THE MAXIMALLY SYMMETRIC SURFACES IN THE 3-TORUS

SHENG BAI, VANESSA ROBINS, CHAO WANG, AND SHICHENG WANG

ABSTRACT. Suppose an orientation-preserving action of a finite group G on the closed surface Σ_g of genus $g > 1$ extends over the 3-torus T^3 for some embedding $\Sigma_g \subset T^3$. Then $|G| \leq 12(g-1)$, and this upper bound $12(g-1)$ can be achieved for $g = n^2 + 1, 3n^2 + 1, 2n^3 + 1, 4n^3 + 1, 8n^3 + 1, n \in \mathbb{Z}_+$. The surfaces in T^3 realizing a maximal symmetry can be either unknotted or knotted. Similar problems in the non-orientable category are also discussed. The connection with minimal surfaces in T^3 is addressed and the situation when the maximally symmetric surfaces above can be realized by minimal surfaces is identified.

CONTENTS

1. Introduction	1
2. Upper and lower bounds of extendable finite groups	4
3. Unknotted examples of the maximally symmetric surfaces	6
4. Knotted examples of the maximally symmetric surfaces	10
5. Minimal surfaces, space groups, proof of the main result	15
References	18

1. INTRODUCTION

Let Σ_g be the closed orientable surface of genus $g > 1$, Π_g be the closed non-orientable surface of genus $g > 2$, and T^3 be the three dimensional torus (3-torus for short). We consider the following question in the smooth category:

Suppose the action of a finite group G on a closed surface S can extend over T^3 . Then what is the maximum order of the group and what does the maximal action look like?

A similar problem has been addressed for surfaces embedded in the 3-sphere, S^3 , which is the simplest compact 3-manifold in the sense that it is a one point compactification of three space and the universal spherical 3-manifold covering all spherical 3-manifolds. See [WWZZ1], [WWZZ2] for orientation-preserving actions, and [WWZ] in the general case. Note that

2010 *Mathematics Subject Classification.* 57M60, 57N10, 57S25, 53A10, 20F65, 05C10;
Key words and phrases. maximal surface symmetry in 3-torus, minimal surface.

The second author is supported by grant No.11501534 of NSFC and the last author is supported by grant No.11371034 of NSFC.

only orientable surfaces Σ_g can be embedded in S^3 . T^3 is another natural and significant 3-manifold for this question. It is the universal compact Euclidean 3-manifold in the sense that it covers all compact Euclidean 3-manifolds; moreover T^3 is covered by 3-space (3-dimensional Euclidean space) and the preimage of a closed surface $S \subset T^3$ under such a covering can be a triply periodic surface, which is an interesting object in the natural sciences and engineering [HBLL+].

Definition 1.1. Let G be a finite group. A G -action on a closed surface S is extendable over T^3 with respect to an embedding $e : S \hookrightarrow T^3$ if G can also act on T^3 such that $h_{T^3} \circ e = e \circ h_S$ for any $h \in G$.

For an embedding of an orientable surface in T^3 we can define whether it is unknotted in the same way as the usual definition for knotting in S^3 .

Definition 1.2. An embedding $e : \Sigma_g \hookrightarrow T^3$ is unknotted if $e(\Sigma_g)$ splits T^3 into two handlebodies. Otherwise it is knotted.

We assume all orientable manifolds in this note are already oriented. According to whether the action preserves the orientation of the surface and T^3 , we can define four classes of maximum orders.

Definition 1.3. Let S be either Σ_g or Π_g . Define $E(S)$, $E^+(S)$, $E_+(\Sigma_g)$ and $E_+^+(\Sigma_g)$ as below:

$E(S)$: the maximum order of all extendable G -actions on S .

$E^+(S)$: the maximum order of all extendable G -actions on S which preserve the orientation of T^3 .

$E_+(\Sigma_g)$: the maximum order of all extendable G -actions on Σ_g which preserve the orientation of Σ_g .

$E_+^+(\Sigma_g)$: the maximum order of extendable G -actions on Σ_g , which preserve the orientations of both T^3 and Σ_g .

We will prove the following Theorem 1.4 and Theorem 1.5. Theorem 1.4 gives the upper bounds of those invariants defined in Definition 1.3, and Theorem 1.5 provides infinitely many g for which the upper bound is realized.

Theorem 1.4. Suppose the genus $g > 1$ for Σ_g and $g > 2$ for Π_g .

- (1) $E_+^+(\Sigma_g) \leq 12(g-1)$.
- (2) $E_+(\Sigma_g) \leq 24(g-1)$, $E^+(\Sigma_g) \leq 24(g-1)$, $E(\Sigma_g) \leq 48(g-1)$.
- (3) $E^+(\Pi_g) \leq 12(g-2)$, $E(\Pi_g) \leq 24(g-2)$.

If an extendable G -action on S realizes one of the above upper bounds, then the corresponding surface $e(S)$ can be thought as a maximally symmetric surface in T^3 .

Theorem 1.5. Suppose n is any positive integer.

- (1) The upper bound of $E_+^+(\Sigma_g)$ in Theorem 1.4 can be achieved by an unknotted embedding for $g = 2n^3 + 1, 4n^3 + 1, 8n^3 + 1$; and by a knotted

embedding for $g = 2n^3 + 1, 4n^3 + 1, 8n^3 + 1$, where 6 does not divide n ; and $g = n^2 + 1, 3n^2 + 1$.

(2) The upper bound of $E_+(\Sigma_g)$, $E^+(\Sigma_g)$ and $E(\Sigma_g)$ in Theorem 1.4 can be achieved for $g = 2n^3 + 1, 4n^3 + 1, 8n^3 + 1$ by an unknotted embedding.

(3) The upper bound of $E^+(\Pi_g)$ and $E(\Pi_g)$ in Theorem 1.4 can be achieved for $g = 2n^3 + 2, 8n^3 + 2$, where n is odd.

The proof of Theorem 1.4, using the Riemann-Hurwitz Formula and the Equivariant Loop Theorem, is given in §2. The proof of Theorem 1.5 occupies the remaining three sections of the paper, which is outlined below:

In §3 and §4 we will construct various examples to realize the upper bounds in Theorem 1.5 (1), §3 for the unknotted case and §4 for the knotted case, therefore Theorem 1.5 (1) follows, up to the verification of the knottiness of those embeddings. Those examples are explicit in the following sense: The 3-torus is obtained by standard opposite face identification of the cube and we define those surfaces in the cube before the identification. Indeed, for each g in Theorem 1.5 (1), we construct all embeddings $\Sigma_g \subset T^3$ realizing $E_+^+(\Sigma_g)$ we can image for the moment, and we expect those are all embeddings realizing $E_+^+(\Sigma_g)$, see a conjecture below. The constructions in §3 and §4 often rely on an understanding of crystallographic space groups.

The knottedness of the examples in §3 and §4 involved in Theorem 1.5 (1), as well as Theorem 1.5 (2) and (3), could be verified by arguments such as those in §3 and §4, and by some topological reasoning; but we present a more convenient and interesting way in §5. In §5 we first link the verification of knottedness of the examples in §3 and §4 with minimal surfaces in T^3 , or equivalently, triply periodic minimal surfaces in R^3 , which itself is an important topic, see [Me] and [FH] for examples. Historically, many minimal surfaces in T^3 were constructed using the symmetry of T^3 and it is known that minimal surfaces in T^3 must be unknotted [Me]. We identify the examples in §3 with known triply periodic minimal surfaces (up to isotopy) [Br], therefore they are unknotted. On the other hand, by a simple criterion from covering space theory and our constructions, the examples in §4 are knotted, therefore can not be realized by minimal surfaces. We also identify the group actions in §3 and §4 with known space groups as well as their index-2 supergroups; then Theorem 1.5 (2) and (3) are proved.

Corollary 1.6. $\Sigma_g \subset T^3$ achieving the upper bound $E_+^+(g) = 12(g - 1)$ can be realized by minimal surfaces for $g = 2n^3 + 1, 4n^3 + 1, 8n^3 + 1$.

We also show in Example 2.3 that for each g it is easy to construct an extendable action of order $4(g - 1)$ on (T^3, Σ_g) .

We end the introduction by the following

Conjecture 1.7. Suppose the genus $g > 1$.

- (1) All g realizing the upper bound $E_+^+(\Sigma_g) = 12(g-1)$ are listed in Theorem 1.5 (1), and moreover, all examples realizing those g are listed in §3 and §4.
- (2) $E_+^+(\Sigma_g) = 4(g-1)$ for $g \in Z_+ \setminus K$, where $K = \{f_i(n) | n \in Z_+\}$ and f_i runs over finitely many quadratic and cubic functions.

2. UPPER AND LOWER BOUNDS OF EXTENDABLE FINITE GROUPS

We need the following two important results, see [Hu], [MY], to prove Theorem 1.4.

Riemann-Hurwitz Formula. $\Sigma_g \rightarrow \Sigma_{g'}$ is a regular branched covering with transformation group G . Let $a_1, a_2, \dots, a_k$ be the branched points in $\Sigma_{g'}$ having indices $q_1 \leq q_2 \leq \dots \leq q_k$. Then

$$2 - 2g = |G|(2 - 2g' - \sum_{i=1}^k (1 - \frac{1}{q_i}))$$

Equivariant Loop Theorem. Let M be a three manifold with a smooth action of a discrete group G . Let F be an equivariant subsurface of ∂M . If F is not π_1 -injective with respect to inclusion into M , then it admits a G -equivariant compression disk.

Here a nonempty subset X of M is G -equivariant if $h(X) = X$ or $h(X) \cap X = \emptyset$, for any $h \in G$. A disk $D \subset M$ is a compression disk of F if $\partial D \subset F$ and in F it does not bound any disk.

Proof of (1). Suppose there is an extendable G -action on Σ_g which preserves the orientations of both T^3 and Σ_g . Cutting T^3 along $e(\Sigma_g)$ we get a three manifold M . Since both Σ_g and T^3 are orientable, Σ_g must be two-sided in T^3 , therefore M contains two boundaries F_1 and F_2 , and $F_1 \cong F_2 \cong \Sigma_g$.

Now clearly G acts on M . Since the G -action preserves the orientations of both T^3 and Σ_g , each of F_1 and F_2 is a G -equivariant surface. Since $g > 1$, $\pi_1(\Sigma_g)$ is not abelian, but $\pi_1(T^3)$ is abelian, the induced homomorphism $\pi_1(\Sigma_g) \rightarrow \pi_1(T^3)$ is not injective. Then at least one of F_1 and F_2 is not π_1 -injective with respect to the inclusion into M . Suppose F_1 is not π_1 -injective, then it admits a G -equivariant compression disk D of F_1 by the Equivariant Loop Theorem.

If there exists an $h \in G$ such that $h(D) = D$ and h reverses an orientation of D , then we can choose a G -equivariant regular neighbourhood $N(D)$ of D such that there is a homeomorphism $i : D \times [-1, 1] \rightarrow N(D)$ and $i(D \times \{0\}) = D$. Then $D' = i(D \times \{1\})$ is also a G -equivariant compression disk of F_1 , and clearly for this D' , if $h'(D') = D'$ for some $h' \in G$, then h' preserves an orientation of D' . Hence we can assume each element of G that preserves D also preserves its orientation, in particular it is fixed point free on ∂D .

Since the G -action on T^3 preserves the orientations of both Σ_g and T^3 , the induced G -action on M preserves F_1 . With the quotient topology F_1/G

is homeomorphic to some $\Sigma_{g'}$, and $p : F_1 \rightarrow F_1/G$ is a regular branched covering. Since the G action is fixed point free on ∂D by previous discussion, $p(\partial D)$ is a simple closed curve in F_1/G .

Let $a_1, a_2, \dots, a_k$ be the branch points having indices $q_1 \leq q_2 \leq \dots \leq q_k$. Note $2 - 2g < 0$. By the Riemann-Hurwitz Formula we have

$$2 - 2g' - \sum_{i=1}^k \left(1 - \frac{1}{q_i}\right) = \frac{2 - 2g}{|G|} < 0.$$

If $g' = 0$ and $k \leq 3$, then $p(\partial D)$ must bound a disk in F_1/G containing at most one branch point, and ∂D will bound a disk in F_1 , a contradiction. Hence either $g' \geq 1$ or $g' = 0$ and $k \geq 4$.

Notice that for each q_i , we have

$$1 - \frac{1}{q_i} \geq 1/2.$$

Then by elementary calculation we have $|G| \leq 12(g-1)$, and moreover the equality holds if and only if $g' = 0$, $k = 4$ and $(q_1, q_2, q_3, q_4) = (2, 2, 2, 3)$. $\square$

Remark 2.1. Zimmermann first proved that the order of an orientation-preserving finite group action on a handlebody of genus g is bounded by $12(g-1)$ [Zi] soon after the work [MY]. As mentioned by one of the referees, May first discovered the sharp $12(g-1)$ upper bound for the automorphism group of a genus- g Klein surface (i.e. the case when S is a non-orientable surfaces, or has non-empty boundary) [Ma1]. It would be interesting to compare the maximal groups from this paper with those in the May [Ma2] and Zimmermann papers.

By the above proof, we actually have the following:

Theorem 2.2. *Suppose Σ_g is embedded in a three manifold M and a finite group G acts on (Σ_g, M) . If G preserves both sides and the orientation of Σ_g and Σ_g is not π_1 -injective in M , then $|G| \leq 12(g-1)$.*

Now (2) and (3) follow from (1).

Proof of (2). Suppose there is an extendable G -action on Σ_g . Let G^o be the normal subgroup of G containing all elements that preserve both the orientations of Σ_g and T^3 . In the case of $E_+(\Sigma_g)$ or $E^+(\Sigma_g)$, the index of G^o in G is at most two, and in the case of $E(\Sigma_g)$, the index of G^o in G is at most four. Hence $E_+(\Sigma_g) \leq 24(g-1)$, $E^+(\Sigma_g) \leq 24(g-1)$, $E(\Sigma_g) \leq 48(g-1)$. $\square$

Proof of (3). Suppose there is an extendable G -action on Π_g and the action preserves the orientation of T^3 . We can choose an equivariant regular neighbourhood $N(\Pi_g)$ of Π_g , then G also acts on $\partial N(\Pi_g)$. Since T^3 is orientable, $N(\Pi_g)$ is homeomorphic to a twisted $[-1, 1]$ -bundle over Π_g and $\partial N(\Pi_g)$ is the orientable double cover of Π_g under the bundle projection. Since $\chi(\Pi_g) = 2 - g$, $\chi(\partial N(\Pi_g)) = 4 - 2g = 2 - 2(g-1)$. It

follows that $\partial N(\Pi_g)$ is homeomorphic to Σ_{g-1} . Clearly the G -action preserves the two sides of $\partial N(\Pi_g)$. Since the action preserves the orientation of T^3 , it also preserves the orientation of $\partial N(\Pi_g)$. Then by the result of (1), we have $E^+(\Pi_g) \leq 12(g-2)$. And similar to the proof of (2), we have $E(\Pi_g) \leq 24(g-2)$. $\square$

Example 2.3. Let $([0, 1]^3, +)$ be the unit cube with a cross properly embedded shown as the right side of Figure 1. For each $g > 1$, put $g-1$ copies of $([0, 1]^3, +)$ to get $([0, 1]^3, +)_{g-1}$ which is a cuboid with properly embedded antennae, as shown in Figure 1. If we identify the opposite faces of the cuboid, we get (T^3, Θ_g) , where Θ_g is a graph of genus g . If we first only identify the right and left faces, we get $([0, 1]^2 \times S^1, A_{g-1})$, where A_{g-1} is a circular antennae with $g-1$ bars. It is easy to see that $G = D_{g-1} \oplus Z_2$ acts on the pair $([0, 1]^2 \times S^1, A_{g-1})$, where D_{g-1} is the dihedral group acts in a standard way on $([0, 1]^2 \times S^1, A_{g-1})$, and Z_2 is π -rotation of $([0, 1]^2 \times S^1, A_{g-1})$ around the center-circle. Clearly this action preserves each pair of faces of $([0, 1]^2 \times S^1, A_{g-1})$ to be identified, therefore induces an action on (T^3, Θ_g) . Let $N(\Theta_g)$ be a G -invariant regular neighborhood of Θ_g . Then $\partial N(\Theta_g) = \Sigma_g$ and G of order $4(g-1)$ acts on (T^3, Σ_g) .

FIGURE 1. Cuboid with embedded antennae

3. UNKNOTTED EXAMPLES OF THE MAXIMALLY SYMMETRIC SURFACES

In this section we give three classes of examples realizing the upper bound of $E_+^+(\Sigma_g) = 12(g-1)$. In each case we first construct a triply periodic graph Γ in the three-dimensional Euclidean space E^3 . There will be a three-dimensional space group $\mathcal{G}$ preserving Γ . Then we choose a rank-three translation normal subgroup T in $\mathcal{G}$. The space E^3/T is our T^3 . The finite group $G = \mathcal{G}/T$ acts on T^3 preserving the graph $\Theta = \Gamma/T$. Finally, we choose an equivariant regular neighbourhood $N(\Theta)$ of Θ , and $\partial N(\Theta)$ is our surface Σ_g . After the discussion of §5 we will see that for each example the complement of $N(\Theta)$ is also a handlebody, hence the surface is unknotted.

Definition 3.1. Let $T_1 = \{(a, b, c) \mid a, b, c \in \mathbb{Z}\}$ be the group of integer translations. Let $t_x = (1, 0, 0), t_y = (0, 1, 0), t_z = (0, 0, 1)$. An element $t = (a, b, c) \in T_1$ acts on E^3 as following:

$$t : (x, y, z) \mapsto (x + a, y + b, z + c)$$

For $n \in \mathbb{Z}_+$, we define three classes of subgroups T_m of T_1 for those integers m that can be presented (uniquely) in the form $n^3, 2n^3$ and $4n^3$ as follows:

$$\begin{aligned} T_{n^3} &= \langle nt_x, nt_y, nt_z \rangle, \\ T_{2n^3} &= \langle nt_y + nt_z, nt_z + nt_x, nt_x + nt_y \rangle, \\ T_{4n^3} &= \langle -nt_x + nt_y + nt_z, nt_x - nt_y + nt_z, nt_x + nt_y - nt_z \rangle. \end{aligned}$$

The subscript m of T_m is equal to the volume $\text{Vol}(E^3/T_m)$, for the above cases with $m = n^3, 2n^3, 4n^3$. We use the subscript m for convenience, and note that the co-volume of a sublattice is not sufficient to uniquely identify a particular sublattice in general.

Clearly T_{2n^3} and T_{4n^3} are subgroups of T_{n^3} . One can easily verify that $T_{(2n)^3} \subset T_{2n^3}$, since $2nt_x, 2nt_y, 2nt_z$ are linear combinations of $nt_y + nt_z, nt_z + nt_x, nt_x + nt_y$, hence $T_{32n^3} = T_{4(2n)^3} \subset T_{(2n)^3} = T_{8n^3} \subset T_{2n^3}$. Similarly one can verify that $T_{(2n)^3} \subset T_{4n^3}$, hence $T_{16n^3} = T_{2(2n)^3} \subset T_{(2n)^3} = T_{8n^3} \subset T_{4n^3}$.

Definition 3.2. We define five isometries of E^3 as follows:

$$\begin{aligned} r_y &: (x, y, z) \mapsto (-x, y, -z) \\ r_z &: (x, y, z) \mapsto (-x, -y, z) \\ r_{xy} &: (x, y, z) \mapsto (y, x, -z) \\ r_{xyz} &: (x, y, z) \mapsto (z, x, y) \\ t_{1/2} &: (x, y, z) \mapsto (x + 1/2, y + 1/2, z + 1/2) \end{aligned}$$

The isometries r_y, r_z and r_{xy} are 2-fold rotations (i.e., rotation by an angle of π) about the y -axis, z -axis and the line $x = y, z = 0$ respectively. The isometry r_{xyz} is a positive 3-fold rotation about the cube body diagonal, $x = y = z$ (i.e., right-hand rule rotation by $2\pi/3$ about the direction $[1, 1, 1]$.)

The notations of space groups used below come from the International Tables for Crystallography [Ha].

Example 3.3. Let Γ_{min}^1 be the one-skeleton of the unit cube $[0, 1]^3$ in E^3 as in Figure 2. Let $\Gamma^1 = \bigcup_{t \in T_1} t(\Gamma_{min}^1)$. Then Γ^1 , the one-skeleton of the tessellation of E^3 by unit cubes, is a triply periodic graph called the simple or primitive cubic lattice (**pcu** in [RCSR]).

Let $H^1 = \langle r_y, r_z, r_{xy}, r_{xyz} \rangle$, which is the well-known orientation-preserving isometric group of $[-1, 1]^3$ of order 24 (with Schönflies symbol O). Let $\mathcal{G}^1 = \langle T_1, H^1 \rangle$, this is the space group $[P432]$. Then $\mathcal{G}^1$ preserves Γ^1 . Now in $T^3 \cong E^3/T_1$ we have a graph $\Theta_1^1 = \Gamma^1/T_1$. It has one vertex and three edges, hence its Euler characteristic $\chi(\Theta_1^1) = -2$ and its genus $g = 1 - \chi(\Theta_1^1) = 3$. Clearly $G_1^1 = \mathcal{G}^1/T_1 \cong H^1$ acts on $T^3 \cong E^3/T_1$ preserving Θ_1^1 , and G_1^1

FIGURE 2. One-skeleton of $[0, 1]^3$

has order 24. Hence when we choose an equivariant regular neighbourhood $N(\Theta_1^1)$ of Θ_1^1 , we get an extendable action of order 24 on $\Sigma_3 \cong \partial N(\Theta_1^1)$.

Similarly in $T^3 \cong E^3/T_{n^3}$ we have a graph $\Theta_{n^3}^1 = \Gamma^1/T_{n^3}$. Since T_{n^3} is a normal subgroup of $\mathcal{G}^1$, $G_{n^3}^1 = \mathcal{G}^1/T_{n^3}$ acts on $T^3 \cong E^3/T_{n^3}$ preserving $\Theta_{n^3}^1$.

$$\begin{aligned}\chi(\Theta_{n^3}^1) &= \chi(\Theta_1^1) \cdot \text{Vol}(E^3/T_{n^3}) = -2n^3 \\ |G_{n^3}^1| &= |G_1^1| \cdot \text{Vol}(E^3/T_{n^3}) = 24n^3\end{aligned}$$

Hence the genus of $\Theta_{n^3}^1$ is $2n^3 + 1$. Then we can get an extendable action of order $24n^3$ on $\Sigma_{2n^3+1} \cong \partial N(\Theta_{n^3}^1)$.

Notice that when $m = 2n^3, 4n^3, n \in \mathbb{Z}_+$, T_m is also a normal subgroup of $\mathcal{G}^1$. Similar to the above construction, we can get an extendable action of order $24m$ on $\Sigma_{2m+1} \cong \partial N(\Theta_m^1)$. Here $\Theta_m^1 = \Gamma^1/T_m$ is a graph in $T^3 \cong E^3/T_m$.

In the above example the superscript 1 in Γ^1 or Θ_m^1 is equal to the volume of the ‘minimal 3-torus’ E^3/T_1 . It is also equal to the number of vertices of the graph Θ_1^1 . In the following examples we use similar notations.

Example 3.4. Let Γ_{min}^2 be the graph in the unit cube $[0, 1]^3$ in E^3 as in Figure 3. It consists of four edges from $(1/2, 1/2, 1/2)$ to $(0, 0, 0)$, $(0, 1, 1)$, $(1, 0, 1)$, $(1, 1, 0)$. Let $\Gamma^2 = \bigcup_{t \in T_2} t(\Gamma_{min}^2)$. One can check that it is connected and is known to scientists as the bonding structure of diamond (**dia** in [RCSR]). Figure 3 shows a fundamental region of T_2 .

Let $H^2 = \langle r_y, r_z, r_{xyz} \rangle$. It is the orientation-preserving isometric group of the regular tetrahedron formed by the convex hull of $(1, 1, 1)$, $(1, -1, -1)$, $(-1, 1, -1)$, and $(-1, -1, 1)$ and has Schönflies symbol T . Let $\mathcal{G}^2 = \langle T_2, H^2, t_{1/2} r_{xy} \rangle$, this is the space group $[F4_132]$. Then $\mathcal{G}^2$ preserves Γ^2 .

Now in $T^3 \cong E^3/T_2$ we have a graph $\Theta_2^2 = \Gamma^2/T_2$. It has two vertices and four edges, hence its Euler characteristic $\chi(\Theta_2^2) = -2$ and its genus is 3. T_2 is a normal subgroup of $\mathcal{G}^2$ and $G_2^2 = \mathcal{G}^2/T_2$ has order 24. It acts on $T^3 \cong E^3/T_2$ preserving Θ_2^2 . Hence when we choose an equivariant regular

FIGURE 3. Γ_{min}^2 in $[0, 2] \times [0, 1] \times [0, 1]$. This is a fundamental region for T_2 , but recall that the translations for T_2 are $y + z$, $z + x$, $x + y$, and these do not glue up directly opposite faces of the rectangular prism illustrated.

neighbourhood $N(\Theta_2^2)$ of Θ_2^2 , we get an extendable action of order 24 on $\Sigma_3 \cong \partial N(\Theta_2^2)$.

Similarly when $m = n^3, 4n^3, 16n^3, n \in \mathbb{Z}_+$, in $T^3 \cong E^3/T_{2m}$ we have a graph $\Theta_{2m}^2 = \Gamma^2/T_{2m}$. One can check that T_{2m} is a normal subgroup of $\mathcal{G}^2$, and the quotient $G_{2m}^2 = \mathcal{G}^2/T_{2m}$ acts on $T^3 \cong E^3/T_{2m}$ preserving Θ_{2m}^2 .

$$\begin{aligned}\chi(\Theta_{2m}^2) &= \chi(\Theta_2^2) \cdot \text{Vol}(E^3/T_{2m})/\text{Vol}(E^3/T_2) = -2m \\ |G_{2m}^2| &= |G_2^2| \cdot \text{Vol}(E^3/T_{2m})/\text{Vol}(E^3/T_2) = 24m\end{aligned}$$

Hence the genus of Θ_{2m}^2 is $2m + 1$. Then we can get an extendable action of order $24m$ on $\Sigma_{2m+1} \cong \partial N(\Theta_{2m}^2)$.

Example 3.5. Let γ be the graph in the unit cube $[0, 1]^3$ in E^3 as in Figure 4. It has three edges from $(1/4, 1/4, 1/4)$ to $(0, 1/2, 1/4)$, $(1/4, 0, 1/2)$ and $(1/2, 1/4, 0)$. Let $\Gamma_{min}^4 = \gamma \cup t_y^2 t_z r_y r_z(\gamma) \cup t_x^2 t_y t_z r_y(\gamma) \cup t_x^2 t_y r_z(\gamma)$. Then let $\Gamma^4 = \bigcup_{t \in T_4} t(\Gamma_{min}^4)$. One can check that it is connected, and is the oft-rediscovered chiral vertex-transitive net of degree-3 known by many names [HOP], including **srs** in [RCSR]. Figure 4 shows a fundamental region of T_4 .

Let $\mathcal{G}^4 = \langle T_4, t_x t_z r_z, t_y t_z r_y, r_{xyz}, t_x t_{1/2} r_{xy} \rangle$, it is the space group $[I4_132]$. Then $\mathcal{G}^4$ preserves Γ^4 .

Now in $T^3 \cong E^3/T_4$ we have a graph $\Theta_4^4 = \Gamma^4/T_4$. It has four vertices and six edges, hence its Euler characteristic $\chi(\Theta_4^4) = -2$ and its genus is 3. T_4 is a normal subgroup of $\mathcal{G}^4$ and $G_4^4 = \mathcal{G}^4/T_4$ has order 24. It acts on $T^3 \cong E^3/T_4$ preserving Θ_4^4 . Hence when we choose an equivariant regular neighbourhood $N(\Theta_4^4)$ of Θ_4^4 , we get an extendable action of order 24 on $\Sigma_3 \cong \partial N(\Theta_4^4)$.

Similarly when $m = n^3, 2n^3, 4n^3, n \in \mathbb{Z}_+$, in $T^3 \cong E^3/T_{4m}$ we have a graph $\Theta_{4m}^4 = \Gamma^4/T_{4m}$. One can check that T_{4m} is a normal subgroup of $\mathcal{G}^4$,

FIGURE 4. Γ_{min}^4 in $[0, 2] \times [0, 2] \times [0, 1]$. Similar to the previous example, this is a fundamental region for T_4 although the T_4 translations do not identify directly opposite faces of the rectangular prism illustrated.

and the quotient $G_{4m}^4 = \mathcal{G}^4/T_{4m}$ acts on $T^3 \cong E^3/T_{4m}$ preserving Θ_{4m}^4 .

$$\begin{aligned}\chi(\Theta_{4m}^4) &= \chi(\Theta_4^4) \cdot \text{Vol}(E^3/T_{4m})/\text{Vol}(E^3/T_4) = -2m \\ |G_{4m}^4| &= |G_4^4| \cdot \text{Vol}(E^3/T_{4m})/\text{Vol}(E^3/T_4) = 24m\end{aligned}$$

Hence the genus of Θ_{4m}^4 is $2m + 1$. Then we can get an extendable action of order $24m$ on $\Sigma_{2m+1} \cong \partial N(\Theta_{4m}^4)$.

4. KNOTTED EXAMPLES OF THE MAXIMALLY SYMMETRIC SURFACES

In the section we give six further classes of examples realizing the upper bound of $E_+^+(\Sigma_g) = 12(g - 1)$. In each case below the embedding is knotted. The surface $e(\Sigma_g)$ does bound a handlebody in T^3 on one side, but the complement of the handlebody is not a handlebody, and this will be clear after the discussion of §5.

Constructions of the extendable actions and surfaces is similar to §4, but here the graph Γ in E^3 can be disconnected. There is a space group $\mathcal{G}$ preserving Γ and a rank three translation normal subgroup T in $\mathcal{G}$ so that the graph $\Theta = \Gamma/T$ is connected. The finite group $G = \mathcal{G}/T$ acts on $T^3 \cong E^3/T$ preserving Θ . Then we choose an equivariant regular neighbourhood $N(\Theta)$ of Θ , and $\partial N(\Theta)$ is our surface Σ_g .

Example 4.1. Let $\Gamma^{1,2} = \Gamma^1 \cup t_{1/2}(\Gamma^1)$. The graph $t_{1/2}(\Gamma^1)$ can be thought of as the dual of Γ^1 in E^3 . $\Gamma^{1,2}$ has two connected components and is named **pcu-c** in [RCSR]. In the unit cube $[0, 1]^3$ it is as in Figure 5.

FIGURE 5. Part of $\Gamma^{1,2}$ in $[0, 1]^3$

Let $\mathcal{G}^{1,2} = \langle \mathcal{G}^1, t_{1/2} \rangle$, it is the space group $[I432]$, and let $T_{1/2} = \langle T_1, t_{1/2} \rangle = \{(a/2, b/2, c/2) \mid (a, b, c) \in T_4\}$. Then $\mathcal{G}^{1,2}$ preserves $\Gamma^{1,2}$ and $T_{1/2}$ is a normal subgroup of $\mathcal{G}^{1,2}$. Since $t_{1/2}$ changes the two components of $\Gamma^{1,2}$, the graph $\Theta_{1/2}^{1,2} = \Gamma^{1,2}/T_{1/2}$ in $T^3 \cong E^3/T_{1/2}$ is connected. Its Euler characteristic $\chi(\Theta_{1/2}^{1,2}) = -2$ and its genus is 3. $G_{1/2}^{1,2} = \mathcal{G}^{1,2}/T_{1/2}$ has order 24. It acts on $T^3 \cong E^3/T_{1/2}$ preserving $\Theta_{1/2}^{1,2}$. Hence when we choose an equivariant regular neighbourhood $N(\Theta_{1/2}^{1,2})$ of $\Theta_{1/2}^{1,2}$, we get an extendable action of order 24 on $\Sigma_3 \cong \partial N(\Theta_{1/2}^{1,2})$.

Similarly let $T_{n^3/2} = \{(a/2, b/2, c/2) \mid (a, b, c) \in T_{4n^3}\}, n \in \mathbb{Z}_+, 2 \nmid n$. Then the graph $\Theta_{n^3/2}^{1,2} = \Gamma^{1,2}/T_{n^3/2}$ in $T^3 \cong E^3/T_{n^3/2}$ is connected. Since $T_{n^3/2}$ is a normal subgroup of $\mathcal{G}^{1,2}$, $G_{n^3/2}^{1,2} = \mathcal{G}^{1,2}/T_{n^3/2}$ acts on $T^3 \cong E^3/T_{n^3/2}$ preserving $\Theta_{n^3/2}^{1,2}$.

$$\begin{aligned} \chi(\Theta_{n^3/2}^{1,2}) &= \chi(\Theta_{1/2}^{1,2}) \cdot \text{Vol}(E^3/T_{n^3/2})/\text{Vol}(E^3/T_{1/2}) = -2n^3 \\ |G_{n^3/2}^{1,2}| &= |G_{1/2}^{1,2}| \cdot \text{Vol}(E^3/T_{n^3/2})/\text{Vol}(E^3/T_{1/2}) = 24n^3 \end{aligned}$$

Hence the genus of $\Theta_{n^3/2}^{1,2}$ is $2n^3 + 1$. Then we can get an extendable action of order $24n^3$ on $\Sigma_{2n^3+1} \cong \partial N(\Theta_{n^3/2}^{1,2})$, here $n \in \mathbb{Z}_+, 2 \nmid n$.

Example 4.2. Let $\Gamma^{2,2} = \Gamma^2 \cup t_x(\Gamma^2)$. The graph $t_x(\Gamma^2)$ can be thought as the dual of Γ^2 in E^3 . $\Gamma^{2,2}$ has two connected components and is called **dia-c** in [RCSR]. In the fundamental region of T_2 it is as in Figure 6.

Let $\mathcal{G}^{2,2} = \langle \mathcal{G}^2, t_x \rangle$, it is the space group $[P4_232]$. Then $\mathcal{G}^{2,2}$ preserves $\Gamma^{2,2}$ and contains T_1 as a normal subgroup. Since t_x changes the two components of $\Gamma^{2,2}$, the graph $\Theta_1^{2,2} = \Gamma^{2,2}/T_1$ in $T^3 \cong E^3/T_1$ is connected. Its Euler characteristic $\chi(\Theta_1^{2,2}) = -2$, and its genus is 3. $G_1^{2,2} = \mathcal{G}^{2,2}/T_1$ has order 24. It acts on $T^3 \cong E^3/T_1$ preserving $\Theta_1^{2,2}$. Hence when we choose an

FIGURE 6. Part of $\Gamma^{2,2}$ in $[0, 2] \times [0, 1] \times [0, 1]$

equivariant regular neighbourhood $N(\Theta_1^{2,2})$ of $\Theta_1^{2,2}$, we get an extendable action of order 24 on $\Sigma_3 \cong \partial N(\Theta_1^{2,2})$.

Similarly if $m = n^3, 4n^3, n \in \mathbb{Z}_+, 2 \nmid n$, the graph $\Theta_m^{2,2} = \Gamma^{2,2}/T_m$ in $T^3 \cong E^3/T_m$ is connected. One can check that T_m is a normal subgroup of $\mathcal{G}^{2,2}$, $G_m^{2,2} = \mathcal{G}^{2,2}/T_m$ acts on $T^3 \cong E^3/T_m$ preserving $\Theta_m^{2,2}$.

$$\chi(\Theta_m^{2,2}) = \chi(\Theta_1^{2,2}) \cdot \text{Vol}(E^3/T_m) = -2m$$

$$|G_m^{2,2}| = |G_1^{2,2}| \cdot \text{Vol}(E^3/T_m) = 24m$$

Hence the genus of $\Theta_m^{2,2}$ is $2m + 1$. Then we can get an extendable action of order $24m$ on $\Sigma_{2m+1} \cong \partial N(\Theta_m^{2,2})$, here $m = n^3, 4n^3, n \in \mathbb{Z}_+, 2 \nmid n$.

Example 4.3. Let $\Gamma^{4,4} = \Gamma^4 \cup t_x(\Gamma^4) \cup t_y(\Gamma^4) \cup t_z(\Gamma^4)$. $\Gamma^{4,4}$ has four connected components. In the unit cube $[0, 1]^3$ in E^3 it is as in Figure 7.

FIGURE 7. Part of $\Gamma^{4,4}$ in $[0, 1]^3$

One can check that $\mathcal{G}^{2,2}$ defined in Example 4.2 (space group $[P4_232]$) preserves $\Gamma^{4,4}$, and the graph $\Theta_1^{4,4} = \Gamma^{4,4}/T_1$ in $T^3 \cong E^3/T_1$ is connected. Its Euler characteristic $\chi(\Theta_1^{4,4}) = -2$, and its genus is 3. $G_1^{2,2} = \mathcal{G}^{2,2}/T_1$ has order 24. It acts on $T^3 \cong E^3/T_1$ preserving $\Theta_1^{4,4}$. Hence when we choose an equivariant regular neighbourhood $N(\Theta_1^{4,4})$ of $\Theta_1^{4,4}$, we get an extendable action of order 24 on $\Sigma_3 \cong \partial N(\Theta_1^{4,4})$.

Similarly if $m = n^3, 2n^3, n \in \mathbb{Z}_+, 2 \nmid n$, the graph $\Theta_m^{4,4} = \Gamma^{4,4}/T_m$ in $T^3 \cong E^3/T_m$ is connected. $G_m^{2,2}$ acts on $T^3 \cong E^3/T_m$ preserving $\Theta_m^{4,4}$.

$$\begin{aligned}\chi(\Theta_m^{4,4}) &= \chi(\Theta_1^{4,4}) \cdot \text{Vol}(E^3/T_m) = -2m \\ |G_m^{2,2}| &= |G_1^{2,2}| \cdot \text{Vol}(E^3/T_m) = 24m\end{aligned}$$

Hence the genus of $\Theta_m^{4,4}$ is $2m + 1$. Then we can get an extendable action of order $24m$ on $\Sigma_{2m+1} \cong \partial N(\Theta_m^{2,2})$, here $m = n^3, 2n^3, n \in \mathbb{Z}_+, 2 \nmid n$.

Example 4.4. Let $\Gamma^{4,8} = \Gamma^{4,4} \cup t_{1/2}(\Gamma^{4,4})$. It has eight connected components. In the unit cube $[0, 1]^3$ it is as in Figure 8.

FIGURE 8. Part of $\Gamma^{4,8}$ in $[0, 1]^3$

One can check that $\mathcal{G}^{1,2}$ defined in Example 4.1 (space group $[I432]$) preserves $\Gamma^{4,8}$, and the graph $\Theta_{1/2}^{4,8} = \Gamma^{4,8}/T_{1/2}$ in $T^3 \cong E^3/T_{1/2}$ is connected. Its Euler characteristic is -2 and its genus is 3. The order 24 group $G_{1/2}^{1,2}$ acts on $T^3 \cong E^3/T_{1/2}$ preserving $\Theta_{1/2}^{4,8}$. Hence when we choose an equivariant regular neighbourhood $N(\Theta_{1/2}^{4,8})$ of $\Theta_{1/2}^{4,8}$, we get an extendable action of order 24 on $\Sigma_3 \cong \partial N(\Theta_{1/2}^{4,8})$.

Similarly for $n \in \mathbb{Z}_+, 2 \nmid n$, the graph $\Theta_{n^3/2}^{4,8} = \Gamma^{4,8}/T_{n^3/2}$ in $T^3 \cong E^3/T_{n^3/2}$ is connected. $G_{n^3/2}^{1,2}$ acts on $T^3 \cong E^3/T_{n^3/2}$ preserving $\Theta_{n^3/2}^{4,8}$.

$$\begin{aligned}\chi(\Theta_{n^3/2}^{4,8}) &= \chi(\Theta_{1/2}^{4,8}) \cdot \text{Vol}(E^3/T_{n^3/2})/\text{Vol}(E^3/T_{1/2}) = -2n^3 \\ |G_{n^3/2}^{1,2}| &= |G_{1/2}^{1,2}| \cdot \text{Vol}(E^3/T_{n^3/2})/\text{Vol}(E^3/T_{1/2}) = 24n^3\end{aligned}$$

Hence the genus of $\Theta_{n^3/2}^{4,8}$ is $2n^3 + 1$. Then we can get an extendable action of order $24n^3$ on $\Sigma_{2n^3+1} \cong \partial N(\Theta_{n^3/2}^{4,8})$, here $n \in \mathbb{Z}_+, 2 \nmid n$.

Example 4.5. Let γ' be the graph in the unit cube $[0, 1]^3$ in E^3 as in Figure 4. There are three edges from $(1/4, 1/4, 1/4)$ to $(0, 1/4, 1/2)$, $(1/2, 0, 1/4)$, $(1/4, 1/2, 0)$, and three edges from $(0, 3/4, 1/2)$ to $(1/2, 3/4, 1)$, $(1/2, 0, 3/4)$ to $(1, 1/2, 3/4)$ and $(3/4, 1/2, 0)$ to $(3/4, 1, 1/2)$ separately.

Let $\Gamma_{min}^{\prime 4} = \gamma' \cup t_y^2 t_z r_y r_z(\gamma') \cup t_x^2 t_y t_z r_y(\gamma') \cup t_x^2 t_y r_z(\gamma')$, and $\Gamma^{\prime 4} = \bigcup_{t \in T_4} t(\Gamma_{min}^{\prime 4})$. One can check that $\Gamma^{\prime 4}$ has 27 connected components, each is similar to a mirror image of Γ^4 . Figure 9 shows the part of $\Gamma^{\prime 4}$ in a fundamental region of T_4 .

FIGURE 9. Part of $\Gamma^{\prime 4}_{min}$ in $[0, 2] \times [0, 2] \times [0, 1]$

By construction, $\mathcal{G}^4$ (space group $[I4_132]$) preserves $\Gamma^{\prime 4}$. And one can check that $\Theta_4^{\prime 4} = \Gamma^{\prime 4}/T_4$ in $T^3 \cong E^3/T_4$ is connected. $\Theta_4^{\prime 4}$ has four vertices and six edges, hence it has Euler characteristic -2 and genus 3. The order-24 group G_4^4 acts on $T^3 \cong E^3/T_4$ preserving $\Theta_4^{\prime 4}$. Hence when we choose an equivariant regular neighbourhood $N(\Theta_4^{\prime 4})$ of $\Theta_4^{\prime 4}$, we get an extendable action of order 24 on $\Sigma_3 \cong \partial N(\Theta_4^{\prime 4})$.

Similarly when $m = n^3, 2n^3, 4n^3, n \in \mathbb{Z}_+, 3 \nmid n$, the graph $\Theta_{4m}^{\prime 4} = \Gamma^{\prime 4}/T_{4m}$ in $T^3 \cong E^3/T_{4m}$ is connected. The quotient $G_{4m}^4 = \mathcal{G}^4/T_{4m}$ acts on $T^3 \cong$

E^3/T_{4m} preserving Θ_{4m}^4 .

$$\begin{aligned}\chi(\Theta_{4m}^4) &= \chi(\Theta_4^4) \cdot \text{Vol}(E^3/T_{4m})/\text{Vol}(E^3/T_4) = -2m \\ |G_{4m}^4| &= |G_4^4| \cdot \text{Vol}(E^3/T_{4m})/\text{Vol}(E^3/T_4) = 24m\end{aligned}$$

Hence the genus of Θ_{4m}^4 is $2m + 1$. Then we can get an extendable action of order $24m$ on $\Sigma_{2m+1} \cong \partial N(\Theta_{4m}^4)$, here $m = n^3, 2n^3, 4n^3, n \in \mathbb{Z}_+, 3 \nmid n$.

Definition 4.6. Let $t_\omega = (-1/2, \sqrt{3}/2, 0)$. Define

$$\begin{aligned}T_{n^2}^\omega &= \langle nt_\omega, nt_x, t_z \rangle, \\ T_{3n^2}^\omega &= \langle 2nt_\omega + nt_x, nt_\omega + 2nt_x, t_z \rangle.\end{aligned}$$

Define a rotation r_ω on E^3 as following:

$$r_\omega : (x, y, z) \mapsto \left(-\frac{1}{2}x - \frac{\sqrt{3}}{2}y, \frac{\sqrt{3}}{2}x - \frac{1}{2}y, z\right)$$

T_1^ω is the hexagonal 3D lattice and r_ω is a rotation by $2\pi/3$ right-handed with respect to the direction $[0, 0, 1]$.

Example 4.7. Let P be a regular hexagon in the xy -plane. It has center $(0, 0, 0)$ and contains $(1/2, \sqrt{3}/6, 0)$ as a vertex, see Figure 10. Let Γ_{min}^P be the boundary of P in E^3 . Let $\Gamma^P = \bigcup_{t \in T_1^\omega} t(\Gamma_{min}^P)$. It contains infinitely many connected components, and for each $n \in \mathbb{Z}$, the horizontal plane $\{(x, y, n) \mid x, y \in E^2\}$ contains exactly one of them. A local picture of Γ^P is as in Figure 11.

FIGURE 10. A regular hexagon

Let $H^\omega = \langle r_\omega, r_x, r_y \rangle$; it is the isometric group of P (Schönflies symbol D_6). Let $\mathcal{G}^\omega = \langle T_1^\omega, H^\omega \rangle$, this is the space group $[P622]$. Then $\mathcal{G}^\omega$ preserves Γ^P , and $\Theta_1^P = \Gamma^P/T_1^\omega$ is a connected graph in $T^3 \cong E^3/T_1^\omega$. It has two

FIGURE 11. A local picture of Γ^P

vertices and three edges, hence its Euler characteristic $\chi(\Theta_1^\omega) = -1$ and its genus is 2. Clearly $G_1^\omega = \mathcal{G}^\omega/T_1^\omega$ acts on $T^3 \cong E^3/T_1^\omega$ preserving Θ_1^P , and $G_1^\omega \cong H^\omega$ has order 12. Hence when we choose an equivariant regular neighbourhood $N(\Theta_1^P)$ of Θ_1^P , we get an extendable action of order 12 on $\Sigma_2 \cong \partial N(\Theta_1^P)$.

Notice that when $m = n^2, 3n^2, n \in \mathbb{Z}_+$, T_m^ω is a normal subgroup of $\mathcal{G}^\omega$. Similarly the graph $\Theta_m^P = \Gamma^P/T_m^\omega$ in $T^3 \cong E^3/T_m^\omega$ is connected, and $G_m^\omega = \mathcal{G}^\omega/T_m^\omega$ acts on $T^3 \cong E^3/T_m^\omega$ preserving Θ_m^P .

$$\begin{aligned}\chi(\Theta_m^P) &= \chi(\Theta_1^P) \cdot \text{Vol}(E^3/T_m^\omega)/\text{Vol}(E^3/T_1^\omega) = -m \\ |G_m^\omega| &= |G_1^\omega| \cdot \text{Vol}(E^3/T_m^\omega)/\text{Vol}(E^3/T_1^\omega) = 12m\end{aligned}$$

Hence the genus of Θ_m^P is $m + 1$. Then we can get an extendable action of order $12m$ on $\Sigma_{m+1} \cong \partial N(\Theta_m^P)$.

5. MINIMAL SURFACES, SPACE GROUPS, PROOF OF THE MAIN RESULT

In this section we will finish the proof of Theorem 1.5. We need some results about triply periodic minimal surfaces, space groups, and a lemma given below.

There are three classical triply periodic minimal surfaces that admit high symmetries: Schwarz's P surface, Schwarz's D [Schw] and Schoen's gyroid surface illustrated in Figure 12 [Br]. Denote them by S_P , S_D and S_G . For each surface there is a side-preserving translational lattice such that the corresponding quotient surface has genus three. A fundamental region for the lattice in the case of S_P is illustrated in Figure 12. A fundamental region

for S_D has volume two in the coordinate system of Section 3. In Figure 12 we draw S_D in the region $[0, 1] \times [0, 2] \times [0, 1]$ to match the domain used in Figure 3. A fundamental region for S_G has volume four, but in Figure 12 we draw S_G in the domain $[0, 2] \times [0, 2] \times [0, 2]$, twice the volume of the fundamental region used in Figure 4 to assist in visualisation, and because this is the standard non-primitive cubic unit cell for the space group $[I4_132]$.

FIGURE 12. Translational unit cells of the minimal surfaces S_P , and S_D match the translational unit cells of their associated graphs, Γ^1 , Γ^2 , and are displayed with the same viewing direction. The picture of S_G above shows the cube $[0, 2] \times [0, 2] \times [0, 2]$, twice the region of that used to depict Γ^4 , but depicted with the same viewing direction.

The following results are known (and can be checked):

- (a) Boundaries of equivariant regular neighbourhoods of Γ^1 , Γ^2 and Γ^4 in Example 3.3, 3.4 and 3.5 can be realized by S_P , S_D and S_G respectively.
- (b) S_P , S_D and S_G are unknotted in T^3 . Actually, any genus $g > 1$ minimal orientable closed surface in T^3 must be unknotted, [Me].

The notations of space groups below come from [Ha].

- (c) $\mathcal{G}^1$, $\mathcal{G}^2$, $\mathcal{G}^4$, $\mathcal{G}^{1,2}$ and $\mathcal{G}^{2,2}$ in Example 3.3, 3.4, 3.5, 4.1 and 4.2 are the space groups $[P432]$, $[F4_132]$, $[I4_132]$, $[I432]$ and $[P4_232]$ respectively.

(d) If we allow isometries that do not preserve orientations of the surface or of space then we find S_P , S_D , S_G are preserved by space groups $[Im\bar{3}m]$, $[Pn\bar{3}m]$, $[Ia\bar{3}d]$ respectively.

(e) We have the following index two subgroup sequences:

- $[P432] \subset [I432] \subset [Im\bar{3}m]$
- $[P432] \subset [Pm\bar{3}m] \subset [Im\bar{3}m]$
- $[F4_132] \subset [P4_232] \subset [Pn\bar{3}m]$
- $[I4_132] \subset [Ia\bar{3}d]$

Lemma 5.1. *Let T be a lattice in E^3 , $p : E^3 \rightarrow T^3 = E^3/T$ be the covering map. Σ_g is an embedded surface in T^3 . If $p^{-1}(\Sigma_g)$ is not connected, then Σ_g is knotted.*

Proof. Note first, from the definition of unknotted embedding $\Sigma_g \subset T^3$, the induced map on the fundamental groups must be surjective. Let F be a connected component of $p^{-1}(\Sigma_g)$. Let $St(F)$ be its stable subgroup in T , i.e., the lattice translations that preserve F . If $p^{-1}(\Sigma_g)$ has more than one component then $St(F) \neq T$, hence $\pi_1(\Sigma_g) \rightarrow \pi_1(T^3)$ is not surjective. Hence Σ_g is knotted in T^3 . $\square$

Proof of Theorem 1.5. By the above results (a) and (b), the surfaces in T^3 described in §3 are unknotted. Since $\Gamma^{1,2}$, $\Gamma^{2,2}$, $\Gamma^{4,4}$, $\Gamma^{4,8}$, Γ'^4 and Γ^P are disconnected, by Lemma 5.1, the surfaces in T^3 described in §4 are knotted. Then by combining the Examples in §3 and §4, we finish the proof of Theorem 1.5 (1).

Now we are going to prove Theorem 1.5 (2) and (3) based on the examples in §3 and §4, and the results (c), (d) and (e) above.

For (2), recall $\mathcal{G}^1 = [P432]$. Example 3.3 told us that when $m = n^3, 2n^3, 4n^3$ we have the action $\mathcal{G}^1/T_m$ on Γ^1/T_m , or equivalently on S_P/T_m by (a), of order $24m$. By (d) $[Im\bar{3}m]$ preserves S_P . By (e), we have an order $96m$ extendable $[Im\bar{3}m]/T_m$ -action on $S_P/T_m \cong \Sigma_{2m+1}$ realizing the upper bound of $E(\Sigma_g)$, where $g = 2m + 1$.

The group $[Im\bar{3}m]/T_m$ contains two elements h_1 and h_2 satisfying: h_1 preserves the orientation of S_P/T_m and reverses the orientation of E^3/T_m (e.g. reflection in a plane transverse to S_P) and h_2 reverses the orientation of S_P/T_m and preserves the orientation of E^3/T_m (e.g. rotation by π about an axis that lies in S_P). Example choices for these elements are

$$\begin{aligned} h_1 : (x, y, z) &\mapsto (1 - x, y, z) \\ h_2 : (x, y, z) &\mapsto (-x + 1/2, -z + 1/2, -y + 1/2) \end{aligned}$$

Then choosing the index two subgroups of $[Im\bar{3}m]/T_m$, $\langle [P432], h_1 \rangle/T_m = [Pm\bar{3}m]/T_m$ and $\langle [P432], h_2 \rangle/T_m = [I432]/T_m$, we get an extendable action on Σ_{2m+1} , realizing the upper bound of $E_+(\Sigma_g)$ and $E^+(\Sigma_g)$ respectively. Now (2) is proved.

For (3), recall $\mathcal{G}^2 = [F4_132]$. Example 3.4 told us that when $m = n^3, 4n^3, 16n^3$ we have the action $\mathcal{G}^2/T_{2m}$ on Γ^2/T_{2m} , or equivalently on

S_D/T_{2m} by (a), of order $24m$. Similar to the case for S_P above, we also have an order $96m$ extendable $[Pn\bar{3}m]/T_{2m}$ -action on $S_D/T_{2m} \cong \Sigma_{2m+1}$ realizing the upper bound of $E(\Sigma_g)$, where $g = 2m + 1$.

Furthermore, if $m = n^3, 4n^3, n \in \mathbb{Z}_+, n$ odd, $[Pn\bar{3}m]/T_{2m}$ contains a translation h reversing an orientation of S_D/T_{2m} and preserving an orientation of E^3/T_{2m} (see Example 4.2 for details, or see S_D in Figure 12 where h is the translation t_x). Modulo this translation we can get an order $48m$ extendable $[Pn\bar{3}m]/T_m$ -action on $S_D/T_m \cong \Pi_{2m+2}$, realizing the upper bound $24(g-2)$ of $E(\Pi_g)$, where $g = 2m+2$. Then choosing the index two subgroup $[P4_232] \subset [Pn\bar{3}m]$, we can get an extendable action on Π_{2m+2} , realizing the upper bound of $E^+(\Pi_g)$. Now (3) is proved. $\square$

Remark 5.2. In the above discussion for (3), the regular neighbourhood $N(S_D/T_m)$ of S_D/T_m is homeomorphic to a twisted $[-1, 1]$ -bundle over S_D/T_m . Its complement in E^3/T_m is essentially the regular neighbourhood $N(\Theta_m^{2,2})$ of $\Theta_m^{2,2}$ in Example 4.2. Similarly for $n \in \mathbb{Z}_+, n$ odd, and $m = n^3/2$, we can get an order $48m$ extendable $[Im\bar{3}m]/T_{n^3/2}$ -action on $S_P/T_{n^3/2} \cong \Pi_{2n^3+2}$, realizing the upper bound of $E(\Pi_g)$. Choosing the index two subgroup $[I432]$, we can get an extendable action on Π_{2n^3+2} , realizing the upper bound of $E^+(\Pi_g)$. The regular neighbourhood $N(S_P/T_{n^3/2})$ is homeomorphic to a twisted $[-1, 1]$ -bundle over $S_P/T_{n^3/2}$. Its complement in $E^3/T_{n^3/2}$ is essentially the regular neighbourhood $N(\Theta_{n^3/2}^{1,2})$ in Example 4.1. There is no similar construction for a regular neighbourhood of the Gyroid minimal surface because there is no translation that reverses the orientation of S_G/T_4 ; the handlebodies on each side of S_G are mirror images of one another.

REFERENCES

- [Br] K. Brakke, *Minimal surface page*,
<http://www.susqu.edu/brakke/evolver/examples/periodic/periodic.html>
- [FH] C. Frohman, J. Hass, *Unstable minimal surfaces and Heegaard splittings*. Invent. Math. 95 (1989), no. 3, 529-540.
- [Ha] T. Hahn (Ed.), *International tables for crystallography*, D. Reidel Publishing Company, (2005).
- [He] J. Hempel, *3-manifolds*, Princeton University Press, (1976).
- [Hu] A. Hurwitz, *Über algebraische Gebilde mit eindeutigen Transformationen in sich*, Mathematische Annalen, 41(3) (1892), 403-442.
- [HBLL+] S.T. Hyde, Z. Blum, T. Landh, S. Lidin, B.W. Ninham, S. Andersson and K. Larsson. *The Language of Shape: The Role of Curvature in Condensed Matter*. New York, USA: Elsevier, (1996).
- [HOP] S.T. Hyde, M. O’Keeffe, D.M. Proserpio, *A Short History of an Elusive Yet Ubiquitous Structure in Chemistry, Materials, and Mathematics*, Angewandte Chemie Int. Ed. 47, (2008), 79968000.
- [Ma1] C. L. May, *Automorphisms of compact Klein surfaces with boundary*, Pacific J. Math. 59 (1975), 199-210.
- [Ma2] C. L. May, *Large automorphism groups of compact Klein surfaces with boundary*, Glasgow Math. J. 1 (1977), 1-10.

- [Me] W. H. Meeks, *The conformal structure and geometry of triply periodic minimal surfaces in R^3* , Bull. Amer. Math. Soc. 83 (1977), no. 1, 134-136.
- [MY] W. H. Meeks, S. T. Yau, *The equivariant Dehn's lemma and loop theorem*, Commentarii Mathematici Helvetici, 56(1) (1981), 225-239.
- [RCSR] M. O'Keeffe, M.A. Peskov, S.J. Ramsden, O.M. Yaghi, *The Reticular Chemistry Structure Resource (RCSR) Database of, and Symbols for Crystal Nets* Accts. Chem. Res. 41, (2008), 1782-1789.
- [Schw] H. A. Schwarz, *Gesammelte Mathematische Abhandlungen*. Bd 1, Berlin: Springer, (1890).
- [Scho] A. H. Schoen, NASA Tech. Note No. D-5541, (1970)
- [WWZZ1] C. Wang, S. C. Wang, Y. M. Zhang, B. Zimmermann, *Extending finite group actions on surfaces over S^3* . Topology Appl. **160**, (2013), no. 16, 2088-2103.
- [WWZZ2] C. Wang, S. C. Wang, Y. M. Zhang, B. Zimmerman, *Maximal Orders of Extendable Finite Group Actions on Surfaces*. Groups Geom. Dyn. 9 (2015), no. 4, 1001-1045.
- [WWZ] C. Wang, S. C. Wang, Y. M. Zhang, *Maximal Orders of Extendable Actions on Surfaces*. Acta Math. Sin. (Engl. Ser.) 32 (2016), no. 1, 54-68.
- [Zi] B. Zimmermann, *Über Homomorphismen n -dimensionaler Henkelkörper und endliche Erweiterungen von Schottky-Gruppen*, Comment. Math. Helv. 56 (1981), 474-486.

SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, CHINA
E-mail address: Bais@math.pku.edu.cn

DEPARTMENT OF APPLIED MATHEMATICS, RESEARCH SCHOOL OF PHYSICS AND ENGINEERING, THE AUSTRALIAN NATIONAL UNIVERSITY, CANBERRA, AUSTRALIA
E-mail address: vanessa.robins@anu.edu.au

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI 230026, CHINA
E-mail address: chao_wang_1987@126.com

SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, CHINA
E-mail address: wangsc@math.pku.edu.cn