THE TOPOLOGY OF ASYMPTOTICALLY EUCLIDEAN
STATIC PERFECT FLUID SPACE-TIME

by

Abul Kasem Muhammad Masood-ul-Alam

A thesis submitted for the degree of

Doctor of Philosophy

of the

Australian National University

July 1985

ACKNOWLEDGEMENTS

I express my gratitude to my supervisor, Professor Leon Simon, for his interest in my work and for his continuous assistance. Also I wish to thank him very much for his many useful comments and suggestions in the preparation of this work. I should like to thank Dr Gary Bunting for informing me of the open problem of spherical symmetry of an asymptotically flat perfect fluid space-time, and my colleague, David Adams, for many helpful discussions. I particularly want to thank Mrs Barbara Geary for her excellent typing of this thesis. She has endured many awkward changes and revisions without complaint, and has transformed my messy drafts and redrafts into a beautifully typed manuscript.

I am grateful to the Australian National University for providing me with an ANU PhD Scholarship during my study.

CONTENTS

ACKNOWLEDGMENTS	1
STATEMENT	1
ABSTRACT	1
CHAPTER I. BASIC DEFINITIONS AND STATEMENT OF THE MAIN THEOREM	1
1.1. Notations and Conventions	1
1.2. The Main Theorem	1
CHAPTER II. THE STABILITY PROBLEM	1
2.1. The Stability Problem	1
2.2. The Stability Problem	1
2.3. Appendix: Some special cases for embedded disks	1
CHAPTER III. PROOF OF THE MAIN THEOREM	1
3.1. The Main Theorem	1
3.2. The Main Theorem	1
3.3. The Main Theorem	1
3.4. The Main Theorem	1
3.5. The Main Theorem	1
3.6. The Main Theorem	1
3.7. The Main Theorem	1
3.8. The Main Theorem	1
3.9. The Main Theorem	1
3.10. The Main Theorem	1
3.11. The Main Theorem	1
3.12. The Main Theorem	1
3.13. The Main Theorem	1
3.14. The Main Theorem	1
3.15. The Main Theorem	1
3.16. The Main Theorem	1
3.17. The Main Theorem	1
3.18. The Main Theorem	1
3.19. The Main Theorem	1
3.20. The Main Theorem	1
3.21. The Main Theorem	1
3.22. The Main Theorem	1
3.23. The Main Theorem	1
3.24. The Main Theorem	1
3.25. The Main Theorem	1
3.26. The Main Theorem	1
3.27. The Main Theorem	1
3.28. The Main Theorem	1
3.29. The Main Theorem	1
3.30. The Main Theorem	1
3.31. The Main Theorem	1
3.32. The Main Theorem	1
3.33. The Main Theorem	1
3.34. The Main Theorem	1
3.35. The Main Theorem	1
3.36. The Main Theorem	1
3.37. The Main Theorem	1
3.38. The Main Theorem	1
3.39. The Main Theorem	1
3.40. The Main Theorem	1
3.41. The Main Theorem	1
3.42. The Main Theorem	1
3.43. The Main Theorem	1
3.44. The Main Theorem	1
3.45. The Main Theorem	1
3.46. The Main Theorem	1
3.47. The Main Theorem	1
3.48. The Main Theorem	1
3.49. The Main Theorem	1
3.50. The Main Theorem	1
3.51. The Main Theorem	1
3.52. The Main Theorem	1
3.53. The Main Theorem	1
3.54. The Main Theorem	1
3.55. The Main Theorem	1
3.56. The Main Theorem	1
3.57. The Main Theorem	1
3.58. The Main Theorem	1
3.59. The Main Theorem	1
3.60. The Main Theorem	1
3.61. The Main Theorem	1
3.62. The Main Theorem	1
3.63. The Main Theorem	1
3.64. The Main Theorem	1
3.65. The Main Theorem	1
3.66. The Main Theorem	1
3.67. The Main Theorem	1
3.68. The Main Theorem	1
3.69. The Main Theorem	1
3.70. The Main Theorem	1
3.71. The Main Theorem	1
3.72. The Main Theorem	1
3.73. The Main Theorem	1
3.74. The Main Theorem	1
3.75. The Main Theorem	1
3.76. The Main Theorem	1
3.77. The Main Theorem	1
3.78. The Main Theorem	1
3.79. The Main Theorem	1
3.80. The Main Theorem	1
3.81. The Main Theorem	1
3.82. The Main Theorem	1
3.83. The Main Theorem	1
3.84. The Main Theorem	1
3.85. The Main Theorem	1
3.86. The Main Theorem	1
3.87. The Main Theorem	1
3.88. The Main Theorem	1
3.89. The Main Theorem	1
3.90. The Main Theorem	1
3.91. The Main Theorem	1
3.92. The Main Theorem	1
3.93. The Main Theorem	1
3.94. The Main Theorem	1
3.95. The Main Theorem	1
3.96. The Main Theorem	1
3.97. The Main Theorem	1
3.98. The Main Theorem	1
3.99. The Main Theorem	1
3.100. The Main Theorem	1
APPENDIX A	1
APPENDIX B	1
APPENDIX C	1
APPENDIX D	1
APPENDIX E	1
APPENDIX F	1
APPENDIX G	1
APPENDIX H	1
APPENDIX I	1
APPENDIX J	1
APPENDIX K	1
APPENDIX L	1
APPENDIX M	1
APPENDIX N	1
APPENDIX O	1
APPENDIX P	1
APPENDIX Q	1
APPENDIX R	1
APPENDIX S	1
APPENDIX T	1
APPENDIX U	1
APPENDIX V	1
APPENDIX W	1
APPENDIX X	1
APPENDIX Y	1
APPENDIX Z	1
REFERENCES	1

I declare that the work contained herein, excepting those parts for which there is an explicit statement to the contrary, is my own. The observation that Theorem 2.2 holds for $C^{1,1}$ metric is due to Professor Leon Simon. The rest of Chapter II, including the alternative method in the appendix at the end of this chapter, is my work.

A.K.H. Masood-ul-Alam

CONTENTS

ACKNOWLEDGEMENTS	(i)
STATEMENT	(ii)
ABSTRACT	(iv)
CHAPTER I: BASIC NOTATION AND STATEMENT OF THE MAIN THEOREM	1
1.1. Notation and Conventions	1
1.2. The Main Theorem	3
CHAPTER II: THE STABILITY INEQUALITY	8
2.0.	8
2.1.	8
2.2. Appendix: Some approximation results for embedded stable minimal surfaces	13
CHAPTER III: PROOF OF THE MAIN THEOREM	45
3.0.	45
3.1.	45
CHAPTER IV: APPLICATION TO GENERAL RELATIVITY	60
4.0.	60
4.1. Static Perfect Fluid Space-time	60
4.2. Gannon's Theorem	62
4.3. Extension to $C^{1,1}$ metric	67
CHAPTER V: MISCELLANEOUS REMARKS AND GENERALIZATIONS	73
5.0.	73
5.1.	73
5.2. Further Applications to General Relativity	76
APPENDIX A	80
APPENDIX B	85
APPENDIX C	86
APPENDIX D	89
APPENDIX E	91
REFERENCES	95

ABSTRACT

In this thesis we prove that a (geodesically) complete, asymptotically Euclidean, static perfect fluid space-time with connected fluid region and satisfying the time-like convergence condition is diffeomorphic to $\mathbb{R}^3 \times \mathbb{R}$. It is believed that such a space-time would be spherically symmetric at least for physically reasonable conditions on the density function ρ and the pressure function p .

The above assertion (that the space-time is diffeomorphic to $\mathbb{R}^3 \times \mathbb{R}$) has been claimed in [LB] provided the Poincaré conjecture is valid. In fact a theorem due to Gannon [G1] says that such a space-time is diffeomorphic to $N \times \mathbb{R}$ where N is a simply connected complete 3-manifold. The asymptotic conditions then imply that N has the same homotopy as $\mathbb{R}^3$ ([LB]). Thus Gannon's result reduces the question to proving the non-existence of fake 3-cells in N (in particular it would give the full result if the 3 dimensional Poincaré conjecture were known to be true).

Proving the non-existence of fake 3-cells in an appropriate class of asymptotically Euclidean Riemannian 3-manifolds is thus the main point of this thesis.

The main arguments needed for this are given in Chapter II and Chapter III of the thesis. In Chapter II we point out that the theorem of Meeks, Simon and Yau (Theorem 1 in [MSY]) applied in the present context, gives (in case there are fake 3-cells) a totally geodesic embedded sphere not intersecting the "fluid region" where $\rho + p > 0$. The argument to show this is similar to an argument of Frankel and Galloway [FG] (who also used the existence theorem of [MSY]), but here we need an additional approximation argument to make the appropriate stability statement. In the appendix to Chapter II we give an alternative approximation argument based on convergence

of sequences $\{\Sigma_k\}$ of smooth compact surfaces in a Riemannian 3-manifold (N, g) , where Σ_k is stable minimal in the smooth metrics k_g approximating the $C^{1,1}$ metric g . We believe the results here are of some independent interest.

In Chapter III we use an argument (inspired partly by an argument of Robinson [R] from his proof of spherical symmetry of the exterior Schwarchild solution) to prove that there cannot exist a totally geodesic embedded sphere which does not intersect the fluid region. Combined with the result of Chapter II, this completes the proof of the main theorem (1.1) for Riemannian 3-manifolds.

In Chapter IV we give Gannon's proof of simple connectedness modified to handle the case of $C^{1,1}$ metric and we complete the proof of the main result for static perfect fluid space-times.

In Chapter V we consider various generalizations of the main theorem (1.1). In particular, the topology of suitable space-like hypersurfaces in a certain class of space-times, not necessarily static or perfect fluid, is investigated.

CHAPTER I

BASIC NOTATION AND STATEMENT OF THE MAIN THEOREM

1.1. Notation and Conventions

With regards to tensor notation we use the following conventions.

Italic capital indices $A, B, C, \dots$ run from 0 to 3, Greek indices $\alpha, \beta, \gamma, \dots$ run from 1 to 3 and Italic indices $a, b, c, \dots$ run from 2 to 3.

In local co-ordinates, for a metric $g_{\alpha\beta}$ the Ricci curvature is $\text{Ric}(g)_{\alpha\beta} = R^{\mu}_{\alpha\mu\beta}$, where $R_{\alpha\beta\mu\nu}$ is the Riemann curvature tensor. For a vector field Z^{β} we have $R^{\alpha}_{\beta\mu\nu} Z^{\beta} = Z^{\alpha}_{;\nu\mu} - Z^{\alpha}_{;\mu\nu}$. Here $;$ denotes covariant differentiation in the g metric.

Let Σ be a smooth surface embedded in the smooth three dimensional Riemannian manifold (N, g) . (By smooth we shall mean C^{∞} .) In local coordinates $\parallel$ indicates covariant differentiation with respect to the induced connection on Σ . For a tensor T belonging to the tensor bundle of Σ we have

$$T^{\alpha\dots}_{\gamma\dots}\parallel_{\varepsilon} = \bar{T}^{\beta\dots}_{\delta\dots};_{\sigma} h^{\alpha}_{\beta} h^{\delta}_{\gamma} \dots h^{\sigma}_{\varepsilon} \text{ on } \Sigma$$

where $\bar{T}$ is a local extension of T in a neighbourhood of Σ in N , and where $h_{\alpha\beta} = g_{\alpha\beta} - n_{\alpha} n_{\beta}$, $h^{\alpha}_{\beta} = g^{\alpha\sigma} h_{\sigma\beta}$ on Σ , n_{α} being the unit normal form on Σ . Clearly $h^{\alpha\beta} n_{\alpha} = 0$, $h^{\alpha\beta}\parallel_{\varepsilon} = 0$.

The second fundamental form A of Σ is the tensor

$$A_{\alpha\beta} = h^{\nu}_{\alpha} h^{\mu}_{\beta} \bar{n}_{\nu;\mu}, \text{ where } \bar{n}_{\alpha} \text{ is any local extension of } n_{\alpha} \text{ in } N. \text{ Mean}$$

curvature $H = A_{\alpha\beta} g^{\alpha\beta}$. $|A|$ denotes the length of the second fundamental form: $|A|^2 = A_{\alpha\beta} A^{\alpha\beta}$.

The Riemann curvature of the induced metric on Σ is denoted by $h_{R\alpha\beta\gamma\delta}$ where

$$h_{R\alpha\beta\gamma\delta} Y^\beta = Y^\alpha_{||\delta\gamma} - Y^\alpha_{||\gamma\delta}$$

for all vector fields Y on Σ . The indices are raised using g . We include the embedding equations for easy reference.

$$\text{Gauss' equation: } h_{R\alpha\beta\gamma\delta} = R^\epsilon_{\rho\mu\nu} h^\alpha_\epsilon h^\rho_\beta h^\mu_\gamma h^\nu_\delta + A^\alpha_\gamma A_{\beta\delta} - A^\alpha_\delta A_{\beta\gamma}.$$

$$\text{Contracted Gauss' equation: } h_R = R - 2 \text{Ric}(g)_{\alpha\beta} n^\alpha n^\beta + H^2 - |A|^2, \quad h_R \text{ and } R \text{ being respectively the scalar curvatures of } \Sigma \text{ and } N.$$

$$\text{Codazzi's equation: } A^\alpha_{\beta||\alpha} - A^\alpha_{\alpha||\beta} = \text{Ric}(g)_{\epsilon\sigma} n^\delta h^\epsilon_\beta.$$

For a vector field Y on N , $Y|_p$ denotes its value at a point $p \in N$. $g(\cdot, \cdot)$ denotes the inner product on the tangent space of N and $\langle \cdot, \cdot \rangle$ denotes the corresponding induced inner product on Σ . In general the induced inner product on the tensor bundles of N and Σ will be denoted by $g(\cdot, \cdot)$ and by $\langle \cdot, \cdot \rangle$ respectively. $|T|^2 \equiv g(T, T)$. For a tensor field T defined on $\zeta_1 \subset N$ and for $\zeta_2 \subset \zeta_1$, $T|_{\zeta_2}$ denotes the restriction of T on ζ_2 .

$|\Sigma|$ denotes the area (that is the usual two dimensional Hausdorff measure H^2 in N) of Σ considered as a submanifold of the Riemannian manifold (N, g) . $|S|_E$ denotes the area of a surface S in $\mathbb{R}^3$. L^n denotes the n -dimensional Lebesgue measure in $\mathbb{R}^n$.

∇ and Δ will respectively denote the gradient operators and the Laplacian on (N, g) . ∇_Σ and Δ_Σ are the corresponding operators on Σ with respect to the induced metric. ${}^4\nabla$ and ${}^4\Delta$ will respectively denote the gradient operators and the Laplacian for the four dimensional Lorentz metric 4g of a spacetime $(M, {}^4g)$.

For a curve $\gamma : [\alpha, \beta] \rightarrow N$, parametrized by t , $(\partial/\partial t)_\gamma|_t$ denotes the tangent vector at $\gamma(t)$.

For subsets ζ_1 and ζ_2 of a topological space T , $\zeta_1 \subset\subset \zeta_2$ means that $\overline{\zeta_1}$ is a compact subset of ζ_2 where $\overline{\zeta_1}$ is the closure of ζ_1 in T . $\zeta_1 \sim \zeta_2$ denotes the set of points in ζ_1 but not in ζ_2 .

$B_R(x)$ is the open ball in $\mathbb{R}^3$ with radius R and centre at $x \in \mathbb{R}^3$. $B_R^2(x)$ denotes the open disk in $\mathbb{R}^2$ with radius R and centre at $x \in \mathbb{R}^2$.

For a bounded, open, connected subset Ω of $\mathbb{R}^n$, $C^{k,\lambda}(\Omega)$, $k \geq 0$, $\lambda \in (0, 1)$, denotes the space of C^k functions on Ω whose k th order partial derivatives are locally Holder continuous in Ω with exponent λ . $C^{k,1}(\Omega)$ denotes the space of C^k functions whose k th order partial derivatives are locally Lipschitz continuous in Ω . For $1 \leq p < \infty$, $W^{k,p}(\Omega)$ denotes the Sobolev space; that is, $W^{k,p}(\Omega)$ is the set of all k times weakly differentiable functions u in Ω such that $D^\alpha u \in L^p(\Omega)$ for all $|\alpha| \leq k$. For a function ψ , $\text{supp } \psi$ denotes its support.

1.2. The Main Theorem

The main theorem we prove is the following.

1.1. THEOREM. *Assumptions: $(N, \{u_\alpha, \varphi_\alpha\}_{\alpha \in A})$ is a C^∞ simply connected three dimensional differentiable manifold with a complete Riemannian metric g satisfying the following conditions:*

(a) *In local co-ordinates given by φ_α , $g_{\alpha\beta} \in C^{1,1}(\varphi_\alpha(u_\alpha))$ and g satisfies*

$$1.2 \quad \text{Ric}(g)_{\alpha\beta} = V^{-1}V_{;\alpha\beta} + \Phi_1 g_{\alpha\beta}$$

and

$$1.3 \quad \Delta V = V\Phi_2 ,$$

where ∇ denotes the covariant differentiation for g , Δ is the Laplacian for g , V is a locally $C^{1,1}$ positive function on N , Φ_1, Φ_2 are bounded measurable functions on N , and $\Phi_2 \geq 0$. (The above equations are assumed to hold almost everywhere in N .)

(b) There exists an open connected set $Q \subset N$ such that $\text{ess inf}_K (\Phi_1 + \Phi_2) > 0$ for all compact $K \subset Q$, and $\Phi_1, \Phi_2 = 0$ in $N \sim \bar{Q}$.

(c) There exists an open connected set $N_0 \subset N$ such that $\bar{N}_0$ is compact and $N \sim \bar{N}_0$ is diffeomorphic to $\mathbb{R}^3 \sim \bar{B}_1$ where $\bar{B}_1$ is the closed unit ball centered at the origin and, with respect to the standard co-ordinate system in $\mathbb{R}^3$, we have, on $N \sim \bar{N}_0$,

$$1.4 \quad g_{\alpha\beta} = \delta_{\alpha\beta} + O(|x|^{-\lambda}) \quad \text{and} \quad \frac{\partial g_{\alpha\beta}}{\partial x^\sigma} = O(|x|^{-1-\lambda})$$

for some $\lambda \in (0, 1)$, where $|x| \equiv \left(\sum_{\alpha=1}^3 (x^\alpha)^2 \right)^{\frac{1}{2}} \rightarrow \infty$.

Assertion: N is topologically $\mathbb{R}^3$ and hence diffeomorphic to $\mathbb{R}^3$.

1.5. Remark. Henceforth the co-ordinate system in (c) above will be referred to as the asymptotic co-ordinate system.

1.6. Remark. On $N \sim \bar{Q}$ we have $\Delta V = 0$. Hence on $N \sim \bar{Q}$, V is locally $C^{2,\mu}$, $\mu \in (0, 1)$ (see the paragraph following 3.9 in Chapter III). It is shown in Chapter III that when $\bar{Q}$ is compact, by virtue of 1.4 we have in the asymptotic co-ordinate system,

$$1.7 \quad V = C_1 - \frac{C_2}{|x|} + \eta, \quad \text{where} \quad \eta = O(|x|^{-1-\beta}), \quad \frac{\partial \eta}{\partial x^\alpha} = O(|x|^{-2-\beta}),$$

and
$$\left(R^{-3} \int_{N_{3R} \sim N_R} \left| \frac{\partial^2 \eta}{\partial x^\alpha \partial x^\tau} \right|^2 \right)^{\frac{1}{2}} = O(R^{-3-\beta}) \quad \text{for some } \beta \in (0, 1),$$

constants $C_1 > 0$ and $C_2 \geq 0$. Here $N_R \subset N$ is such that in the

asymptotic co-ordinate system, $N \sim N_R = \mathbb{R}^3 \sim B_R(0)$. Without loss of generality we write 1.7 as

1.8
$$V = 1 - \frac{m}{|x|} + \eta \quad \text{where } m \geq 0 \quad \text{and } \eta \text{ is as above.}$$

It is well-known that when $m = 0$ and Φ_1 tends to 0 at infinity, then

g is flat. To see this we note that in this case 1.3 and 1.8 imply

$V \equiv \text{constant}$. Hence by 1.2 we have $\text{Ric}(g)_{\alpha\beta} = \Phi_1 g_{\alpha\beta}$. The

contracted Bianchi identity for g (see (A1) in Appendix A) then

implies that $\Phi_1 \equiv \text{constant}$. Thus $\Phi_1 \equiv 0$ and g is Ricci flat and hence flat.

1.9. Remark. The contracted Bianchi identity for g implies that $\Phi_2 - \Phi_1$ is a Lipschitz function on N and

1.10
$$(\Phi_2 - \Phi_1)_{;\beta} = -2V^{-1}(\Phi_1 + \Phi_2)V_{;\beta}$$

(see Appendix A). For simplicity we put

1.11
$$\begin{cases} \Phi_2 - \Phi_1 = 16\pi p, \\ 3\Phi_1 + \Phi_2 = 16\pi \rho, \end{cases}$$

where the factor 16π has been used in accordance with the usual convention of Physics Literature and has no significance here. It follows from 1.10 that p and ρ are functions of V and

1.12
$$\frac{dp}{dV} = -V^{-1}(\rho(V) + p(V))$$

at least when $V^{-1}(t)$ and $V^{-1}(t) \sim \{x \in N \mid \nabla V(x) = 0\}$ are connected sets

for each $t \in \text{Range } V$ and the critical set $C \equiv \{x \in N \mid \nabla V(x) = 0\}$ has H^3 measure zero (see Appendix A for details).

It is shown in [KS] that if $\bar{Q}$ is compact and p is non-negative then ρ cannot be a Lipschitz function of p unless $p \equiv 0$. In [KS] it is assumed that V has one critical point, and the argument is as follows: we put $V = e^u$. Then using $-Kp \leq \rho(0) - \rho(p)$ for some constant K we get, from 1.12,

$$\frac{d}{du} (pe^{(1+K)u}) \geq -\rho(0)e^{(1+K)u}.$$

This implies, for $u \leq u_0$,

$$p(u)e^{(1+K)u} \leq p(u_0)e^{(1+K)u_0} + \rho(0) \int_u^{u_0} e^{(1+K)s} ds.$$

Now suppose $u = u_0$ on the boundary ∂Q of Q . Then $p(u_0) = 0$ since $p \equiv 0$ on $N \sim \bar{Q}$ and, by 1.10, p is continuous. Thus

$$1.13 \quad p(u)e^{(1+K)u} \leq \rho(0) \int_u^{u_0} e^{(1+K)s} ds.$$

Also by applying the maximum principle to 1.3 we get $e^u \leq e^{u_0}$ in Q .

Hence if $\rho(0) = 0$ we have, by 1.13, $p \equiv 0$ in Q since p is assumed to be non-negative. Thus if $d\rho/dp$ is bounded then either ρ must have a jump discontinuity at ∂Q or (N, g) is flat by virtue of 1.10.

We give an alternative argument which does not need the assumptions on the critical points of V . If ρ is a Lipschitz function of p then 1.10 and 1.3 give

$$\Delta p + \left(2 + \frac{d\rho}{dp}\right) V^{-1} V_{;\beta} p_{;\beta} = -4\pi(\rho+p)(\rho+3p) \leq 0.$$

Now p vanishes outside $\bar{Q}$ and $N \sim \bar{Q} \neq \emptyset$. Hence by the (generalized) strong maximum principle (Theorem 8.19 in [GT]), $p \equiv 0$.

1.14. Remark. The assumption $\Phi_2 \geq 0$ in (b) is not necessary in case

Q is unbounded. This is because in this case, if a sphere $S \subset N$ with $S \cap Q = \emptyset$ bounds the compact manifold-with-boundary $N_1 \subset N$ then $N_1 \cap Q = \emptyset$ (since, by hypothesis Q is connected; for details see Remark 3.1 in Chapter III). Hence Case II in the proof of Theorem 3.22 cannot occur automatically. The proof of Case II uses 1.8 and that $V < 1$ on $N \sim \bar{Q}$. The condition $\Phi_2 \geq 0$ enables us to use the maximum principle to deduce that $V < 1$ on N and also to determine the sign of C_2 in 1.7.

1.15. Remark. A physical example where g and V are piecewise C^3 and ρ is discontinuous across ∂Q is the static stellar model described in [KS] and [L]. Our asymptotic conditions are slightly more general than those given in [KS] or [K]. In particular (when $\bar{Q}$ is compact) we have deduced the asymptotic expression 1.7 for V instead of assuming it.

1.16. Remark. Further generalizations of Theorem 1.1 are given in Chapter V. In particular alternative definitions for " Q " are given in Section 5.1.

CHAPTER II

THE STABILITY INEQUALITY

2.0.

In this chapter we use a theorem of Meeks, Simon and Yau (Theorem 1 in [MSY]) regarding the existence of minimal surfaces in 3-manifolds to prove that either N is topologically Euclidean or there exists an embedded totally geodesic sphere in $N \sim Q$. The proof is essentially a straightforward modification of a result due to Frankel and Galloway (corollary to Theorem 1 in [FG]). However, since the metric is only $C^{1,1}$ in our case, we have to deduce a suitable form of the "stability inequality" involving the Ricci curvature which, in our case, is only defined almost everywhere and hence does not, in general, make sense on the minimal surface.

2.1.

Let $\mathcal{C}$ denote the collection of all connected compact C^2 2-dimensional surfaces-without-boundary embedded in N , and let $\mathcal{C}_1$ denote the collection of compact embedded surfaces Σ such that each component of Σ is an element of $\mathcal{C}$. Given $\Sigma \in \mathcal{C}$ we let $I(\Sigma)$ be the collection of all $\hat{\Sigma}$ such that $\hat{\Sigma}$ is isotopic to Σ via a smooth isotopy:

$\psi : [0, 1] \times N \rightarrow N$ where

(i) $\psi_0(x) \equiv \psi(0, x) = 1_N(x)$, 1_N being the identity map on

N ;

(ii) ψ_t , defined by $\psi_t(x) = \psi(t, x)$, $(t, x) \in [0, 1] \times N$,

is a diffeomorphism of N onto N ;

(iii) $\psi_t|_{N \sim K} = 1_{N \sim K}$ for $t \in [0, 1]$ for some compact set

$K \subset N$ independent of t .

We shall say that a two-sided surface $S \in \mathcal{C}$ is area minimizing if there exists $d > 0$ such that

$$2.1 \quad |S| \leq \inf_{\substack{\hat{\Sigma} \in \mathcal{I}(S) \\ \hat{\Sigma} \subset \{x \in N \mid \text{dist}(x, S) < d\}}} |\hat{\Sigma}|.$$

Now from the asymptotic conditions (c) in the hypothesis of the main Theorem 1.1 it follows that there exists a smooth sphere S_r in $N - \bar{Q}$ given by $|x| = r$ in the asymptotic co-ordinate system such that the mean curvature of S_r in (N, g) with respect to the outward normal is strictly positive. Let $\bar{N}_1$ be the simply connected compact submanifold of N with boundary $\partial \bar{N}_1 = S_r$ (that is, in asymptotic co-ordinate system $N \sim \bar{N}_1 = \mathbb{R}^3 \sim \bar{B}_r(0)$). Then

2.2 THEOREM [MSY]. *Either $\bar{N}_1$ is diffeomorphic to a closed unit ball in $\mathbb{R}^3$ or there exists a $C^{2,\alpha}$, $\alpha \in (0, 1)$, embedded area minimizing minimal sphere S in the interior N_1 of $\bar{N}_1$.*

Proof of 2.2. It is a particular case of the more general fundamental existence theorem (Theorem 1) in [MSY]. We note that the only places in the proof of the latter theorem where the smoothness of the metric is used are in the definition of homogeneous regularity and in the arguments relating to homothetic expansion on page 639. (As an alternative method we can avoid the use of the above theorem for $C^{1,1}$ metric altogether and can directly apply the smooth metric version of this theorem (which is proved in [MSY]) to a sequence of smooth metrics approximating the $C^{1,1}$ metric to prove our final theorem (2.8) of this chapter. This alternative approach is given in the appendix at the end of this chapter.) On page 639 in [MSY] it is assumed that N can be locally isometrically embedded in the Euclidean space. This can be avoided by considering local co-ordinates representation for N in the neighbourhood of a point x_0 with the $C^{1,1}$

metric $g_{\alpha\beta}$ satisfying $g_{\alpha\beta}(0) = \delta_{\alpha\beta}$ and $\left(\partial g_{\alpha\beta}/\partial x^\tau\right)(0) = 0$ where 0 is the image of x_0 . The definition of homogeneous regularity can be modified by having a countable atlas with uniformly bounded $C^{1,1}$ norm of the metric in each chart. By hypothesis N has such an atlas. Finally, we note that since N is simply connected there cannot be any embedded projective space in N (by Theorem 4.7 on page 108 in [H]). $\square$

We shall now deduce a suitable form of the "stability inequality" from 2.1 and use this inequality to prove the existence of a totally geodesic embedded sphere $S \subset N \sim Q$ in case N is not topologically Euclidean.

We take an 1-parameter family $\{\varphi_t\}_{0 \leq t \leq 1}$ of diffeomorphisms $N_2 \rightarrow N_2$ of some neighbourhood N_2 of S in N such that

- (i) $\varphi(t, x) \equiv \varphi_t(x)$ is a C^2 map: $(-1, 1) \times N_2 \rightarrow N_2$;
- (ii) $\varphi_0(x) \equiv x$, $x \in N_2$;
- (iii) $\partial\varphi(t, x)/\partial t|_{0,S} = \xi \hat{n}$,

where ξ is a smooth function with bounded C^1 norm in N_2 and $\hat{n}$ is a smooth vector field on S such that for some given $\varepsilon > 0$, $|\hat{n} - n_0|^2 < \varepsilon$ and $|\nabla_Y(\hat{n} - n_0)|^2 < \varepsilon|Y|^2$ for all vectors $Y \in T_x N$, $x \in S$, n_0 being a unit normal vector field on S .

Now let $f(t) = |S_t|$ where $S_t = \varphi_t(S)$. Then $df/dt|_0 = 0$ since S is area minimizing. Also given any interval $I_j = (-1/j, 1/j)$ (however small) about $t = 0$ and given any $\bar{t} \in I_j$ we have

$$0 \leq f(\bar{t}) - f(0) = \int_0^{\bar{t}} \frac{df}{dt} dt = \int_0^{\bar{t}} \int_0^t \frac{d^2f}{ds^2} ds dt .$$

Thus $d^2f/ds^2 \geq 0$ on a subset $A_j \subset I_j$ with A_j having positive measure.

Now there exist $C_j \subset A_j$ with C_j having positive measure such that, for all $t \in C_j$, $\partial^2 g_{\alpha\beta} / \partial x^\sigma \partial x^\tau$ exists almost everywhere on S_t . To see this we consider the local chart (U_i, φ_i) . Let B be the set

$$\left\{ x \in U_i \mid x \in \varphi_t(S), t \in \left[-\frac{1}{j}, \frac{1}{j} \right] \text{ for sufficiently large } j \right.$$

$$\left. \text{and some } \frac{\partial^2 g_{\alpha\beta}}{\partial x^\sigma \partial x^\tau} \text{ is not defined at } x \right\}.$$

Then

$$0 = \int_{\varphi_i(B)} |\nabla t| dL^3 = \int_{-\bar{t}}^{\bar{t}} H^2(\varphi_i(B \cap S_t)) dt$$

by the co-area formula. Thus for almost all $t \in [-\bar{t}, \bar{t}]$, $H^2(B \cap S_t) = 0$.

For any $t \in C_j$ we have the following second variation formula on

S_t :

$$\left(\frac{d^2}{ds^2} |S_s| \right)_{s=t} = \int_{S_t} \left(\operatorname{div}_{S_t} Z + (\operatorname{div}_{S_t} X)^2 + \sum_{a=1}^2 |(\nabla_{\tau_a} X)^\perp|^2 \right. \\ \left. - \sum_{b,a=1}^2 \langle \tau_a, \nabla_{\tau_b} X \rangle \langle \tau_b, \nabla_{\tau_a} X \rangle - \sum_{a=1}^2 \langle R(\tau_a, X)X, \tau_a \rangle \right)$$

where τ_a is an orthonormal vector basis on S_t , $X = (d/ds)(\varphi_s(x))|_{s=t}$,

$x \in S_t$; $Z = (d^2/ds^2)(\varphi_s(x))|_{s=t}$, $x \in S_t$; $(\nabla_{\tau_a} X)^\perp$ is the part of $\nabla_{\tau_a} X$

normal to S_t and $(R(\cdot, \cdot)\cdot)$ is the Riemann tensor.

Restricting t to a subset D_j of C_j with D_j having positive measure, we have $X = \xi(n + \eta)$ where n is the unit normal vector to S_t

and where $|\eta|^2 < \varepsilon$, $|\nabla_Y \eta|^2 < \varepsilon|Y|^2$ for $Y \in T_x N$, $x \in S_t$. Hence we

have, for $t \in D_j$,

$$\begin{aligned}
2.3 \quad 0 &\leq \left(\frac{d^2}{ds^2} |S_s| \right)_{s=t} \\
&= \int_{S_t} \left(g(Z, n) H_t + \xi^2 H_t^2 + |\nabla_{S_t} \xi|^2 - |A_t|^2 \xi^2 - \xi^2 \text{Ric}(g)(n, n) \right) + c\epsilon
\end{aligned}$$

where A_t is the second fundamental form of S_t and H_t is the mean curvature of S_t . c is a constant depending only on C^1 norm of ξ .

This implies

$$\int_{S_t} \left(\xi^2 \left(|A_t|^2 + \text{Ric}(g)(n, n) \right) - |\nabla_{S_t} \xi|^2 \right) \leq \int_{S_t} \left(g(Z, n) H_t + \xi^2 H_t^2 \right) + c\epsilon.$$

Now there exists $E_j \subset D_j$ also with positive measure such that 1.2 and 1.3 hold almost everywhere on S_t for all $t \in E_j$. Hence taking a smooth ξ such that the C^1 norm of $(\xi - V)$ in N_2 is less than ϵ we get, for all $t \in E_j$,

$$2.4 \quad \int_{S_t} \left(V^2 \left(|A_t|^2 + \text{Ric}(g)(n, n) \right) - |\nabla_{S_t} V|^2 \right) \leq \int_{S_t} \left(g(Z, n) H_t + V^2 H_t^2 \right) + c\epsilon.$$

This implies, since $V|_{S_t}$ is $W^{2,p}(S_t)$,

$$2.5 \quad \int_{S_t} \left(V^2 \left(|A_t|^2 + \text{Ric}(g)(n, n) \right) + V \Delta_{S_t} V \right) \leq \int_{S_t} \left(g(Z, n) H_t + V^2 H_t^2 \right) + c\epsilon.$$

Now using the following well known formula derived in Appendix B,

$$2.6 \quad \Delta \psi = \Delta_{S_t} \psi + \psi_{;\alpha\beta} n^\alpha n^\beta + H_t \nabla \psi(n) \quad \text{for any } C^2 \psi$$

which also holds for V on S_t , $t \in E_j$, we have

$$\begin{aligned}
2.7 \quad \int_{S_t} \left(V^2 \left(|A_t|^2 + \text{Ric}(g)(n, n) \right) + V \Delta V - V V_{;\alpha\beta} n^\alpha n^\beta \right) \\
\leq \int_{S_t} \left(g(Z, n) H_t + V^2 H_t^2 + V H_t \nabla V(n) \right) + c\epsilon.
\end{aligned}$$

But $V^2 \text{Ric}(g)(n, n) + V\Delta V - VV_{;\alpha\beta}n^\alpha n^\beta = V^2(\Phi_1 + \Phi_2) \geq V^2 h$ where h is a continuous function in N with $h > 0$ in Q . Hence

$$\int_{S_t} V^2(|A_t|^2 + h) \leq \int_{S_t} \left(g(Z, n)H_t + V^2 H_t^2 + VH_t \nabla V(n) \right) + c\varepsilon$$

for all $t \in E_j$.

But $E_j \subset (-1/j, 1/j)$. Hence letting $j \rightarrow \infty$ and noting that $S = S_0$ is a C^2 minimal surface we have

$$\int_S V^2(|A|^2 + h) \leq 0$$

where A is the second fundamental form of S . Since $h > 0$ in Q we get that $|A| = 0$ and $|S \cap Q| = 0$. Since S is C^2 we have proved the following theorem:

2.8 THEOREM. *Either N is diffeomorphic to $\mathbb{R}^3$ or there exists a $C^{2,\alpha}$, $\alpha \in (0, 1)$, embedded totally geodesic sphere S in $N \sim Q$.*

2.2. APPENDIX: Some approximation results for embedded stable minimal surfaces

In this appendix we discuss convergence of sequences $\{\Sigma_k\}$ of stable minimal surfaces in N , where Σ_k is stable minimal relative to a smooth metric k_g approximating the $C^{1,1}$ metric g . This approximation is done in such a way that the relevant stability inequalities behave well under the limiting operation. We believe the results here are of some independent interest, and they also provide an alternative method for deducing the results of the above chapter. In particular we prove Theorem 2.8 by appealing only to the original (smooth metric) version of Theorem 2.2 which was proved in [MSY].

At first, following Schoen, Simon and Yau [SSY], we prove an L^5

estimate for the length of the second fundamental form of a compact stable minimal surface in a smooth Riemannian 3-manifold. Our aim is to get a uniform estimate for a sequence of such surfaces Σ_k with uniformly bounded area in the smooth Riemannian manifolds $(N, {}^k g)$ having uniformly bounded curvature and with ${}^k g \rightarrow g$ in the $C^{1,\alpha}$ norm. The proof differs slightly from that of [SSY] in that we get rid of terms involving the third derivatives of the metrics by suitable integration by parts over the compact surfaces. Using the above curvature estimate a compactness result (Lemma 2.90) for the sequence $\{\Sigma_k\}$ is obtained. Finally the compactness result is used to give an alternative proof of Theorem 2.8.

Curvature estimate

2.9 LEMMA. *Suppose $\tilde{\Sigma}$ is an embedded compact, stable minimal surface in a 3-dimensional smooth Riemannian manifold $(N, \tilde{g})$. Let $|A|$ be the length of the second fundamental form of $\tilde{\Sigma}$, $R_{\alpha\beta\delta\gamma}$ be the curvature tensor of $(N, \tilde{g})$ and $|\tilde{\Sigma}|_{\tilde{g}}$ be the area of $\tilde{\Sigma}$ with respect to the induced metric. Then*

$$2.10 \quad \int_{\tilde{\Sigma}} |A|^5 \leq \beta_1 |\tilde{\Sigma}|_{\tilde{g}}$$

where β_1 is a constant depending only on the bound

$$\sup_{\tilde{\Sigma}} |R_{\alpha\beta\delta\gamma} R^{\alpha\beta\delta\gamma}| \leq \beta_0.$$

Proof of 2.9. We reproduce the proof in [SSY] except for the modification necessary in our case.

Let n_α be the unit normal form on $\tilde{\Sigma}$ and $\bar{n}_\alpha$ be an extension of n_α in a neighbourhood of $\tilde{\Sigma}$. Differentiating covariantly the second fundamental form $A_{\alpha\beta} = \bar{n}_{\nu;\rho} h^\nu_\alpha h^\rho_\beta$ with respect to the induced connection on $\tilde{\Sigma}$ we get

$$2.11 \quad A_{\alpha\beta||\mu} - A_{\alpha\mu||\beta} = h^\nu_\alpha h^\rho_\beta h^\sigma_\mu R_{\nu\epsilon\sigma\rho} n^\epsilon.$$

Here ; and || denote covariant differentiations with respect to the connection for $\tilde{g}$ and the induced connection respectively, and

$h^{\alpha\beta} = \tilde{g}^{\alpha\beta} - n^\alpha n^\beta$. The calculation becomes easy by noting that on $\tilde{\Sigma}$ we have $h^{\alpha\beta} n_\alpha = 0$ and $\bar{n}_{\alpha;\rho} n^\alpha h^\rho_\beta = 0$, so that

$$\begin{aligned} A_{\alpha\beta||\mu} &= \left[\bar{n}_{\nu;\rho} (\delta^\nu_\alpha, -\bar{n}^\nu \bar{n}_\alpha,) (\delta^\rho_\beta, -\bar{n}^\rho \bar{n}_\beta,) \right]_{;\mu} h^{\alpha'}_\alpha h^{\beta'}_\beta h^{\mu'}_\mu \\ &= \bar{n}_{\nu;\rho\mu} h^\nu_\alpha h^\rho_\beta h^{\mu'}_\mu - h^\nu_\alpha \bar{n}_{\nu;\rho} n^\rho A_{\beta\mu}. \end{aligned}$$

Differentiating covariantly 2.11 we get

$$2.12 \quad A_{\alpha\beta||\mu\pi} = \left(h^\nu_\alpha h^\rho_\beta h^\sigma_\mu R_{\nu\epsilon\sigma\rho} n^\epsilon \right)_{||\pi} + A_{\alpha\mu||\beta\pi}.$$

Now let $h^\alpha_{R\beta\gamma\delta}$ be the curvature tensor of the induced metric. Then the commutation rule gives

$$2.13 \quad A_{\alpha\beta||\mu\pi} - A_{\alpha\beta||\pi\mu} = h^\nu_{R\alpha\nu\pi\mu} A^\nu_\beta - h^\nu_{R\beta\pi\mu} A_{\alpha\nu}.$$

Hence from 2.12 we have

$$\begin{aligned} 2.14 \quad A_{\alpha\beta||\mu\pi} &= \left(h^\nu_\alpha h^\rho_\beta h^\sigma_\mu R_{\nu\epsilon\sigma\rho} n^\epsilon \right)_{||\pi} + A_{\alpha\mu||\pi\beta} + h^\nu_{R\alpha\nu\pi\beta} A^\nu_\mu - h^\nu_{R\mu\pi\beta} A_{\alpha\nu} \\ &= \left(h^\nu_\alpha h^\rho_\beta h^\sigma_\mu R_{\nu\epsilon\sigma\rho} n^\epsilon \right)_{||\pi} + A_{\alpha\mu||\pi\beta} + R_{\sigma\tau\omega\rho} h^\sigma_\alpha h^\omega_\pi h^\rho_\beta A^\tau_\mu + A_{\alpha\pi} A_{\nu\beta} A^\nu_\mu \\ &\quad - A_{\alpha\beta} A_{\nu\pi} A^\nu_\mu - R_{\sigma\tau\omega\rho} h^\sigma_\nu h^\tau_\mu h^\omega_\pi h^\rho_\beta A^\nu_\alpha - A_{\nu\pi} A_{\mu\beta} A^\nu_\alpha + A_{\nu\beta} A_{\mu\pi} A^\nu_\alpha \end{aligned}$$

where we have used the Gauss' equation

$$2.15 \quad h^\nu_{R\alpha\nu\pi\beta} = R_{\sigma\tau\omega\rho} h^\sigma_\alpha h^\tau_\nu h^\omega_\pi h^\rho_\beta + A_{\alpha\pi} A_{\nu\beta} - A_{\alpha\beta} A_{\nu\pi}.$$

2.14 implies

$$\begin{aligned} 2.16 \quad h^{\mu\pi} A_{\alpha\beta||\mu\pi} &\equiv \Delta_{\tilde{\Sigma}} A_{\alpha\beta} = \left(h^\nu_\alpha h^\rho_\beta h^{\sigma\pi} R_{\nu\epsilon\sigma\rho} n^\epsilon \right)_{||\pi} + A_{\alpha\mu||}^\mu{}_\beta + R_{\sigma\tau\omega\rho} h^\sigma_\alpha h^{\omega\mu} h^\rho_\beta A^\tau_\mu \\ &\quad + A_{\alpha\pi} A_{\nu\beta} A^\nu_\mu - A_{\alpha\beta} |A|^2 - R_{\sigma\tau\omega\rho} h^{\tau\omega} h^\rho_\beta A^\sigma_\alpha - A_{\nu\pi} A^\pi_\beta A^\nu_\alpha + A_{\nu\beta} A^\nu_\alpha A^\mu_\mu, \end{aligned}$$

where we have used $h^{\mu\pi}_{||\alpha} = 0$ and $A_{\alpha\mu||\pi\beta} h^{\mu\pi} = A_{\alpha\mu||\beta}{}^{\mu}$. Thus using

$A_{\mu}{}^{\mu} = 0$ and Codazzi's equation

$$2.17 \quad A_{\alpha}{}^{\mu}{}_{||\mu} - A_{\mu}{}^{\mu}{}_{||\alpha} = R_{\rho\tau} n^{\tau} h^{\rho}{}_{\alpha}$$

we get

$$2.18 \quad \Delta_{\tilde{\Sigma}} A_{\alpha\beta} = \left(h^{\nu}{}_{\alpha} h^{\rho}{}_{\beta} h^{\sigma\pi} R_{\nu\epsilon\sigma\rho} n^{\epsilon} \right)_{||\pi} + \left(R_{\rho\tau} n^{\tau} h^{\rho}{}_{\alpha} \right)_{||\beta} + R_{\sigma\tau\omega\rho} A^{\tau\omega} h^{\sigma}{}_{\alpha} h^{\rho}{}_{\beta} \\ - A_{\alpha\beta} |A|^2 - R_{\sigma\tau\omega\rho} A^{\sigma}{}_{\alpha} h^{\tau\omega} h^{\rho}{}_{\beta}.$$

Contracting with $A^{\alpha\beta}$ and writing the terms involving the third derivatives of the metric $\tilde{g}$ in divergence form we have

$$2.19 \quad A^{\alpha\beta} \Delta_{\tilde{\Sigma}} A_{\alpha\beta} = \left(A^{\nu\rho} h^{\sigma\pi} R_{\nu\epsilon\sigma\rho} n^{\epsilon} \right)_{||\pi} - A^{\nu\rho}{}_{||}{}^{\sigma} R_{\nu\epsilon\sigma\rho} n^{\epsilon} \\ + \left(A^{\rho\beta} R_{\rho\tau} n^{\tau} \right)_{||\beta} - A^{\rho\beta}{}_{||\beta} R_{\rho\tau} n^{\tau} + R_{\sigma\tau\omega\rho} \left(A^{\tau\omega} A^{\sigma\rho} - A^{\sigma}{}_{\alpha} A^{\alpha\rho} h^{\tau\omega} \right) - |A|^4.$$

Now from 2.11 we have

$$A^{\nu\rho}{}_{||}{}^{\sigma} R_{\nu\epsilon\sigma\rho} = A^{\nu\sigma}{}_{||}{}^{\rho} R_{\nu\epsilon\sigma\rho} + h^{\nu\tau} h^{\sigma\mu} h^{\rho\alpha} R_{\tau\pi\mu\alpha} n^{\pi} R_{\nu\epsilon\sigma\rho} \\ = A^{\nu\rho}{}_{||}{}^{\sigma} R_{\nu\epsilon\sigma\rho} + h^{\nu\tau} h^{\sigma\mu} h^{\rho\alpha} R_{\tau\pi\mu\alpha} n^{\pi} R_{\nu\epsilon\sigma\rho}.$$

Hence using 2.17 we obtain, from 2.19,

$$2.20 \quad A^{\alpha\beta} \Delta_{\tilde{\Sigma}} A_{\alpha\beta} = \left(A^{\nu\rho} h^{\sigma\pi} R_{\nu\epsilon\sigma\rho} n^{\epsilon} \right)_{||\pi} + \left(A^{\rho\beta} R_{\rho\tau} n^{\tau} \right)_{||\beta} - \frac{1}{2} n^{\pi} n^{\epsilon} h^{\nu\tau} h^{\sigma\mu} h^{\rho\alpha} R_{\tau\pi\mu\alpha} R_{\nu\epsilon\sigma\rho} \\ - R_{\rho\tau} n^{\tau} R_{\pi\mu} n^{\mu} h^{\pi\rho} + R_{\sigma\tau\omega\rho} \left(A^{\tau\omega} A^{\sigma\rho} - A^{\sigma}{}_{\alpha} A^{\alpha\rho} h^{\tau\omega} \right) - |A|^4.$$

2.20 is equation 1.21 of [SSY] except that we have written the terms involving the third derivatives of the metric $\tilde{g}$ in divergence form on $\tilde{\Sigma}$.

Now at points where $|A| \neq 0$ we have, for any $\epsilon > 0$,

$$2.21 \quad |\nabla_{\tilde{\Sigma}} |A||^2 \left(1 + \frac{1}{1+\epsilon} \right) \leq \frac{1}{2} \Delta_{\tilde{\Sigma}} |A|^2 - A^{\alpha\beta} \Delta_{\tilde{\Sigma}} A_{\alpha\beta} + \frac{4\beta_0}{\epsilon}.$$

This inequality is derived as follows.

Without loss of generality we may assume that at a particular point

$p \in \tilde{\Sigma}$, $g_{\alpha\beta} = \delta_{\alpha\beta}$, A_{ab} is diagonal for $a, b \in \{2, 3\}$; and locally about p in $\tilde{\Sigma}$, $g^{1\alpha} = g_{1\alpha} = \delta_{\alpha}^1$, $A_{1\alpha} = 0$ for $\alpha \in \{1, 2, 3\}$. Then, at p ,

$$\begin{aligned}
 4|A|^2 \left(\frac{1}{2} \Delta_{\tilde{\Sigma}} |A|^2 - A^{\alpha\beta} \Delta_{\tilde{\Sigma}} A_{\alpha\beta} \right) - |\nabla_{\tilde{\Sigma}} |A|^2|^2 \\
 = 4 \sum_{c=2}^3 (A_{cc})^2 \sum_{a,b,m=2}^3 (A_{ab||m})^2 - 4 \sum_{m=2}^3 \left(\sum_{a=2}^3 A_{aa} A_{aa||m} \right) \left(\sum_{b=2}^3 A_{bb} A_{bb||m} \right) \\
 = 8 \sum_{c=2}^3 (A_{cc})^2 \sum_{m=2}^3 (A_{32||m})^2 \\
 = 8 \sum_{c=2}^3 (A_{cc})^2 \sum_{a \neq b} (A_{ab||a})^2.
 \end{aligned}$$

Now by triangle inequality,

$$\begin{aligned}
 v \left(\sum_{a \neq b} (A_{ab||a})^2 \right) &\geq v \left(\sum_{a \neq b} (A_{aa||b})^2 \right) - v \left(\sum_{a \neq b} (A_{aa||b} - A_{ab||a})^2 \right) \\
 &= v \left(\sum_{a \neq b} (A_{aa||b})^2 \right) - v \left(\sum_{a,b=2}^3 (A_{aa||b} - A_{ab||a})^2 \right) \\
 &= v \left(\sum_{a \neq b} (A_{aa||b})^2 \right) - v \left(\sum_{a,b=2}^3 (R_{a1ba})^2 \right)
 \end{aligned}$$

where we have used 2.11. Thus

$$v \left(\sum_{a \neq b} (A_{ab||a})^2 \right) \geq v \left(\sum_{a \neq b} (A_{aa||b})^2 \right) - \sqrt{2\beta_0}.$$

Hence for any $\varepsilon > 0$ we have

$$\sum_{a \neq b} (A_{ab||a})^2 \geq \left(\sum_{a \neq b} (A_{aa||b})^2 \right) (1+\varepsilon)^{-1} - 2\beta_0 \varepsilon^{-1}.$$

In the above we have used the fact that for $A, B, C \geq 0$,

$\sqrt{A} \geq \sqrt{B} - \sqrt{C} \Rightarrow A \geq (1+\varepsilon)^{-1} B - C\varepsilon^{-1}$ for any $\varepsilon > 0$. Hence we get

$$\begin{aligned}
 2.22 \quad 4|A|^2 \left(\frac{1}{2} \Delta_{\tilde{\Sigma}} |A|^2 - A^{\alpha\beta} \Delta_{\tilde{\Sigma}} A_{\alpha\beta} \right) - |\nabla_{\tilde{\Sigma}} |A|^2|^2 \\
 \geq 8|A|^2 \left((1+\varepsilon)^{-1} \sum_{a \neq b} (A_{aa||b})^2 - 2\beta_0 \varepsilon^{-1} \right).
 \end{aligned}$$

Again

$$\begin{aligned}
|\nabla_{\tilde{\Sigma}}|A|^2|^2 &= 4 \sum_{m=2}^3 \left(\sum_{\alpha=2}^3 A_{\alpha\alpha} A_{\alpha\alpha} \parallel m \right)^2 \\
&\leq 4 \sum_{m=2}^3 \left(\sum_{\alpha=2}^3 (A_{\alpha\alpha})^2 \right) \left(\sum_{b=2}^3 (A_{bb} \parallel m)^2 \right) \\
&= 4|A|^2 \left(\sum_{b \neq m} (A_{bb} \parallel m)^2 + \sum_b (A_{bb} \parallel b)^2 \right).
\end{aligned}$$

But since $\tilde{\Sigma}$ is minimal we have, for $c \neq b$,

$$A_{cc} \parallel b + A_{bb} \parallel b = \sum_{\alpha=2}^3 A_{\alpha\alpha} \parallel b = 0.$$

Thus

$$\begin{aligned}
|\nabla_{\tilde{\Sigma}}|A|^2|^2 &\leq 4|A|^2 \left(\sum_{b \neq m} (A_{bb} \parallel m)^2 + \sum_{b \neq c} (A_{cc} \parallel b)^2 \right) \\
&= 8|A|^2 \sum_{b \neq m} (A_{bb} \parallel m)^2.
\end{aligned}$$

Putting the above inequality in 2.22 we get

$$4|A|^2 \left(\frac{1}{2} \Delta_{\tilde{\Sigma}} |A|^2 - A^{\alpha\beta} \Delta_{\tilde{\Sigma}} A_{\alpha\beta} \right) \geq (1 + (1+\epsilon)^{-1}) |\nabla_{\tilde{\Sigma}}|A|^2|^2 - 16|A|^2 \beta_0 \epsilon^{-1}.$$

Finally dividing by $4|A|^2 \neq 0$ we get 2.21. Hence using the estimate

$$\left| R_{\alpha\beta\gamma\delta} R^{\alpha\beta\gamma\delta} \right| < \beta_0 \quad \text{on } \tilde{\Sigma} \quad \text{and 2.20 we have at points where } |A| \neq 0,$$

$$\begin{aligned}
2.23 \quad |\nabla_{\tilde{\Sigma}}|A|^2|^2 \left(1 + \frac{1}{1+\epsilon} \right) &\leq \frac{1}{2} \Delta_{\tilde{\Sigma}} |A|^2 + |A|^4 + C_1 \sqrt{\beta_0} |A|^2 + C_2 \beta_0 + \frac{4}{\epsilon} \beta_0 \\
&\quad - \left(A^{\nu\rho} h^{\sigma\pi} R_{\nu\epsilon\sigma\rho} n^\epsilon \right) \parallel_\pi - \left(A^{\rho\beta} R_{\rho\tau} n^\tau \right) \parallel_\beta
\end{aligned}$$

where C_1 and C_2 are absolute positive constants.

This inequality continues to hold globally in the distribution sense even if $|A|$ vanishes at various points. Hence multiplying 2.23 with

$|A|^{2q}$ ($q > 0$), integrating over $\tilde{\Sigma}$ and integrating by parts the last two terms we get

$$\begin{aligned}
2.24 \quad & \int_{\tilde{\Sigma}} |A|^{2q} |\nabla_{\tilde{\Sigma}} |A||^2 \left(1 + \frac{1}{1+\varepsilon}\right) \\
& \leq \frac{1}{2} \int_{\tilde{\Sigma}} |A|^{2q} \Delta_{\tilde{\Sigma}} |A|^2 + \int_{\tilde{\Sigma}} |A|^{2q} \left(|A|^4 + c_1 \sqrt{\beta_0} |A|^2 + c_2 \beta_0 + \frac{4\beta_0}{\varepsilon} \right. \\
& \quad \left. + \int_{\tilde{\Sigma}} (|A|^{2q}) \|\pi^{A\nu\rho h\sigma\pi} R_{\nu\varepsilon\sigma\rho} n^\varepsilon + \int_{\tilde{\Sigma}} (|A|^{2q}) \|\beta^{A\rho\beta} R_{\rho\tau} n^\tau \right).
\end{aligned}$$

Hence using

$$\left| (|A|^{2q}) \|\pi^{A\nu\rho h\sigma\pi} R_{\nu\varepsilon\sigma\rho} n^\varepsilon \right| \leq c_3 |\nabla_{\tilde{\Sigma}} |A|^{2q}| |A| \sqrt{\beta_0}$$

and so on, we obtain

$$\begin{aligned}
2.25 \quad & \int_{\tilde{\Sigma}} |A|^{2q} |\nabla_{\tilde{\Sigma}} |A||^2 \left(1 + \frac{1}{1+\varepsilon}\right) \leq - \int_{\tilde{\Sigma}} 2q |A|^{2q} |\nabla_{\tilde{\Sigma}} |A||^2 \\
& + \int_{\tilde{\Sigma}} |A|^{2q} \left(|A|^4 + c_4 \sqrt{\beta_0} |A|^2 + c_5 \beta_0 + \frac{4\beta_0}{\varepsilon} \right) + \int_{\tilde{\Sigma}} c_6 \sqrt{\beta_0} 2q |A|^{2q} |\nabla_{\tilde{\Sigma}} |A||.
\end{aligned}$$

In the above and in what follows c_m , $m \in \mathbb{Z}$, are absolute positive constants.

Next we consider the stability inequality

$$2.26 \quad \int_{\tilde{\Sigma}} \left(|A|^{2+R_{\alpha\beta} n^\alpha n^\beta} \right) \xi^2 - \int_{\tilde{\Sigma}} |\nabla_{\tilde{\Sigma}} \xi|^2 \leq 0 \quad \text{for all } \xi \in C^{0,1}(\tilde{\Sigma}).$$

We put $\xi = |A|^{1+q}$; this implies $|\nabla_{\tilde{\Sigma}} \xi|^2 = (1+q)^2 |A|^{2q} |\nabla_{\tilde{\Sigma}} |A||^2$. Hence from 2.26 we get

$$2.27 \quad \int_{\tilde{\Sigma}} |A|^{2q+4} \leq \int_{\tilde{\Sigma}} (1+q)^2 |A|^{2q} |\nabla_{\tilde{\Sigma}} |A||^2 + \int_{\tilde{\Sigma}} c_7 \sqrt{\beta_0} |A|^{2+2q}.$$

We add 2.25 and 2.27 and use

$$2.28 \quad \int_{\tilde{\Sigma}} c_6 \sqrt{\beta_0} 2q |A|^{2q} |\nabla_{\tilde{\Sigma}} |A|| \leq \int_{\tilde{\Sigma}} \varepsilon q^2 |A|^{2q} |\nabla_{\tilde{\Sigma}} |A||^2 + \int_{\tilde{\Sigma}} c_6^2 \beta_0 |A|^{2q} \varepsilon^{-1}$$

to get

$$\begin{aligned}
2.29 \quad \int_{\tilde{\Sigma}} |A|^{2q} |\nabla_{\tilde{\Sigma}} |A||^2 (1+(1+\varepsilon)^{-1}) &\leq \int_{\tilde{\Sigma}} (1+q^2) |A|^{2q} |\nabla_{\tilde{\Sigma}} |A||^2 \\
&+ \varepsilon q^2 \int_{\tilde{\Sigma}} |A|^{2q} |\nabla_{\tilde{\Sigma}} |A||^2 + \int_{\tilde{\Sigma}} \frac{C_8 \beta_0}{\varepsilon} |A|^{2q} + \int_{\tilde{\Sigma}} C_7 \sqrt{\beta_0} |A|^{2+2q} \\
&+ \int_{\tilde{\Sigma}} |A|^{2q} \left(C_4 \sqrt{\beta_0} |A|^2 + C_5 \beta_0 + 4\beta_0/\varepsilon \right).
\end{aligned}$$

This implies, for some constant $C > 0$,

$$2.30 \quad ((1+\varepsilon)^{-1} - q^2(1+\varepsilon)) \int_{\tilde{\Sigma}} |A|^{2q} |\nabla_{\tilde{\Sigma}} |A||^2 \leq \int_{\tilde{\Sigma}} |A|^{2q} \left(C \sqrt{\beta_0} |A|^2 + C(1+\varepsilon)^{-1} \beta_0 \right).$$

From 2.27 and 2.30 we find L^5 bound for $|A|$. We put $q = \frac{1}{2}$. Then

$((1+\varepsilon)^{-1} - q^2(1+\varepsilon)) = \frac{1}{4}(1+\varepsilon)^{-1}(3-2\varepsilon-\varepsilon^2) > 0$ for any $\varepsilon < 1$. Hence 2.30 implies

$$2.31 \quad \int_{\tilde{\Sigma}} |A| |\nabla_{\tilde{\Sigma}} |A||^2 \leq C'_1 (|A|^3 + |A|)$$

for some positive number $C'_1 = C'_1(\beta_0, \varepsilon)$.

Putting $q = \frac{1}{2}$ in 2.27 yields

$$2.32 \quad \int_{\tilde{\Sigma}} |A|^5 \leq C'_2 \int_{\tilde{\Sigma}} (|A| |\nabla_{\tilde{\Sigma}} |A||^2 + |A|^3)$$

where $C'_2 = C'_2(\beta_0)$ is a constant.

2.31 and 2.32 give

$$2.33 \quad \int_{\tilde{\Sigma}} |A|^5 \leq C'_3 \int_{\tilde{\Sigma}} (|A|^3 + |A|)$$

with new constant $C'_3 = C'_3(\beta_0)$.

Now for each $\varepsilon_0 > 0$ Young's inequality gives

$$|A|^p \leq \varepsilon_0 |A|^5 + C'_4, \quad p = 1, 3,$$

for some positive constants $C'_4 = C'_4(\varepsilon_0, p)$. Thus 2.33 implies

$$2.34 \quad \int_{\tilde{\Sigma}} |A|^5 \leq \beta_1 |\tilde{\Sigma}|_{\tilde{g}}$$

for some constant $\beta_1 = \beta_1(\beta_0)$. $\square$

Compactness result

Let (N, g) be a Riemannian manifold having the following properties:

2.35 (i) (N, g) admits a sequence $\{q_i\}$ of points and a countable

atlas $\{U_i, \varphi_i\}$ such that, for some fixed real number

$R > 0$, $\varphi_i(U_i) \supset \supset B_{2R}(0) \subset \mathbb{R}^3$, $\varphi_i^{-1}(0) = q_i$ for all

i and the sets $\{\varphi_i^{-1}(B_{R/2}(0))\}$ also cover N ;

(ii) in the intersection $U_i \cap U_j$ the map $\varphi_i \circ \varphi_j^{-1}$ and their derivatives (say up to order 4) are uniformly bounded for all i and j ;

(iii) there exists a constant C_0 independent of i such that

$\sup_{\alpha, \beta \in \{1, 3\}} |g_{\alpha\beta}(x)|_{1,1;\varphi_i(U_i)} < C_0$ for all i , where $g_{\alpha\beta}$

are the components of g in the chart (U_i, φ_i) and

$|\cdot|_{1,1;\varphi_i(U_i)}$ is the usual $C^{1,1}$ norm on $\varphi_i(U_i)$; and

(iv) there exists a sequence of smooth metrics k_g on N such that $(N, {}^k_g)$ has uniformly bounded curvature independent of k and

$$2.36 \quad \lim_{k \rightarrow \infty} \left[\sup_i \sup_{\alpha, \beta \in \{1, 3\}} |g_{\alpha\beta}^{-k} g_{\alpha\beta}|_{1,\lambda;\varphi_i(U_i)} \right] = 0 \quad \text{for each } \lambda \in (0, 1)$$

where k_g are the components of k_g in the chart (U_i, φ_i) ,

and where $|\cdot|_{1,\lambda;\varphi_i(U_i)}$ is the $C^{1,\lambda}$ norm on $\varphi_i(U_i)$.

For example any smooth complete Riemannian manifold with injectivity

radius bounded from below and with bounded curvature satisfies the above properties (ref. Lemma 2.25 and 2.26 on page 48 in [A]).

We denote $\varphi_i^{-1}(B_{2R}(0))$ by B_i and assume that on the chart (B_i, φ_i) we have

$$2.37 \quad \begin{cases} |k_{g_{\alpha\beta}}|, |k_g^{\alpha\beta}|, |k_{\Gamma_{\beta\sigma}}^{\alpha}| \leq c_0, \\ \lambda_0 \sum_{\alpha=1}^3 x_{\alpha}^2 \leq k_g^{\alpha\beta} x_{\alpha} x_{\beta} \leq \Lambda_0 \sum_{\alpha=1}^3 x_{\alpha}^2 \text{ for all } (x_1, x_2, x_3) \in \mathbb{R}^3, \end{cases}$$

where c_0, λ_0 and Λ_0 are positive constants independent of k and i ,

and $k_{\Gamma_{\beta\sigma}}^{\alpha}$ are Christoffel symbols for k_g .

Let $\{\Sigma_k\}$ be a sequence of smooth, embedded compact surfaces in N such that each Σ_k is stable minimal with respect to the metric k_g and the area $|\Sigma_k|_{k_g}$ is uniformly bounded. It follows from Lemma 2.9 that

$$2.38 \quad \int_{\Sigma_k} |A(\Sigma_k, k_g)|^5 \leq c_1$$

where $|A(\Sigma_k, k_g)|$ is the length of the second fundamental form of Σ_k in (N, k_g) and c_1 is a constant independent of k .

Now we shall show that 2.38 implies that $|A(\Sigma_k, k_g)|$ is uniformly bounded. Let $A_E(\varphi_i(\Sigma_k \cap B_i))$ denote the second fundamental form of $\varphi_i(\Sigma_k \cap B_i)$ considered as a surface in $\mathbb{R}^3$ given by the Cartesian coordinates of the chart (B_i, φ_i) . We need the following lemma.

$$2.39. \text{ LEMMA. } |A(\Sigma_k \cap B_i, k_g)(p)|^2 \leq c_2 \left(|A_E(\varphi_i(\Sigma_k \cap B_i))(\varphi_i(p))|^2 + 1 \right),$$

$|A_E(\varphi_i(\Sigma_k \cap B_i))(\varphi_i(p))|^2 \leq c_3 \left(\left| A(\Sigma_k \cap B_i, {}^k_g)(p) \right|^2 + 1 \right)$ where $p \in B_i$ and c_2 and c_3 are positive constants independent of p, k and i .

Proof of 2.39. We suppose v_α is the unit normal to $\varphi_i(\Sigma_k \cap B_i)$ with respect to the Euclidean metric. Let $\bar{v}_\alpha$ be an extension of v_α

about $\varphi_i(\Sigma_k \cap B_i)$ such that $\sum_{\alpha=0}^2 v_\alpha^2 = 1$. We define

$$2.40 \quad \bar{n}_\alpha = \varphi \bar{v}_\alpha$$

where

$$2.41 \quad \varphi = \left(k_g^{\alpha\beta} \bar{v}_\alpha \bar{v}_\beta \right)^{-\frac{1}{2}}.$$

Then n_α , the restriction of $\bar{n}_\alpha$ on $\Sigma_k \cap B_i$ will be the unit normal form in the k_g metric. Let $h^{\alpha\beta} = k_g^{\alpha\beta} - n^\alpha n^\beta$ where $n^\alpha = k_g^{\alpha\beta} n_\beta$. Let k_∇ be the connection for the k_g metric. Then using

$$0 = \left(k_{\nabla n} \right)_\beta = \varphi^{-1} \frac{\partial \varphi}{\partial x^\beta} + \varphi^2 \frac{\partial \bar{v}_\tau}{\partial x^\beta} v_\alpha k_g^{\tau\alpha} - k_{\Gamma \rho\beta}^\tau n_\tau n^\rho$$

we get

$$2.42 \quad \left| A(\Sigma_k \cap B_i, {}^k_g) \right|^2 = \left(\varphi \frac{\partial \bar{v}_\varepsilon}{\partial x^\sigma} h^\varepsilon_\rho - k_{\Gamma \sigma\rho}^\tau n_\tau \right) \left(\varphi \frac{\partial \bar{v}_\mu}{\partial x^\beta} h^\mu_\alpha - k_{\Gamma \alpha\beta}^\tau n_\tau \right) h^{\alpha\rho} h^{\beta\sigma}.$$

Since Cauchy-Schwarz inequality holds for the inner product

$$2.43 \quad \langle P, Q \rangle := P_{\alpha\beta} Q_{\sigma\rho} h^{\alpha\rho} h^{\beta\sigma}$$

we can use $\langle P-Q, P-Q \rangle \leq 2\langle P, P \rangle + 2\langle Q, Q \rangle$ and

$$\langle P-Q, P-Q \rangle + \frac{1}{2}\langle P, P \rangle + 8\langle Q, Q \rangle \geq \langle P-Q, P-Q \rangle + 2\langle P, Q \rangle \geq \langle P, P \rangle + \langle Q, Q \rangle$$

to obtain

$$\begin{aligned}
2.45 \quad \left| A \left(\Sigma_k \cap B_i, k_g \right) \right|^2 &\leq 2 \left(\varphi^2 \frac{\partial \bar{v}_\varepsilon}{\partial x^\sigma} \frac{\partial \bar{v}_\mu}{\partial x^\beta} h^\varepsilon_\rho h^\mu_\alpha h^{\alpha\rho} h^{\beta\sigma} + k_{\Gamma_{\sigma\rho} n_\tau}^\tau k_{\Gamma_{\alpha\beta} n_\tau}^{\tau'} h^{\alpha\rho} h^{\beta\sigma} \right) \\
&= 2 \left(\varphi^2 \frac{\partial \bar{v}_\varepsilon}{\partial x^\sigma} \frac{\partial \bar{v}_\mu}{\partial x^\beta} h^{\varepsilon\mu} h^{\beta\sigma} + k_{\Gamma_{\sigma\rho} n_\tau}^\tau k_{\Gamma_{\alpha\beta} n_\tau}^{\tau'} h^{\alpha\rho} h^{\beta\sigma} \right)
\end{aligned}$$

and

$$2.46 \quad \left| A \left(\Sigma_k \cap B_i, k_g \right) \right|^2 + 7 \left(k_{\Gamma_{\sigma\rho} n_\tau}^\tau k_{\Gamma_{\alpha\beta} n_\tau}^{\tau'} h^{\alpha\rho} h^{\beta\sigma} \right) \geq \frac{1}{2} \varphi^2 \frac{\partial \bar{v}_\varepsilon}{\partial x^\sigma} \frac{\partial \bar{v}_\mu}{\partial x^\beta} h^{\varepsilon\mu} h^{\beta\sigma} .$$

Now we choose a new Euclidean co-ordinate system obtained by rotation from the original co-ordinate system such that at $\varphi_i(p)$, $v_\alpha = \delta_\alpha^1$. This implies, at $\varphi_i(p)$,

$$\begin{aligned}
&\varphi = (k_g^{11})^{-\frac{1}{2}}, \quad n^\alpha = k_g^{1\alpha} / (k_g^{11})^{\frac{1}{2}}, \\
2.47 \quad h^{\alpha\beta} &= k_g^{\alpha\beta} - k_g^{1\alpha} k_g^{1\beta} / k_g^{11} = \begin{cases} k_g^{ab} - k_g^{1a} k_g^{1b} / k_g^{11}, & a, b \in \{2, 3\}, \\ 0 & \text{otherwise.} \end{cases}
\end{aligned}$$

Also

$$2.48 \quad \left. \frac{\partial \bar{v}_1}{\partial x^\beta} \right|_{\varphi_i(p)} = 0 \quad \text{since} \quad \bar{v}_\alpha \frac{\partial \bar{v}_\alpha}{\partial x^\beta} = 0 .$$

Hence

$$2.49 \quad |A_E(\varphi_i(\Sigma_k \cap B_i))(\varphi_i(p))|^2 = \sum_{a,b=2}^3 \left(\frac{\partial \bar{v}_a}{\partial x^b} \right)^2 .$$

Since the orthonormal transformation with respect to the Euclidean metric does not change the bounds on $|k_g^{\alpha\beta}|$, $|k_{\Gamma_{\beta\rho}}^\alpha|$ and the eigenvalues of $k_g^{\alpha\beta}$ we have, at $\varphi_i(p)$,

$$2.50 \quad 0 < \lambda_0 \leq k_g^{\alpha\beta} v_\alpha v_\beta = \frac{1}{\varphi^2} \leq \Lambda_0$$

and from 2.47 we have

$$2.51 \quad 0 < c_4 \leq \det h^{ab} = \det k_g^{\alpha\beta} \leq c_5$$

where $\lambda_0, \Lambda_0, c_4$ and c_5 are positive constants independent of p, k, i .

2.51 and the bound on $|\text{Trace } h^{ab}|$ implies

$$2.52 \quad c_6 \sum_{a=2}^3 X_a^2 \leq h^{ab} X_a X_b \leq c_7 \sum_{a=2}^3 X_a^2 \quad \text{for all } (X_2, X_3) \in \mathbb{R}^2$$

where c_6 and c_7 are positive constants independent of k, p, i . Hence

using 2.47-2.52 and bounds on $|{}^k g_{\alpha\beta}|$, $|{}^k g^{\alpha\beta}|$ and $|{}^k \Gamma_{\rho\beta}^\alpha|$ in 2.45 and 2.46

we get

$$\left| A\left(\Sigma_k \cap B_i, {}^k g\right)(p) \right|^2 \leq c_8 |A_E(\varphi_i(\Sigma_k \cap B_i))(\varphi_i(p))|^2 + c_9$$

and

$$\left| A\left(\Sigma_k \cap B_i, {}^k g\right)(p) \right|^2 + c_{10} \geq c_{11} |A_E(\varphi_i(\Sigma_k \cap B_i))(\varphi_i(p))|^2$$

respectively, where c_8, c_9, c_{10} and c_{11} are positive constants

independent of p, k and i .

Hence the lemma follows. $\square$

From the uniform L^5 bound for the length of the second fundamental form of the sequence of surfaces $\{\Sigma_k\}$ follows the following lemma. We shall be in a particular chart (φ_i, B_i) . Let

$$x_0 \in B_R(\varphi_i(q_i)) \cap \varphi_i(\Sigma_k \cap B_i)$$

where we recall $B_R(\varphi_i(q_i)) = B_R(0) \subset B_{2R}(0) = \varphi_i(B_i)$. We use the notation

$[\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)]^*$ to denote the component of $\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)$,

$0 < \rho \leq R$ which contains x_0 . Let v_α be the components of the unit

normal form on $\varphi_i(\Sigma_k \cap B_i)$ with respect to the Euclidean metric on

$\varphi_i(B_i)$.

2.53. LEMMA. For all $\rho < R/4$,

$$\sup_{x \in [\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)]^*} |v_\alpha(x) - v_\alpha(x_0)| \leq c_{12}(\rho/R)^{3/5}$$

where c_{12} is a constant independent of k , x_0 and i .

Proof of 2.53. Let $x_1 \in \varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)$. In the following c_m , $m \in \mathbb{Z}$, will always denote a positive constant independent of k , i , x_0 and x_1 . By Lemma 2.39 and the fact that the area measures on $\Sigma_k \cap B_i$ in the k_g metric and in the Euclidean metric have bounded ratio independent of k and depending only on c_0 , λ_0 and Λ_0 occurring in 2.37 we have (for $\rho < R/4$)

$$\begin{aligned} 2.54 \quad & \int_{\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_1)} |A_E(\varphi_i(\Sigma_k \cap B_i))|^5 d\alpha \\ & \leq c_{13} \int_{\Sigma_k \cap \varphi_i^{-1}(B_\rho(x_1))} |A(\Sigma_k \cap B_i, k_g)|^5 + c_{14} |\Sigma_k \cap \varphi_i^{-1}(B_\rho(x_1))|_{k_g}. \end{aligned}$$

By virtue of 2.38 and the fact that the area $|\Sigma_k|_{k_g}$ are uniformly bounded, 2.54 yields

$$2.55 \quad \int_{\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_1)} |A_E(\varphi_i(\Sigma_k \cap B_i))|^5 d\alpha \leq c_{15}.$$

Then using the Hölder inequality we have

$$2.56 \quad \int_{\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_1)} |A_E(\varphi_i(\Sigma_k \cap B_i))| d\alpha \leq c_{16} |\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_1)|_E^{4/5}.$$

Now it is well-known (see Theorem 17.7 and Remark (3) of 17.9 in [LS1]) that

$$\begin{aligned} 2.57 \quad & \rho^{-2} |\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_1)|_E \\ & \leq c_{17} + c_{18} \int_{\varphi_i(\Sigma_k \cap B_i) \cap B_R(x_1)} |H_E(\varphi_i(\Sigma_k \cap B_i))|^4 \end{aligned}$$

where $H_E(\varphi_i(\Sigma_k \cap B_i))$ is the mean curvature of $\varphi_i(\Sigma_k \cap B_i)$ as a surface in

$\mathbb{R}^3$.

Hence 2.56 yields

$$2.58 \quad \int_{\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_1)} |A_E(\varphi_i(\Sigma_k \cap B_i))| d\alpha \leq c_{19} \rho^{8/5}$$

where we have used the fact $H^2 \leq 2|A|^2$ so that

$$\int |H_E(\varphi_i(\Sigma_k \cap B_i))|^4 \leq c_{20} \int |A_E(\varphi_i(\Sigma_k \cap B_i))|^4 \leq c_{21} \quad \text{by 2.55.}$$

Hence using

$$2.59 \quad |A_E(\varphi_i(\Sigma_k \cap B_i))|^2 = \sum_{\alpha, \beta} \left(\frac{\partial v_\alpha}{\partial x^\beta} \right)^2 - \sum_\alpha \left(\sum_\beta v_\beta \frac{\partial v_\alpha}{\partial x^\beta} \right) \equiv \sum_\alpha |\partial v_\alpha|^2$$

where ∂ denotes tangential gradient operator, we have

$$2.60 \quad \int_{\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_1)} |\partial v_\alpha|^2 \leq c_{22} \rho^{8/5}$$

for all $x_1 \in \varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)$ and for all $\rho \leq R/4$.

Hence the lemma follows from the following analogue of the Morrey estimate for two dimensional surfaces due to Simon [LS2].

(We simply quote the appropriate result without giving the proof which can be found in [LS2]. See also Lemma 16.4 in [GT].)

Let M be a 2-dimensional C^∞ submanifold of $\mathbb{R}^3$, suppose h is a C^1 function on M and suppose $K > 0$, $\beta \in (0, 1)$, are such that

$$\int_{G_\rho(x_1)} |\partial h| d\alpha \leq K \rho (\rho/R)^\beta$$

for all $x_1 \in G_{R/4}(x_0) \equiv \{x \in M \mid |x - x_0| < R/4\}$ and for all $\rho \in (0, R/4)$.

Here R is such that $(\bar{M} \sim M) \cap \{x \in \mathbb{R}^3 \mid |x - x_0| \leq R\} = \emptyset$ and $x_0 \in M$.

Then $\sup_{x \in G_\rho^*(x_0)} |h(x) - h(x_0)| \leq cK(\rho/R)^\beta$ where $G_\rho^*(x_0)$ denotes the

component of $G_\rho(x_0)$ which contains x_0 , c is a constant depending on

the constants Λ_3 and Λ_4 defined as follows:

$$2.61 \quad \int_{G_{R/2}(x_0)} H^2 d\alpha \leq \Lambda_4 ,$$

$$2.62 \quad (3R/4)^{-2} |G_{3R/4}(x_0)|_E \leq \Lambda_3 . \quad \square$$

2.63. Remark. Since v is unit normal $|v_\alpha(x) - v_\alpha(x_0)| \leq 2$. This implies

$$2.64 \quad \sup_{x \in [\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)]^*} |v_\alpha(x) - v_\alpha(x_0)| \leq c_{23}(\rho/R)^{3/5}$$

for all $\rho \in (0, R)$. Hence

$$2.65 \quad |v_\alpha(x_1) - v_\alpha(x_2)| \leq c_{24}(|x_1 - x_2|/R)^{3/5}$$

for all $x_1, x_2 \in [\varphi_i(\Sigma_k \cap B_i) \cap B_{R/4}(x_0)]^*$.

The $3/5$ Hölder estimate for the unit normal can be improved using the minimal surface equation.

Let P be the rotation taking the co-ordinates with respect to the $\{x^\alpha\}_{\alpha=1,2,3}$ co-ordinate system in $\varphi_i(B_i)$ to those with respect to the new co-ordinate system $\{y^\alpha\}_{\alpha=1,2,3}$ where y^1 is in the direction of $v(x_0)$ and the (y^2, y^3) plane is parallel to the tangent plane of $\varphi_i(\Sigma_k \cap B_i)$ at $x_0 = P^{-1}(y_0) \in B_R(0)$. Then there exists a $\rho^* > 0$ such that

$$\varphi_i(\Sigma_k \cap B_i) \cap B_{\rho^*}(x_0) = \text{graph } k_u = \{(y^1, y) : y^1 = k_u(y)\}$$

where $y = (y^2, y^3)$. At $y_0 = (y_0^2, y_0^3)$ we have

$$2.66 \quad \frac{\partial^k u}{\partial y^\alpha} \equiv D_\alpha(k_u) = 0, \quad \alpha = 2, 3.$$

We suppose ρ^* is the maximum number less than or equal to $R/4$ such that $|D(k_u)| \leq 1$ for each k . Then from 2.65 we get

$$2.67 \quad 1/\sqrt{2} \leq c_{25}(\rho^*/R)^{3/5}.$$

Thus ρ^* can be chosen independent of k , x and i such that

$$2.68 \quad |D({}^k u)| \leq 1.$$

The unit normal form n of $\Sigma_k \cap \varphi_i^{-1}(B_{\rho^*}(y_0))$ with respect to the metric k_g is given by

$$2.69 \quad n = \frac{\xi}{v}.$$

Here ξ is a one-form on $\Sigma_k \cap \varphi_i^{-1}(B_{\rho^*}(y_0))$ namely,

$$2.70 \quad \xi({}^k u(y), y) = (1, -D({}^k u)(y))$$

and v is a scalar,

$$2.71 \quad v({}^k u(y), y) = \left({}^k_g(\xi, \xi) \Big|_{{}^k u(y), y} \right)^{\frac{1}{2}} = \left({}^k_{\tilde{g}}{}^{\alpha\beta} \xi_\alpha \xi_\beta \right) \Big|_{{}^k u(y), y},$$

${}^k_{\tilde{g}}{}^{\alpha\beta}$ being the components of k_g in $\{y^\alpha\}$ co-ordinate system.

The function ${}^k_u : B_{\rho^*}^2(y_0) \rightarrow \mathbb{R}$ satisfies the following elliptic partial differential equation on the disc $B_{\rho^*}^2(y_0)$:

$$2.72 \quad {}^k_{\psi}{}^{ab} D_a D_b ({}^k u) = {}^k_f$$

where

$$2.73 \quad {}^k_{\psi}{}^{ab}(y) = {}^k_{\tilde{g}}{}^{ab} - n^\alpha n^\beta$$

and

$$2.74 \quad {}^k_f(y) = \left(-{}^k_{\tilde{\Gamma}}{}^1_{\alpha\beta} + {}^k_{\tilde{\Gamma}}{}^c_{\alpha\beta} D_c ({}^k u) \right) ({}^k_{\tilde{g}}{}^{\alpha\beta} - n^\alpha n^\beta).$$

In equations 2.73 and 2.74, ${}^k_{\tilde{g}}{}^{\alpha\beta}$ and ${}^k_{\tilde{\Gamma}}{}^\sigma_{\alpha\beta}$ are evaluated at $(u(y), y)$

and $n^\alpha = {}^k_{\tilde{g}}{}^{\alpha\beta} n_\beta$.

Now we have

$$2.75 \quad \lambda_0 (1 + |D(k_u)|^2) \leq v^2 \leq \Lambda_0 (1 + |D(k_u)|^2) ,$$

$$2.76 \quad 0 < \tau_0 \leq \det \left(\tilde{g}^{ab} - n^a n^b \right) = \frac{\det \tilde{g}^{\alpha\beta}}{v^2} \leq \tau_1$$

and

$$2.77 \quad |\text{Trace } k_\psi^{ab}| \leq \tau_2 \quad \text{in } B_{\rho^*}^2(y)$$

where τ_0, τ_1 and τ_2 are constants independent of k, x_0 and i .

But k_ψ^{ab} is nonnegative semidefinite. Hence there exists a constant τ_4 independent of k, x_0 , and i such that

$$2.78 \quad 0 < \tau_4 \sum_{a=2}^3 X_a^2 \leq k_\psi^{ab} X_a X_b \quad \text{in } B_{\rho^*}^2 \quad \text{for all } x \in \mathbb{R}^2 .$$

Now using the fact that, for each $\lambda \in (0, 1)$, $\tilde{g}^{\alpha\beta}$ is uniformly bounded in $C^{1,\lambda}(\overline{B}_{2R}(0))$ for all k and $|D(k_u)| \leq 1$ and using 2.65 with $\lambda' = 3/5$ we get

$$2.79 \quad |k_u|_{1,\lambda'; B_{\rho^*}^2(y_0)} < c_{25} ,$$

$$2.80 \quad |k_\psi|_{0,\lambda'; B_{\rho^*}^2(y_0)} < c_{26} ,$$

$$2.81 \quad |k_f|_{0,\lambda'; B_{\rho^*}^2(y_0)}^{(2)} < c_{27} ,$$

where c_{25}, c_{26} and c_{27} are constants independent of k, x and i , and

$$\begin{aligned} |f|_{0,\lambda; B_{\rho^*}^2}^{(2)} &\equiv \sup_{y \in B_{\rho^*}^2} \left[\text{dist}(y, \partial B_{\rho^*}^2) \right]^2 |f(y)| \\ &+ \sup_{y_1, y_2 \in B_{\rho^*}^2} \left\{ \frac{|f(y_1) - f(y_2)|}{|y_1 - y_2|^\lambda} \left[\min(\text{dist}(y_1, \partial B_{\rho^*}^2), \text{dist}(y_2, \partial B_{\rho^*}^2)) \right]^2 \right\}. \end{aligned}$$

Since k_u is smooth, by Theorem 6.2 of [GT] (and also using the relation 4.17" on page 61 in [GT]) we get

$$\begin{aligned}
2.82 \quad |^k u|_{2,\lambda'; B_{\rho^*/2}^2(y_0)} &\leq c_{28} \left(1 + |^k f|_{0,\lambda'; B_{\rho^*}^2(y_0)}^{(2)} \right)^{\min(1, (\rho^*/2)^{2+\lambda'})} \\
&\leq c_{29}
\end{aligned}$$

where $c_{28} = c_{28}(\lambda')$, $c_{29} = c_{29}(\lambda')$ are constants independent of k, x

and i . 2.82 yields local $C^{0,\lambda}$ estimate for k_{ψ}^{ab} and k_f^{ab} with any $\lambda \in (0, 1)$. Hence we ultimately get

$$2.83 \quad \text{for all } \lambda \in (0, 1), \quad |^k u|_{2,\lambda; B_{\rho^*/4}^2(y_0)} \leq c_{30},$$

$c_{30} = c_{30}(\lambda)$ being a constant independent of k, x , and i .

Thus we arrive at the following lemma:

2.84. LEMMA. Let $x_0 \in \varphi_j(\Sigma_k \cap B_i) \cap B_R(0)$. Then considering

$\varphi_i(\Sigma_k \cap B_i)$ locally as graph $^k \tilde{u}$ over the tangent plane at x_0 we have, for some positive ρ^* independent of k, x and i ,

$$2.85 \quad |^k u|_{2,\lambda; B_{\rho^*/4}^2(y_0)} \leq c_{31} \quad \text{for all } \lambda \in (0, 1),$$

where $c_{31} = c_{31}(\lambda)$ is a constant independent of k, x_0 and i .

Proof of 2.84. Immediately follows from 2.83. The constant c_{31} is different from c_{30} because of the change of the plane over which $\varphi_i(\Sigma_k \cap B_i)$ is locally considered as graph. $\square$

Now we prove a lower bound for the ratio

$$|[\varphi_i(\Sigma_k \cap B_i) \cap B_{r'}(x_0)]^*|_{E/r'}^2 \quad \text{for } x_0 \in \varphi_i(\Sigma_k \cap B_i) \cap B_R(0).$$

2.86. LEMMA. Let $x_0 \in B_R(0) \cap \varphi_i(\Sigma_k \cap B_i)$. There exists a positive number $\bar{\rho}$ independent of k, x_0 and i such that, for all $\rho < \bar{\rho}$,

$$2.87 \quad |[\varphi_i(\Sigma_k \cap B_i) \cap B_{\rho}(x_0)]^*|_E \geq \pi \rho^2 / 2.$$

Proof of 2.86. The lemma follows from the well known area estimate for the two-dimensional graphs in $\mathbb{R}^3$; for example the inequality 16.31 in [GT]. In our case, for $\rho < \rho^*/4$ we have

$$\pi\rho^2 \leq |[\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)]^*|_E + (\rho^2/2) \int_{\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)} |A_E(\varphi_i(\Sigma_k \cap B_i))|^2 da.$$

This implies, by the Hölder inequality,

$$\pi\rho^2 \leq |[\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)]^*|_E + (\rho^2/2) |[\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)]_E|^{3/5} \left(\int_{\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)} |A_E(\varphi_i(\Sigma_k \cap B_i))|^5 da \right)^{2/5}.$$

Hence using the inequalities 2.55 and 2.57 respectively we get

$$\pi\rho^2 \leq |[\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)]^*|_E + \rho^{2+6/5} c_{32}.$$

In the above and in what follows all the constants c_m , $m \in \mathbb{Z}$, are independent of k , x_0 and i . Thus we have

$$\rho^2 \left[\pi - (1+c_{32})\rho^{6/5} \right] \leq |[\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)]^*|_E.$$

Hence provided $(1+c_{32})\rho^{6/5} \leq \pi/2$ (that is $\rho \leq |\pi/2(1+c_{32})|^{5/6} \equiv c_{33} > 0$) we have

$$\pi\rho^2/2 \leq |[\varphi_i(\Sigma_k \cap B_i) \cap B_\rho(x_0)]^*|_E. \quad \square$$

For later use we define

$$2.89 \quad r = \frac{1}{4} \min \left(c_{33}, \frac{\rho^*}{4} \right).$$

Now we can prove the following compactness lemma. N , g and Σ_k are as in the paragraph preceding 2.38.

2.90. LEMMA. Suppose for all k the surfaces Σ_k are contained in a fixed compact subset of N . Then there exists a subsequence $\{\Sigma_{k_j}\}$ of

$\{\Sigma_k\}$ converging locally in the $C^{2,\alpha}$, $\alpha \in (0, 1)$, sense to a $C^{2,\alpha}$ compact, embedded minimal surface in (N, g) .

Proof of 2.90. Suppose all the surfaces Σ_k are contained in the connected open subset N of N such that $\bar{N}$ is compact. We cover $\bar{N}$ with a finite number of sets B_i , $i \in I$. Let r be as in 2.89. We cover each $B_{3R/5}(\varphi_i(q_i)) \subset \varphi_i(B_i)$, $i \in I$, with finite number J of Euclidean balls $B_{r/4}(x_j)$ such that $B_r(x_j) \subset B_R(\varphi_i(q_i))$ for all $j \in J$.

We first note that there exists a subsequence of $\{\Sigma_k\}$ (henceforth the subsequence is also denoted by $\{\Sigma_k\}$) such that each ball $B_{r/2}(x_j)$ either intersects an infinite number of members of $\{\Sigma_k\}$ or it does not intersect any member of $\{\Sigma_k\}$. This is because if any of the finite number of balls intersects only a finite number of surfaces then these finite number of surfaces can be deleted from the sequence.

Now let $F(i, j, k)$ be the collection of components of $\varphi_i(\Sigma_k \cap B_i) \cap B_r(x_j)$ such that their intersection with $B_{r/2}(x_j)$ is non-empty and let T be the number of elements in $F(i, j, k)$. For the l th element of $F(i, j, k)$ we have, from 2.87,

$$|[\varphi_i(\Sigma_k \cap B_i) \cap B_{r/2}(\hat{x})]^*|_E \leq \pi r^2/8$$

where $\hat{x} \in B_{r/2}(x_j)$ and $\hat{x}$ lies on the l th element of $F(i, j, k)$.

Summing over all the T components we have,

$$T \pi r^2/8 \leq c_{34} |\Sigma_k|_{k_g} \leq c_{35}.$$

Hence T is bounded independent of i, j, k . Let $\{S_k^1\}$ be the infinite sequence of surfaces $\{\varphi_i(\Sigma_k \cap B_i) \cap B_r(x_j)\}$ where only the first element of $F(i, j, k)$ has been considered. Then there exists a

subsequence $\{S_{k_1}^1\}$ of $\{S_k^1\}$ such that $\{S_{k_1}^1\}$ converges to a $C^{2,\alpha'}$, $\alpha' > 0$, surface (or to null set) in $B_{r/2}(x_j)$. This is because, for each k , $S_k^1 \cap B_r(x_j) \subset \text{graph } k_u^1$ where k_u^1 is a function on some two dimensional disc $B_r^2(x_j)$. Hence there exists a subsequence (also denoted by) $\{S_k^1\}$ such that, for each k , $S_k^1 \cap B_r(x_j) \subset \text{graph } k_{\tilde{u}}^1$ where $k_{\tilde{u}}^1$ are functions on the same disc ${}^{(1)}B_r^2(x_j)$. Now, because of Lemma 2.84, $\{k_{\tilde{u}}^1\}$ have uniformly bounded $C^{2,\alpha''}({}^{(1)}B_r^2(x_j))$ norm, $\alpha'' \in (0, 1)$. This implies there exists a subsequence $\{k_{\tilde{u}}^1\}$ such that $k_{\tilde{u}}^1 \rightarrow u^1$ in $C^{2,\alpha'}({}^{(1)}B_r^2(x_j))$, $0 < \alpha' < \alpha''$. Hence there exists a subsequence $\{S_{k_1}^1\}$ of $\{S_k^1\}$ such that $S_{k_1}^1 \cap B_r(x_j) \rightarrow \text{graph } u^1 \cap B_r(x_j)$. (In particular $\text{graph } u^1 \cap B_{r/2}(x_j)$ may be empty if the limiting surface, $\text{graph } u^1$, just touches $B_{r/2}(x_j)$. But if nonempty it is always connected.)

Now we consider the subsequence $\{\Sigma_{k_1}\}$ of $\{\Sigma_k\}$. We may suppose that infinite number of Σ_{k_1} have got components in $B_{r/2}(x_j)$ other than $\{S_{k_1}^1\}$. This is because otherwise finite number of surfaces can be removed to have a sequence with only one component in $B_{r/2}(x_j)$. Let $\{S_{k_1}^2\}$ be the infinite sequence of surfaces $\{\varphi_i(\Sigma_{k_1} \cap B_i) \cap B_r(x_j)\}$ where only the second element of $F(i, j, k_1)$ has been considered. As before we get a subsequence $\{S_k^2\}$ such that $\{S_k^2 \cap B_r(x_j)\}$ converges to a $C^{2,\alpha'}$ surface

(or null set) in $B_{r/2}(x_j) : S_k^2 \cap B_{r/2}(x_j) \rightarrow \text{graph } u^2 \cap B_{r/2}(x_j)$ where $u^2 \in C^{2,\alpha'}\left((2)_{B_r^2}(x_j)\right)$ for some disc $(2)_{B_r^2}(x_j)$.

Since there are only finite numbers of components to consider we find that there exists a subsequence $\{\Sigma_k\}$ such that on $B_{r/2}(x_j)$, $\varphi_i(\Sigma_k \cap B_i)$

converges to the finite union $\bigcup_l \text{graph } u^l \cap B_{r/2}(x_j)$ where

$u^l \in C^{2,\alpha'}\left((l)_{B_r^2}(x_j)\right)$ for some discs $(l)_{B_r^2}(x_j)$. We note that two pieces

$\text{graph } u^{l_1}$ and $\text{graph } u^{l_2}$ cannot cut each other in $B_{r/2}(x_j)$. This is

because otherwise Σ_k will be self-intersecting for sufficiently large k .

Also if they touch at a point $x \in B_{r/2}(x_j)$ then they will coincide in

$B_{r/2}(x_j)$. To prove this first we note (by taking the limit as $k \rightarrow \infty$ in

2.79-2.81) that $\text{graph } u^{l_1}$ and $\text{graph } u^{l_2}$ are minimal in the g metric.

Hence on some disc $B^2 \equiv \tilde{B}_{r/2}^2(x_j)$ they satisfy the partial differential

equation $D_a D_b (\tilde{u}^l) \psi^{ab}(\tilde{u}^l(y), y) = f(\tilde{u}^l(y), y)$ for $l = l_1, l_2$. The " $\sim$ "

sign has been used to express $\text{graph } u^l$ over $(l)_{B_{r/2}^2}(x_j)$ as $\text{graph } \tilde{u}^l$

over B^2 . Now for some positive constants c_m , $m \in \mathbb{Z}$, we have, for

$y \in B^2$,

$$|\psi^{ab}(\tilde{u}^{l_1}(y), y) - \psi^{ab}(\tilde{u}^{l_2}(y), y)| \leq c_{36} |\tilde{u}^{l_1}(y) - \tilde{u}^{l_2}(y)|$$

and

$$|f(\tilde{u}^{l_1}(y), y) - f(\tilde{u}^{l_2}(y), y)| \leq c_{37} |\tilde{u}^{l_1}(y) - \tilde{u}^{l_2}(y)|.$$

Hence

$$\left[D_a D_b (\tilde{u}^{l_1} - \tilde{u}^{l_2}) \right] \psi^{ab}(\tilde{u}^{l_1}(y), y) \leq c_{38} |\tilde{u}^{l_1}(y) - \tilde{u}^{l_2}(y)|.$$

Now we can assume $\tilde{u}^{l_1} \geq \tilde{u}^{l_2}$. Then $u = \tilde{u}^{l_1} - \tilde{u}^{l_2} \geq 0$ satisfies

$$(D_a D_b u) \psi^{ab}(\tilde{u}^{l_1}(y), y) - c_{39} u \leq 0.$$

But $u = 0$ is a nonpositive interior minimum. Hence, by strong maximum principle $u = 0$, that is, $\tilde{u}^{l_1} = \tilde{u}^{l_2}$. Thus graph $\tilde{u}^{l_1}$ and graph $\tilde{u}^{l_2}$ coincide on B^2 .

We now repeat the same argument with the subsequence $\{\Sigma_k\}$ so obtained in the second ball $B_r(x_{j'})$, $j' \neq j$, and so on. Since the balls $\{B_{r/4}(x_j)\}_{j \in J}$ cover $B_{3R/5}(\varphi_i(q_i))$ and since the sets $\{\varphi_i^{-1}(B_{R/2}(q_i))\}_{i \in I}$ cover N , the repetition of the above procedure in balls $B_r(x_j) \subset \varphi_i(B_i)$ for all $j \in J$, $i \in I$, gives the existence of a subsequence $\{\Sigma_k\}$ such that Σ_k converges to the finite union of $C^{2,\alpha'}$ graphs, namely,

$$2.91 \quad \Sigma = \bigcup_{i \in I} \bigcup_{j \in J} \bigcup_l \varphi_i^{-1} \left[\text{graph } u^l(j) \cap \bar{B}_{r/4}(x_j) \right]$$

where each $[\varphi_i(\Sigma_k \cap B_i) \cap B_r(x_j)]_l$ tends to $\text{graph } u^l(j)$ on $B_{r/2}(x_j)$ in $C^{2,\alpha'}$ sense, $[\varphi_i(\Sigma_k \cap B_i) \cap B_r(x_j)]_l$ being the l th element of $F(i, j, k')$ (that is, the l th component of $\varphi_i(\Sigma_k \cap B_i) \cap B_r(x_j)$ such that $\varphi_i(\Sigma_k \cap B_i) \cap B_{r/2}(x_j)$ is nonempty).

Σ is compact because it is a finite union of compact sets. Σ is

without boundary because in all $B_{r/4}(x_j)$, $B_{r/4}(x_j) \cap \phi_i(\Sigma_k, \cap B_i)$ converge to smooth pieces without boundary in $B_{r/4}(x_j)$ and these pieces coincide on $B_{r/2}(x_j) \cap B_{r/2}(x_{j'}) \cap \phi_i(\Sigma_k, \cap B_i) \neq \emptyset$ for all $j \neq j'$. As the balls $B_{r/4}(x_j)$ cover the whole of the region we find that Σ is without boundary. Σ is non self-intersecting since the pieces are either pairwise disjoint or they coincide. Σ is thus a $C^{2,\alpha}$ compact embedded surface without boundary and Σ is minimal in the g metric. This completes the proof of 2.90. $\square$

2.92. Remark. In case each Σ_k is a sphere using the Gauss-Bonnet theorem and the fact that Σ_k converges to the limiting surface in $C^{2,\alpha}$, $\alpha \in (0, 1)$, sense we get

$$\int_{\Sigma} nK = \lim_{k' \rightarrow \infty} \int_{\Sigma_{k'}} K_{k'} = 4\pi,$$

where K and $K_{k'}$ are respectively the Gauss curvatures of Σ and $\Sigma_{k'}$, in the metric induced from a smooth metric on N , and n is the number of sheets of $\Sigma_{k'}$ converging to Σ . Hence

$$\int_{\Sigma} K > 0$$

so that either Σ is a sphere ($n = 1$) or Σ is a projective space ($n = 2$). In particular when N is simply connected Σ is a sphere since embedded projective space cannot occur in N (by Theorem 4.7 on page 108 in [H]).

Alternative proof of Theorem 2.8. As before, from the asymptotic conditions (c) in the hypothesis of the main Theorem 1.1, it follows that there exists a smooth sphere S_{α_0} in $N \sim \overline{Q}$ given by $x = \alpha_0$ in the asymptotic coordinate system such that the mean curvature of S_{α_0} in

(N, g) with respect to the outward normal is strictly positive. Let N_1 and N_2 be bounded, connected open subsets of N containing S_{α_0} as well as its interior. We suppose $N_1 \subset\subset N_2$ and $\{(u_i, \varphi_i)\}_{i \in I}$ is a finite atlas for $\bar{N}_2$ satisfying the following properties:

- (i) in the intersection $u_i \cap u_j$ the maps $\varphi_i^{-1} \circ \varphi_j$ and their derivatives are uniformly bounded for all $i, j \in I$;
- (ii) there exists positive constants c_{49} , λ' and Λ' such that

$$2.93 \quad \sup_{i \in I; \alpha, \beta \in \{1, 2, 3\}} |g_{\alpha\beta}^i|_{1,1;\varphi_i(u_i)} < c_{49},$$

$g_{\alpha\beta}^i$ being the components of g in the chart (u_i, φ_i) , and

$$2.94 \quad \lambda' \sum_{\alpha=1}^3 (x^\alpha)^2 \leq g_{\alpha\beta}^i x^\alpha x^\beta \leq \Lambda' \sum_{\alpha=1}^3 (x^\alpha)^2 \quad \text{for all } (x^1, x^2, x^3) \in \mathbb{R}^3,$$

and for all $i \in I$.

Let $\{\psi_i\}_{i \in I}$ be a C^∞ partition of unity subordinate to the atlas $\{(u_i, \varphi_i)\}_{i \in I}$ such that $N_2 \subset \bigcup_{i \in I} \text{supp } \psi_i$. Let

$$2.95 \quad \rho(x) = \begin{cases} c \exp(1/(|x|^2 - 1)) & , \quad |x| \leq 1, \\ 0 & , \quad |x| \geq 1, \end{cases}$$

c being the constant such that $\int_{\mathbb{R}^3} \rho dx = 1$, be a mollifier on $\mathbb{R}^3$.

For any function f on $\varphi_j(u_j) \subset \mathbb{R}^3$ we define $f_{h(j)}$ by

$$2.96 \quad f_{h(j)}(\varphi_j(p)) = h^{-3} \int_{\varphi_j(u_j)} f(x) \rho((x - \varphi_j(p))/h) dx$$

where $p \in \text{supp } \psi_j$ and $h > 0$. Then we define

$$2.97 \quad f_h(\varphi_j(p)) = \sum_i \psi_i(p) \left(f \circ \varphi_j \circ \varphi_i^{-1} \right)_{h(i)}(\varphi_i(p)) .$$

For any tensor T on N_2 we define the tensor T_h as follows:

$$2.98 \quad T_h|_p = \sum_{i \in I} \psi_i(p) T_{h(i)}|_p , \quad p \in N_2 ,$$

where $T_{h(i)}$ are the tensors on $\text{supp } \psi_i$ such that in the chart (u_i, φ_i)

the components $(T_{h(i)})_{\alpha\beta\dots}^i$ and $T_{\alpha\beta\dots}^i$ of $T_{h(i)}$ and T respectively are related by

$$2.99 \quad (T_{h(i)})_{\alpha\beta\dots}^i(\varphi_i(p)) = \left(T_{\alpha\beta\dots}^i \right)_{h(i)}(\varphi_i(p)) .$$

We note that when T is a scalar on U_j , 2.98 is identical to 2.97

with $T_h = f_h \circ \varphi_j$.

Clearly for any sequence of $h \downarrow 0$, $\{g_h\}$ is a sequence of smooth Riemannian metrics converging to g in $C^{1,\lambda}$ norm on $\bar{N}_1$ for each $\lambda \in (0, 1)$. Furthermore

$$2.100 \quad (f_1 + f_2)_h = (f_1)_h + (f_2)_h$$

and if u is $C^{0,1}(\varphi_j(U_j))$ and f is bounded then

$$2.101 \quad u_h f_h - (uf)_h = O(h) .$$

Also we have

$$(T_h)_{\alpha\beta}^j - \left(T_{\alpha\beta}^j \right)_h = O(h) ,$$

provided $T_{\alpha\beta}^i$, $i \in I$, are uniformly bounded.

With these preliminaries we prove the following lemma.

2.102. LEMMA. Let $\hat{\nabla}$ and $\hat{\Delta}$ denote respectively the gradient operator and the Laplacian for the metric g_h . Then, on N_1 ,

$$2.103 \quad \text{Ric}(g_h) = V_h^{-1} \hat{\nabla} \hat{\nabla} V_h + (\Phi_1)_h g_h + F^h ,$$

$$2.104 \quad \hat{\Delta}V_h = (\Phi_2)_h V_h + G^h ,$$

where F^h are type $(0, 2)$ tensors and G^h are scalars such that

$$2.105 \quad \sup_{i \in I} \sup_{p \in U_i \cap N_1} \left| F^h_{\alpha\beta}(\varphi_i(p)) \right| \leq c_{50} h$$

and

$$2.106 \quad \sup_{i \in I} \sup_{p \in U_i \cap N_1} \left| G^h(\varphi_i(p)) \right| \leq c_{50} h ,$$

c_{50} being a constant independent of α, β and h .

Proof of 2.102. The proof follows by mollifying 1.2 and 1.3 and by straight forward calculation. In the following we only show that

$\text{Ric}(g_h)^i_{\alpha\beta} - \left(\text{Ric}(g)^i_{\alpha\beta} \right)_h$ is of $O(h)$, that is

$$2.107 \quad \sup_{i \in I} \sup_{p \in U_i \cap N_1} \left| \text{Ric}(g_h)^i_{\alpha\beta}(\varphi_i(p)) - \left(\text{Ric}(g)^i_{\alpha\beta} \right)_h(\varphi_i(p)) \right| < c_{51} h$$

and

$$2.108 \quad (\Phi_1)_h(g_h)^i_{\alpha\beta} - \left(\Phi_1 g^i_{\alpha\beta} \right)_h = O(h) .$$

Here and in what follows c_m , $m \in \mathbb{Z}$, denote positive constants independent of $i \in I$, $p \in U_i \cap N_1$ and $\alpha, \beta, \dots \in \{1, 2, 3\}$.

We suppose $\{x_{(j)}^\alpha\}$ denotes the local co-ordinate system in the chart (U_i, φ_i) . Then, by virtue of 2.97, 2.98 and 2.99, we have

$$2.109 \quad (g_h)^j_{\alpha\beta}(x_{(j)}) = \sum_i \psi_i \left(\varphi_j^{-1}(x_j) \right) \left(g^i_{\mu\nu} \right)_{h(i)}(x_{(i)}(x_{(j)})) \frac{\partial x_{(i)}^\mu}{\partial x_{(j)}^\alpha} \frac{\partial x_{(i)}^\nu}{\partial x_{(j)}^\beta}$$

which yields

$$2.110 \quad \left| \frac{\partial (g_h)^j_{\alpha\beta}}{\partial x^\tau}(x) - \frac{\partial g^j_{\alpha\beta}}{\partial x^\tau}(x) \right| < c_{52} h$$

and

$$2.111 \quad \left| \frac{\partial^2 (g_h)^j_{\alpha\beta}}{\partial x^\sigma \partial x^\tau} (x) - \left(\frac{\partial^2 g^j_{\alpha\beta}}{\partial x^\sigma \partial x^\tau} \right)_h (x) \right| < c_{53} h .$$

Now

$$\text{Ric}(g_h)^j_{\alpha\beta} = \frac{1}{2} (g_h)^{\rho\sigma}_j \left[- \frac{\partial^2 (g_h)^j_{\alpha\beta}}{\partial x^\rho \partial x^\sigma} + \frac{\partial^2 (g_h)^j_{\sigma\alpha}}{\partial x^\rho \partial x^\beta} - \frac{\partial^2 (g_h)^j_{\sigma\rho}}{\partial x^\alpha \partial x^\beta} + \frac{\partial^2 (g_h)^j_{\rho\beta}}{\partial x^\alpha \partial x^\sigma} \right] + R \left((g_h)^j_{\alpha\beta} \right)$$

where $R \left((g_h)^j_{\alpha\beta} \right)$ denotes the remaining terms involving only products and sums of $(g_h)^{\mu\nu}_j$, $(g_h)^j_{\mu\nu}$, $\partial (g_h)^{\mu\nu}_j / \partial x^\tau$ and $\partial (g_h)^j_{\mu\nu} / \partial x^\tau$. Hence, by 2.111,

$$\begin{aligned} \text{Ric}(g_h)^j_{\alpha\beta} &= \frac{1}{2} g^{\rho\sigma}_j \left[- \left(\frac{\partial^2 g^j_{\alpha\beta}}{\partial x^\rho \partial x^\sigma} \right)_h + \left(\frac{\partial^2 g^j_{\sigma\alpha}}{\partial x^\rho \partial x^\beta} \right)_h - \left(\frac{\partial^2 g^j_{\sigma\rho}}{\partial x^\alpha \partial x^\beta} \right)_h + \left(\frac{\partial^2 g^j_{\rho\beta}}{\partial x^\alpha \partial x^\sigma} \right)_h \right] + O(h) + R \left((g_h)^j_{\alpha\beta} \right) \\ &= \frac{1}{2} g^{\rho\sigma}_j \left[- \frac{\partial^2 g^j_{\alpha\beta}}{\partial x^\rho \partial x^\sigma} + \frac{\partial^2 g^j_{\sigma\alpha}}{\partial x^\rho \partial x^\beta} - \frac{\partial^2 g^j_{\sigma\rho}}{\partial x^\alpha \partial x^\beta} + \frac{\partial^2 g^j_{\rho\beta}}{\partial x^\alpha \partial x^\sigma} \right]_h + O(h) + R \left((g_h)^j_{\alpha\beta} \right) \end{aligned}$$

where in the last step we have used 2.101 and 2.100. Hence

$$\begin{aligned} \text{Ric}(g_h)^j_{\alpha\beta} &= \left[(\text{Ric}(g))^j_{\alpha\beta} \right]_h - \left[R(g^j_{\alpha\beta}) \right]_h + R \left((g_h)^j_{\alpha\beta} \right) + O(h) \\ &= \left[(\text{Ric}(g))^j_{\alpha\beta} \right]_h + O(h) . \end{aligned}$$

Thus 2.107 is established. To prove 2.108 we note that

$$\left(\Phi_1 g^j_{\alpha\beta} \right)_h = (\Phi_1)_h \left(g^j_{\alpha\beta} \right)_h + O(h) \quad \text{by 2.101.}$$

$$\text{Hence } \left(\Phi_1 g^j_{\alpha\beta} \right)_h = (\Phi_1)_h (g_h)^j_{\alpha\beta} + O(h) . \quad \square$$

Proof of 2.8. For sufficiently small h , S_{α_0} has strictly positive mean curvature with respect to the outward normal for all metrics g_h .

By hypothesis N is simply connected and hence N cannot contain embedded projective space. Now by the smooth metric version of Theorem 2.2 either $\overline{N}_1$ is diffeomorphic to a closed unit ball in $\mathbb{R}^3$ or there exists a stable minimal sphere h_S in the g_h metric embedded in N_1 . By the compactness lemma, 2.90, and Remark 2.92, there exists a subsequence of $\{g_h\}$ (which we also denote by $\{g_h\}$) such that h_S converges to a $C^{2,\alpha}$, $\alpha \in (0, 1)$, embedded minimal sphere S in (N, g) in locally $C^{2,\alpha}$ sense. Each h_S satisfies the following stability inequality

$$2.112 \quad \int_{{}^h_S} \left(\left(|A({}^h_S, g_h)|^2 + \text{Ric}(g_h)({}^h_n, {}^h_n) \right) \xi^2 - |\hat{\nabla}_{{}^h_S} \xi|^2 \right) \leq 0, \quad \xi \in C^1({}^h_S),$$

where h_n is the unit normal vector on h_S in the g_h metric and $\hat{\nabla}_{{}^h_S}$ is the gradient operator on h_S in the metric induced from g_h .

We put $\xi = V_h$ in 2.112 to obtain

$$2.113 \quad \int_{{}^h_S} \left(\left(|A({}^h_S, g_h)|^2 + \text{Ric}(g_h)({}^h_n, {}^h_n) \right) V_h^2 + V_h \hat{\Delta}_{{}^h_S} V_h \right) \leq 0$$

where $\hat{\Delta}_{{}^h_S}$ is the Laplacian on h_S with respect to metric induced g_h .

Now h_S being minimal on h_S we have (see Appendix B)

$$2.114 \quad \hat{\Delta} V_h = \hat{\Delta}_{{}^h_S} V_h + (\hat{\nabla} \hat{\nabla} V_h)({}^h_n, {}^h_n).$$

2.114 gives

$$\begin{aligned} 2.115 \quad & \text{Ric}(g_h)({}^h_n, {}^h_n) V_h^2 + V_h \hat{\Delta}_{{}^h_S} V_h \\ & = \text{Ric}(g_h)({}^h_n, {}^h_n) V_h^2 + V_h \hat{\Delta} V_h - V_h (\hat{\nabla} \hat{\nabla} V_h)({}^h_n, {}^h_n). \end{aligned}$$

Using 2.115 and Lemma 2.102 in 2.113 we get

$$2.116 \quad \int_{h_S} \left(\left| A \left({}^h S, g_h \right) \right|^2 V_h^2 + V_h^2 (\Phi_1 + \Phi_2)_h + V_h^2 F^h ({}^h n, {}^h n) + V_h G^h \right) \leq 0 .$$

Now we have ${}^h n \leq \left(\left(g_h ({}^h n, {}^h n) \right) / \lambda' \right)^{\frac{1}{2}}$ and V_h uniformly bounded for all h .

Thus using 2.105 and 2.106 in 2.116 we get

$$2.117 \quad \left| \int_{h_S} \left(\left| A \left({}^h S, g_h \right) \right|^2 V_h^2 + V_h^2 (\Phi_1 + \Phi_2)_h \right) \right| \leq c_{64} h |{}^h S|_{g_h}$$

where c_{64} is a constant independent of h .

Finally since each ${}^h S$ is obtained from a minimizing sequence from S_{α_0} we have

$$2.118 \quad |{}^h S|_{g_h} \leq |S_{\alpha_0}|_{g_h} \leq c_{65}$$

for some constant c_{65} independent of h .

2.117, 2.118 and the fact that $\Phi_1 + \Phi_2$ is nonnegative yield

$$2.119 \quad \begin{cases} \lim_{h \downarrow 0} \int_{h_S} \left| A \left({}^h S, g_h \right) \right|^2 V_h^2 = 0 , \\ \lim_{h \downarrow 0} \int_{h_S} (\Phi_1 + \Phi_2)_h V_h^2 = 0 . \end{cases}$$

Now V is strictly positive on N . Hence V_h is greater than some positive constant on $\bar{N}_1$ for all sufficiently small h . Hence 2.119 implies

$$2.120 \quad \begin{cases} \lim_{h \downarrow 0} \int_{h_S} \left| A \left({}^h S, g_h \right) \right|^2 = 0 , \\ \lim_{h \downarrow 0} \int_{h_S} (\Phi_1 + \Phi_2)_h = 0 . \end{cases}$$

Thus the $C^{2,\alpha}$ limiting surface S is totally geodesic in (N, g) and

and since $(\Phi_1 + \Phi_2)_h(x) > f(x) > 0$ for $x \in Q$ where $f(x)$ is independent of h , we have $S \cap Q = \emptyset$. $\square$

CHAPTER III

PROOF OF THE MAIN THEOREM

3.0.

In this chapter we shall prove that no embedded totally geodesic sphere exists in $N \sim Q$ and thereby we shall complete the proof of the main theorem.

3.1.

3.1 Remark. Since N is simply connected and S is compact co-dimension 1 embedded submanifold of N , S separates N and $N \sim S$ has exactly two closed components, say N_1 and N_2 having boundary S (see Lemma 4.4 and Theorem 4.6 on page 107 in [H]). It follows from the asymptotic condition (c) that exactly one of the components, say N_1 , is bounded. Thus $\overline{N}_1$ is a compact manifold with boundary S so that Stoke's formula holds. Similar is the case with the set obtained from N_2 by deleting the exterior of an asymptotic sphere S_R given by $|x| = R$ in the asymptotic coordinate system for sufficiently large R .

In the following lemma we deduce some formulae we shall need later.

3.2 LEMMA. Let S be a (C^2) totally geodesic embedded sphere in (N, g) such that $S \subset N \sim Q$. We suppose n is a continuous unit normal form on S . Then

3.3 (i) $g(n, \nabla V) = m'$, a constant on S ;

3.4 (ii) $\int_S \frac{|\nabla_S V|^2}{V^2} = 4\pi$, where ∇_S is the gradient operator on S

with respect to the metric induced from g ;

and provided $V < 1$,

(iii) for a sequence T_l of smooth spheres in $N \sim \bar{Q}$ converging to S in the C^2 sense

$$3.5 \quad \lim_{l \rightarrow \infty} \int_{T_l} \frac{cV^2+a}{V(1-V^2)^3} g(\tilde{n}, \nabla w) = \int_S 2m' \left\langle \nabla_S \left(\frac{cV^2+a}{V(1-V^2)^3} \right), \nabla_S V \right\rangle$$

where $w = |\nabla V|^2$, $\tilde{n} = \tilde{n}(l)$ is the smooth unit normal form on T_l consistent in direction with n , $\langle \cdot, \cdot \rangle$ denotes induced inner product on T_l and c, a are arbitrary constants to be specified later.

3.6 Remark. If $S \subset N \sim \bar{Q}$, we may replace 3.5 by the pointwise relation $g(n, \nabla w) = -m'RV$ on S , R being the scalar curvature of S . However S may touch ∂Q where the metric is not C^2 . The extra term $(cV^2+a)/(V(1-V^2)^3)$ has been introduced for later application. In 3.3 the sign of the constant m' depends on which normal direction is considered.

Before proceeding with the proof of Lemma 3.2, we want to make a remark about the differentiability of the metric g on $N \sim \bar{Q}$.

3.7 Remark. Müller zum Hagen [M] has shown that when the metric g is C^3 , there is an analytic atlas for $N \sim \bar{Q}$ in which the components of the metric and V are analytic functions of coordinates. The proof can be easily modified to the case when g is originally assumed to be locally $C^{1,1}$, although the isometry between the original atlas and the new atlas for N may be of lower regularity: that is, C^2 and locally $W^{3,p}$, $1 \leq p < \infty$. However we shall make no use of this fact in what follows.

Proof of 3.2. We first note that there exists a sequence T_l of smooth spheres in $N \sim \bar{Q}$ converging to S in the C^2 sense. This is because S separates N (see Remark 3.1) and by hypothesis Q is

connected.

Now in $N \sim \overline{Q}$, 1.2 and 1.3 become

$$3.8 \quad \text{Ric}(g)_{\alpha\beta} = V^{-1}V_{;\alpha\beta}$$

and

$$3.9 \quad \Delta V = 0$$

respectively.

Standard PDE regularity theorems (for example, Theorem 8.10 in [GT]) applied to 3.9 gives V to be locally $W^{3,p}$, $1 < p < \infty$, on $N \sim \overline{Q}$. In particular V is C^2 on $N \sim \overline{Q}$. From 3.8 it follows that $\text{Ric}(g)$ is locally $W^{1,p}$ on $N \sim \overline{Q}$; that is, in particular $\text{Ric}(g)$ is locally C^λ , $\lambda \in (0, 1)$ on $N \sim \overline{Q}$.

Now for the smooth compact embedded surface T_ℓ in $N \sim \overline{Q}$ we can use the following weak form of Codazzi's equation:

$$3.10 \quad \int_{T_\ell} b^\mu \text{Ric}(g)_{\alpha\beta} \left(\delta_\mu^\alpha - \tilde{n}^\alpha \tilde{n}_\mu \right) \tilde{n}^\beta = \int_{T_\ell} \left(b^\mu_{||\mu} H - b^\mu_{||\beta} A^\beta_{\mu} \right)$$

for any C^1 vector field b^μ on T_ℓ . Here $A_{\alpha\beta} = A_{\alpha\beta}(\ell)$ is the second fundamental form of T_ℓ and $H = H(\ell)$ is the mean curvature of T_ℓ . 3.10 can be obtained by approximating g with smooth metrics in $W^{2,p}$ norm in a neighbourhood of T_ℓ in N .

Now let f be any C^2 function on S and let $\bar{f}$ be a C^2 extension of f in a fixed neighbourhood of S such that $\bar{f}$ restricted to T_ℓ is a C^2 function on T_ℓ for all sufficiently large ℓ .

On any such T_ℓ in local co-ordinates we have

$$\begin{aligned}
3.11 \quad \bar{f}^{\|\sigma} \left(V; \alpha \tilde{n}^\alpha \right)_{\|\sigma} &= \bar{f}^{\|\sigma} \left(V; \alpha \tilde{n}^\alpha \right)_{;\beta} \left(\delta_\sigma^\beta - \tilde{n}^\beta \tilde{n}_\sigma \right) \\
&= \bar{f}^{\|\sigma} \left(V; \alpha \beta \left(\delta_\sigma^\beta - \tilde{n}^\beta \tilde{n}_\sigma \right) \tilde{n}^\alpha + V; \alpha \tilde{n}^\alpha; \beta \left(\delta_\sigma^\beta - \tilde{n}^\beta \tilde{n}_\sigma \right) \right)
\end{aligned}$$

where $\tilde{n}$ is any local extension of $\tilde{n}$ in N .

On using 3.8 and $\left(\tilde{n}_\alpha \tilde{n}^\alpha \right)_{;\beta} \left(\delta_\sigma^\beta - \tilde{n}^\beta \tilde{n}_\sigma \right) = 0$, 3.11 yields

$$3.12 \quad \bar{f}^{\|\sigma} \left(V; \alpha \tilde{n}^\alpha \right)_{\|\sigma} = \bar{f}^{\|\sigma} \left(V \operatorname{Ric}(g)_{\alpha\beta} \left(\delta_\sigma^\beta - \tilde{n}^\beta \tilde{n}_\sigma \right) \tilde{n}^\alpha + V^{\|\mu} A_{\mu\sigma} \right).$$

Hence integrating and using 3.10 with $b^\sigma = V \bar{f}^{\|\sigma}$ we have

$$3.13 \quad \int_{T_l} \bar{f}^{\|\sigma} \left(V; \alpha \tilde{n}^\alpha \right)_{\|\sigma} = \int_{T_l} \left((V \bar{f}^{\|\sigma})_{\|\sigma} H - (V \bar{f}^{\|\sigma})_{\|\alpha} A^\alpha_{\sigma} + A_{\mu\sigma} V^{\|\mu} \bar{f}^{\|\sigma} \right)$$

which on simplifying the right hand side and integrating the left hand side by parts yields

$$3.14 \quad - \int_{T_l} g(\tilde{n}, \nabla V) \Delta_{T_l} \bar{f} = \int_{T_l} (V H \Delta_{T_l} \bar{f} + H \langle \nabla_{T_l} V, \nabla_{T_l} \bar{f} \rangle - V \langle A, \nabla_{T_l} \nabla_{T_l} \bar{f} \rangle).$$

Now $\nabla_{T_l} \nabla_{T_l} \bar{f}$, $\Delta_{T_l} \bar{f}$ and $\langle \nabla_{T_l} V, \nabla_{T_l} \bar{f} \rangle$ are uniformly bounded

independent of l . Hence letting $l \rightarrow \infty$ and recalling that S is totally geodesic we get

$$3.15 \quad \int_S g(n, \nabla V) \Delta_S f = 0.$$

Since this holds for arbitrary C^2 function f on S and since $g(n, \nabla V)$ is continuous we have $g(n, \nabla V) = \text{constant}$ on S . Thus (i) is proved.

To prove (ii) first we note that the scalar curvature of g in $N \sim \bar{Q}$ vanishes. Now using the Gauss-Bonnet theorem and the contracted Gauss equation for a sequence of smooth metrics approaching g in $W^{2,p}$ norm in a neighbourhood of T_l in N , we get

$$3.16 \quad \int_{T_l} \left(-\operatorname{Ric}(g)_{\alpha\beta} \tilde{n}^\alpha \tilde{n}^\beta + H^2/2 - |A|^2/2 \right) = 4\pi.$$

This implies, by 3.8,

$$3.17 \quad \int_{T_l} \left(-V^{-1} V_{;\alpha\beta} \tilde{n}^\alpha \tilde{n}^\beta + H^2/2 - |A|^2/2 \right) = 4\pi .$$

Now using 2.6 with $\psi = V$ and 3.9 in 3.17 we get

$$3.18 \quad \int_{T_l} \left(\frac{|\nabla_{T_l} V|^2}{V^2} + g(\tilde{n}, \nabla V)H/V + H^2/2 - |A|^2/2 \right) = 4\pi .$$

Hence letting l tend to ∞ we get (ii).

Finally to prove (iii) we note that

$$\begin{aligned} 3.19 \quad g(\tilde{n}, \nabla w)/V &= 2V^{-1} V_{;\alpha\beta} V^{;\beta} \tilde{n}^\alpha \\ &= 2V^{-1} V_{;\alpha\rho} \left(\delta_\beta^\rho - \tilde{n}^\rho \tilde{n}_\beta \right) V^{;\beta} \tilde{n}^\alpha + 2V^{-1} \left(V^{;\beta} \tilde{n}_\beta \right) V_{;\alpha\rho} \tilde{n}^\alpha \tilde{n}^\rho \\ &= 2 \operatorname{Ric}(g)_{\alpha\rho} \left(\delta_\beta^\rho - \tilde{n}^\rho \tilde{n}_\beta \right) V^{;\beta} \tilde{n}^\alpha + 2V^{-1} g(\tilde{n}, \nabla V) \left(-\Delta_{T_l} V - g(\tilde{n}, \nabla V)H \right) . \end{aligned}$$

In the last step above we have used 3.8, 2.6 and 3.9 in addition to replacing $V^{;\beta}$ in the first term by $V^{\parallel\beta}$.

Hence using 3.10 with $b^\beta = \frac{cV^2+a}{(1-V^2)^3} V^{\parallel\beta}$ we get

$$\begin{aligned} 3.20 \quad \int_{T_l} \frac{cV^2+a}{V(1-V^2)^3} g(\tilde{n}, \nabla w) &= -2 \int_{T_l} \left(\frac{cV^2+a}{(1-V^2)^3} V^{\parallel\beta} \right)_{\parallel\alpha} A^\alpha_\beta + 2 \int_{T_l} \left(\frac{cV^2+a}{(1-V^2)^3} V^{\parallel\beta} \right)_{\parallel\beta} H \\ &\quad + 2 \int_{T_l} \frac{cV^2+a}{V(1-V^2)^3} g(\tilde{n}, \nabla V) \left(-\Delta_{T_l} V - g(\tilde{n}, \nabla V)H \right) . \end{aligned}$$

Now we have

$$\begin{aligned}
3.21 \quad 2 \int_{T_l} \frac{cV^2+a}{V(1-V^2)^3} g(\tilde{n}, \nabla V) (-\Delta_{T_l} V) &= 2 \int_{T_l} g(\tilde{n}, \nabla V) \left\langle \nabla_{T_l} \left(\frac{cV^2+a}{V(1-V^2)^3} \right), \nabla_{T_l} V \right\rangle \\
&+ 2 \int_{T_l} \frac{cV^2+a}{V(1-V^2)^3} \langle \nabla_{T_l} V, \nabla_{T_l} g(\tilde{n}, \nabla V) \rangle
\end{aligned}$$

where the second term in the right hand side equals $-2 \int g(\tilde{n}, \nabla V) \Delta_{T_l} f$ for

$$f = \int \left(\frac{cV^2+a}{V(1-V^2)^3} \right) dV \quad \text{and can be evaluated using 3.14. Hence letting } l \text{ tend}$$

to ∞ we get, from 3.20,

$$\lim_{l \rightarrow \infty} \int_{T_l} \frac{cV^2+a}{V(1-V^2)^3} g(\tilde{n}, \nabla w) = 2m' \int_S \left\langle \nabla_S \left(\frac{cV^2+a}{V(1-V^2)^3} \right), \nabla_S V \right\rangle.$$

This completes the proof of the lemma. $\square$

3.22 THEOREM. *There cannot be any totally geodesic embedded C^2 sphere S in (N, g) such that $S \subset N \sim Q$.*

Proof of 3.22. Let us suppose to the contrary that there is a S . Since by hypothesis Q is connected we have by Remark 3.1 either of the following cases:

Case I: $Q \subset \bar{N}_2$ or

Case II: $Q \subset \bar{N}_1$.

We consider Case I first. On N_1 we have $\Delta V = 0$ and on $\partial N_1 = S$ we have $g(n, \nabla V) = \text{constant}$ by 3.3. This implies $g(n, \nabla V) = 0$ on S .

Hence

$$\int_{N_1} |\nabla V|^2 = - \int_{N_1} V \Delta V + (1/2) \int_{N_1} \Delta V^2 = \frac{1}{2} \int_S V g(n, \nabla V) = 0$$

giving $|\nabla V|^2 = 0$ on N_1 . Thus V is constant in the interior of N_1

and hence, by continuity, on S . This implies

$$\int_S |\nabla_S V|^2 / V^2 = 0$$

which contradicts 3.4. Thus $Q \notin \overline{N}_2$.

Now we show that Case II also does not occur and hence get a contradiction. We shall use Robinson's divergence form inequality ([R]) on $N \sim \overline{Q}$. First we show that in the asymptotic co-ordinate system we have

$$V = C_1 - C_2/|x| + \eta$$

where $\eta = O(|x|^{-1-\beta})$, $\partial\eta/\partial x^\alpha = O(|x|^{-2-\beta})$ and

$$\left(R^{-3} \int_{N_{3R} \sim N_R} \left| \frac{\partial^2 \eta}{\partial x^\alpha \partial x^\tau} \right|^2 \right)^{1/2} = O(R^{-3-\beta}) \quad \text{for some } \beta \in (0, 1),$$

constants $C_1 > 0$ and $C_2 \geq 0$.

To see this first we note that $\overline{Q}$ is compact because $Q \subset \overline{N}_1$. Hence

$$\Delta V = 0 \quad \text{on } N \sim N_{R_0} \quad \text{for some } R_0 \text{ sufficiently large.}$$

In the asymptotic co-ordinate system V therefore satisfies the following divergence form PDE:

$$3.23 \quad \frac{\partial}{\partial x^\alpha} \left(g_{\alpha\beta} \sqrt{|g|} \frac{\partial V}{\partial x^\beta} \right) = 0$$

where $|g| = \det(g_{\alpha\beta})$. Hence by the Harnack inequality (Theorem 8.20 in [GT]) we have, for all $R > 2R_0$,

$$3.24 \quad \sup_{S_R} V \leq C_3 \inf_{S_R} V$$

where the constant C_3 is independent of R (this can be seen for example by covering S_R by fixed numbers of balls of radius $R/8$).

We note for later reference that 3.23 can be written

$$3.23' \quad \Delta_{\mathbb{R}^3} V = (\delta^{\alpha\beta} - g^{\alpha\beta}) \frac{\partial^2 V}{\partial x^\alpha \partial x^\beta} - \frac{\partial}{\partial x^\alpha} (g^{\alpha\beta} \sqrt{|g|}) \frac{\partial V}{\partial x^\beta} / \sqrt{|g|}$$

where $\Delta_{\mathbb{R}^3}$ denotes the usual Laplacian on $\mathbb{R}^3$.

Now let

$$3.25 \quad \psi(r) = C_4 \left(R_1^{-\varepsilon} - r^{-\varepsilon} \right) \quad \text{and} \quad \xi = \psi - V$$

where $r \geq R_1 > R_0$, $1 > \varepsilon > 0$ and C_4 is a constant to be specified later. Then

$$3.26 \quad \Delta \xi = \Delta \psi = \varepsilon(1-\varepsilon)C_4 r^{-2-\varepsilon} + \varepsilon C_4 \times O(r^{-2-\varepsilon-\alpha})$$

where we have used the assumption 1.4 concerning asymptotic behaviour of $g_{\alpha\beta}$. Thus $\Delta \xi > 0$ for sufficiently large R_1 . On S_{R_1} , $\xi = -V$.

Hence we can choose C_4 such that $\xi = 0$ at some point in $N \sim \bar{N}_{R_1}$. But

by maximum principle (Theorem 8.19 in [GT]) ξ cannot have non-negative maximum in the interior of any compact set in $N \sim \bar{N}_{R_1}$. Hence

$\limsup_{R \rightarrow \infty} \xi > 0$. So on every S_R , R sufficiently large, there exists a

point p such that $\psi(p) \geq V(p)$. Hence by 3.24 and 3.25 we get that V is bounded on N .

Now applying Theorems 8.8 and 8.32 in [GT] to 3.23 (after scaling) we have the following $W^{2,2}$ estimate

$$3.27 \quad \sup_{N_{3R} \sim N_{2R}} V + R \sup_{N_{3R} \sim N_{2R}} |DV| + \left\{ R^{-3} \int_{N_{3R} \sim N_{2R}} R^4 |D^2 V|^2 \right\}^{1/2} \\ \leq C \left\{ R^{-3} \int_{N_{4R} \sim N_R} V^2 \right\}^{1/2}$$

for all $R \geq 2R_0$. Here D denotes the usual gradient operator in $\mathbb{R}^3$.

Hence from 3.23' and 1.4 we have

$$3.28 \quad \Delta_{\mathbb{R}^3} V = f$$

where

$$\left(\frac{1}{R^3} \int_{N_{3R} \sim N_{2R}} f^2 \right)^{1/2} \leq CR^{-2-\alpha} \quad \text{for all } R \geq R_0 .$$

Taking smaller α if necessary, we may assume $\alpha \neq k^{-1}$ for all $k \in \mathbb{Z}$.

There exists a solution ζ of $\Delta_{\mathbb{R}^3} \zeta = f$ (ref. page 6 in [CHS]) with

$$3.29 \quad \left(\frac{1}{R^3} \int_{N_{3R} \sim N_{2R}} \zeta^2 \right)^{1/2} \leq CR^{-\alpha} .$$

Then we have

$$3.30 \quad \Delta_{\mathbb{R}^3} (V - \zeta) = 0 .$$

Hence $V - \zeta$ has the eigenfunction expansion

$$3.31 \quad (V - \zeta)(r\omega) = C_1 - \frac{C_2}{r} + \sum_{j=1}^{\infty} \left(a_j r^{\mu_j} - b_j r^{-\gamma_j} \right) \phi_j(\omega)$$

in terms of the eigenfunctions ϕ_j of the Laplacian on S^2 . Here

$C_1, C_2, a_j, b_j, \mu_j, \gamma_j$ are constants and $\mu_{j+1} \geq \mu_j > 0$, $\gamma_{j+1} \geq \gamma_j > 1$ (see [CHS]). Since V is bounded $a_j = 0$ for all j . Thus

$$3.32 \quad (V - \zeta)(r\omega) = C_1 - \frac{C_2}{r} + \sum_{j=1}^{\infty} b_j r^{-\gamma_j} \phi_j(\omega) .$$

This implies, by 3.29,

$$3.33 \quad \left(\frac{1}{R^3} \int_{N_{3R} \sim N_{2R}} (V - C_1)^2 \right)^{1/2} = O(R^{-\alpha})$$

for all sufficiently large R .

Hence using 3.27 with $V - C_1$ in place of V , and the arbitrariness of R above we have, for sufficiently large R ,

$$3.34 \quad \sup_{N_{2R} \sim N_R} |(V - C_1)| = O(R^{-\alpha}) .$$

That is, $V = C_1 + O(r^{-\alpha})$.

By the maximum principle $C_1 > 0$. Without loss of generality we assume $C_1 = 1$.

Now we write 3.27 with V replaced by $V - 1$:

$$3.35 \quad \sup_{N_{3R} \sim N_{2R}} (V-1) + R \sup_{N_{3R} \sim N_{2R}} |DV| + \left(R^3 \int_{N_{3R} \sim N_{2R}} R^4 |D^2 V|^2 \right)^{1/2} \\ \leq \left(R^{-3} \int_{N_{4R} \sim N_R} (V-1)^2 \right)^{1/2}.$$

Using 3.34 this time we get

$$3.36 \quad \left(\frac{1}{R^3} \int_{N_{3R} \sim N_{2R}} f^2 \right)^{1/2} \leq CR^{-2-2\alpha}.$$

Hence 3.29 is improved to

$$\left(\frac{1}{R^3} \int_{N_{3R} \sim N_R} \zeta^2 \right)^{1/2} \leq CR^{-2\alpha}.$$

Now we select $k_0 \in \mathbb{Z}$ such that $k_0 \alpha < 1$, $(k_0+1)\alpha > 1$. Then repeating the procedure (k_0+1) -times we have

$$V = 1 - \frac{C_2}{r} + O(r^{-1-\beta}) \quad \text{where } \beta > 0.$$

Finally writing $\eta = V - 1 + C_2/r$ we have

$$\Delta_{\mathbb{R}^3} \eta = f.$$

Hence applying the estimate 3.27 with V replaced by η we get

$$R \sup_{N_{3R} \sim N_R} |D\eta| + \left(R^{-3} \int_{N_{3R} \sim N_R} R^4 |D^2 \eta|^2 \right)^{1/2} \leq CR^{-1-\beta}.$$

Now by maximum principle $C_2 \geq 0$ (see Remark 1.6 for discussion on the case when $C_2 = 0$). Hence,

$$3.37 \quad V = 1 - m/|x| + \eta$$

where m is a nonnegative constant, $\eta = O(|x|^{-1-\beta})$, $|D\eta| = O(|x|^{-2-\beta})$ and

$$\left(R^{-3} \int_{N_{3R} \sim N_R} |D^2 \eta|^2 \right)^{1/2} = O(R^{-3-\beta}).$$

Clearly in case $m \neq 0$, $V < 1$ on N .

Now on each bounded open subset N_4 of $N \sim \overline{Q}$ we have the following inequality (provided $V < 1$ on $N \sim \overline{Q}$ ([R]))

$$3.38 \quad (FV^{-1}w^{;\alpha} + GV^{;\alpha})_{;\alpha} \geq 0$$

where

$$3.39 \quad F = (cV^2 + \alpha) / (1 - V^2)^3$$

and

$$3.40 \quad G = -2c / (1 - V^2)^3 + 6(cV^2 + \alpha) / (1 - V^2)^4,$$

c and α being constants such that $F > 0$ on N_2 .

For the sake of completeness the derivation of 3.38 is included in Appendix C. Here we only show that w is locally C^2 on $N \sim \overline{Q}$ so that 3.38 makes sense pointwise. Differentiating 3.9 we get $V^{;\alpha}_{;\alpha\beta} = 0$. This

implies by commutation law, $V^{;\alpha}_{;\alpha\beta} = -\text{Ric}(g)_{\tau\beta} V^{;\tau}$. Hence,

$$\begin{aligned} 3.41 \quad \Delta w &= 2 \left(V^{;\alpha}_{;\alpha\beta} V^{;\beta} \right)_{;\alpha} = 2 V^{;\alpha}_{;\alpha\beta} V^{;\beta} + 2 V^{;\alpha}_{;\alpha\beta} V^{;\beta\alpha} \\ &= -2 \text{Ric}(g)_{\tau\beta} V^{;\tau} V^{;\beta} + 2 V^{;\alpha}_{;\alpha\beta} V^{;\beta\alpha}. \end{aligned}$$

Since $\text{Ric}(g)$ is locally C^α , $\alpha \in (0, 1)$, we get that w is locally $C^{2,\alpha}$.

Now Stoke's formula applied to 3.38 gives

$$3.42 \quad \lim_{L \rightarrow \infty} \int_{T_L} (FV^{-1}g(\tilde{n}, \nabla w) + Gwg(\tilde{n}, \nabla V)) \\ + \lim_{R \rightarrow \infty} \int_{S_R} (FV^{-1}g({}^R n, \nabla w) + Gwg({}^R n, \nabla V)) \geq 0$$

where T_L and $\tilde{n}$ are as in Lemma 3.2. S_R is the asymptotic sphere

$|x| = R$ and ${}^R n$ is the outward normal to S_R . $\tilde{n}$ is directed outward

with respect to the volume bounded by T_L and S_R . Hence by 3.9 and 3.37

we have

$$\lim_{L \rightarrow \infty} \int_{T_L} g(\tilde{n}, \nabla V) = - \lim_{R \rightarrow \infty} \int_{S_R} g({}^R n, \nabla V) = -4\pi m.$$

Thus by 3.3 we have

$$3.43 \quad m' = g(n, \nabla V)|_S = -4\pi m/|S|,$$

n being the unit normal form on S consistent in direction with $\tilde{n}$. We need to consider only the case $m > 0$ (for $m = 0$, arguments of Case I apply in the domain exterior to S). Hence we assume $V < 1$.

The integral at infinity in 3.42 is evaluated in Appendix D. We have

$$3.44 \quad \lim_{R \rightarrow \infty} \int_{S_R} FV^{-1}g({}^R n, \nabla w) + Gwg({}^R n, \nabla V) = -\pi(c+a)/2m.$$

Using 3.39, 3.40, 3.43 and 3.44 in 3.42 we get

$$3.45 \quad \lim_{L \rightarrow \infty} \int_{T_L} \frac{cV^2+a}{V(1-V^2)^3} g(\tilde{n}, \nabla w) + \int_S \frac{-2cw}{(1-V^2)^3} m' + \int_S \frac{6(cV^2+a)}{(1-V^2)^4} wm' \\ \geq \pi(c+a)/2m.$$

Hence by 3.5 we get

$$3.46 \quad m' \int_S \left\langle 2\nabla_S \left(\frac{cV^2+a}{V(1-V^2)^3} \right), \nabla_S V \right\rangle - m' \int_S \frac{2cw}{(1-V^2)^3} + m' \int_S \frac{6(cV^2+a)w}{(1-V^2)^4} \geq \pi(c+a)/2m.$$

Dividing by $|m'|$ we get

$$3.47 \quad \int_S \left[\left\langle -2\nabla_S \left(\frac{cV^2+a}{V(1-V^2)^3} \right), \nabla_S V \right\rangle + \frac{2cw}{(1-V^2)^3} - \frac{6(cV^2+a)w}{(1-V^2)^4} \right] \geq \pi(c+a)/(2m|m'|).$$

Following Robinson [R] we shall choose two different set of values for c and α to obtain two inequalities contradicting each other. We put $c = -1$, $\alpha = 1$. We check that $F > 0$ on $N \sim \bar{Q}$. Then 3.47 becomes

$$3.48 \quad \int_S \left[\left\langle -2\nabla_S \left(\frac{1}{V(1-V^2)^2} \right), \nabla_S V \right\rangle - \frac{2w}{(1-V^2)^3} - \frac{6w}{(1-V^2)^3} \right] \geq 0.$$

This implies

$$3.49 \quad \int_S \left[2 \frac{|\nabla_S V|^2}{V^2(1-V^2)^2} + \frac{(-2)(-2)(-2V)}{V(1-V^2)^3} |\nabla_S V|^2 \right] \geq \int_S \frac{8|\nabla_S V|^2}{(1-V^2)^3} + \int_S \frac{8|g(n, \nabla V)|^2}{(1-V^2)^3}$$

where we have used $w = |\nabla_S V|^2 + |g(n, \nabla V)|^2$. Thus

$$3.50 \quad \int_S \left[2 \frac{|\nabla_S V|^2}{V^2(1-V^2)^2} - \frac{16|\nabla_S V|^2}{(1-V^2)^3} \right] \geq 8m'^2 \int_S \frac{1}{(1-V^2)^3} > 8m'^2 |S|.$$

The last inequality follows because we have $1/(1-V^2)^3 > 1$ on S as $V > 0$. Thus by adding the terms in the left hand side we get

$$3.51 \quad \int_S \frac{2(1-9V^2)}{V^2(1-V^2)^3} |\nabla_S V|^2 > 8m'^2 |S|.$$

Now using

$$3.52 \quad \frac{1-9V^2}{(1-V^2)^3} < 1 \quad \text{on } S$$

we have

$$3.53 \quad \int_S \frac{|\nabla_S V|^2}{V^2} > 4m'^2 |S|.$$

To check 3.52 we suppose $f(V) = (1-9V^2)/(1-V^2)^3$. Then

$$\frac{df}{dV} = - \frac{2+6V^2}{(1-V^2)^4} 6V < 0$$

for $V > 0$. Thus $f(V)$ is a strictly decreasing function of V . Hence $f(V) < f(0) = 1$ for $V > 0$.

Finally using 3.4 we get

$$4\pi > 4m'^2 |S| .$$

Thus by 3.43 we have

$$3.54 \quad |S| > 16\pi m'^2 .$$

We now consider $c = 1$ and $\alpha = 0$ in 3.47. Clearly F is positive. Thus we have

$$3.55 \quad \int_S \left[\left\langle -2\nabla_S \left(\frac{V}{(1-V^2)^3} \right), \nabla_S V \right\rangle + \frac{2w}{(1-V^2)^3} - \frac{6V^2 w}{(1-V^2)^4} \right] \geq \pi / (2m|m'|) .$$

This implies

$$3.56 \quad \int_S \left[-\frac{2|\nabla_S V|^2}{(1-V^2)^3} + \frac{(-2)(-3)(-2V)V|\nabla_S V|^2}{(1-V^2)^4} + \frac{2w}{(1-V^2)^4} (1-V^2-3V^2) \right] \geq \pi / (2m|m'|) .$$

Using $w = |\nabla_S V|^2 + m'^2$ we get

$$3.57 \quad \int_S \left[-\frac{2|\nabla_S V|^2}{(1-V^2)^3} - \frac{12V^2|\nabla_S V|^2}{(1-V^2)^4} + \frac{2|\nabla_S V|^2(1-4V^2)}{(1-V^2)^4} + \frac{2m'^2(1-4V^2)}{(1-V^2)^4} \right] \geq \pi / (2m|m'|) .$$

Now adding the first three terms we have

$$3.58 \quad \int_S \left[\frac{-18V^2}{(1-V^2)^4} |\nabla_S V|^2 + \frac{2m'^2(1-4V^2)}{(1-V^2)^4} \right] \geq \pi / (2m|m'|) .$$

Thus

$$3.59 \quad 2m'^2 \int_S \frac{1-4V^2}{(1-V^2)^4} \geq \pi / (2m|m'|) .$$

Then using

$$3.60 \quad \frac{1-4V^2}{(1-V^2)^4} < 1 \quad \text{on } S$$

we get

$$3.61 \quad 2m'^2 |S| > \pi / (2m|m'|) .$$

To check 3.60 we suppose $f(V) = (1-4V^2)/(1-V^2)^4$. Then

$$\frac{df}{dV} = \frac{-3V^2}{(1-V^2)^5} \cdot 8V < 0$$

for $V > 0$. Thus $f(V) < f(0) = 1$ for $V > 0$.

Using 3.43 in 3.61 we get

$$32\pi^2 m^2 / |S| > |S| / 8m^2$$

giving $16\pi m^2 > |S|$ which contradicts 3.54. $\square$

We can now complete the proof of the main Theorem 1.1.

Proof of 1.1. By Theorem 2.2 and Theorem 2.8 either N is diffeomorphic to $\mathbb{R}^3$ or there exists a C^2 embedded totally geodesic sphere S in $N - Q$. But by Theorem 3.22 such a sphere cannot exist. Hence the theorem follows. $\square$

CHAPTER IV

APPLICATION TO GENERAL RELATIVITY

4.0.

In this chapter we apply our main theorem to prove that a complete, asymptotically Euclidean, static perfect fluid space-time with connected fluid region and satisfying "timelike convergence condition" is diffeomorphic to $\mathbb{R}^3 \times \mathbb{R}$, without using the Poincaré conjecture. The first section deals with the preliminaries regarding the static perfect fluid space-time. In Section 4.2 after discussing some standard preliminary results from Lorentzian geometry, we describe Gannon's proof for simple connectedness of the Cauchy surface when the metric is smooth. In Section 4.3 we modify the proof for $C^{1,1}$ metric. This enables us to apply our main theorem and hence complete the proof of our assertion.

4.1. Static perfect fluid space-time

By a static perfect fluid spacetime we mean a geodesically complete space-time $(M, {}^4g)$ such that:

(i) M is a C^∞ manifold diffeomorphic to $N \times \mathbb{R}$ where for each $t \in \mathbb{R}$, $N_t = N \times \{t\}$ is a spacelike three-manifold.

(ii) The Lorentz metric 4g can be written as

$$4.1 \quad {}^4g = -V^2(dt \otimes dt) + g$$

where V is a positive $C^{1,1}$ function and g is a tensor such that g restricted to N is a Riemannian metric on N , and V and g are independent of t . We assume that 4g is at least $C^{1,1}$.

(iii) $(M, {}^4g)$ satisfies Einstein's equation

$$4.2 \quad \text{Ric}({}^4g)_{AB} - \frac{1}{2} \text{Scalar}({}^4g) {}^4g_{AB} = 8\pi((\rho+p)u_A u_B + p {}^4g_{AB})$$

where ρ and p are bounded measurable functions and u_A is a unit timelike vector field on M .

Clearly in the co-ordinate system $\{t, x^1, x^2, x^3\}$, where $\{x^1, x^2, x^3\}$ is a co-ordinate system on N , the unit timelike normal form n on each N_t is given by $n_0 = V^{-1}$, $n_\alpha = 0$, $\alpha = 1, 2, 3$. Thus n_A is independent of t . Hence each N_t is totally geodesic in M and by virtue of the Gauss-Codazzi embedding equations for the Lorentzian metric 4g , 4.2 decomposes into

$$4.3 \quad \text{Ric}(g)_{\alpha\beta} = V^{-1}V_{;\alpha\beta} + 4\pi(\rho-p)g_{\alpha\beta},$$

and

$$4.4 \quad \Delta V = 4\pi V(\rho+3p) \text{ on } N,$$

where $;$ denotes the covariant derivative with respect to g and Δ denotes the Laplacian with respect to g .

It is clear that ρ and p are independent of t . It follows from 4.2 that if 4g satisfies the timelike convergence condition, namely,

$$4.5 \quad \text{Ric}({}^4g)(W, W) \geq 0$$

for all timelike vectors W , then $\rho + 3p \geq 0$. By continuity 4.5 implies the null convergence condition, namely, $\text{Ric}({}^4g)(K, K) \geq 0$ for all null vectors K . By virtue of 4.2 the latter condition is satisfied if and only if $\rho + p \geq 0$.

We also assume that there exists an open connected region $Q \subset N$ such that $\text{ess inf}_K (\rho+p) > 0$ for all compact $K \subset Q$ and $\rho = p = 0$ in $N \sim \overline{Q}$.

The functions ρ and p are respectively called the density and the pressure of the fluid. The assumption that the fluid region Q is connected is needed here in order to apply our main theorem. However the

null convergence condition $\rho + p \geq 0$ is crucial both in the proof of Gannon's theorem and in the proof of our main theorem. We also assume that 4g satisfies the timelike convergence condition so that by 4.4, ΔV is non-negative. However when $\mathcal{Q}$ is unbounded, the null convergence condition will be sufficient for our purpose. Remark 1.14 is relevant here.

We say that $(M, {}^4g)$ is asymptotically Euclidean if (N, g) satisfies condition (c) of the main Theorem 1.1 and for some $\mu \in (0, 1)$, V satisfies (by 1.7, this is automatic if $\overline{\mathcal{Q}}$ is compact)

$$4.6 \quad 1 - V = O(|x|^{-\mu}) \quad , \quad \frac{\partial V}{\partial x^\alpha} = O(|x|^{-1-\mu}) \quad \text{as } |x| \rightarrow \infty .$$

4.2. Gannon's theorem

First we recall some standard definitions and results from Lorentzian geometry. We shall follow mostly Hawking and Ellis [HE]. Let $(\tilde{M}, {}^4\tilde{g})$ be a smooth spacetime not necessarily complete. For sets $\zeta \subset \tilde{M}$ the chronological future $I^+(\zeta)$ is the set of all points in $\tilde{M}$ which can be reached from ζ by a future-directed timelike curve (of nonzero extent) in $\tilde{M}$. The causal future $J^+(\zeta)$ is the union of ζ with the set of all points in $\tilde{M}$ which can be reached from ζ by a future-directed non-space-like curve in $\tilde{M}$. A set $\zeta \subset \tilde{M}$ is said to be achronal if $I^+(\zeta) \cap \zeta = \emptyset$, that is, if there are no two points of ζ with timelike separation. A set ζ is said to be a future set if $\zeta \supset I^+(\zeta)$. For any set ζ , $J^+(\zeta)$ is a future set. We shall use the following well-known proposition.

4.7 PROPOSITION. *If ζ is a future set then its boundary $\partial\zeta$ is a closed, embedded, achronal three-dimensional Lipschitz (that is $C^{0,1}$) manifold.*

Proof of 4.7. We refer to Proposition 6.3.1 in [HE]. □

We need to consider the congruences of null geodesics orthogonal to a smooth spacelike embedded two-dimensional manifold A . Let W and X be two null vector fields orthogonal to A and normalized such that ${}^4\tilde{g}(W, X) = -1$. At any point $p \in A$, $W|_p$ and $X|_p$ span the timelike complementary subspace orthogonal to the two dimensional tangent space of A at p . We complete the pseudo-orthonormal basis by introducing two unit mutually orthogonal spacelike vector fields Y and Z on A . Now we define two null second fundamental form tensors of A by

$$4.8 \quad {}^{(W)}\chi_{AB} = h^C_A h^D_B {}^4\nabla_D W_C \text{ and } {}^{(X)}\chi_{AB} = h^C_A h^D_B {}^4\nabla_D X_C$$

where h^C_A is the projection operator from $T_p \tilde{M}$ to $T_p A$ and is given by

$$h^C_A = \left(Y^C Y_A + Z^C Z_A \right).$$

There will be two families of null geodesics normal to A corresponding to the two null normals W and X . Let $\gamma(\omega)$ be a null geodesic along W , ω being an affine parameter such that $W|_p = (\partial/\partial\omega)_\gamma|_0$ (we recall that ω is an affine parameter if the covariant derivative of $(\partial/\partial\omega)_\gamma$ with respect to $(\partial/\partial\omega)_\gamma$ vanishes). We may fix the pseudo-orthonormal basis (E_1, E_2, E_3, E_4) by taking $E_1 = Y$, $E_2 = Z$, $E_3 = X$ and $E_4 = W$ at A , and using parallel propagation along the null geodesics. For a null geodesic $\gamma(\omega)$ a point q on $\gamma(\omega)$ is said to be conjugate to A along $\gamma(\omega)$ if there is a Jacobi field J along $\gamma(\omega)$ normal to $(\partial/\partial\omega)_\gamma$ and not identically zero, which satisfies with respect to the above pseudo-orthonormal basis $J^m = 0$, $m = 1, 2$, at q and the initial conditions $dJ^m/d\omega = {}^{(W)}\chi_{mn} J^n$, $m, n \in \{1, 2\}$, at $p = \gamma(0)$ at A . We note that J^3 vanishes all along γ . We discuss J^4 in Appendix E (after (E3)).

(Some authors, for example [BE], call q a focal point.) The following lemma is well-known.

4.9 LEMMA (Focusing Lemma). *Let $\text{Ric}({}^4\tilde{g})(W, W) \geq 0$ everywhere. If the "expansion" ${}^{(W)}\chi_{AB} {}^4\tilde{g}^{AB}$ is negative on A , there will be a point conjugate to A along $\gamma(\omega)$ within an affine distance*

$$\omega_0 = 2 / \left[- {}^{(W)}\chi_{AB} {}^4\tilde{g}^{AB} \right] \text{ from } A, \text{ provided } \gamma(\omega) \text{ can be extended that far.}$$

Moreover, points on a null geodesic beyond any point conjugate to A would lie in $I^+(A)$.

Proof of 4.9. For the first part we refer to Proposition 4.4.6 and for the second part we refer to Proposition 4.5.14 in [HE]. $\square$

The spacetime $(\tilde{M}, {}^4\tilde{g})$ is said to be causally simple if for every compact set $K \subset \tilde{M}$, $\partial J^+(K) = J^+(K) \sim I^+(K)$ and $\partial J^-(K) = J^-(K) \sim I^-(K)$. Here $J^-(K)$ is the causal past of K . In a causally simple spacetime $\partial J^+(\zeta)$ for any set ζ must lie on some future directed null geodesics from ζ . This follows from Proposition 4.5.10 in [HE] which says that if p and q are joined by a non-spacelike curve $\tau(t)$ which is not a null geodesic they can also be joined by a timelike curve.

A Cauchy surface is a spacelike hypersurface which every non-spacelike curve intersects exactly once. We shall need the following proposition.

4.10 PROPOSITION. *Suppose $(\tilde{M}, {}^4\tilde{g})$ has a Cauchy surface. Then $(\tilde{M}, {}^4\tilde{g})$ is causally simple.*

Proof of 4.10. By Proposition 6.6.3 in [HE] existence of Cauchy surface implies "global hyperbolicity" and by Proposition 6.6.1 in [HE] "global hyperbolicity" implies that the spacetime is causally simple. $\square$

4.11 Remark. Thus if A is a two dimensional smooth embedded submanifold of a Cauchy surface then the points of $\partial J^+(A)$ are contained in the null geodesics with past endpoints on A and orthogonal to A . There

are two such families of null geodesics. The following is an important lemma due to Gannon [G1].

4.12 LEMMA. *Let $\hat{N}$ be a simply connected Cauchy surface and A be a two dimensional smooth embedded connected submanifold of $\hat{N}$. Then, with reference to the above remark, a null geodesic of one family cannot meet a null geodesic of the other family in $\partial J^+(A) \sim A$.*

Proof of 4.12 (References [G1], [G2]). Suppose to the contrary that there is a null geodesic γ belonging to one family meeting a null geodesic σ belonging to the other family in $\partial J^+(A)$ off A . Let p and q be the endpoints of γ and σ respectively on A and let τ be the path on A joining p and q . $\gamma \cup \sigma \cup \tau$ form a closed loop δ in $\partial J^+(A)$. Then, by pushing τ a little off A into the family containing σ and joining the resulting endpoints in a suitable manner, we can deform δ to a closed loop $\bar{\delta}$ in $\partial J^+(A)$ meeting A only at the point p and transversely there. By choosing a global timelike vector field Y we project $\bar{\delta}$ along the integral curves of Y to a loop ω in $\hat{N}$ meeting A transversely at p . Since $\hat{N}$ is simply connected the intersection number of ω with A is zero mod 2. Hence ω must meet A in some other point r . The integral curve of Y through r is then a timelike path from A to $\partial J^+(A)$ which is impossible as $\partial J^+(A)$ is achronal. $\square$

Now we shall state and prove Proposition 1.2 in Gannon's paper [G1]. We state the proposition as a theorem with a view to application to the static perfect fluid spacetime.

4.13 THEOREM (Gannon [G1]). *Let $(\tilde{M}, {}^4\tilde{g})$ be a geodesically complete smooth spacetime with Cauchy surface $\tilde{N}$ so that $\tilde{M}$ is homeomorphic to $\tilde{N} \times \mathbb{R}$. Suppose on $\tilde{M}$, for all null vectors W , $\text{Ric}({}^4\tilde{g})_{AB} W^A W^B \geq 0$. Suppose for some compact submanifold-with-boundary $\bar{N} \subset \tilde{N}$, $\tilde{N} \sim \bar{N}$ is homeomorphic to $\mathbb{R}^3$ minus the closed unit ball and suppose $\partial \bar{N}$ is a smooth*

sphere with the inward directed null geodesics orthogonal to $\partial\bar{N}$ converging (that is, the expansion $\chi_{AB} {}^4\tilde{g}^{AB}$ is negative), everywhere on $\partial\bar{N}$. (Here inward means those geodesics which locally project orthogonally into N .)

Then $\tilde{N}$ is simply connected.

Proof of 4.13. Suppose to the contrary $\tilde{N}$ is nonsimply connected. Let $\hat{M}$ be the universal covering space for $\tilde{M}$ and $\pi : \hat{M} \rightarrow \tilde{M}$ be the corresponding covering map. Since $\tilde{M}$ is diffeomorphic to $\tilde{N} \times \mathbb{R}$, $\pi^{-1}(\tilde{N}) \equiv \hat{N}$ is the universal covering space for $\tilde{N}$ and $\hat{M}$ is diffeomorphic to $\hat{N} \times \mathbb{R}$. Let ${}^4\hat{g}$ be the lift of ${}^4\tilde{g}$ for $\hat{M}$. Then clearly $\hat{N}$ is a Cauchy surface for the spacetime $(\hat{M}, {}^4\hat{g})$.

Let A be one component of $\pi^{-1}(\partial\bar{N})$. Then π restricted to A , $\pi|_A$ is a covering map for $\partial\bar{N}$. But $\partial\bar{N}$ is a 2-sphere which is simply connected. Hence $\pi|_A$ is a diffeomorphism. But for a point $p \in \partial\bar{N}$, $\pi^{-1}(p)$ contains more than one point because $\pi_1(\tilde{N}) \neq 0$. Hence we have more than one component of $\pi^{-1}(\partial\bar{N})$.

Now we have $\partial\pi^{-1}(\bar{N}) = \pi^{-1}(\partial\bar{N})$. But $\bar{N}$ is a retract of $\tilde{N}$; hence $\pi^{-1}(\bar{N})$ is the universal covering space $\hat{N}$ for $\bar{N}$. Since A is a component of $\partial\pi^{-1}(\bar{N})$ and since $\partial\pi^{-1}(\bar{N})$ has at least two components, A is not the boundary of a compact three dimensional manifold in $\hat{N}$. But $\hat{N}$ is a retract of $\hat{M}$ so A does not bound a compact three dimensional manifold in $\hat{M}$.

Now, by Proposition 4.7, $\partial J^+(A)$ is an acronal, embedded, closed, three dimensional Lipschitz manifold in $\hat{M}$. Since $\hat{M}$ has a Cauchy surface by Proposition 4.10, $\hat{M}$ is causally simple. Hence, by Remark 4.11, $\partial J^+(A)$ is contained in the null geodesics with past end points on A and meeting

A orthogonally. By Lemma 4.12 the inward directed null geodesics do not meet the outgoing null geodesics off A . Hence both families form proper closed submanifolds of $\hat{M}$ with boundary A . Since $\tilde{M}$ is null geodesically complete so is $\hat{M}$. Hence, by the Focussing Lemma 4.9 the inward directed null geodesics (which are everywhere converging on A by hypothesis) must leave $\partial J^+(A)$ within a finite affine distance. That implies that the closed submanifold with boundary A consisting of the family of inward directed null geodesics is compact. This contradicts our earlier statement that A does not bound any compact three dimensional manifold in $\hat{M}$. Hence $\tilde{N}$ must be simply connected. $\square$

4.3. Extension to $C^{1,1}$ metric

We now extend Gannon's Theorem for $C^{1,1}$ metric 4g for the case when $(M, {}^4g)$ is the asymptotically Euclidean static perfect fluid space-time of Section 4.1. This can be done in a way similar to the extension of singularity theorems to the case when the metric is $C^{1,1}$, that is, by taking a smooth sequence of metrics and using a sharpened version of the Focusing Lemma (ref. page 285 of [HE]).

Let N be as in Section 4.1 and let $\partial N_R \subset N$ denote the sphere $|x| = R$ in the asymptotic co-ordinate system. In the co-ordinate system $\{t, x^1, x^2, x^3\}$, the two null vectors orthogonal to ∂N_R can be given by

$$\begin{aligned}
 4.14 \quad \pm W^A &= {}^4g^{AB} \left(f \frac{\partial t}{\partial x^B} \mp \frac{\partial |x|}{\partial x^B} \right) |x| (2g^{\pi\mu} x^\pi x^\mu)^{-1/2} \\
 &= \begin{cases} V^{-1} & , \quad A = 0 , \\ \mp g^{\alpha\beta} x^\beta (2g^{\pi\mu} x^\pi x^\mu)^{-1/2} & , \quad A = \alpha = 1, 2, 3 , \end{cases}
 \end{aligned}$$

where

$$4.15 \quad f < 0, \quad f^2 = \left[{}^4g^{AB} \frac{\partial |x|}{\partial x^A} \frac{\partial |x|}{\partial x^B} \right] / \left[{}^4g^{AB} \frac{\partial t}{\partial x^A} \frac{\partial t}{\partial x^B} \right] = V^2 (g^{\alpha\beta} x^\alpha x^\beta) / |x|^2$$

and the factors $|x| / (2g^{\pi\mu} x^\pi x^\mu)^{1/2}$ are included so that $\pm W^A$ are normalized to give $g({}^+W^A, {}^-W^A) = -1$. Clearly ${}^+W \equiv W$ is a future-directed, inward directed null vector orthogonal to ∂N_R .

Now let Y_a , $a \in \{2, 3\}$, be a fixed smooth local basis vector field on ∂N_R and Z_a , $a \in \{2, 3\}$ be a local orthonormal (in the 4g metric) basis vector field on ∂N_R . We have

$$4.16 \quad Z_a = \Pi_{ab} Y_b \quad \text{where } \Pi_{ab} \text{ is a } 2 \times 2 \text{ matrix.}$$

Clearly

$$4.17 \quad (\Pi^T \Pi)^{-1}_{bc} = g(Y_b, Y_c) .$$

Hence the expansion of the inward directed null geodesics on ∂N_R in the 4g metric is given by

$$4.18 \quad \begin{aligned} (W) \chi_{AB} {}^4g^{AB} &= \sum_{a=2}^3 {}^4g(Z_a, {}^4\nabla_{Z_a} W) \\ &= \sum_{b,c=2}^3 ({}^4\nabla_B W_A) Y_b^A Y_c^B (\Pi^T \Pi)_{bc} . \end{aligned}$$

By virtue of the asymptotic conditions we have in $\{t, x^1, x^2, x^3\}$ the co-ordinate system

$$4.19 \quad \begin{aligned} (W) \chi_{AB} {}^4g^{AB} &= \sum_{\alpha=2}^3 \left[\frac{\partial W_\alpha}{\partial x^\beta} - \Gamma_{\alpha\beta}^\tau W_\tau \right] Z_\alpha^\alpha Z_\alpha^\beta \\ &= \sum_{\alpha=2}^3 \left[-\delta_\beta^\alpha (2g^{\pi\mu} x^\pi x^\mu)^{-1/2} + 2x^\alpha (2g^{\pi\mu} x^\pi x^\mu)^{-3/2} g^{\beta\tau} x^\tau \right] Z_\alpha^\alpha Z_\alpha^\beta \\ &\quad + O(1/R^2) = -\frac{\sqrt{2}}{R} + O(1/R^2) . \end{aligned}$$

Thus for sufficiently large R , $^{(W)}\chi_{AB} {}^4g^{AB}$ is everywhere negative on ∂N_R . We now fix a particular ∂N_R , say ∂N , such that $^{(W)}\chi_{AB} {}^4g^{AB}$ is strictly negative everywhere on ∂N .

4.20 Remark. It is also clear from 4.18 that if $\{^k g\}$ is a sequence of smooth metrics approaching to g in $C^{1,\lambda}$ norm on an open set in N containing ∂N then for sufficiently large k the expansion of ∂N in the 4g_k metric on M defined by

$$4.21 \quad {}^4g_k = -{}^k V^2 dt^2 + {}^k g, \quad {}^k V \text{ independent of } t,$$

is also negative everywhere on ∂N .

We now give a sharpened version of the first part of the Focusing Lemma 4.9. In fact for null geodesic congruences we have the following result.

4.22 LEMMA. Let $(\tilde{M}, {}^4\tilde{g})$ be a smooth complete spacetime, A be a smooth embedded two dimensional spacelike submanifold, $\gamma : [0, \omega^*] \rightarrow \tilde{M}$ be a null geodesic with past end point on A , meeting A orthogonally at p . Let $(\partial/\partial\omega)_\gamma|_0 = W|_p$ where W is a given null vector field orthogonal to A and satisfying ${}^4\nabla_W W = 0$ in a neighbourhood of p . Then if

$$\theta_0 = {}^{(W)}\chi_{AB} {}^4\tilde{g}^{AB}|_p \text{ is negative, } \omega^* > 4|\theta_0|^{-1} \text{ and}$$

$$\text{Ric}({}^4\tilde{g})\left(\left(\frac{\partial}{\partial\omega}\right)_\gamma\Big|_\omega, \left(\frac{\partial}{\partial\omega}\right)_\gamma\Big|_\omega\right) \geq -\epsilon > -\frac{1}{4}|\theta_0|^2$$

for $\omega \in [0, \omega^*]$ then there will be a point on $\gamma(\omega)$ conjugate to A along γ within an affine distance $\omega_0 = 2\left(\left(1-2\epsilon\theta_0^{-2}\right)|\theta_0|\right)^{-1}$.

Proof of 4.22. The proof is similar to the proof of Lemma 4.9. For the sake of completeness we include the proof in Appendix E. $\square$

4.13' THEOREM (Extended Gannon's Theorem). Let $(M, {}^4g)$ be the asymptotically Euclidean complete $C^{1,1}$ static perfect fluid space-time

satisfying the null convergence condition and let N be the $t = 0$ hypersurface. Then N is simply connected.

Proof of 4.13'. We put $\omega^* = 8 \sup_{\partial N} \left(\left| {}^{(W)}\chi_{AB} {}^4g^{AB} \right| \right)^{-1}$. Let

$\gamma_p : [0, \omega^*] \rightarrow M$ denote the inward directed null geodesics with past ends p at ∂N and orthogonal to ∂N and having parameter length ω^* . Then $\gamma_p(\omega)$ satisfies the following ordinary differential equations in the co-ordinate system $\{t, x^1, x^2, x^3\}$,

$$4.23 \quad \left. \begin{aligned} \frac{d^2 t}{d\omega^2} &= 0 \end{aligned} \right\}$$

$$4.24 \quad \left. \begin{aligned} \frac{d^2 x^\alpha}{d\omega^2} + \Gamma_{\beta\sigma}^\alpha \frac{dx^\beta}{d\omega} \frac{dx^\sigma}{d\omega} &= 0 \end{aligned} \right\}$$

with initial condition $\gamma_p(0) = p$, $(\partial/\partial\omega)|_p = W|_p$, where $W|_p$ is the inward directed null vector orthogonal to ∂N and is given by 4.14. $\Gamma_{\beta\sigma}^\alpha$ are the Christoffel symbols of the metric g .

We take an open set $N_5 \subset N$ such that $\bar{N}_5$ is compact and

$\bigcup_{p \in \partial N} \gamma_p \subset N_5 \times [0, 2\omega^* W^0(p)]$ where $W^0(p)$ is the 0th component of $W|_p$.

Now let $\{^k g\}$ be a sequence of smooth complete Riemannian metrics on N and $\{^k V\}$ be a sequence of smooth functions on N such that $\{^k g\}$ and $\{^k V\}$ converge to g and V respectively in $C^{1,\lambda}$, $\lambda \in (0, 1)$, norm on $\bar{N}_5$. Let 4g_k be as in 4.21. By Remark 4.20 we may assume that for each k the expansion $\theta_k(p)$ of ∂N in the 4g_k metric is negative everywhere on ∂N and say,

$$4.25 \quad \frac{3}{2} \inf_{\partial N} \left| {}^{(W)}\chi_{AB} {}^4g^{AB} \right| \geq \inf_{\partial N} |\theta_k| \geq \frac{1}{2} \inf_{\partial N} \left| {}^{(W)}\chi_{AB} {}^4g^{AB} \right|.$$

Let ${}^k\gamma_p : [0, \omega^*] \rightarrow M$ denote the ingoing null geodesics with past ends at ∂N and orthogonal to ∂N in the 4g_k metric. From 4.14 it follows that the initial tangent vectors ${}^k W|_p$ in the 4g_k metric approaches $W|_p$ as k tends to infinity. Thus for all sufficiently large k we have

$$4.26 \quad \bigcup_{p \in \partial N} {}^k\gamma_p \subset N_5 \times [0, 2\omega^* W^0(p)] .$$

This is because the solution of the geodesic equations depend continuously on the initial positions, initial tangent vectors and the coefficients by the standard theorems of ordinary differential equations (see, for example, Theorem 2.1 on page 94 in [HA]). Also since $\bar{N}_5 \times [0, 2\omega^* W^0(p)]$ is compact there is an upper bound for the components of ${}^k W$ and W in $N_5 \times [0, 2\omega^* W^0(p)]$. Hence for sufficiently large k we find $\varepsilon > 0$ with

$$4.27 \quad \text{Ric}({}^4g_k)({}^k W, {}^k W) \geq -\varepsilon > -\frac{1}{16} \inf_{\partial N} \left| (W)_{\chi_{AB}} {}^4g^{AB} \right|^2 .$$

We now fix a particular k , say k' , such that 4.26 and 4.27 hold for ${}^4g_{k'}$. As in the proof of 4.13 let A be a component of $\pi^{-1}(\partial N)$ where $\pi : \hat{M} \rightarrow M$ is the universal covering map for M . 4.25 and 4.27 give

$$\text{Ric}({}^4g_{k'})({}^{k'} W, {}^{k'} W) \geq -\varepsilon > -\frac{1}{4} \inf_{\partial N} |\theta_{k'}|^2$$

along ${}^{k'}\gamma_p(\omega)$ for $\omega \leq \omega^*$.

Hence by Lemma 4.22 the inward directed null geodesics will leave $\partial J^+(A)$ within parameter length

$$4 \sup_{\partial N} |\theta_{k'}|^{-1} \leq 8 \sup_{\partial N} \left| (W)_{\chi_{AB}} {}^4g^{AB} \right|^{-1} = \omega^* .$$

Hence, as in the proof of 4.13, $\partial J^+(A)$ in the ${}^4g_{k'}$ metric bounds a

compact set in $\hat{M}$ so that N must be simply connected. $\square$

As a consequence of the extended Gannon's Theorem 4.13' and our main theorem we prove the following theorem. This result has been claimed in [LB] assuming the Poincaré conjecture to be true. In fact the asymptotic conditions and Gannon's Theorem imply that N has the same homotopy as $\mathbb{R}^3$ ([LB]). What we have shown here is that a fake 3-cell cannot occur in N and hence N is diffeomorphic to $\mathbb{R}^3$.

4.28 THEOREM. *A complete asymptotically Euclidean static perfect fluid space-time having connected fluid region and satisfying the timelike convergence condition is diffeomorphic to $\mathbb{R}^3 \times \mathbb{R}$.*

Proof of 4.28. By the extended Gannon's Theorem N is simply connected and hence our main Theorem 1.1 applies. $\square$

CHAPTER V

MISCELLANEOUS REMARKS AND GENERALIZATIONS

5.0.

In this chapter we discuss various generalizations to Theorem 1.1 and some further applications to general relativity. In particular in Section 5.2 we show how Theorem 1.1 can be generalized to suitable space-like hypersurfaces in a certain class of space-times, not necessarily static or perfect fluid. As before the results are different from those in the paper of Frankel and Galloway [FG]; for example, using the asymptotic conditions we can allow certain terms $(\Phi_1 \text{ and } \Phi_2)$ to vanish identically outside a connected set, whereas the results in [FG] require that $\text{ess inf}_K (\Phi_1 + \Phi_2) > 0$ for all compact subsets K of the 3-manifold whose topology is to be investigated.

5.1.

Here we give an alternative definition of $Q \subset N$:

Q is connected, $\text{ess inf}_K (\Phi_1 + \Phi_2) > 0$ for all compact sets

$K \subset Q$ and $\Phi_2 \equiv 0$ on $N \sim \overline{Q}$.

Because of the following lemma the main Theorem 1.1 continues to hold with Q defined as above.

5.1 LEMMA. *Let Q be defined as above. Suppose $\Phi_1 + \Phi_2 \geq 0$ on N .*

If there exists an embedded area minimizing (defined in 2.1) minimal C^2 sphere S on $N \sim Q$ then $\Phi_1 \equiv 0$ on each component of $N \sim \overline{Q}$ with closure intersecting S .

5.2 Remark. In the above lemma we continue to assume that N is

simply connected so that S separates N . The assumption that V is a positive $C^{1,1}$ function on N is also used.

Proof of 5.1. Since $\Phi_2 \equiv 0$ on $N \sim Q$ the equation (A1) in Appendix A implies that Φ_1 is a Lipschitz function on $N \sim \overline{Q}$ and

$$5.3 \quad \Phi_{1;\beta} = 2V^{-1}\Phi_1 V_{;\beta}.$$

Hence $\Phi_1 = CV^2$ on $N \sim \overline{Q}$ where by virtue of $\Phi_1 + \Phi_2 \geq 0$, C (which is constant on each component of $N \sim \overline{Q}$) is non-negative.

Now as in the proof of Theorem 2.8 we can approximate S by a sequence of C^2 spheres S_t , $t \downarrow 0$. We can take $S_t \subset N \sim \overline{Q}$ because S separates N and Q is connected. The fact that S is area-minimizing and Φ_1 is continuous on (each component of) $N \sim Q$ gives (using 2.7) that

$$\lim_{t \rightarrow 0} \int_{S_t} \Phi_1 V^2 = 0. \quad \text{Hence} \quad \int_S CV^4 = 0.$$

But $V > 0$. Hence on S , $C = 0$ giving $\Phi_1 \equiv 0$ on each component of $N \sim \overline{Q}$ with closure intersecting S . $\square$

5.4 Remark. If we use an extra asymptotic condition that the scalar curvature R of g is $o(1)$ then it follows that $\Phi_1 \equiv 0$ on each unbounded component of $N \sim \overline{Q}$ without the assumption that an area minimizing sphere S exists in $N \sim Q$. This is because from 1.2 we have $R = \Delta V + 3\Phi_1$ and by hypothesis $\Delta V = 0$. However by Theorems 2.2 and 2.8 if such S does not exist then N is topologically Euclidean and hence the conclusion of the main theorem follows without the above mentioned asymptotic decay of R .

5.5 Remark. Arguments similar to those used in the proof of Lemma 5.1 show that we can also define Q as follows: Q is connected, $\text{ess inf}_K (\Phi_1 + \Phi_2) > 0$ for all compact sets $K \subset Q$ and $\Phi_1 \equiv 0$ in $N \sim \overline{Q}$.

Then (A1) in Appendix A implies that $\Phi_{2;\beta} = -2V^{-1}\Phi_2 V_{;\beta}$ giving $\Phi_2 = CV^{-2}$ where C is constant on each component of $N \sim \overline{Q}$. The same arguments as before then imply that $\Phi_2 \equiv 0$ on each component of $N \sim \overline{Q}$ with closure intersecting S .

We shall now discuss some generalizations of equation 1.2.

5.6 LEMMA. *The main Theorem 1.1 continues to hold with the usual conditions on V, g, Φ_1 and Φ_2 if we replace 1.2 by*

$$5.7 \quad C_1 \Phi_1 g_{\alpha\beta} \geq \text{Ric}(g)_{\alpha\beta} - V^{-1}V_{;\alpha\beta} \geq \Phi_1 g_{\alpha\beta} \quad \text{almost everywhere}$$

where C_1 is a constant.

The first inequality in 5.7 is added to ensure that $\Phi_1 \equiv 0$ on $N \sim \overline{Q}$ implies

$$5.8 \quad \text{Ric}(g)_{\alpha\beta} = V^{-1}V_{;\alpha\beta}, \quad \text{on } N \sim \overline{Q}.$$

Proof of 5.6. 5.8 enables us to apply the reasoning of Chapter III (in particular we refer to Theorem 3.22) to prove that no embedded C^2 totally geodesic sphere exists in $N \sim \overline{Q}$. To prove our assertion that Theorem 1.1 holds with 1.2 replaced by 5.7 we now simply need to use the inequality

$$V^2 \text{Ric}(g)(n, n) + V\Delta V - VV_{;\alpha\beta}n^\alpha n^\beta \geq V^2(\Phi_1 + \Phi_2)$$

in 2.7 instead of the equality. The rest of the arguments in the proof of Theorem 2.8 then apply and since there is no embedded C^2 totally geodesic sphere in $N \sim \overline{Q}$, it follows that N must be diffeomorphic to $\mathbb{R}^3$. $\square$

5.9 Remark. It is important to note that when Φ_1 satisfies the inequality 5.7 the contracted Bianchi identity for g may no longer be equivalent to

$$(\Phi_2 - \Phi_1)_{;\beta} = -2V^{-1}(\Phi_1 + \Phi_2)V_{;\beta}.$$

Hence we should be careful in using the alternative definition of Q given in the paragraph preceding Lemma 5.1. However in an open set where $\Phi_1 \equiv 0$, the above relation continues to hold because in this case 5.7 becomes an equality.

5.2. Further applications to general relativity

We shall now apply Theorem 1.1 to a complete, asymptotically Euclidean (that is, the induced metric satisfies the asymptotic conditions 1.4) simply connected space-like hyper-surface N in a suitable class of space-times $(M, {}^4g)$, not necessarily static. In particular we shall show that Theorem 4.28 holds for any static space-time, not necessarily perfect fluid, under similar asymptotic and energy conditions (here by energy conditions we mean the conditions on the energy-momentum tensors which will be equivalent to the usual conditions on Φ_1 and Φ_2 in condition (b) of Theorem 1.1).

Let $(M, {}^4g)$ and N be as above. In a neighbourhood of N in M we write

$${}^4g = -V^2 dt^2 + g_{\alpha\beta} dx^\alpha dx^\beta, \quad \alpha, \beta \in \{1, 2, 3\},$$

where $\{x^\alpha\}$, $\alpha = 1, 2, 3$, is a co-ordinate system on N ;

$g_{\alpha\beta} = g_{\alpha\beta}(t, x^\tau)$ and $V = V(t, x^\tau)$. Suppose g and V restricted to N are respectively $C^{1,1}$ Riemannian metric and $C^{1,1}$ positive function.

The unit time-like vector field u^A on N is given in the above co-ordinate system, by

$$5.10 \quad u^0 = V^{-1} \frac{\partial}{\partial t}; \quad u^\alpha = 0, \quad \alpha = 1, 2, 3.$$

Using the Gauss-Codazzi embedding equations we get

$$5.11 \quad \text{Ric}(g)_{\alpha\beta} = V^{-1} V_{;\alpha\beta} + \text{Ric}({}^4g)_{\alpha\beta} + 2\Omega_{\alpha\sigma}\Omega_{\beta}^{\sigma} - \Omega\Omega_{\alpha\beta} - \frac{\partial\Omega_{\alpha\beta}}{\partial t},$$

$$5.12 \quad V^{-1} \Delta V = \text{Ric}({}^4g)_{00} V^{-2} + g^{\alpha\beta} \frac{\partial \Omega_{\alpha\beta}}{\partial t} - |\Omega|^2 ,$$

where $\Omega_{\alpha\beta} = \frac{1}{2} V^{-1} \frac{\partial g_{\alpha\beta}}{\partial t}$ is the second fundamental form of N in M ,

$\Omega = \Omega_{\alpha\beta} g^{\alpha\beta}$, and $|\Omega|^2 = \Omega_{\alpha\beta} \Omega_{\rho\sigma} g^{\alpha\rho} g^{\beta\sigma}$. As usual Δ and ∇ denote

respectively the Laplacian and covariant differentiation relative to the g metric and the indices are raised by $g^{\alpha\beta}$.

For $p \in N$ there exist $\lambda(p)$ and $\Lambda(p)$ such that

$$5.13 \quad \lambda(p)g(X, X)|_p \leq (\text{Ric}({}^4g)(X, X) + 2\Omega_{\alpha\sigma} \Omega_{\beta}^{\sigma} X^{\alpha} X^{\beta} - \Omega \Omega_{\alpha\beta} X^{\alpha} X^{\beta} - \frac{\partial \Omega_{\alpha\beta}}{\partial t} X^{\alpha} X^{\beta})(p) \\ \leq \Lambda(p)g(X, X)|_p$$

for all vectors X tangent to N at p . We shall take

$$5.14 \quad \Phi_2 = \text{Ric}({}^4g_{00}) V^{-2} + g^{\alpha\beta} \frac{\partial \Omega_{\alpha\beta}}{\partial t} - |\Omega|^2$$

and

$$5.15 \quad \Phi_1 = \lambda .$$

Now we suppose that Φ_1 and Φ_2 satisfies the following conditions:

$$5.16 \quad \text{(i) on } N, \quad \Phi_1 + \Phi_2 \geq 0, \quad \Phi_2 \geq 0 \quad \text{and} \quad C\Phi_1 \geq \Lambda \quad \text{for some} \\ \text{constant } C; \text{ and}$$

(ii) there exists a connected set $Q \subset N$ with

$$\text{ess inf}_K (\Phi_1 + \Phi_2) > 0 \quad \text{for all compact } K \subset Q \quad \text{and}$$

$$\Phi_1 = \Phi_2 = 0 \quad \text{on } N \sim \overline{Q} .$$

Then by virtue of Lemma 5.6 our main Theorem 1.1 applies and gives N to be diffeomorphic to $\mathbb{R}^3$. In particular if N is a Cauchy surface, then by a well known theorem due to Geroch (see Proposition 6.6.8 in [HE]) M will be diffeomorphic to $\mathbb{R}^3 \times \mathbb{R}$.

In case the Cauchy surface N is not *a priori* known to be simply connected, Gannon's Theorem (see Theorems 4.13, 4.13' and the paragraph preceding 4.13) can be used to prove that N is simply connected provided $(M, {}^4g)$ is geodesically complete and satisfies the null convergence condition, and on N , V satisfies the asymptotic condition 4.6.

As an example we now consider the case of a static space-time $(M, {}^4g)$ not necessarily perfect fluid. We assume $(M, {}^4g)$ to be asymptotically Euclidean in the sense of 1.4 and 4.6; and to satisfy

$$5.17 \quad \text{Ric}({}^4g)(K, K) \geq 0$$

for all null vectors K . Hence any $t = \text{constant}$ hyper-surface N is simply connected (by Gannon's Theorem 4.13').

Since the metric is static we have $\Omega_{\alpha\beta} = 0$. Hence 5.11 and 5.12 become

$$\text{Ric}(g)_{\alpha\beta} = V^{-1}V_{;\alpha\beta} + \text{Ric}({}^4g)_{\alpha\beta}$$

and

$$V^{-1}\Delta V = \text{Ric}({}^4g)_{00}V^{-2}.$$

Let

$$\Phi_2 = \text{Ric}({}^4g)_{00}V^{-2}$$

and

$$\Phi_1 = \lambda,$$

where

$$\lambda(p) = \inf \left(\left(\text{Ric}({}^4g)_{\alpha\beta} X^\alpha X^\beta \right)(p) / g(X, X)|_p \right),$$

the infimum is over all vectors X tangent to N at p .

Let n be any unit vector tangent to N . Putting $K^A = u^A + n^A$ in 5.17 we have

$$\text{Ric}(g)_{\alpha\beta} n^\alpha n^\beta + V^{-2} \text{Ric}(g)_{00} \geq 0$$

where we have used the fact $\text{Ric}(g)_{0\beta} = 0$ by Codazzi's equation since N is totally geodesic in M .

Thus $\Phi_1 + \Phi_2 \geq 0$. If we strengthen the assumption 5.17 to

$$5.18 \quad \text{Ric}({}^4g)(W, W) \geq 0$$

for all time-like vectors W we get in particular (taking $W^A = u^A$)

$$\text{Ric}({}^4g)_{00} V^{-2} \geq 0.$$

Let Q be a connected set in N such that on Q , $\text{ess inf}_K (\Phi_1 + \Phi_2) > 0$ for all compact sets $K \subset Q$; and on $N \sim \bar{Q}$, $\text{Ric}({}^4g) \equiv 0$. We shall call Q the "non-vacuum region". Since the conditions 5.16 (i) and (ii) are satisfied, the main Theorem 1.1 applies. Hence we have proved the following theorem (compare with Theorem 4.28).

5.19 THEOREM. *Let $(M, {}^4g)$ be an asymptotically Euclidean (in the sense of 1.4 and 4.6), geodesically complete static space-time satisfying the time-like convergence condition (5.18). If $(M, {}^4g)$ has connected non-vacuum region (defined above) then M is diffeomorphic to $\mathbb{R}^3 \times \mathbb{R}$.*

5.20 Remark. Since for a static space-time N is totally geodesic, it follows from 4.19 that we do not need to assume the asymptotic condition 4.6 to apply Gannon's theorem. Hence condition 4.6 can be dropped from the hypothesis of the above theorem.

APPENDIX A

In this appendix we give the proof of the well-known fact that the "contracted Bianchi identity" for g implies that p is a Lipschitz function on N and almost everywhere

$$(A1) \quad p_{;\beta} = -V^{-1}(\rho+p)V_{;\beta}$$

where $p = (\Phi_2 - \Phi_1)/16\pi$ and $\rho = (\Phi_2 + 3\Phi_1)/16\pi$.

We shall also show (as mentioned in Remark 1.9) that p and ρ are functions of V with

$$\frac{dp}{dV} = -V^{-1}(\rho(V)+p(V))$$

when $V^{-1}(t)$ and $V^{-1}(t) \sim \{x \in N \mid \nabla V(x) = 0\}$ are connected sets for each $t \in \text{Range } V$ and the set $C \equiv \{x \in N \mid \nabla V(x) = 0\}$ has H^3 measure zero. The latter two assumptions are made for the sake of simplicity; it may be they can be dropped.

Our assumptions however cover a much more complicated situation than that described in [KS] where it is assumed that V has only one critical point.

For any $W^{1,p}$ vector field b^β on N having compact support in N we have

$$(A2) \quad \int_N \text{Ric}(g)^\alpha_\beta b^\beta_{;\alpha} = \frac{1}{2} \int_N R b^\beta_{;\beta},$$

where R is the scalar curvature of g .

The above equation is obtained by approximating g with a sequence of smooth metrics in $W^{2,p}$, contracting the contracted Bianchi identities of the smooth metrics with b^β , integrating by parts to get rid of the terms involving the third derivatives of the metric and finally taking the limit.

We note that the integration by parts is possible since the contracted Bianchi identity is in the divergence form. We have

$$(A3) \quad R = \Phi_2 + 3\Phi_1 .$$

Putting 1.2 and (A3) in (A2) we get

$$(A4) \quad \int_N V^{-1} V^{;\alpha}_{\beta} b^{\beta}_{;\alpha} + \Phi_1 b^{\beta}_{;\beta} = \frac{1}{2} \int_N (\Phi_2 + 3\Phi_1) b^{\beta}_{;\beta} .$$

Now V satisfies the following weak form of the commutation rule

$$(A5) \quad - \int_N \Delta V b^{\beta}_{;\beta} + \int_N V^{;\alpha}_{\beta} b^{\beta}_{;\alpha} = - \int_N \text{Ric}(g)_{\alpha\beta} V^{;\alpha} b^{\beta}$$

which can be obtained by using the fact that V is $W^{2,p}$ and the corresponding pointwise relation. Replacing b^{β} in (A5) by $V^{-1} b^{\beta}$ which is also a valid test function we have

$$(A6) \quad - \int_N V^{-1} \Delta V b^{\beta}_{;\beta} + \int_N V^{-2} \Delta V b^{\beta} V_{;\beta} + \int_N V^{-1} V^{;\alpha}_{\beta} b^{\beta}_{;\alpha} - \int_N V^{-2} V^{;\alpha}_{\beta} b^{\beta} V_{;\alpha} \\ = - \int_N \text{Ric}(g)_{\alpha\beta} V^{;\alpha} V^{;\beta} V^{-1} .$$

Hence using 1.3, 1.2 and (A4) in (A6) and simplifying we get

$$\int_N (\Phi_1 - \Phi_2) b^{\beta}_{;\beta} + 2 \int_N V^{-1} (\Phi_1 + \Phi_2) b^{\beta} V_{;\beta} = 0 .$$

Thus $p = (\Phi_2 - \Phi_1)/16\pi$ has bounded weak derivative. Hence p is Lipschitz on N and almost everywhere we have

$$p_{;\beta} = -V^{-1}(\rho + p)V_{;\beta} .$$

Thus we have proved (A1).

Now we can show that p is constant on level sets of V . We have

$$V^{-1}(t) = (V^{-1}(t) \sim C) \cup (V^{-1}(t) \cap C) .$$

By C^1 Sard-type theorem (ref. page 56 in [LS1]) for L^1 -almost every $t \in \mathbb{R}$ we have

$$(A7) \quad H^2(V^{-1}(t) \cap C) = 0$$

and $V^{-1}(t) \sim C$ is a (connected by hypothesis) two dimensional C^1 submanifold. Thus for such t given any $q \in V^{-1}(t) \sim C$ there exists a neighbourhood U of q in N with (V, x^2, x^3) as a co-ordinate system in it. By (A1) on $U \cap (V^{-1}(t) \sim C)$ we have $\partial p / \partial x^a = 0$, $a \in \{2, 3\}$. Hence p is constant on $V^{-1}(t) \sim C$ since by hypothesis it is connected.

Now we consider an arbitrary $t \in \text{Range } V$. We have sequences $\{t_j^+\}$ and $\{t_j^-\}$ with

$$\lim_{j \rightarrow \infty} t_j^\pm = t \quad \text{and} \quad t_j^- < t < t_j^+,$$

where for each $t_j^\pm$, $V^{-1}(t_j^\pm) \sim C$ is a connected 2-dimensional C^1 submanifold. We claim

$$(A8) \quad \partial V^{-1}(t) = \left(\lim_{j \rightarrow \infty} V^{-1}(t_j^+) \sim C \right) \cup \left(\lim_{j \rightarrow \infty} V^{-1}(t_j^-) \sim C \right).$$

Here, for $A_j \subset N$, $\lim_{j \rightarrow \infty} A_j$ means

$$\{x \in N \mid x = \lim x_j \text{ for some sequence } \{x_j\} \text{ with } x_j \in A_j\}.$$

To see this let $x \in \partial V^{-1}(t)$. Then there exists $\sigma > 0$ and $y \in B_\sigma(x)$ with $V(y) = s \neq t$. Without loss of generality we assume $s > t$. Since V is continuous there exists $\varepsilon > 0$ such that for all $w \in B_\varepsilon(y)$, $V(w) \geq (s+t)/2$ and $B_\varepsilon(y) \subset B_\sigma(x)$. We take j large enough so that $t_j^+ < (s+t)/2$. Now there exists a $r_j > 0$ such that $V > t_j^+$ on $B_{r_j}(x)$. We consider the solid cone formed by joining the vertex x with points on $B_\varepsilon(y)$. On each ray from x to $B_\varepsilon(y)$ there exists a point z with $V(z) = t_j^+$. Since $z \notin B_{r_j}(x)$ we have $H^2\left(V^{-1}(t_j^+) \cap B_\sigma(x)\right) > 0$. But $H^2\left(V^{-1}(t_j^+) \cap C\right) = 0$. Hence, since $\sigma > 0$ is arbitrary, x is a limit

point of $V^{-1}(t_j^+) \sim C$. Thus

$$\partial V^{-1}(t) \subset \left(\lim_{j \rightarrow \infty} V^{-1}(t_j^+) \sim C \right) \cup \left(\lim_{j \rightarrow \infty} V^{-1}(t_j^-) \sim C \right).$$

The reverse inclusion is trivial. Hence we have proved (A8).

Let

$$\Gamma^\pm = \lim_{j \rightarrow \infty} V^{-1}(t_j^\pm) \sim C.$$

Then p is constant on Γ^+ and on Γ^- . Also $\partial V^{-1}(t) = \Gamma^+ \cup \Gamma^-$.

Suppose the interior of $V^{-1}(t)$ is nonempty. Then, by (A1), p is constant on each component of the interior of $V^{-1}(t)$. Since the interior of $V^{-1}(t)$ must have closure intersecting either Γ^+ or Γ^- we find that p can have only two values on $V^{-1}(t)$. But by hypothesis $V^{-1}(t)$ is connected. Also p is continuous. Hence p is constant on $V^{-1}(t)$.

For almost all $t \in \mathbb{R}$, p is constant on $V^{-1}(t) \sim C$ because (A1) implies $dp/dV = -V^{-1}(\rho+p)$ in the co-ordinate system (V, x^2, x^3) in some neighbourhood U of $q \in V^{-1}(t) \sim C$. Since, for such t , $H^2(V^{-1}(t) \cap C) = 0$ we can take p to be constant on $V^{-1}(t)$ if necessary by redefining p on a set of measure zero. For arbitrary t if $V^{-1}(t)$ has got nonempty interior then, by 1.3, $p = 3p = \text{constant}$ there. Now we show that the remaining set S has H^3 -measure zero so that p need not be defined on S . Let $T \subset \mathbb{R}$ be the set of all t where (A7) does not hold. Then $L^1(T) = 0$ and $S = \bigcup_{t \in T} V^{-1}(t)$. By the co-area formula and definition of C we have, on any local chart (B_i, φ_i) ,

$$\int_{S \cap B_i} |\nabla V| = \int_{S \cap B_i} |\nabla V| = \int_T H^2(S \cap B_i \cap V^{-1}(t)) dt = 0,$$

since $L^1(T) = 0$. On $S \sim C$, $|\nabla V| > 0$. Hence $H^3(S \sim C) = 0$. But by hypothesis $H^3(C) = 0$. Hence $H^3(S) = 0$. This completes the proof of the second assertion. $\square$

APPENDIX B

The following well-known formula has been used often. Let S be a C^2 surface in (N, g) . Then we have, for any C^2 function ψ on N ,

$$(B1) \quad \Delta\psi = \Delta_S\psi + \psi_{;\rho\epsilon} n^\rho n^\epsilon + H\nabla\psi(n)$$

where as usual Δ and ∇ are respectively the Laplacian and the gradient operator in the g metric, Δ_S is the Laplacian on S in the induced metric on S , n is the unit normal vector on S and H is the mean curvature of S .

To prove (B1) let $h^{\alpha\beta} = g^{\alpha\beta} - \bar{n}^\alpha \bar{n}^\beta$, where $\bar{n}$ is a local extension of n , and let $\parallel$ denote the covariant differentiation on S in the induced metric. Then

$$\begin{aligned} \Delta_S\psi &= \psi_{\parallel\alpha\beta} h^{\alpha\beta} = \left(\psi_{;\rho} h^\rho_\alpha \right)_{\parallel\beta} h^{\alpha\beta} \\ &= \left(\psi_{;\rho} h^\rho_\tau \right)_{;\epsilon} h^{\alpha\beta} h^\tau_\alpha h^\epsilon_\beta \\ &= \psi_{;\rho\epsilon} h^{\rho\epsilon} + \psi_{;\rho} \left(-\bar{n}^\rho_{;\epsilon} n_\tau - \bar{n}_\tau_{;\epsilon} n^\rho \right) h^{\tau\epsilon} \\ &= \psi_{;\rho\epsilon} h^{\rho\epsilon} - \left(\psi_{;\rho} n^\rho \right) \bar{n}_\tau_{;\epsilon} h^{\tau\epsilon} \quad \text{since } n_\tau h^{\tau\epsilon} = 0. \end{aligned}$$

Hence we get $\Delta_S\psi = \Delta\psi - \psi_{;\rho\epsilon} n^\rho n^\epsilon - H\nabla\psi(n)$ which implies (B1). $\square$

APPENDIX C

Here we deduce Robinson's inequality 3.38 on $N \sim \overline{Q}$ (see also [R]).

(Notice that our computation avoids the assumption that V is analytic used in the argument of [R].)

$$(C1) \quad (FV^{-1}w^{;\alpha} + GwV^{;\alpha})_{;\alpha} \geq 0$$

where

$$(C2) \quad F = (cV^2 + \alpha) / (1 - V^2)^3$$

and

$$(C3) \quad G = -2c / (1 - V^2)^3 + 6(cV^2 + \alpha) / (1 - V^2)^4.$$

Here $w = |\nabla V|^2$, $0 < V < 1$, $\Delta V = 0$ (on $N \sim \overline{Q}$) and c, α are constants. We note that $\text{Ric}(g)_{\alpha\beta} = V^{-1}V_{;\alpha\beta}$ on $N \sim \overline{Q}$.

We recall that w is (locally) C^2 on $N \sim \overline{Q}$. Hence

$$\begin{aligned} (FV^{-1}w^{;\alpha} + GwV^{;\alpha})_{;\alpha} &= FV^{-1}\Delta w - FV^{-2}V_{;\alpha}w^{;\alpha} + 2cV_{;\alpha}w^{;\alpha} / (1 - V^2)^3 \\ &\quad + 6(cV^2 + \alpha)V_{;\alpha}w^{;\alpha} / (1 - V^2)^4 + Gw\Delta V + Gw_{;\alpha}V^{;\alpha} \\ &\quad - 12cVw^2 / (1 - V^2)^4 + 48(cV^2 + \alpha)w^2V / (1 - V^2)^5 + 12cVw^2 / (1 - V^2)^4 \\ &= FV^{-1}\Delta w - FV^{-2}V_{;\alpha}w^{;\alpha} + 12Fw_{;\alpha}V^{;\alpha} / (1 - V^2) + 48Fw^2V / (1 - V^2)^2 \end{aligned}$$

where we have used $\Delta V = 0$.

At points where $w \neq 0$ we add and subtract $3F|\nabla w|^2 / 4Vw$ to obtain

$$(C4) \quad (FV^{-1}w^{;\alpha} + GwV^{;\alpha})_{;\alpha} = \frac{1}{4}FV^{-1} \left[4V^{-4}w\Delta w - 4V^{-5}wV_{;\alpha}w^{;\alpha} - 3V^{-4}|\nabla w|^2 \right] + \frac{3}{4}V^{-1}w^{-1}F|\nabla w + 8wVV / (1 - V^2)|^2.$$

Now we show that at points where $w \neq 0$ we have

$$(C5) \quad 4V^{-4}w\Delta w - 4V^{-5}wV_{;\alpha}w^{;\alpha} - 3V^{-4}|\nabla w|^2 = R_{\alpha\beta\gamma}R^{\alpha\beta\gamma} \geq 0$$

where $R_{\alpha\beta\gamma} = 2R_{\alpha[\beta;\gamma]}$. Here and in what follows we have used the notation

$$\text{Ric}(g)_{\alpha\beta} = R_{\alpha\beta} \quad \text{and}$$

$$T_{\alpha\dots[\beta\gamma]\dots\tau} = \frac{1}{2}(T_{\alpha\dots\beta\gamma\dots\tau} - T_{\alpha\dots\gamma\beta\dots\tau}) .$$

Differentiating $R_{\alpha\beta} = V^{-1}V_{;\alpha\beta}$ we obtain that, almost everywhere in $N \sim \overline{Q}$,

$$R_{\alpha\beta\gamma} = 2V^{-1}V_{;\alpha[\beta\gamma]} - 2V^{-2}V_{;[\gamma^V;\beta]\alpha} = V^{-1}R_{\alpha\tau\gamma\beta}V^{;\tau} - 2V^{-2}V_{;[\gamma^V;\beta]\alpha} .$$

Since N is three dimensional and $R = 0$ on $N \sim \overline{Q}$ we get, almost everywhere in $N \sim \overline{Q}$,

$$\begin{aligned} R_{\alpha\beta\gamma} &= 2V^{-1}(g_{\alpha[\gamma^R\beta]\tau} + g_{\tau[\beta^R\gamma]\alpha})V^{;\tau} - 2V^{-2}V_{;[\gamma^V;\beta]\alpha} \\ &= 2V^{-2}\left(g_{\alpha[\gamma^V;\beta]\tau}V^{;\tau} + 2V_{;[\beta^V;\gamma]\alpha}\right) \end{aligned}$$

where we have used $R_{\beta\tau} = V^{-1}V_{;\beta\tau}$ and $V_{;[\gamma^V;\beta]\alpha} = -V_{;[\beta^V;\gamma]\alpha}$. Hence,

using $w_{;\beta} = 2V_{;\beta\tau}V^{;\tau}$ we get, almost everywhere,

$$R_{\alpha\beta\gamma} = V^{-2}(g_{\alpha[\gamma^w;\beta]} + 4V_{;[\beta^V;\gamma]\alpha}) .$$

This implies

$$\begin{aligned} R_{\alpha\beta\gamma}R^{\alpha\beta\gamma} &= V^{-4}\left(g_{\alpha[\gamma^w;\beta]}g^{\alpha[\gamma^w;\beta]} + 8g_{\alpha[\gamma^w;\beta]}V^{;[\beta^V;\gamma]\alpha} + 16V_{;[\beta^V;\gamma]\alpha}V^{;[\beta^V;\gamma]\alpha}\right) \\ &= V^{-4}\left(\delta_{\gamma}^{[\gamma^w;\beta]}w_{;\beta} + 8w_{;\beta}V^{;[\beta^V;\gamma]\gamma} + 16V_{;\beta^V;\gamma\alpha}V^{;[\beta^V;\gamma]\alpha}\right) \\ &= V^{-4}\left(|\nabla w|^2 - 4w_{;\beta}V^{;\gamma}V^{;\beta}_{\gamma} + 8wV_{;\gamma\alpha}V^{;\gamma\alpha} - 8\left(V_{;\beta^V;\beta\alpha}\right)\left(V^{;\gamma}V_{;\gamma\alpha}\right)\right) \end{aligned}$$

where we have used $\Delta V = 0$. Thus almost everywhere on $N \sim \overline{Q}$ we have

$$(C6) \quad R_{\alpha\beta\gamma}R^{\alpha\beta\gamma} = V^{-4}\left(8wV_{;\gamma\alpha}V^{;\gamma\alpha} - 3|\nabla w|^2\right) .$$

Now, as in 3.41, we have $2V_{;\alpha\gamma}V^{;\alpha\gamma} = \Delta w + 2R_{\tau\beta}V^{;\tau}V^{;\beta}$. Hence using

$$2R_{\tau\beta}V^{;\tau} = V^{-1}w_{;\beta} \quad \text{we get}$$

$\Delta w = V^{-1} w_{;\gamma} V^{;\gamma} + 2V_{;\gamma\alpha} V^{;\gamma\alpha}$. Putting this expression in (C6) we get (C5).

From (C4) and (C5) we get

$$(C7) \quad (FV^{-1} w^{;\alpha} + GwV^{;\alpha})_{;\alpha} \geq 0 \quad \text{at points where } w \neq 0.$$

But the left hand side vanishes identically on any open set where $w \equiv 0$.

Hence our claim follows because of the continuity of $(FV^{-1} w^{;\alpha} + GwV^{;\alpha})_{;\alpha}$. $\square$

APPENDIX D

We show that (as claimed in Chapter III, 3.44)

$$(D1) \quad \lim_{R \rightarrow \infty} \int_{S_R} (FV^{-1}g(R_n, \nabla w) + Gwg(R_n, \nabla V)) = -\pi(c+a)/2m .$$

From 3.37 we have in the asymptotic co-ordinate system for some $\beta \in (0, \lambda)$ where λ is as in 1.4,

$$V_{;\mu} = m\dot{x}^\mu/|x|^3 + O(|x|^{-2-\beta}) ,$$

$$w = m^2/|x|^4 + O(|x|^{-4-\beta})$$

and

$$g(R_n, \nabla w) = 2V_{;\mu\tau}V^\mu(R_n^\tau) = -4m^2/|x|^5 + \tilde{\eta}$$

where

$$(D2) \quad \left(r^{-3} \int_{N_{2r} \sim N_r} \tilde{\eta}^2 \right)^{\frac{1}{2}} = O(r^{-5-\beta}) .$$

Also we have

$$(1-V^2)^{-3} = R^3/8m^3 + O(R^2)$$

and

$$(1-V^2)^{-4} = R^4/16m^4 + O(R^2) .$$

Hence

$$\begin{aligned} \lim_{R \rightarrow \infty} \int_{S_R} FV^{-1}g(R_n, \nabla w) \\ &= \lim_{R \rightarrow \infty} \left[4\pi R^2(c+a) (R^3/8m^3) (-4m^2/R^5) + \int_{S_R} \tilde{\eta}(c+a) \times O(R^3) \right] \\ &= -2\pi(c+a)/m , \end{aligned}$$

since by (D2) there exists at least one $R \in (r, 2r)$ such that

$$\int_{S_R} \tilde{\eta} = O(R^{-3-\beta}) .$$

Similarly

$$\begin{aligned}
 \lim_{R \rightarrow \infty} \int_{S_R} G \omega g(R_n, \nabla V) \\
 &= \lim_{R \rightarrow \infty} [4\pi R^2 (6(c+a)R^4/16m^4 + O(R^3)) ((m^2/R^4)(m/R^2) + O(R^{-6-\beta}))] \\
 &= 3\pi(c+a)/2m.
 \end{aligned}$$

Hence (D1) follows. $\square$

APPENDIX E

Proof of Lemma 4.22. We choose the pseudo-orthonormal basis vectors along $\gamma(\omega)$ as follows: at A , $E_4 = W$, E_1 and E_2 are mutually orthogonal unit tangent vectors of A , E_3 is null and ${}^4g(E_1, E_3) = 0 = {}^4g(E_2, E_3)$, ${}^4g(E_3, E_4) = -1$. We then use parallel propagation along the null geodesics with tangent vectors W . We shall first deduce the ("vorticity free") Raychaudhuri equation for null geodesics (ref: [HE], pages 86-88, 100-101).

Let J be the Jacobi field along $\gamma(\omega)$ with initial conditions

$$(E1) \quad \left(\frac{\partial J^n}{\partial \omega} \right) \Big|_{\gamma|_{\omega}} = \chi_{mn} J^m(\gamma(\omega)), \quad m, n \in \{1, 2\},$$

and

$$(E2) \quad J^3(\gamma(0)) = 0 = \left(\frac{\partial J^3}{\partial \omega} \right) \Big|_0,$$

where $\chi_{mn} = {}^{(W)}\chi_{mn}$ (see 4.8 in Chapter IV) is a symmetric matrix. We note that (E2) and the Jacobi equation

$$(E3) \quad \left(\frac{\partial^2 J^A}{\partial \omega^2} \right) \Big|_{\gamma|_{\omega}} = -R^A{}_{4B4} J^B(\gamma(\omega))$$

implies that J^3 vanishes all along γ (note that $R^3{}_{4B4} = 0$). Also for null geodesics we are not interested in the tangential component J^4 , that is, we identify the vector fields differing only in the tangential direction. (Note that J^4 satisfying $(\partial^2 J^4 / \partial \omega^2) \Big|_{\gamma|_{\omega}} = - \sum_{n=1}^2 R^4{}_{4n4} J^n(\gamma(\omega))$ is determined by J^1 and J^2 up to addition of a term linear in ω .)

Thus (E3) becomes

$$(E4) \quad \left. \left(\frac{\partial^2 J^n}{\partial \omega^2} \right) \right|_{\gamma} \Big|_{\omega} = - \sum_{m=1}^2 R_{n4m4} J^m(\gamma(\omega)) , \quad n, m = \{1, 2\} .$$

Now J^n also satisfies the following ordinary differential equation

$$(E5) \quad \left. \left(\frac{\partial J^n}{\partial \omega} \right) \right|_{\gamma} \Big|_{\omega} = \left({}^4\nabla_m W^n \right) J^m(\gamma(\omega)) .$$

Hence $J^n(\gamma(\omega))$ can be expressed in terms of their values at some point say $\gamma(0)$, by (see pages 68-69 in [CL])

$$(E6) \quad J^n(\gamma(\omega)) = A_{nm}(\omega) J^m(\gamma(0))$$

where $A_{nm}(\omega)$ is a smooth 2×2 matrix function of ω and

$$A_{nm}(0) = \delta_{mn} , \quad \frac{dA_{nm}}{d\omega}(0) = \chi_{mn} .$$

A_{nm} will become singular when the first point conjugate to A along γ is reached. Putting (E6) in (E4) we get

$$(E7) \quad \frac{d^2 A_{nm}}{d\omega^2} = - \sum_{l=1}^2 R_{n4l4} A_{lm} .$$

Multiplying (E7) with $(A^{-1})_{ml}$ and using

$$\sum_{n=1}^2 R_{n4n4} = \text{Ric}({}^4g)(W, W)$$

we get (as long as A_{ml} is non-singular)

$$(E8) \quad \sum_{m,n} (A^{-1})_{mn} \frac{d^2 A_{nm}}{d\omega^2} = - \text{Ric}({}^4g)(W, W) .$$

Now we define

$$\theta_{mn} \equiv \sum_l \frac{1}{2} \left[(A^{-1})_{ln} \frac{dA_{ml}}{d\omega} + (A^{-1})_{lm} \frac{dA_{nl}}{d\omega} \right] ,$$

$$\theta = \text{Trace } \theta_{mn} ,$$

$$\sigma_{mn} = \theta_{mn} - \frac{1}{2} \theta \delta_{mn} ,$$

$$2\sigma^2 = \sum_{m,n} \sigma_{mn}^2$$

and

$$w_{mn} = \sum_l \frac{1}{2} \left[(A^{-1})_{ln} \frac{dA_{ml}}{d\omega} - (A^{-1})_{lm} \frac{dA_{nl}}{d\omega} \right].$$

Then

$$\begin{aligned} 2\sigma^2 + \frac{1}{2}\theta^2 &= \sum_{m,n} \left(\frac{1}{2} \sum_l \left[(A^{-1})_{ln} \frac{dA_{ml}}{d\omega} + (A^{-1})_{lm} \frac{dA_{nl}}{d\omega} \right] \right)^2 \\ &= \sum_{m,n} w_{mn}^2 + \sum_{l,m,k,n} (A^{-1})_{ln} \frac{dA_{ml}}{d\omega} (A^{-1})_{km} \frac{dA_{nk}}{d\omega} \\ &= \sum_{m,n} w_{mn}^2 - \sum_{m,n} \frac{d(A^{-1})_{mn}}{d\omega} \frac{dA_{nm}}{d\omega}. \end{aligned}$$

Now we show that $w_{mn} = 0$. We have

$$\begin{aligned} \sum_{m,n} \frac{d}{d\omega} (A_{mp} w_{mn} A_{nk}) &= \frac{1}{2} \sum_m \left[A_{mp} \frac{d^2 A_{mk}}{d\omega^2} - \frac{d^2 A_{mp}}{d\omega^2} A_{mk} \right] \\ &= \frac{1}{2} \sum_{m,l} [-A_{mp} R_{m4l4} A_{lk} + A_{mk} R_{m4l4} A_{lp}] \\ &= 0. \end{aligned}$$

But initially $dA_{nm}/d\omega = \chi_{pn} A_{pm}$ giving $A_{mp} w_{mn} A_{nk} = 0$ initially.

Hence, all along γ , $A_{mp} w_{mn} A_{nk} = 0$. Hence $w_{mn} = 0$ (as long as A_{mp} is nonsingular) so that

$$(E9) \quad 2\sigma^2 + \frac{1}{2}\theta^2 = - \sum_{m,n} \frac{d(A^{-1})_{mn}}{d\omega} \frac{dA_{nm}}{d\omega}.$$

Now from (E8) we get

$$(E10) \quad \frac{d\theta}{d\omega} = -\text{Ric}({}^4g)(W, W) + \sum_{m,n} \frac{d(A^{-1})_{mn}}{d\omega} \frac{dA_{nm}}{d\omega}.$$

Hence using (E9) we get

$$(E11) \quad \frac{d\theta}{d\omega} = -\text{Ric}({}^4g)(W, W) - 2\sigma^2 - \frac{1}{2}\theta^2.$$

(E11) is called the ("vorticity free") Raychaudhuri equation for null geodesics. From (E11) we have, using $\text{Ric}({}^4g)(W, W) \geq -\varepsilon$ for $\omega \in [0, \omega^*]$ where $\omega^* > 4|\theta_0|^{-1}$,

$$\frac{d\theta^{-1}}{d\omega} \geq \frac{1}{2} - \varepsilon\theta^{-2}.$$

Initially $|\theta(\gamma(0))|^{-1} = \theta_0^{-1} < 0$ and also $\varepsilon < \frac{1}{4}\theta_0^2$ for $[0, \omega^*]$.

Hence θ^{-1} will become zero within an affine distance

$$\omega_0 = 2 \left(\left(1 - 2\varepsilon\theta_0^{-2} \right) |\theta_0| \right)^{-1} < \omega^*. \quad \text{But } \theta = (\det A)^{-1} (d/d\omega)(\det A) \text{ where}$$

$(d/d\omega)(\det A)$ is finite by (E7). Hence there will be a point of γ within affine distance $\omega_0 < \omega^*$ conjugate to A along γ . $\square$

REFERENCES

- [A] T. Aubin, *Nonlinear Analysis on Manifolds, Monge-Ampère Equations*, Springer-Verlag, New York, Heidelberg, Berlin, 1982.
- [BE] J. Beem, P. Ehrlich, *Global Lorentzian Geometry* (Pure and Applied Mathematics, 67), Marcel Dekker, New York, 1981.
- [CHS] L. Cafarelli, R. Hardt, L. Simon, "Minimal surfaces with isolated singularities", *Manuscripta Math.* 48, 1984, 1-18.
- [CL] E. Coddington, N. Levinson, *Theory of Ordinary Differential Equations*, McGraw-Hill, New York, 1955.
- [FG] T. Frankel, G. Galloway, "Stable minimal surfaces and spatial topology in general relativity", *Math. Z.* 181, 1982, 395-406.
- [G1] D. Gannon, "Singularities in nonsimply connected space-times", *J. Math. Phys.* 16, 1975, 2364-2367.
- [G2] D. Gannon, "On the topology of space-like hypersurfaces, singularities, and black holes", *Gen. Relativity Gravitation* 7, 1976, 219-232.
- [GT] D. Gilberg, N. Trudinger, *Elliptic Partial Differential Equations of Second Order* (2nd Edition), Springer-Verlag, Berlin, Heidelberg, New York, 1984.
- [H] M. Hirsch, *Differential Topology* (Graduate Texts in Mathematics, 33), Springer-Verlag, New York, Heidelberg, Berlin, 1976.
- [HA] P. Hartman, *Ordinary Differential Equations*, Baltimore, 1973.
- [HE] S. Hawking, G. Ellis, *The Large Scale Structure of Space-time*, Cambridge University Press, Cambridge, 1973.
- [K] H. Künzle, "On the spherical symmetry of a static perfect fluid", *Comm. Math. Phys.* 20, 1971, 85-100.
- [KS] H. Künzle, J. Savage, "A global analysis approach to the general relativistic fluid ball problem", *Gen. Relativity Gravitation* 12, 1980, 155-174.

- [L] L. Lindblom, "Some properties of static general relativistic stellar models", *J. Math. Phys.* **21**, 1980, 1455-1459.
- [LB] L. Lindblom, D. Brill, "Comments on the topology of nonsingular stellar models", *Essays in General Relativity, a Festschrift for Abraham Taub* (F.J. Tipler, ed.), 13-19, Academic Press, New York, San Francisco, London, 1980.
- [LS1] L. Simon, *Lectures on Geometric Measure Theory*, Proceedings of the C.M.A., Australian National University, Canberra, **3**, 1983.
- [LS2] L. Simon, "A Hölder estimate for quasiconformal maps between surfaces in Euclidean space", *Acta Math.* **139**, 1977, 19-51.
- [M] H. Müller zum Hagen, "On the analyticity of static vacuum solutions of Einstein's equations", *Proc. Cambridge Philos. Soc.* **67**, 1970, 415-421.
- [MSY] W. Meeks III, L. Simon, S.T. Yau, "Embedded minimal surfaces, exotic spheres, and manifolds with positive Ricci curvature", *Ann. Math.* **116**, 1982, 621-659.
- [R] D. Robinson, "A simple proof of the generalization of Israel's theorem", *Gen. Relativity Gravitation* **8**, 1977, 695-698.
- [SSY] R. Schoen, L. Simon, S.T. Yau, "Curvature estimates for minimal hypersurfaces", *Acta Math.* **134**, 1975, 276-288.