MULTIPLIER PROBLEMS FOR SPACES
OF CONTINUOUS FUNCTIONS WITH
 p -SUMMABLE TRANSFORMS

by

Lynette Myra Bloom (née Butler)

Except where specifically stated otherwise the results in this
thesis are the product of my own research.

A thesis submitted for the degree of
Doctor of Philosophy
at the
Australian National University
Canberra
February, 1978

ACKNOWLEDGEMENTS

My greatest thanks are due to my supervisor, Professor Robert Edwards, for his patient and expert guidance during the research for this thesis.

I should like to express my gratitude to my husband,

STATEMENT

Dr. Walter Bloom, for his encouragement, and to him, Professor Yitzhak Katznelson and Dr. Richard Loy go my thanks

for me. Except where specifically stated otherwise the results in this thesis are the product of my own research. via Hodgkinson (who typed the bulk of this thesis) and Ms. Helen Carwardine (who typed Chapter 4) for their excellent typing.

L M Bloom

Lynette M. Bloom

CONTENTS

STATEMENT

(i)

ACKNOWLEDGEMENTS

(ii)

CONTENTS

(iii)

ABSTRACT

(iv)

CHAPTER 1 NOTATION AND PRELIMINARIES

1.1 ACKNOWLEDGEMENTS

(v)

1.2 Definitions and notation

(vi)

My greatest thanks are due to my supervisor, Professor

1.3 Multiplier spaces

(vii)

Robert Edwards, for his patient and expert guidance during the research for this thesis.

2.1 Introduction

(viii)

I should like to express my gratitude to my husband,

2.2 A^p is not an algebra for $p \in (1, 2)$

(ix)

Dr. Walter Bloom, for his constant encouragement; and to him,

2.3 Extension of Construction 2.2.1

(x)

Professor Yitzhak Katznelson and Dr. Richard Loy go my thanks

2.4 Construction of $f \in A^p$ with Fourier

(xi)

for many interesting helpful discussions.

2.5 Series divergent at a point

(xii)

Finally, I should like to thank Mrs. Sylvia Hodgkinson (who

2.6 Strict inclusion results for A^p , $p \in (1, 2)$

(xiii)

typed the bulk of this thesis) and Ms. Helen Carwardine (who typed

2.6 Some further results for A^p spaces

(xiv)

Chapter 4) for their excellent typing.

CHAPTER 3 THE FOURIER MULTIPLIER PROBLEM FOR A^p , $p \in (1, 2]$

3.1 Introduction

(xv)

3.2 The multipliers (A^p, A^p) , $p \in (1, 2]$

(xvi)

3.3 Strict inclusion results for (A^p, A^q) , $p \in (1, 2]$

(xvii)

3.4 The multipliers (A^p, A^q) , $p \in (1, 2]$, $q \in [p, 2]$

(xviii)

3.5 The multipliers (A^p, A^q) , $p \in (1, 2]$, $q \in (1, p)$

(xix)

CONTENTS

STATEMENT	(i)
ACKNOWLEDGEMENTS	(ii)
CONTENTS	(iii)
ABSTRACT	1
CHAPTER 1 NOTATION AND PRELIMINARIES	
1.1 General references	2
1.2 Definitions and notation	3
1.3 Multiplier spaces	12
CHAPTER 2 THE SPACES A^p , $p \in [1, \infty)$	
2.1 Introduction	18
2.2 A^p is not an algebra for $p \in (1, 2)$	20
2.3 Extension of Construction 2.2.1	26
2.4 Construction of $f \in A^\#$ with Fourier series divergent at a point	31
2.5 Strict inclusion results for A^p , $p \in [1, 2]$	34
2.6 Some further results for A^p spaces	38
CHAPTER 3 THE FOURIER MULTIPLIER PROBLEM FOR A^p , $p \in [1, 2]$	
3.1 Introduction	45
3.2 The multipliers (A^p, A^p) , $p \in [1, 2]$	46
3.3 Strict inclusion results for (A^p, A^q) , $p \in [1, 2]$	50
3.4 The multipliers (A^p, A^q) , $p \in [1, 2]$, $q \in [p, 2]$	57
3.5 The multipliers (A^p, A^q) , $p \in (1, 2]$, $q \in [1, p)$	58

CHAPTER 4	POINTWISE MULTIPLIERS OF A^p , $p \in [1,2]$	
4.1	Introduction	66
4.2	Some inclusion results for pointwise multipliers	66
4.3	$M(A^p, A^q) = \{0\}$, if $2 \geq p > q \geq 1$	72
4.4	Some sufficient conditions for membership of $M(A^p)$	75
4.5	An alternative approach	87
APPENDIX A	BOUNDS FOR TWO SPECIAL BILINEAR FORMS	
A.1	Introduction	93
A.2	Bounds for Hilbert's B form	94
A.3	Bounds for Stěckin's bilinear form	97
APPENDIX B	SOME ADDITIONAL RESULTS	
B.1	Idempotent Fourier Multipliers from A^p to A^q , $p, q \in [1,2]$	100
B.2	$M_p \cap C = M(A^p)$, $p \in (1,2)$	106
REFERENCES		110
FURTHER REFERENCES		114
NOTES		115
INDEX OF SYMBOLS		116

In Chapter 4 we consider the pointwise multiplier-spaces $M(A^p, A^q)$, written $M(A^p)$ when $q = p$. It is proved there that $M(A^p, A^q) = \{0\}$ when $2 \geq p > q \geq 1$. Here, and in Appendix B, we establish various inclusion results for these multiplier spaces and prove several results involving the translation-continuity of functions in $M(A^p)$. With the aid of the results in Appendix A we obtain sufficient conditions for membership of $M(A^p)$ in such a way as to show that the functions so obtained are translation-continuous. In Appendix B we prove that $M(A^p) = M_p \cap C$, $p \in [1,2]$ and discuss some consequences of this result.

The material of 2.2 and that of 2.5 and Chapter 3 appear in [2] and [4] respectively.

ABSTRACT

The space A^p , $p \in [1, \infty)$, is defined as the space of those functions on the circle group T which are continuous and have p -summable Fourier transforms. Each A^p space will be normed by $N_p: f \mapsto \|f\|_\infty + \|\hat{f}\|_p$, under which it is a Banach space. In this thesis we shall be concerned with these spaces and both their Fourier (or convolution) and pointwise multiplier spaces.

In Chapter 2 we consider the spaces A^p in detail. In particular, we prove that A^p , $p \in (1, 2)$ is not an algebra and establish constructively the strict inclusion results

$$\bigcup_{p \in [1, q)} A^p \subsetneq A^q, \text{ if } q \in (1, 2]$$

and

$$A^q \subsetneq \bigcap_{p \in (q, 2]} A^p, \text{ if } q \in [1, 2).$$

In Appendix B we identify the idempotent elements of (A^p, A^q) .

In Chapter 3 we consider the spaces (A^p, A^q) of Fourier multipliers from A^p to A^q . We identify $M + \mathcal{FL}^{p'}$ as the strong dual of A^p , and prove that $(A^p, A^q) = M + \mathcal{FL}^{p'}$, if $p \in [1, 2]$, $q \in [p, 2]$. In the case $p \in (1, 2]$, $q \in [1, p)$ we give a sufficient and several necessary conditions for membership of (A^p, A^q) . In addition, we establish constructively the strict inclusion results

$$(A^q, A^q) \subsetneq \bigcap_{p \in [1, q)} (A^p, A^p), \text{ if } q \in (1, 2]$$

and

$$\bigcup_{p \in (q, 2]} (A^p, A^p) \subsetneq (A^q, A^q), \text{ if } q \in [1, 2).$$

In Chapter 4 we consider the pointwise multiplier spaces $M(A^p, A^q)$, written $M(A^p)$ when $q = p$. It is proved there that $M(A^p, A^q) = \{0\}$ when $2 \geq p > q \geq 1$. Here, and in Appendix B, we establish various inclusion results for these multiplier spaces and prove several results involving the translation-continuity of functions in $M(A^p)$: With the aid of the results in Appendix A we obtain sufficient conditions for membership of $M(A^p)$ in such a way as to show that the functions so obtained are translation-continuous. In Appendix B we prove that $M(A^p) = M_p \cap C$, $p \in [1, 2]$ and discuss some consequences of this result.

The material of 2.2 and that of 2.5 and Chapter 3 appear in [2] and [4] respectively.

CHAPTER 1

NOTATION AND PRELIMINARIES

1.1 General references

The symbol G will always denote an LCA (locally compact abelian Hausdorff) group and will possess typical elements $a, x, y, \dots$, and identity element 0 . The dual group of G will be denoted by Γ . We shall write λ for a Haar measure on G . When G is compact λ will be normalised to have total variation equal to 1. Where no confusion arises we abbreviate $\int_G f(x) d\lambda(x)$ by $\int_G f(x) dx$.

In what follows we reserve the symbols $\mathbb{R}, \mathbb{C}, \mathbb{Z}$ for the additive groups of the real numbers, the complex numbers, and the integers respectively. We also reserve the symbol $\mathbb{T}$ to denote the circle group, by which we mean here the quotient group $\mathbb{R}/2\pi\mathbb{Z}$. For details of this identification and its implications we refer the reader to either [5] or [12]. For simplicity we shall replace $a \in \mathbb{T}$ with that element x of $[0, 2\pi)$ for which a is the coset $x + 2\pi\mathbb{Z}$.

We shall henceforth be concerned almost exclusively with analysis on the compact group $\mathbb{T}$ and its (discrete) dual group $\mathbb{Z}$; the Haar measure on $\mathbb{T}$ is given by $\frac{1}{2\pi} \mu$, where μ is Lebesgue measure on $\mathbb{R}$ restricted to $[0, 2\pi)$ and the appropriate identification is made as mentioned previously; that on $\mathbb{Z}$ is just the usual counting measure. With these choices of Haar measures Parseval's theorem holds.

For Fourier analysis our main references will be [5] and [23], with our notation in most cases following the former. For general functional analysis our background references will be [6] and [17].

1.2 Definitions and notation

In this section we collect together some of the most frequently used basic definitions and notation ; these will remain standard throughout.

1.2.1 SPACES OF FUNCTIONS, MEASURES AND DISTRIBUTIONS

We denote by $C, C^\infty, \mathcal{D}$ respectively the space of continuous functions, the space of infinitely differentiable functions, and the space of distributions on T . We consider that C is equipped with the uniform topology; C^∞ has the topology defined by the metric $d : (f, g) \rightarrow \|f - g\|_{(\infty)}$, where

$$\|f - g\|_{(\infty)} = \sum_{k=0}^{\infty} 2^{-k} \|f - g\|_{(k)} \{1 + \|f - g\|_{(k)}\}^{-1}$$

with

$$\|f - g\|_{(k)} = \sup_{0 \leq h \leq k} \|f^{(h)} - g^{(h)}\|_{\infty} ;$$

and $\mathcal{D}$ has its usual topology as the strong dual of C^∞ .

When $p \in [1, \infty]$, $L^p(G)$ will denote the normed Lebesgue space of equivalence classes of functions; denote its norm by $\|\cdot\|_p$. As is customary, we write L^p and ℓ^p in the cases $G = \mathbb{T}$ and $G = \mathbb{Z}$ respectively. Also, p' will denote the exponent conjugate to p ; that is $1' = \infty$, $\infty' = 1$ and $\frac{1}{p} + \frac{1}{p'} = 1$ for $p \in (1, \infty)$.

If $f \in L^1(G)$ we write $\hat{f}$ for the Fourier transform of f , defined on Γ by

$$\hat{f}(\gamma) = \int_G f(x) \overline{\gamma(x)} dx.$$

We shall give $\hat{}$ a dual role. If the function g on Γ is a Fourier transform then $\hat{g}$ will denote that function on G whose transform is g ; in other words, the "inverse" transform of g .

The Fourier transform can also be defined for distributions. Letting e_k denote the character of $\mathbb{T}$ given by $e_k(x) = e^{ikx}$, we define the Fourier transform of $\sigma \in \mathcal{D}'$ via the duality relation

$$\hat{\sigma}(n) = \langle \sigma, e_{-n} \rangle.$$

A will denote the subspace of C consisting of those functions f in C for which $\sum_{n \in \mathbb{Z}} |\hat{f}(n)| < \infty$. Under the norm

$$\|\cdot\|_A : f \mapsto \|\hat{f}\|_1, A \text{ is a Banach algebra isometrically isomorphic to}$$

ℓ^1 .

M and PM will denote respectively the space of complex Radon measures and the space of complex pseudomeasures on T , and they will be equipped with their usual topologies as the strong duals of C and A respectively.

For $p \in [1, \infty]$ we use Fl^p to denote the space of those distributions whose Fourier transforms belong to ℓ^p . Each Fl^p is a Banach space under the norm $\sigma \rightarrow \|\hat{\sigma}\|_p$. Note here that we can make the identifications $Fl^1 = A$ and $Fl^\infty = PM$. Also, for $p \in [1, 2]$, the Hausdorff-Young inequality ensures that $Fl^p \subseteq L^{p'}$.

We are now in a position to introduce the spaces A^p with which we shall be primarily concerned throughout this thesis. These spaces are defined by

$$A^p = C \cap Fl^p, \quad p \in [1, \infty).$$

It is not difficult to verify that A^p is a Banach space under the norm N_p , given by

$$N_p : f \rightarrow \|f\|_\infty + \|\hat{f}\|_p;$$

and that we can make the identifications $A^1 = A$ and $A^p = C$ for $p \in [2, \infty)$. Note also that TP , the space of trigonometric polynomials on T , is dense in A^p . Because of its frequent occurrence in what is to follow we shall find it convenient to write $A^\# = \bigcap_{p>1} A^p$. The

space $A^\#$ may be regarded as a Fréchet space with defining seminorms $f \rightarrow N_{p_k}(f)$, where $(p_k)_{k=1}^\infty$ is any sequence satisfying $p_k > 1$ and $p_k \downarrow 1$.

We take this opportunity to note that for $f \in A^p$ and $n \in \mathbb{Z}$,

$$\| (e_n f)^\wedge \|_p = \| \hat{f} \|_p, \quad \text{for all } n \in \mathbb{Z}. \quad (1.2.1)$$

and

$$N_p(e_n f) = N_p(f). \quad (1.2.2)$$

Also, ξ_E will denote the characteristic function of the set E .

1.2.2 TRANSLATIONS AND REFLECTIONS

For $\alpha \in G$ and a function f on G we define the α -translate $\tau_\alpha f$ of f by

$$\tau_\alpha f : x \mapsto f(x - \alpha), \quad \text{for all } x \in G;$$

and the reflection $\check{f}$ of f by

$$\check{f} : x \mapsto f(-x), \quad \text{for all } x \in G.$$

For $\sigma \in \mathcal{D}$ and $\alpha \in \mathbb{T}$ the corresponding definitions for $\tau_\alpha \sigma$ and $\check{\sigma}$ are given by the duality relations

$$\langle \tau_\alpha \sigma, g \rangle = \langle \sigma, \tau_\alpha g \rangle$$

and

$$\langle \check{\sigma}, g \rangle = \langle \sigma, \check{g} \rangle,$$

1.2.4 CONVOLUTION

where these are to hold for all $g \in C^\infty$.

A normed space S of distributions will be called translation - continuous if $\lim_{\alpha \rightarrow 0} \|\tau_\alpha \sigma - \sigma\| = 0$, for all $\sigma \in S$.

1.2.3 LIPSCHITZ SPACES

Suppose $\alpha \in (0, 1]$. Then we define the following Lipschitz spaces:

$$\Lambda_\alpha = \{f \in C : \sup_{\alpha \neq 0} \frac{\|\tau_\alpha f - f\|_\infty}{|\alpha|^\alpha} < \infty\} \quad (1.2.3)$$

and

$$\Lambda_\alpha^1 = \{f \in L^1 : \sup_{\alpha \neq 0} \frac{\|\tau_\alpha f - f\|_1}{|\alpha|^\alpha} < \infty\}. \quad (1.2.4)$$

The spaces Λ_α and $C \cap \Lambda_\alpha^1$ are Banach algebras under the topologies defined by the respective norms

$$\|\cdot\|_{\Lambda_\alpha} : f \mapsto \|f\|_\infty + \sup_{\alpha \neq 0} \frac{\|\tau_\alpha f - f\|_\infty}{|\alpha|^\alpha}$$

and

$$\|\cdot\|_{\Lambda_\alpha^1} : f \mapsto \|f\|_\infty + \sup_{\alpha \neq 0} \frac{\|\tau_\alpha f - f\|_1}{|\alpha|^\alpha}.$$

It is worth noting here that TP is not a dense subspace of either Λ_α or $C \cap \Lambda_\alpha^1$; and hence that neither Λ_α nor $C \cap \Lambda_\alpha^1$ is translation-continuous.

1.2.4 CONVOLUTION

For measurable functions f, g on G their convolution $f * g$ is given by

$$f * g = \int_G \tau_y f \cdot g(y) dy.$$

Among the properties of convolution it is worth particular note here that, if either of f, g is in C or C^∞ , then so is $f * g$. For further details the reader is referred to [5].

We shall need the notion of convolution also for distributions. We shall introduce this, as in [5], 12.6, by first defining $\sigma * u \in C^\infty$ for $\sigma \in \mathcal{D}$, $u \in C^\infty$ and using this to define $\sigma * \mu$ for $\sigma, \mu \in \mathcal{D}$. To this end

$$\sigma * u(x) = \langle \sigma, \tau_x^\vee u \rangle;$$

and then $\sigma * \mu \in \mathcal{D}$ is given by

$$\langle \sigma * \mu, u \rangle = \langle \sigma, (\mu * u)^\vee \rangle, \text{ for all } u \in C^\infty.$$

1.2.5 DIRICHLET AND FEJÉR KERNELS

For $f \in L^1$ the trigonometric series $\sum_{n \in \mathbb{Z}} \hat{f}(n) e_n$ is called

the Fourier series of f . Then, for any non-negative integer N , the N -th symmetric partial sum of the Fourier series of f is

$$s_N f = \sum_{|n| \leq N} \hat{f}(n) e_n. \quad (1.2.5)$$

This N -th partial sum can be expressed as the convolution

$s_N f = D_N * f$, where

$$D_N(x) = \sum_{|n| \leq N} e^{inx} = \begin{cases} \sin(N + \frac{1}{2})x / \sin \frac{1}{2}x, & \text{if } x \not\equiv 0 \pmod{2\pi}, \\ 2N + 1, & \text{otherwise.} \end{cases}$$

The sequence $(D_N)_{N=0}^{\infty}$ is called the Dirichlet kernel; although we shall sometimes refer to D_N itself as the Dirichlet kernel of order N .

The N -th Cesàro sum $\sigma_N f$ of the Fourier series of f is given by

$$\sigma_N f = (N + 1)^{-1} \sum_{n=0}^N s_n f.$$

This Cesàro sum can be expressed as the convolution $\sigma_N f = F_N * f$, where

$$\begin{aligned} F_N(x) &= (N + 1)^{-1} \sum_{n=0}^N D_n(x) \\ &= \sum_{|n| \leq N} \left(1 - \frac{|n|}{N + 1}\right) e^{inx} \end{aligned}$$

$$= \begin{cases} [\sin \frac{1}{2} (N+1)x / \sin \frac{1}{2} x]^2 (N+1)^{-1}, & x \not\equiv 0 \pmod{2\pi}, \\ N+1, & \text{otherwise,} \end{cases} \quad (1.2.9)$$

constitute the Fejér kernels. Remarks on nomenclature similar to those made for the Dirichlet kernel will also apply to the Fejér kernel.

There are many "listed" properties both for the Dirichlet and the Fejér kernel. We shall conclude this section by obtaining some norm estimates related to the Dirichlet kernel, but first we shall introduce some more notation. If ϕ and ψ are positive functions on $\{1, 2, \dots\}$ we write $\phi \sim \psi$ if and only if $0 < \inf \psi \phi^{-1} \leq \sup \psi \phi^{-1} < \infty$. We now have the following results.

1.2.6 LEMMA. Suppose $p \in (1, 2)$. Then

$$\|D_N\|_{\infty} = 2N + 1 \sim N, \quad (1.2.6)$$

$$\|\hat{D}_N\|_p = (2N + 1)^{\frac{1}{p}} \sim N^{\frac{1}{p}} \quad (1.2.7)$$

and

$$\|(D_N^2)^{\wedge}\|_p \sim N^{1+\frac{1}{p}}. \quad (1.2.8)$$

Proof. Both (1.2.6) and (1.2.7) are immediately apparent from the definition of D_N . We shall prove (1.2.8) here.

$$\|(D_N^2)^{\wedge}\|_p^p = \|\hat{D}_N * \hat{D}_N\|_p^p$$

$$= \sum_{|n| \leq 2N+1} \left| 2N + 1 - |n| \right|^p$$

$$= (2N + 1)^p + 2 \sum_{n=1}^{2N} n^p . \quad (1.2.9)$$

Making use in (1.2.9) of the result

$$(p + 1)^{-1} (k - 1)^{p+1} \leq \sum_{n=1}^k n^p \leq (p + 1)^{-1} k^{p+1} , \quad (1.2.11)$$

we obtain

$$2(p + 1)^{-1} (2N - 1)^{p+1} \leq \| (D_N^2)^{\wedge} \|_p^p \leq 2(p + 2)(p + 1)^{-1} (2N + 1)^{p+1} ,$$

from which (1.2.8) follows. //

1.2.7 RUDIN-SHAPIRO POLYNOMIALS

The Rudin-Shapiro polynomials were first introduced for T (see [29]) and were then defined for all infinite *LCA* groups (see [16], (37.19) (6)). Recently Price and McMullen in [25] have introduced analogous Rudin-Shapiro sequences for infinite compact non-abelian groups.

We shall be concerned here with the Rudin-Shapiro polynomials on T . These will prove to be a key tool for some of the work of Chapter 2. In [23], p.33 the Rudin-Shapiro polynomials P_m , $m = 0, 1, 2, \dots$, are defined by

$$P_m = \sum_{n=0}^{2^m-1} \varepsilon_m(n) e_n ,$$

where $\varepsilon_m(n) \in \{-1, 1\}$ are chosen in such a way that

$$\|P_m\|_\infty \leq 2^{\frac{1}{2}(m+1)} \quad \text{and} \quad \|\hat{P}_m\|_p = 2^{m/p}. \quad (1.2.10)$$

We define here $P_v^* = P_m$ for the unique m for which $2^m \leq v < 2^{m+1}$.

Then

$$\|P_v^*\|_\infty \leq 2^{\frac{1}{2}(m+1)} \leq v^{\frac{1}{2}}/2 \quad (1.2.11)$$

and

$$v^{\frac{1}{p}} \geq \|\hat{P}_v^*\|_p = 2^{\frac{m}{p}} > v^{\frac{1}{p}} 2^{-\frac{1}{p}}. \quad (1.2.12)$$

1.3 Multiplier spaces

In the following chapters we shall consider two different types of multipliers, namely Fourier (or convolution) multipliers and point-wise multipliers.

1.3.1 FOURIER MULTIPLIER SPACES

Suppose here that E and F denote Banach or Fréchet spaces of distributions on $\mathbb{T}$ which are invariant under translations. By a Fourier multiplier from E to F we mean a continuous linear operator which commutes with translations. The problem of identifying the Fourier multipliers for particular spaces E and F began with Fekete in [11] and has been considered since by many others. The interested reader is referred here to [24] and to [5], Chapter 16 and the many references given in each of these. In [19], Hörmander showed that such operators can be written as convolution with an

appropriate distribution. We shall denote the space of such distributions by (E, F) and refer to $\phi \in (E, F)$ as a Fourier multiplier from E to F . We shall of course be primarily concerned with the cases where E and F (not necessarily equal) are A^p spaces for some $p \in [1, 2]$.

We denote by M_p the space of all $f \in L^\infty$ for which $\hat{f} * g \in \ell^p$ whenever $g \in \ell^p$; for $f \in M_p$ we express its norm $\|f\|_{p,p}$ by

$$\|f\|_{p,p} = \sup\{\|\hat{f} * g\|_p : g \in \ell^p \text{ and } \|g\|_p \leq 1\}.$$

M_p here is the special case (for $G = \mathbb{Z}$) of the multiplier space $M_p(\Gamma)$ consisting of those $f \in L^\infty(\Gamma)$ for which $T_f g \in L^p(G)$ whenever $g \in L^2(G) \cap L^p(G)$, where $(T_f g)^\wedge = f \cdot \hat{g}$. In the general case

$$\|f\|_{p,p} = \sup\{\|T_f g\|_p : g \in L^p(G) \cap L^2(G) \text{ and } \|g\|_p \leq 1\}.$$

By applying certain results from the (at present unpublished) book [9] we obtain sufficient conditions for certain functions to belong to $M_p(\Gamma)$. For the sake of completeness we include here (without proof) the results required, in a form more suited to this thesis; see Theorem 1.3.4 and Corollary 1.3.5 below. We need first the following definitions.

1.3.2 DEFINITION. $(U_k)_{k \in \mathbb{Z}}$ is a covering family in (or for) the LCA group G if and only if the following conditions are satisfied:

(i) $(U_k)_{k \in \mathbb{Z}}$ is a base of relatively compact measurable neighbourhoods of 0 in G such that $\bigcup_{k \in \mathbb{Z}} U_k = G$ and, for $k_1 \leq k_2$, $U_{k_1} \supseteq U_{k_2}$.

(ii) there exists a function $\theta : \mathbb{Z} \rightarrow \mathbb{Z}$ and a positive number b such that, for every $k \in \mathbb{Z}$,

$$\theta(k) < k,$$

$$U_k - U_k \subseteq U_{\theta(k)},$$

and

$$\lambda(U_{\theta(k)}) \leq b \lambda(U_k).$$

1.3.3 DEFINITION. Suppose $f \in L^1(G)$. Then we write $J(f) = \sup_{\beta} J_{\beta}(f)$, where

$$J_{\beta}(f) = \sup_{y \in U_{\beta}} \int_{G \setminus U_{\theta(\beta)}} |f(x-y) - f(x)| dx.$$

1.3.4 THEOREM. Suppose $f \in L^1(G)$. Then, for $p \in (1, \infty)$,

$$\|\hat{f}\|_{p,p} \leq b^{2/p^*} B_p \max(J(f), \|\hat{f}\|_{\infty}),$$

where $p^* = \min(p, p')$ and

$$B_p = \begin{cases} [2p(16 - 7p)(p - 1)^{-1}(2 - p)^{-1}]^{1/p}, & p \in (1, 2) \\ 1, & p = 2 \\ B_{p'}, & p \in (2, \infty) \end{cases} \quad (1.3.1)$$

. //

1.3.5 COROLLARY. Suppose $\phi \in L^\infty(\Gamma)$ and there exists a sequence $(f_\nu)_{\nu=1}^\infty$ of integrable functions on G such that

$$\kappa = \sup_\nu \max \{J(f_\nu), \|\hat{f}_\nu\|_\infty\} < \infty$$

1.3.8 LEMMA. Suppose $p, q \in [1, 2]$. Then

and

$$\phi = \lim_{\nu \rightarrow \infty} \hat{f}_\nu \quad \text{l.a.e. on } \Gamma.$$

Then, for $p \in (1, \infty)$, $\phi \in M_p(\Gamma)$ and

$$\|\phi\|_{p,p} \leq b^{2/p^*} B_p \kappa. \quad (1.3.2)$$

1.3.6 REMARK. Note here that in the case $G = \mathbb{Z}$ an appropriate covering family is formed by the neighbourhoods U_k , where

$$U_k = \begin{cases} \{0\} & \text{if } k \in \mathbb{Z} \text{ and } k > 0, \\ \{n \in \mathbb{Z} : |n| < 2^{-k}\} & \text{if } k \in \mathbb{Z} \text{ and } k \leq 0. \end{cases}$$

In this case we can take $\theta(k) = k - 1$ and $b = 3$.

1.3.7 POINTWISE MULTIPLIER SPACES

We shall suppose here that E and F are subspaces of L^1 . Then we say that $f \in L^1$ is a pointwise multiplier from E to F whenever $f \cdot E \subseteq F$, where by $f \cdot E$ we mean $\{fg : g \in E\}$. We denote by $M(E, F)$ the set of all pointwise multipliers from E to F . When $E = F$ we abbreviate $M(E, F)$ to $M(E)$ and speak of the

pointwise multipliers of E . Again we shall be primarily concerned with the cases when E and F (not necessarily equal) are A^p spaces for some $p \in [1, 2]$. Before proceeding to the main body of the thesis we shall prove the following result.

1.3.8 LEMMA. Suppose $p, q \in [1, 2]$. Then

$$f \cdot A^p \subseteq A^q \text{ if and only if } \|f\|_{M(p,q)} < \infty, \quad (1.3.1)$$

which, combined with (1.3.5), gives us that $h = fg$. Thus, by the closed graph theorem, V_f is continuous; hence

$$\|f\|_{M(p,q)} = \sup\{N_q(fg) : g \in TP \text{ and } N_p(g) \leq 1\}. \quad (1.3.2)$$

Proof. (i) "only if". Suppose $f \cdot A^p \subseteq A^q$. Since $1 \in A^p$ it follows immediately that $f \in C$. We may identify f with the linear operator $V_f : A^p \rightarrow A^q$ defined by

$$V_f : g \rightarrow fg, \text{ for all } g \in A^p.$$

Also, for each k ,

Now consider a sequence $(g_k)_{k=1}^\infty$ from A^p for which there are functions $g \in A^p$ and $h \in A^q$ satisfying

$$\lim_{k \rightarrow \infty} N_p(g_k - g) = 0 \text{ and } \lim_{k \rightarrow \infty} N_q(fg_k - h) = 0. \quad (1.3.3)$$

Since $\|\cdot\|_\infty \leq N_r(\cdot)$ for all $r \in [1, 2]$, it follows from (1.3.3) that

follows by use of Fatou's lemma (see [34], p.26) that $(fg) \in A^q$ and hence, since $fg \in C$, that $fg \in A^q$. //

$$\lim_{k \rightarrow \infty} \|g_k - g\|_{\infty} = 0 \quad (1.3.4)$$

and

$$\lim_{k \rightarrow \infty} \|fg_k - h\|_{\infty} = 0 \quad (1.3.5)$$

But $f \in C$ so that (1.3.4) implies that $\lim_{k \rightarrow \infty} \|fg_k - fg\|_{\infty} = 0$

which, combined with (1.3.5), gives us that $h = fg$. Thus, by the closed graph theorem, V_f is continuous; hence

$$\|f\|_{M(p,q)} \leq \|V_f\| = \sup\{N_q(fg) : g \in A^p \text{ and } N_p(g) \leq 1\} < \infty.$$

(ii) "if". Suppose $\|f\|_{M(p,q)} < \infty$. Then it follows immediately that $f \in A^q$. Now consider $g \in A^p$. We can, without loss of generality, consider $g \neq 0$. Then, since TP is dense in A^p , there exists a sequence $(g_k)_{k=1}^{\infty}$ from TP satisfying $\lim_{k \rightarrow \infty} N_p(g - g_k) = 0$.

Also, for each k ,

$$N_q(fg_k) \leq \|f\|_{M(p,q)} N_p(g_k). \quad (1.3.6)$$

Since $\lim_{k \rightarrow \infty} N_p(g_k) = N_p(g)$, (1.3.6) ensures that $(fg_k)_{k=1}^{\infty}$ is norm

bounded in A^q . Since also $\lim_{k \rightarrow \infty} (fg_k)^{\wedge} = (fg)^{\wedge}$ pointwise on Z it

follows by use of Fatou's lemma (see [34], p.26) that $(fg)^{\wedge} \in \ell^q$;

and hence, since $fg \in C$, that $fg \in A^q$. //

CHAPTER 2

THE SPACES A^p , $p \in [1, \infty)$

2.1 Introduction

These spaces, defined previously in 1.2.1, are very interesting in their own right. In addition, they have a connection (which we shall not exploit here) with p -Sidon sets (see [10] and [21]) and p -Helson sets (see [13]).

As already noted, $A^1 = A$ and $A^p = \mathbb{C}$ for $p \in [2, \infty)$ so that A^p is an algebra whenever $p = 1$ or $p \in [2, \infty)$. This leads one to ask whether or not A^p is an algebra when $p \in (1, 2)$. In this context see [7], p.183. In private correspondence with us, Professor Y. Katznelson suggested in outline an *existential* proof that for $p \in (1, 2)$, A^p is not an algebra. We have concentrated here on a *constructive* approach, the details of which will be given in 2.2 and may be found also in [4]. In this construction use is made of the Dirichlet kernel, the Rudin-Shapiro polynomials and the various estimates for these obtained in 1.2.6 and 1.2.7 respectively.

In 2.4 we give a construction to show that the spaces A^p , $p \in (1, 2)$ are not among those subspaces of L^∞ which were shown not to be algebras by Leon Brown in [3]. In 2.5 we again make use of the Rudin-Shapiro polynomials to prove two strict inclusion results for the spaces A^p , $p \in [1, 2]$. We then, in 2.6, give some miscellaneous results concerning A^p , $p \in (1, 2)$. These include a sufficient condition for both f and f^{-1} to belong to A^p and a proof that, for $p \in [1, 2]$, A^p is not reflexive.

In Construction 2.2 we need the following (known) result. The proof given here involves the notion of spectrum which, for $f \in L^1$, is defined by

$$sp(f) = \{n \in \mathbb{Z} : \hat{f}(n) \neq 0\}.$$

2.1.1 LEMMA. Let s be a positive integer, ϕ a trigonometric polynomial of degree less than s , and ψ any trigonometric polynomial. Write $\psi_{(2s)}$ for the function $x \rightarrow \psi(2sx)$ on $\mathbb{T}$. Then

$$\|(\phi \psi_{(2s)})^\wedge\|_p = \|\hat{\phi}\|_p \|\psi_{(2s)}^\wedge\|_p.$$

Proof. We can write

$$\theta = \phi \psi_{(2s)} = \sum_{m \in \mathbb{Z}} \hat{\psi}(m) e_{2sm} \phi, \quad (2.2.1)$$

a finite sum. Also, $e_{2sm} \phi$ and $e_{2sm'} \phi$ have disjoint spectra whenever $m \neq m'$. Thus, by (1.2.1),

$$\|\hat{\theta}\|_p^p = \sum_{m \in \mathbb{Z}} |\hat{\psi}(m)|^p \|(e_{2sm} \phi)^\wedge\|_p^p \quad (2.2.2)$$

$$= \sum_{m \in \mathbb{Z}} |\hat{\psi}(m)|^p \|\hat{\phi}\|_p^p$$

$$= \|\hat{\phi}\|_p^p \|\hat{\psi}\|_p^p.$$

2.2 A^p is not an algebra for $p \in (1,2)$.

Given $q_0 \in [1,2)$, functions $f = f_{q_0} \in A^\#$ and $g_q = g_{q,q_0} \in A^q$ are constructed such that $fg_q \notin A^q$ for every $q \in [1, q_0]$. In particular, if $p \in (1,2)$, A^p is not an algebra.

Throughout the construction we assume that $p \in (1,2)$, $q \in [1,2)$, and $a_1, a_2, \dots$ will denote positive constants.

2.2.1 CONSTRUCTION. Let $f_{n,p} = D_n^{P^*} m_1(2s_1)$ and $g_{n,p,q} = D_n^{P^*} m_2(2s_2)$, where

$$m_1 = \left[n^{\frac{2(p-1)}{2-p}} \right], \quad m_2 = \left[n^{\frac{2(q-1)}{2-q}} \right], \quad s_1 = 4n \quad \text{and} \quad s_2 = 4n + 16nm_1.$$

Since $s_1 > n$, Lemma 2.1.1, together with Lemma 1.2.6, (1.2.11) and (1.2.12), shows that

$$N_p(f_{n,p}) \leq a_1 \left(nm_1^{\frac{1}{2}} + n^{\frac{1}{p}} m_1^{\frac{1}{p}} \right) \leq a_2 n^{\frac{1}{(2-p)}}. \quad (2.2.1)$$

Similarly, since $s_2 > n$,

$$N_q(g_{n,p,q}) \leq a_3 n^{\frac{1}{(2-q)}}. \quad (2.2.2)$$

Again, since $s_2 > 2n + 2s_1m_1$ and $s_1 > 2n$, we can use Lemma 2.1.1 to obtain

$$\begin{aligned} \left\| \left(f_{n,p} g_{n,p,q} \right)^\wedge \right\|_q &= \left\| \left(D_n^2 P_{m_1}^*(2s_1) P_{m_2}^*(2s_2) \right)^\wedge \right\|_q \\ &= \left\| (D_n^2)^\wedge \right\|_q \left\| \left(P_{m_1}^*(2s_1) \right)^\wedge \right\|_q \left\| \left(P_{m_2}^*(2s_2) \right)^\wedge \right\|_q. \end{aligned}$$

It now follows that

$$\| \left\{ f_{n,p} g_{n,p,q} \right\}^{\wedge} \|_q \geq a_4 n^{\kappa}, \quad (2.2.3)$$

where

$$\kappa = \frac{8q - 4 - 2q^2 + pq^2 - 3pq + 4p}{q(2 - p)(2 - q)}$$

From (2.2.1), (2.2.2) and (2.2.3),

$$\frac{\| \left\{ f_{n,p} g_{n,p,q} \right\}^{\wedge} \|_q}{N_p(f_{n,p}) N_q(g_{n,p,q})} \geq a_5 n^{\sigma}, \quad (2.2.4)$$

where

$$\sigma = \frac{(p - 1)(2 - q)}{q(2 - p)}$$

Then (2.2.4) gives
We next estimate the size of $sp\left\{f_{n,p}\right\}$ and $sp\left\{g_{n,p,q}\right\}$:

$$sp\left\{f_{n,p}\right\} \subseteq [-n, n + 2s_1m_1], \quad (2.2.6)$$

$$sp\left\{g_{n,p,q}\right\} \subseteq [-n, n + 2s_2m_2].$$

Thus, according to the choice of s_1 and s_2 specified above,

$$sp\left\{f_{n,p}\right\} \cup sp\left\{g_{n,p,q}\right\} \subseteq [-n, n(1 + 8n^p + 32n^{\tau})], \quad (2.2.5)$$

where

$$\rho = 2(q - 1)/(2 - q) \quad \text{and} \quad \tau = \{6(p+q) - 4(pq+2)\}/(2 - q)(2 - p) .$$

Now define

$$f_n^q = \frac{n^{-2} f_{k(n), p(n)}}{N_{p(n)} \left[f_{k(n), p(n)} \right]} ,$$

and

$$g_{n,q}^{\circ} = \frac{\beta(n,q) g_{k(n), p(n), q}}{N_q \left[g_{k(n), p(n), q} \right]} ,$$

where

$$k(n) = 2^{n^2} , \quad p(n) = 1 + \frac{1}{2n} \quad \text{and} \quad \beta(n,q) = 2^{-\frac{1}{2}(2-q)n/2q} .$$

Then (2.2.4) gives

$$\left\| \left(f_n^{\circ} \ g_{n,q}^{\circ} \right)^{\wedge} \right\|_q \geq \alpha_5 n^{-2} 2^{\psi} , \quad (2.2.6)$$

where

$$\psi = \frac{(2n + 1)(2 - q)}{4q(2 - 1/n)}$$

$$\text{and so} \quad \left\| \left(f_n^{\circ} \ g_{n,q}^{\circ} \right)^{\wedge} \right\|_q \rightarrow \infty \quad \text{as} \quad n \rightarrow \infty .$$

For each $f \in \mathcal{C}$ the function $p \rightarrow N_p(f)$ is decreasing for $p \geq 1$, so that the formula

$$N_p(f_n^\circ) = \frac{n^{-2} N_p(f_{k(n), p(n)})}{N_{p(n)}(f_{k(n), p(n)})}$$

shows that to any $p > 1$ corresponds an integer $n_0(p)$ such that

$$N_p(f_n^\circ) \leq n^{-2}, \text{ for all } n \geq n_0(p). \quad (2.2.7)$$

Also

$$N_q(g_{n,q}^\circ) = \beta(n,q). \quad (2.2.8)$$

Finally, let $f_n = e_{v_n} f_n^\circ$ and $g_{n,q} = g_{n,q}^\circ$, where the integers $v_n = v_{n,q_0}$ will be chosen appropriately to ensure that the f_n have pairwise disjoint spectra, and consider

$$f = f_{q_0} = \sum_{n=1}^{\infty} f_n \text{ and } g_p = g_{q,q_0} = \sum_{n=1}^{\infty} g_{n,q}. \quad (2.2.9)$$

By (1.2.2) and (2.2.7)

$$\begin{aligned} N_p(f) &\leq \sum_{n=1}^{\infty} N_p(f_n) \\ &= \sum_{n=1}^{\infty} N_p(f_n^\circ) \\ &\leq \sum_{n=1}^{n_0(p)} N_p(f_n^\circ) + \sum_{n > n_0(p)} n^{-2} \\ &< \infty \text{ for all } p > 1, \end{aligned}$$

and so $f \in A^\#$. Similarly, by (1.2.2) and (2.2.8),

$$N_q(g_q) \leq \sum_{n=1}^{\infty} N_q(g_{n,q})$$

Moreover (2.2.5) and (2.2.11) show that S_{n,q_0} is finite.

Now suppose $\{v_{n,q}\}_{n=1}^{\infty}$ is chosen to satisfy

$$= \sum_{n=1}^{\infty} N_q(g_{n,q}^\circ)$$

$$\leq \sum_{n=1}^{\infty} \beta(n,q)$$

$$< \infty$$

whenever $(r,k) \neq (n,m)$. (2.2.12)

Then, for every $q \leq q_0$ and every $n = 1, 2, 3, \dots$, $\{f g_q\}$ and

so that $g_q \in A^q$ for $q \in [1, 2)$. the latter, and (2.2.10) shows that

By (2.2.9),

$$\begin{aligned} f g_q &= \sum_{r,k=1}^{\infty} f_r g_{k,q} \\ &= \sum_{m=1}^{\infty} f_m g_{m,q} + \sum_{(r,k) \neq (m,m)} f_r g_{k,q}. \end{aligned} \quad (2.2.10)$$

Let

$$S_{n,q_0} = \cup \{sp(f_n^\circ) \cup sp(g_{n,q}^\circ) : q \leq q_0\} \quad (2.2.11)$$

Reverting to (2.2.12) it is not difficult to check that (omitting

explicit reference to q) it suffices to choose $v_i \in \mathbb{Z}$ freely, and to make a choice of $v_n, n = 2, 3, \dots$, by recursion to satisfy

$$= \cup \{sp(f_{k(n),p(n)}) \cup sp(g_{k(n),p(n),q}) : q \leq q_0\}.$$

Then, for $q \leq q_0$,

$$sp(f_n) \cup sp(g_{n,q}) \subseteq v_{n,q_0} + S_{n,q_0},$$

and

$$sp(f_r g_{k,q}) \subseteq v_{r,q_0} + S_{r,q_0} + v_{k,q_0} + S_{k,q_0}.$$

Moreover (2.2.5) and (2.2.11) show that S_{n,q_0} is finite.

Now suppose $\left\{v_{n,q_0}\right\}_{n=1}^{\infty}$ to be chosen to satisfy

$$\left(2 v_{m,q_0} + S_{m,q_0} + S_{m,q_0}\right) \cap \left(v_{r,q_0} + S_{r,q_0} + v_{k,q_0} + S_{k,q_0}\right) = \emptyset$$

$$\text{whenever } (r,k) \neq (m,m). \quad (2.2.12)$$

Then, for every $q \leq q_0$ and every $m = 1, 2, 3, \dots$, $(fg_q)^\wedge$ and $(f_m g_{m,q})^\wedge$ agree on the support of the latter, and (2.2.10) shows that

$$\|(fg_q)^\wedge\|_q^q \geq \|(f_m g_{m,q})^\wedge\|_q^q$$

$$= \|(f_m^\circ g_{m,q}^\circ)^\wedge\|_q^q,$$

the last step by (1.2.1). Hence, by (2.2.6),

$$\|(fg_q)^\wedge\|_q^q = \infty, \text{ for all } q \in [1, q_0].$$

Reverting to (2.2.12) it is not difficult to check that (omitting explicit reference to q_0) it suffices to choose $v_1 \in Z$ freely, and to make a choice of $v_n, n = 2, 3, \dots$, by recurrence to satisfy

$$v_{n+1} \in Z \setminus S'_n(v_1, \dots, v_n; q_0)$$

and

$$2v_{n+1} \in Z \setminus S''_n(v_1, \dots, v_n; q_0),$$

where

$$S'_n(u_1, \dots, u_n; q_0) = \bigcup_{i \leq n} \{u_i + S_{i, q_0} + S_{n+1, q_0} - S_{n+1, q_0} - S_{n+1, q_0}\} \\ \bigcup_{\substack{i \leq n \\ j \leq n}} \{2u_i - u_j + S_{n+1, q_0} + S_{n+1, q_0} - S_{i, q_0} - S_{j, q_0}\},$$

$$S''_n(u_1, \dots, u_n; q_0) = \bigcup_{i \leq j \leq n} \{u_i + u_j + S_{i, q_0} + S_{j, q_0} - S_{n+1, q_0} - S_{n+1, q_0}\}$$

for every $n = 1, 2, \dots$, and every $(u_1, \dots, u_n) \in \mathbb{Z}^n$.

2.2.2 REMARK. The preceding simple construction encounters difficulties if one tries to handle all $q < 2$ in one move. This is because the sets $S_n = \bigcup \{sp(f_n^\circ) \cup sp(g_{n, q}^\circ) : q < 2\}$ are infinite, and it is no longer clear that integers v_n can be chosen so that the analogue of (2.2.12) is satisfied.

However, in 2.3 we shall combine the result in Construction 2.2.1 with category arguments to indicate for $q \in [1, 2)$ the existence of $f \in A^\#$ and $g_q \in A^q$ for which $fg_q \notin A^q$.

2.3 Extension of Construction 2.2.1.

From Construction 2.2.1 we have that, for $r = 1, 2, \dots$,

$$\exists f_r \in A^\# \text{ such that } f_r \notin M(A^q), \forall q \in I_r = [1, 2 - r^{-1}]. \quad (2.3.1)$$

We wish to make use of (2.3.1) to show that

$$\exists f \in A^\# \text{ such that } f \notin M(A^q), \forall q \in I_\infty = [1, 2) = \bigcup_{r=1}^{\infty} I_r. \quad (2.3.2)$$

Using Lemma 1.3.8 we see that (2.3.1) is equivalent to saying that, for $r = 1, 2, \dots$,

$$\exists f_r \in A^\# \text{ such that } \inf_{q \in I_r} \|f_r\|_{M(q, q)} = \infty; \quad (2.3.3)$$

and that (2.3.2) is equivalent to

$$\exists f \in A^\# \text{ such that } \inf_{q \in I_\infty} \|f\|_{M(q,q)} = \infty. \quad (2.3.4)$$

What we shall actually prove is that if (2.3.2) is false then (2.3.1) is false. In our proof we need the following lemma.

2.3.1 LEMMA. *Let r, s denote positive integers. Then*

$$V_{r,s} = \{f \in A^\# \mid \inf_{q \in I_r} \|f\|_{M(q,q)} \leq s\}$$

is closed in $A^\#$.

Proof. Consider a sequence $(f_j)_{j=1}^\infty$ from $V_{r,s}$ which converges to f in $A^\#$. For each $j = 1, 2, \dots$, $\inf_{q \in I_r} \|f_j\|_{M(q,q)} \leq s$ so that given $\delta > 0$ we can choose a sequence $(q_j)_{j=1}^\infty$ for which

$$\|f_j\|_{M(q_j, q_j)} \leq s + \delta. \quad (2.3.5)$$

By considering a subsequence of (q_j) if necessary we can, and shall, suppose that $q_j \rightarrow q$ in I_r . Now for all $g \in TP, g \neq 0$,

$$\frac{N_{q_j}(f_j g)}{N_{q_j}(g)} \leq \|f_j\|_{M(q_j, q_j)}$$

and so by (2.3.5)

$$N_{q_j}(f_j g) \leq (s + \delta) N_{q_j}(g). \quad (2.3.6)$$

Since

$$N_{q_j}(f_j g) = \|f_j g\|_\infty + \|(f_j g)^\wedge\|_{q_j} \quad (2.3.7)$$

and $(f_j) \rightarrow f$ in $A^\#$, it follows that, in C , $(f_j) \rightarrow f$ and hence $(f_j g) \rightarrow fg$. We then have

$$\lim_j \|f_j g\|_\infty = \|fg\|_\infty. \quad (2.3.8)$$

By (2.3.6) and (2.3.7)

$$\left(\sum_{n \in \mathbb{Z}} |(f_j g)^\wedge(n)|^{q_j} \right)^{\frac{1}{q_j}} \leq (s + \delta) N_{q_j}(g) - \|f_j g\|_\infty.$$

Thus, for $K = 1, 2, \dots$,

$$\left(\sum_{|n| \leq K} |(f_j g)^\wedge(n)|^{q_j} \right)^{\frac{1}{q_j}} \leq (s + \delta) N_{q_j}(g) - \|f_j g\|_\infty,$$

which is

$$\begin{aligned} & \left(\sum_{|n| \leq K} \left| \sum_{m \in sp(g)} \hat{f}_j(n-m) \hat{g}(m) \right|^{q_j} \right)^{\frac{1}{q_j}} \\ & \leq (s + \delta) N_{q_j}(g) - \|f_j g\|_\infty. \end{aligned} \quad (2.3.9)$$

Letting $j \rightarrow \infty$ and noting that $\lim_j N_{q_j}(g) = N_q(g)$ we see by (2.3.8) and (2.3.9) that

$$\begin{aligned} & \left(\sum_{|n| \leq K} \left| \sum_{m \in sp(g)} \hat{f}(n-m) \hat{g}(m) \right|^q \right)^{\frac{1}{q}} \\ & \leq (s + \delta) N_q(g) - \|fg\|_\infty. \end{aligned} \quad (2.3.10)$$

By letting $K \rightarrow \infty$ in (2.3.10) we obtain

$$\| (fg)^{\sim} \|_q \leq (s + \delta) N_q(g) - \|fg\|_{\infty},$$

which says that

$$N_q(fg) \leq (s + \delta) N_q(g);$$

consequently

$$\|f\|_{M(q,q)} \leq (s + \delta). \quad (2.3.11)$$

Because (2.3.11) is true for any $\delta > 0$ it follows that

$$\|f\|_{M(q,q)} \leq s \text{ which, since } q \in I_r, \text{ gives us that } f \in V_{r,s}$$

as required. //

We are now in a position to prove that if (2.3.2) is false then (2.3.1) is also false.

Proof. If (2.3.2) is false then by (2.3.4) we have that

$$\inf_{q \in I_{\infty}} \|f\|_{M(q,q)} < \infty, \text{ for all } f \in A^{\#}$$

which, since $I_{\infty} = \bigcup_{r=1}^{\infty} I_r$, is equivalent to

$$\inf_r \inf_{q \in I_r} \|f\|_{M(q,q)} < \infty, \text{ for all } f \in A^{\#}. \quad (2.3.12)$$

For $V_{r,s}$ as in Lemma 2.3.1, (2.3.12) gives us that

$A^{\#} \subseteq \bigcup_{r,s=1}^{\infty} V_{r,s}$. Clearly the reverse inclusion holds so that

$$A^{\#} = \bigcup_{r,s=1}^{\infty} V_{r,s}. \quad (2.3.13)$$

By (2.3.13), Lemma 2.3.1 and Baire's Category theorem, at least one of the sets $V_{r,s}$ has non-empty interior relative to $A^\#$. Hence there exist natural numbers r_0 and s_0 , a function $f_0 \in A^\#$, and real numbers $\alpha > 0$ and $p_0 > 1$ satisfying

$$\left\{ f \in A^\# \text{ and } N_{p_0}(f - f_0) \leq \alpha \right\} \Rightarrow f \in V_{r_0, s_0}. \quad (2.3.14)$$

Now choose $g_0 \in TP$ satisfying $N_{p_0}(g_0 - f_0) \leq \frac{1}{2}\alpha$. By the triangle inequality, (2.3.14) and the definition of V_{r_0, s_0} ,

$$\left\{ f \in A^\# \text{ and } N_p(f - g_0) \leq \frac{1}{2}\alpha \right\} \Rightarrow \inf_{q \in I_{r_0}} \|f\|_{M(q, q)} \leq s_0. \quad (2.3.15)$$

We now see that for $g \in TP$,

$$\begin{aligned} N_q(g_0 g) &= N_q \left(\sum_{n \in sp(g_0)} \hat{g}_0(n) e_n g \right) \\ &\leq \sum_{n \in sp(g_0)} N_q(\hat{g}_0(n) e_n g) \\ &= \sum_{n \in sp(g_0)} |\hat{g}_0(n)| N_q(e_n g) \\ &= N_q(g) \sum_{n \in sp(g_0)} |\hat{g}_0(n)|, \end{aligned}$$

the final step by (1.2.2); hence

$$\|g_0\|_{M(q, q)} \leq \sum_{n \in sp(g_0)} |\hat{g}_0(n)| < \infty. \quad (2.3.16)$$

Writing $\sum_{n \in sp(g_0)} |\hat{g}_0(n)| = \gamma$, we have by (2.3.16) that

$$\sup_{q \in I_{r_0}} \|g_0\|_{M(q, q)} \leq \gamma < \infty. \quad (2.3.17)$$

If $h \in A^\#$ satisfies $N_{p_0}(h) \leq \frac{1}{2}\alpha$ then, by (2.3.15),

$$\inf_{q \in I_{r_0}} \|g_0 + h\|_{M(q,q)} \leq s_0. \quad (2.3.18)$$

For $q \in I_{r_0}$, the triangle inequality together with (2.3.17) gives us that

$$\begin{aligned} \|h\|_{M(q,q)} &\leq \|g_0 + h\|_{M(q,q)} + \|g_0\|_{M(q,q)} \\ &\leq \|g_0 + h\|_{M(q,q)} + \gamma; \end{aligned}$$

and so, by (2.3.18),

$$\inf_{q \in I_{r_0}} \|h\|_{M(q,q)} \leq s_0 + \gamma. \quad (2.3.19)$$

Replacing h by $\frac{1}{2}\alpha h/N_{p_0}(h)$ in (2.3.19) we have that

$$\inf_{q \in I_{r_0}} \|h\|_{M(q,q)} \leq 2\alpha^{-1}(s_0 + \gamma)N_{p_0}(h) < \infty, \text{ for all } h \in A^\#,$$

which provides a contradiction to (2.3.3) and hence to (2.3.1). //

2.4 Construction of $f \in A^\#$ with Fourier series divergent at a point.

For $f \in L^\infty$ we shall write $\|f\|_U = \sup_N \|s_N f\|_\infty$ and denote by U_0 the Banach space of those $f \in L^\infty$ for which $\|f\|_U < \infty$. We shall denote by U the space of functions with uniformly convergent Fourier series. U is the closed subspace of U_0 generated by the trigonometric polynomials; it is a Banach space with norm $\|\cdot\|_U$. Finally, we denote by U^T the subspace of U consisting of those $f \in U$ for which $\hat{f}(n) = 0$ for all $n < 0$. (The T here indicates

that each $f \in U^T$ can be written as a Taylor series.) In [30] R. Salem showed that there exists $f \in U$ for which $f^2 \notin U$. He then asked if U here could be replaced by U^T and this question was answered affirmatively by J-P. Kahane and Y. Katznelson in [22]. Salem's result proves simultaneously that U is not an algebra and that the space $M(U)$ of pointwise multipliers of U is not the whole of U . In [3] Leon Brown used Salem's result to show that any closed subspace of U_0 that contains $1, e_1$ and e_{-1} is not an algebra. We now make use of a construction due to Fejér (see [34] p.299, or [5], 10.3.1) to show that there exists $f \in A^\# \setminus U_0$; and hence that Brown's result cannot be used to prove that $A^{p,p} \in (1,2)$, is not an algebra.

2.4.1 CONSTRUCTION (Fejér). For integers r and q satisfying $r > q \geq 1$ let $Q(x, r, q)$ denote the trigonometric polynomial defined by

$$\begin{aligned} Q(x, r, q) &= 2 \sin rx \sum_{k=1}^q k^{-1} \sin kx \\ &= \frac{\cos(r-q)x}{q} + \dots + \frac{\cos(r-1)x}{1} - \frac{\cos(r+1)x}{1} \\ &\quad - \dots - \frac{\cos(r+q)x}{q}. \end{aligned}$$

Since the partial sums of $\sum_{k=1}^{\infty} k^{-1} \sin kx$ are uniformly bounded (see [5], 7.2.2) the $Q(x, r, q)$ are uniformly bounded.

Suppose now that $(r_k)_{k=1}^{\infty}$ and $(q_k)_{k=1}^{\infty}$ are sequences of integers satisfying

$$1 \leq q_k \leq r_k \quad \text{and} \quad r_k + q_k < r_{k+1} - q_{k+1}. \quad (2.4.1)$$

Suppose further that $(\alpha_k)_{k=1}^{\infty}$ is any sequence of complex numbers for which

$$\sum_{k=1}^{\infty} |\alpha_k| < \infty \quad (2.4.2)$$

and

$$\liminf_{k \rightarrow \infty} |\alpha_k| \log q_k > 0 \quad (2.4.3)$$

One possible choice satisfying (2.4.1), (2.4.2) and (2.4.3) is

$$r_k = 2^{k^3+1}, \quad q_k = 2^{k^3}, \quad \alpha_k = k^{-2} \quad (2.4.4)$$

Now consider the function f defined by

$$f(x) = \sum_{k=1}^{\infty} f_k(x) = \sum_{k=1}^{\infty} \alpha_k Q(x, r_k, q_k) \quad (2.4.5)$$

Because of (2.4.2) and the uniform boundedness of the $Q(x, r, q)$, the series (2.4.5) is uniformly convergent; and hence $f \in C$. Writing $S_k = \text{sp}\{f_k\}$, (2.4.1) ensures that the S_k are pairwise disjoint.

We can combine this with the uniform convergence of (2.4.5) to obtain

$$\hat{f}(n) = \begin{cases} \hat{f}_k(n) & , \quad n \in S_k \\ 0 & , \quad n \notin \bigcup_k S_k \end{cases} \quad (2.4.6)$$

From this it follows that

$$s_{r_k+q_k} f(x) - s_{r_k} f(x) = \sum_{v=r_k+1}^{r_k+q_k} \frac{\alpha_k \cos vx}{r_k - v} \quad (2.5.2)$$

Hence

$$\begin{aligned} |s_{r_k+q_k} f(0) - s_{r_k} f(0)| &= |\alpha_k| |1 + \frac{1}{2} + \dots + q_k^{-1}| \\ &> |\alpha_k| \log q_k \end{aligned}$$

Thus, by (2.4.3), the Fourier series of f diverges at 0; and

$$\|f\|_U \geq \sup_N |s_N f(0)| = \infty,$$

which means that $f \notin U_0$.

We shall show now that if $f \in \mathcal{C}$ is constructed as just described then $f \in A^\#$. By (2.4.6), for $p > 1$,

$$\begin{aligned} \sum_{n \in \mathbb{Z}} |\hat{f}(n)|^p &= \sum_{k=1}^{\infty} \sum_{n \in S_k} |\hat{f}_k(n)|^p \\ &= 2 \sum_{k=1}^{\infty} |\alpha_k|^p \sum_{m=1}^{q_k} m^{-p}. \end{aligned} \quad (2.4.7)$$

By (2.4.7) and the convergence, for $p > 1$, of both $\sum_{m=1}^{\infty} m^{-p}$ and $\sum_{k=1}^{\infty} |\alpha_k|^p$, $\sum_{n \in \mathbb{Z}} |\hat{f}(n)|^p < \infty$; and hence $f \in A^\#$. Making the choice indicated in (2.4.4) we have $f \in A^\# \setminus U_0$, as required.

2.5 Strict inclusion results for A^p , $p \in [1, 2]$.

As one might expect, the strict inclusion result

$$A^p \subsetneq A^q, \quad 1 \leq p < q \leq 2,$$

holds for these spaces. In fact this inclusion can be strengthened to give both

$$\bigcup_{p \in [1, q)} A^p \subsetneq A^q \quad \text{if } q \in (1, 2], \quad (2.5.1)$$

and

$$A^q \subsetneq \bigcap_{p \in (q, 2]} A^p \quad \text{if } q \in [1, 2). \quad (2.5.2)$$

In this section we give constructive proofs to establish both (2.5.1) and (2.5.2).

2.5.1 CONSTRUCTION. The strict inclusion (2.5.1).

Consider a given $q \in (1, 2]$. Define $f_k \in TP$ by

$$f_k = \beta_{k,q} P_k e_{v_k}, \quad k = 0, 1, 2, \dots, \quad (2.5.3)$$

where the sequences $(\beta_{k,q})_{k=0}^{\infty}$ and $(v_k)_{k=0}^{\infty}$ will be chosen appropriately, the latter in such a way to ensure that the $S_k = \text{sp}(f_k)$ are pairwise disjoint.

From (1.2.2), (1.2.10) and (2.5.3) we have

$$N_q(f_k) \sim \beta_{k,q} 2^{k/q}, \quad q \in (1, 2]. \quad (2.5.4)$$

Now consider $f = \sum_{k=0}^{\infty} f_k$. Since $N_q(f) \leq \sum_{k=0}^{\infty} N_q(f_k)$ it follows from

(2.5.4) that a sufficient condition for $f \in A^q$ is that

$$\sum_{k=0}^{\infty} \beta_{k,q} 2^{k/q} < \infty, \quad q \in (1, 2]. \quad (2.5.5)$$

Put

$$\beta_{k,q} = (k+1)^{-2} 2^{-k/q}, \quad k = 0, 1, 2, \dots \quad (2.5.6)$$

With this choice of $\beta_{k,q}$, (2.5.5) is satisfied since

$$\sum_{k=0}^{\infty} \beta_{k,q} 2^{k/q} = \sum_{k=0}^{\infty} (k+1)^{-2} < \infty.$$

We shall now show that, with $(\beta_{k,q})_{k=0}^{\infty}$ as in (2.5.6), the associated function $f \notin \bigcup_{p \in [1, q)} A^p$. Since the series defining f converges in A^q ,

$$\hat{f}(n) = \begin{cases} \hat{f}_k(n), & n \in S_k, \quad k = 0, 1, 2, \dots, \\ 0 & , \quad n \notin \bigcup_k S_k, \end{cases} \quad (2.5.7)$$

where

$$\hat{f}_k(n) = \begin{cases} \varepsilon_k(n) (k+1)^{-2} 2^{-k/q}, & n \in S_k, \\ 0 & , \quad \text{otherwise} \end{cases} \quad (2.5.8)$$

Also, for f_k defined as in (2.5.3),

$$S_k = \{n \in \mathbb{Z} : v_k \leq n \leq v_k + 2^k - 1\}, \quad (2.5.9)$$

so each S_k is a finite set with cardinality

$$|S_k| = 2^k. \quad (2.5.10)$$

Thus, making use of (2.5.7), (2.5.8) and (2.5.10),

$$\begin{aligned} \|\hat{f}\|_p^p &= \sum_{n \in \mathbb{Z}} |\hat{f}(n)|^p \\ &= \sum_{k=0}^{\infty} \sum_{n \in S_k} |\hat{f}_k(n)|^p \\ &= \sum_{k=0}^{\infty} (k+1)^{-2p} 2^{-kp/q} 2^k \\ &= \sum_{k=0}^{\infty} (k+1)^{-2p} 2^{k(1-p/q)} \\ &= \infty \quad \text{for } q \in (1, 2], p \in [1, q); \end{aligned} \quad (2.5.13)$$

and so $f \notin \bigcup_{p \in [1, q)} A^p$.

We still need to choose $\{v_k\}$ appropriately. By (2.5.9) it is sufficient to choose $\{v_k\}$ to be a strictly increasing sequence satisfying

$$v_{k+1} > v_k + 2^k - 1, \quad k = 0, 1, 2, \dots, \quad (2.5.11)$$

to ensure that the S_k , $k = 0, 1, 2, \dots$, are pairwise disjoint.

The condition given in (2.5.11) is satisfied when

$$v_k = 2^{k+1}, \quad k = 0, 1, 2, \dots,$$

and our construction is completed.

2.5.2 CONSTRUCTION. *The strict inclusion (2.5.2).*

The method employed here is similar to that used in 2.5.1.

Consider a given $q \in [1, 2)$ and define $f_k \in TP$ by

$$f_k = (k+1)^{-1/q} 2^{-k/q} p_k e_{2^{k+1}}, \quad k = 0, 1, 2, \dots \quad (2.5.12)$$

For each such k write $S_k = sp\{f_k\}$; it is not hard to see that the S_k are pairwise disjoint. From (1.2.2), (1.2.11), (1.2.12) and (2.5.12).

$$N_p(f_k) \sim (k+1)^{-1/q} 2^{k(1/p-1/q)}, \quad p \in (q, 2]. \quad (2.5.13)$$

Now put $f = \sum_{k=0}^{\infty} f_k$. Since $N_p(f) \leq \sum_{k=0}^{\infty} N_p(f_k)$ we have from (2.5.13)

that $N_p(f) < \infty$ for all $p \in (q, 2]$, and so $f \in \bigcap_{p \in (q, 2]} A^p$. Since

the series defining f converges in A^p for $p \in (q, 2]$,

$$\hat{f}(n) = \begin{cases} \hat{f}_k(n), & n \in S_k, \\ 0 & , n \notin \bigcup_k S_k \end{cases} \quad (2.5.14)$$

where

$$\hat{f}_k(n) = \begin{cases} \epsilon_k(n) (k+1)^{-1/q} 2^{-k/q}, & n \in S_k, \\ 0 & \text{otherwise} \end{cases} \quad (2.5.15)$$

Thus, making use of (2.5.14), (2.5.15) and the fact that the S_k are pairwise disjoint,

Proof. That $f \in C \cap BV$ implies $f \in A^p$ follows directly from (2.6.2) and the definition of A^p . Using (2.6.1) and the fact that

$$\sum_{n=1}^{\infty} n^{-p} < p/(p-1) = p', \quad \text{for } p \in (1, \infty),$$

$$\begin{aligned}
\|\hat{f}\|_q^q &= \sum_{n \in \mathbb{Z}} |\hat{f}(n)|^q \\
&= \sum_{k=0}^{\infty} \sum_{n \in S_k} |\hat{f}_k(n)|^q \\
&= \sum_{k=0}^{\infty} (k+1)^{-1} \\
&= \infty ;
\end{aligned}$$

so that $f \notin A^q$ and our construction is completed.

2.6 Some further results for A^p spaces.

We shall denote by $V[f]$ the total variation of the function f on T ; and by BV the space of functions of bounded variation. Now (see [5], 2.3.6),

$$|n\hat{f}(n)| \leq \frac{1}{2\pi} V[f] \quad \text{for all } n \in \mathbb{Z}, \quad (2.6.1)$$

and hence, if $p > 1$,

$$\hat{f} \in \ell^p \quad \text{for all } f \in BV. \quad (2.6.2)$$

2.6.1 LEMMA. If $f \in C \cap BV$ then $f \in A^\#$ and, for $p \in (1, \infty)$,

$$N_p(f) \leq \|f\|_\infty + \left(\|f\|_\infty^p + \frac{p'}{\pi} V[f] \right)^{1/p},$$

where $\frac{1}{p} + \frac{1}{p'} = 1$.

Proof. That $f \in C \cap BV$ implies $f \in A^\#$ follows directly from (2.6.2) and the definition of $A^\#$. Using (2.6.1) and the fact that

$$\sum_{n=1}^{\infty} n^{-p} \leq p/(p-1) = p', \quad \text{for } p \in (1, \infty),$$

we have

$$\|\hat{f}\|_p^p = |\hat{f}(0)|^p + \sum_{n \in \mathbb{Z} \setminus \{0\}} |\hat{f}(n)|^p$$

$$\leq |\hat{f}(0)|^p + \frac{1}{\pi} V[f] \sum_{n=1}^{\infty} n^{-p}$$

$$\leq |\hat{f}(0)|^p + \frac{p'}{\pi} V[f]$$

$$\leq \|f\|_{\infty}^p + \frac{p'}{\pi} V[f] .$$

Hence

$$N_p(f) = \|f\|_{\infty} + \|\hat{f}\|_p$$

$$\leq \|f\|_{\infty} + \left(\|f\|_{\infty}^p + \frac{p'}{\pi} V[f] \right)^{1/p} ,$$

as required. //

Now consider $p \in (1,2)$. If $f \in \Lambda_{\alpha}^1$ with $\alpha \in (p^{-1},1]$ then (see [34], p.47) $\hat{f}(n) = O(|n|^{-\alpha})$ and so $f \in \mathcal{FL}^p$. Hence, if $f \in C \cap \Lambda_{\alpha}^1$ with $\alpha \in (p^{-1},1]$, then $f \in A^p$. In [1] Alpar proved (using different notation and terminology) the following result.

2.6.2 THEOREM. Suppose that $p \in (1,2)$, $f \in C \cap \Lambda_{\alpha}^1$ with $\alpha \in (p^{-1},1]$ and that the complex function F is holomorphic on an open set containing the range of f . Then $F \circ f \in A^p$. In particular, if f does not vanish on T then the choice of $F: z \rightarrow z^{-1}$ gives $f^{-1} \in A^p$. //

Actually, with our formulation, Theorem 2.6.2 can be proved directly by the use of Banach algebra techniques.

We shall now prove some inclusion results which were communicated privately to the author by Professor Y. Katznelson.

2.6.3 THEOREM. Let $p, q \in (1, 2)$. Then

$Fl^p \cdot Fl^q \subseteq Fl^r$ if and only if $r \geq pq/(p+q-pq)$.

Proof. Since $p \in (1, 2)$ and $q \in (1, 2)$ we have $\frac{1}{p} + \frac{1}{q} > 1$ and so by [16], (20.18)

$$Fl^p \cdot Fl^q \subseteq Fl^{pq/(p+q-pq)};$$

hence

$$Fl^p \cdot Fl^q \subseteq Fl^r \quad \text{if } r \geq pq/(p+q-pq).$$

Conversely, suppose that r is such that

$$Fl^p \cdot Fl^q \subseteq Fl^r. \quad (2.6.3)$$

By (1.2.7) and (1.2.8), $\|\hat{D}_n\|_p \sim n^{1/p}$, $\|\hat{D}_n\|_q \sim n^{1/q}$ and $\|(D_n^2)^\wedge\|_r \sim n^{1+1/r}$. Hence for (2.6.3) to hold we must have

$$n^{1+1/r} = O\left(n^{\frac{1}{p} + \frac{1}{q}}\right) \quad \text{and so } r \geq pq/(p+q-pq). \quad //$$

2.6.4 THEOREM. Let $p, q \in (1, 2)$. Then

$A^q \cdot Fl^p \subseteq Fl^r$ if and only if $r \geq pq/(2+pq-p-q)$.

Proof. If $f \in A^q$, $q \in (1, 2)$, then $f \cdot Fl^1 \subseteq Fl^q$ and $f \cdot Fl^2 \subseteq Fl^2$. By appeal to the Riesz-Thorin interpolation theorem we then have that for all $\alpha \in (0, 1)$,

$$f \cdot Fl^{2/(2-\alpha)} \subseteq Fl^{2q/(q\alpha+2(1-\alpha))}. \quad (2.6.4)$$

If $\alpha = 2(p-1)/p$ then (2.6.4) gives

$$f \cdot Fl^p \subseteq Fl^{pq/(2+pq-p-q)}$$

and so

$$A^q \cdot Fl^p \subseteq Fl^r \quad \text{if } r \geq pq/(2+pq-p-q).$$

Conversely, suppose that r is such that

$$A^q.F\ell^p \subseteq F\ell^r. \quad (2.6.5)$$

By (1.2.7), $\|\hat{D}_n\|_p \sim n^{1/p}$. Also, combination of Lemmas 1.2.6 and 2.1.1 with the estimates (1.2.11) and (1.2.12) gives us that, for $s > 2n$,

$$\left\| \left(D_n^2 P_{m(2s)}^* \right)^\wedge \right\|_r \sim n^{1+1/r} m^{1/r}, \quad (2.6.6)$$

and

$$N_q \left(D_n P_{m(2s)}^* \right) \sim n^{1/q} m^{1/q} + rm^{1/2}. \quad (2.6.7)$$

Choose $m = \left\lceil n^{2(q-1)/(2-q)} \right\rceil$. Then, by (2.6.6) and (2.6.7) respectively,

$$\left\| \left(D_n^2 P_{m(2s)}^* \right)^\wedge \right\|_r \sim n^{1+q/r(2-q)}$$

and

$$N_q \left(D_n P_{m(2s)}^* \right) \sim n^{1/(2-q)}.$$

Thus for (2.6.5) to hold we must have that

$$n^{1+qr^{-1}(2-q)^{-1}} = o \left(n^{(p+2-q)/p(2-q)} \right)$$

and so $r \geq pq/(2+pq - p-q)$.

2.6.5 REMARK. The condition on r given in Theorem 2.6.4 is symmetric in p and q even though $A^q.F\ell^p$ is not.

2.6.6 COROLLARY. Let $p, q \in (1, 2)$. Then

$$A^p.A^q \subseteq A^r \text{ if } r \geq pq/(2+pq - p-q). \quad (2.6.8)$$

Proof. This result follows directly from Theorem 2.6.4 and the definition of A^p . //

2.6.7 THEOREM. Let $p \in (1,2)$ and $q \in (1,2)$. Then

$$A^p \cdot A^q \subseteq \mathcal{FL}^r \text{ only if } r \geq \frac{4(p+q-1) - 3pq}{p+q - pq}.$$

Proof. For $p, q \in (1,2)$ put

$$f = D_n P_{m_1}^*(2s_1) \quad \text{and} \quad g = D_n P_{m_2}^*(2s_2),$$

where

$$m_1 = \left[n^{2(p-1)/(2-p)} \right] \quad \text{and} \quad m_2 = \left[n^{2(q-1)/(2-q)} \right],$$

and s_1, s_2 satisfy

$$s_1 > 2n \quad \text{and} \quad s_2 > 2n + 2s_1 m_1.$$

Then, by Lemma 1.2.6, Lemma 2.1.1 and the estimates (1.2.11) and (1.2.12),

$$N_p(f) \sim n^{1/(2-p)} \quad \text{and} \quad N_q(g) \sim n^{1/(2-q)}; \quad (2.6.9)$$

and also

$$\begin{aligned} \|(fg)^\wedge\|_r &= \left\| \left(D_n^2 P_{m_1}^*(2s_1) P_{m_2}^*(2s_2) \right)^\wedge \right\|_r \\ &\sim n^{1+r^{-1}} + 2(3p+3q-2pq-4)r^{-1}(2-p)^{-1}(2-q)^{-1}. \end{aligned} \quad (2.6.10)$$

Hence if $A^p \cdot A^q \subseteq \mathcal{FL}^r$ then, by (2.6.9) and (2.6.10),

$$n^{1+r^{-1}} + 2(3p+3q-2pq-4)r^{-1}(2-p)^{-1}(2-q)^{-1} = O\left(n^{(4-p-q)/(2-p)(2-q)}\right)$$

and so $r \geq \frac{4(p+q-1) - 3pq}{p+q - pq}$.

2.6.8 REMARK. If $p \in (1,2)$ and $A^p \cdot A^p \subseteq A^r$ then, by Theorem 2.6.7, $r \geq 3 - 2/p > p$; it follows from this that A^p is not an algebra for $p \in (1,2)$. //

We shall conclude this chapter with two theorems, the first of which shows that A^p , $p \in (1,2)$, is translation continuous; this will be used in Chapter 4.

2.6.9 THEOREM. For each $f \in A^p$ the function $\alpha \rightarrow \tau_\alpha f$ is continuous from T to A^p , $p \in (1,2)$.

Proof. Without loss of generality we need show continuity only at 0. Consider $f \in A^p$, $p \in (1,2)$. Since, for all $\alpha \in T$,

$$N_p(\tau_\alpha f - f) = \|\tau_\alpha f - f\|_\infty + \|(\tau_\alpha f - f)^\wedge\|_p$$

and

$$\lim_{\alpha \rightarrow 0} \|\tau_\alpha f - f\|_\infty = 0,$$

we need prove only that

$$\lim_{\alpha \rightarrow 0} \|(\tau_\alpha f - f)^\wedge\|_p = 0;$$

which follows immediately by the Lebesgue dominated convergence theorem. //

In 3.2 we show that the strong dual of A^p may be identified with the space $M + F\ell^{p'}$. We anticipate this result in the proof of the following theorem.

2.6.10 THEOREM. A^p , $p \in [1,2]$, is not reflexive.

Proof. Consider the sequence $(\frac{1}{2}e_n)_{n=1}^\infty$ of functions in the unit ball of A^p , $p \in [1,2]$. Now suppose that some subsequence

$(\frac{1}{2}e_{n_k})_{k=1}^\infty$ of $(\frac{1}{2}e_n)_{n=1}^\infty$ is weakly convergent in A^p , to $g \in A^p$ say.

Then, for all $\sigma \in M + \mathcal{FL}^{p'}$,

$$\lim_{k \rightarrow \infty} \langle \sigma, \frac{1}{2} e_{n_k} \rangle = \langle \sigma, g \rangle ; \quad (1.2)$$

and, in particular (since $A^p \subseteq M + \mathcal{FL}^{p'}$),

$$\lim_{k \rightarrow \infty} \langle \bar{g}, \frac{1}{2} e_{n_k} \rangle = \langle \bar{g}, g \rangle . \quad p \in [1, 2] , \quad (2.6.13)$$

By (2.6.13) and the Riemann-Lebesgue lemma it follows that

$\langle \bar{g}, g \rangle = 0$ and hence $g = 0$. However, there exist

$\sigma \in M + \mathcal{FL}^{p'}$ which do not satisfy $\lim_{k \rightarrow \infty} \langle \sigma, \frac{1}{2} e_{n_k} \rangle = 0$ (the

Dirac measure at 0 for instance) so we have a contradiction. Hence

the unit ball of A^p is not weakly compact and so, by [28], Corollary

3, p.73 together with Smúlian's theorem, A^p is not reflexive. //

CHAPTER 3

THE FOURIER MULTIPLIER PROBLEM FOR A^p , $p \in [1,2]$

3.1 Introduction

By a Fourier multiplier from A^p to A^q , $p \in [1,2]$, $q \in [1,2]$, we mean a continuous linear operator $T: A^p \rightarrow A^q$ which commutes with translations. As can be seen from [5], 16.3.1, to each multiplier $T: A^p \rightarrow A^q$ there corresponds a unique distribution ϕ such that T is (the restriction to A^p of) the operator T_ϕ defined by

$$T_\phi f = \phi * f.$$

We denote the space of such distributions ϕ by (A^p, A^q) and refer to $\phi \in (A^p, A^q)$ as a multiplier from A^p to A^q . Since TP is dense in A^p , a distribution ϕ belongs to (A^p, A^q) if and only if

$$N_q(\phi * f) \leq \text{const. } N_p(f), \text{ for all } f \in TP. \quad (3.1.1)$$

In particular, a distribution ϕ belongs to $(A^p, C) = (A^p, A^2)$ if and only if

$$\|\phi * f\|_\infty \leq \text{const. } N_p(f), \text{ for all } f \in TP; \quad (3.1.2)$$

or (using the translation invariance of N_p) what is equivalent, if and only if

$$|\phi * f(0)| \leq \text{const. } N_p(f), \text{ for all } f \in TP. \quad (3.1.3)$$

We define $(A^p)'$ to be the set of linear functionals ℓ on TP such that

$$|\ell(f)| \leq \text{const. } N_p(f), \text{ for all } f \in TP. \quad (3.1.4)$$

Since TP is dense in A^p , the restriction to TP gives a one-to-one map of the dual of A^p onto $(A^p)'$.

3.2 The multipliers (A^p, A^p) , $p \in [1, 2]$.

In order to identify these spaces we first prove two lemmas.

3.2.1 LEMMA. $(A^p, A^p) = (A^p, C)$, $p \in [1, 2]$.

Proof. Since, for $p \in [1, 2]$, $A^p \subseteq C$ with a continuous injection, we have that $(A^p, A^p) \subseteq (A^p, C)$. Conversely, suppose $\phi \in (A^p, C)$, $p \in [1, 2]$. Then, by (3.1.2), for all $f \in TP$

$$\|\phi * f\|_\infty \leq \text{const. } N_p(f); \quad (3.2.1)$$

hence for all $n \in \mathbb{Z}$

$$|\hat{\phi}(n) \cdot \hat{f}(n)| \leq \|\phi * f\|_\infty \leq \text{const. } N_p(f). \quad (3.2.2)$$

Put $f = e_n$ in (3.2.2) to get

$$|\hat{\phi}(n)| \leq \text{const. for all } n \in \mathbb{Z}.$$

Thus $\phi \in PM$ and, for all $f \in TP$,

$$\|\hat{\phi} \cdot \hat{f}\|_p \leq \|\hat{\phi}\|_\infty \|\hat{f}\|_p. \quad (3.2.3)$$

Combination of (3.2.2) and (3.2.3) shows that

$$N_p(\phi * f) \leq \text{const. } N_p(f), \text{ for all } f \in TP;$$

by (3.1.1), $\phi \in (A^p, A^p)$. Thus $(A^p, C) \subseteq (A^p, A^p)$ which completes our proof. //

3.2.2 REMARK. In view of what was said in 3.1 there is a one-to-one correspondence $\ell \leftrightarrow \phi$ between $(A^p)'$ and (A^p, C) under which

$$\ell(\tau_{-x}f) = \phi * f(x), \text{ for all } f \in TP, x \in T;$$

which is equivalent to

$$\ell(f) = \phi * f(0), \text{ for all } f \in T.$$

3.2.3 LEMMA. To every $\ell \in (A^p)'$ corresponds $\mu \in M$ and $\sigma \in Fl^p$ such that

$$\ell(f) = \mu * f(0) + \sigma * f(0), \text{ for all } f \in TP.$$

The converse is also true.

Proof. Define

$$S = \{(f, \hat{f}) \in C \times \ell^p : f \in TP\}.$$

Take $\ell \in (A^p)'$ and define a map ℓ' on S by

$$\ell' : (f, \hat{f}) \rightarrow \ell(f).$$

ℓ' is clearly linear; and since

$$|\ell'((f, \hat{f}))| = |\ell(f)|$$

$$\leq \text{const. } N_p(f)$$

$$= \text{const. } (\|f\|_\infty + \|\hat{f}\|_p),$$

ℓ' is continuous on S as a subspace of $C \times \ell^p$. Thus, by the Hahn-Banach Theorem, ℓ' can be extended to a continuous linear functional on the whole of $C \times \ell^p$. Denote this extension by ℓ' also.

The mapping $f \rightarrow \ell'((f, 0))$ is a continuous linear functional on C , so it can be represented by a measure, $\mu \in M$, such that

$$\ell'((f, 0)) = \langle \mu, f \rangle = \mu * f(0), \text{ for all } f \in C. \quad (3.2.4)$$

Also, $\theta \rightarrow \ell'((0, \theta))$ is a continuous linear functional on ℓ^p so it can be represented by $\alpha \in \ell^{p'}$, such that

$$\ell'((0, \theta)) = \sum_{n \in \mathbb{Z}} \alpha(n) \theta(n). \quad (3.2.5)$$

Define $\hat{\sigma} \in F\ell^{p'}$ by

$$\hat{\sigma}(n) = \alpha(n), \text{ for all } n \in \mathbb{Z}.$$

Then, for $f \in TP$,

$$\sigma * f = \sum_{n \in \mathbb{Z}} \hat{\sigma}(n) \hat{f}(n) e_n;$$

thus, using (3.2.5), we can write

$$\sigma * f(0) = \sum_{n \in \mathbb{Z}} \alpha(n) \hat{f}(n) = \ell'((0, \hat{f})) \quad (3.2.6)$$

Combination of (3.2.4) and (3.2.6) gives, for all $f \in TP$,

$$\begin{aligned}\ell(f) &= \ell'((f, 0)) + \ell'((0, \hat{f})) \\ &= \mu * f(0) + \sigma * f(0),\end{aligned}$$

where $\mu \in M$ and $\sigma \in Fl^{p'}$.

Conversely, suppose $\mu \in M$ and $\sigma \in Fl^{p'}$ and consider the map defined on TP by

$$\ell: f \rightarrow \mu * f(0) + \sigma * f(0).$$

Then $\ell \in (A^p)'$, since for all $f \in TP$,

$$|\ell(f)| \leq |\mu * f(0)| + |\sigma * f(0)|$$

$$\leq \|\mu * f\|_{\infty} + \|\sigma * f\|_{\infty}$$

$$\leq \|\mu\|_M \|f\|_{\infty} + \|\hat{\sigma}\|_p \|\hat{f}\|_p$$

$$\leq \text{const. } N_p(f) \quad //$$

3.2.4 REMARK. Lemma 3.2.3 provides the identification of the dual of A^p with $M + Fl^{p'}$, which was promised in 2.6.10. This dual space was considered independently, and in a different context, by Gregory in [13]. //

We are now in a position to prove the following result.

3.2.5 THEOREM. Suppose $p \in [1, 2]$. Then

$$(A^p, C) = (A^p, A^p) = M + Fl^{p'}.$$

Proof. By Lemma 3.2.1, $(A^p, A^p) = (A^p, \mathbb{C})$. From Remark 3.2.2 it follows that $\phi \in (A^p, \mathbb{C})$ if and only if

$$\phi * f(x) = \ell(\tau_{-x} f), \text{ for all } f \in TP, x \in T,$$

for some $\ell \in (A^p)'$. Thus, by Lemma 3.2.3, $\phi \in (A^p, \mathbb{C})$ if and only if there exist $\mu \in M$ and $\sigma \in F\ell^p'$ such that, for every $f \in TP$ and every $x \in T$,

$$\begin{aligned} \phi * f(x) &= \mu * \tau_{-x} f(0) + \sigma * \tau_{-x} f(0) \\ &= \mu * f(x) + \sigma * f(x). \end{aligned}$$

This signifies that $\phi = \mu + \sigma$. //

3.3 Strict inclusion results for (A^p, A^p) , $p \in [1, 2]$.

Since N_r is stronger than N_s if $r < s$, it follows easily that $(A^q)' \subseteq (A^p)'$ whenever $1 \leq p < q \leq 2$. What we shall do here is first prove constructively two (strong) strict inclusion results for the spaces $(A^p)'$, $p \in [1, 2]$, and then use these to prove two corresponding strict inclusion results for the multiplier spaces (A^p, A^p) , $p \in [1, 2]$. The strict inclusion results we shall prove for the spaces $(A^p)'$, $p \in [1, 2]$, are the following:

$$(A^q)' \subsetneq \bigcap_{p \in [1, q)} (A^p)' \text{ if } q \in (1, 2] \quad (3.3.1)$$

and

$$\bigcup_{p \in (q, 2]} (A^p)' \subsetneq (A^q)' \text{ if } q \in [1, 2). \quad (3.3.2)$$

3.3.1 CONSTRUCTION. The strict inclusion (3.3.1).

Consider a given $q \in (1, 2]$. We wish to construct a linear functional, ℓ say, on the space TP , such that

$$\ell(f) = \sum_{n \in \mathbb{Z}} c_n \hat{f}(n), \text{ for all } f \in TP,$$

where $(c_n)_{n \in \mathbb{Z}}$ is chosen so that ℓ is not continuous in the topology induced by A^q , but is continuous in the topology induced by A^p for every $p \in [1, q)$. For the case $p \in (1, q)$, to ensure that $\ell \in (A^p)'$ it is sufficient to choose $(c_n) \in \ell^p$; for then, for every $f \in TP$,

$$\begin{aligned} |\ell(f)| &= \left| \sum_{n \in \mathbb{Z}} c_n \hat{f}(n) \right| \\ &\leq \| (c_n) \|_p \| \hat{f} \|_p \\ &\leq \| (c_n) \|_p N_p(f). \end{aligned}$$

Now define

$$f_k = \beta_{k,q}^p e_{v_k}, \quad k = 0, 1, 2, \dots, \quad (3.3.3)$$

where the sequences $(\beta_{k,q})$ and (v_k) will be chosen appropriately, the latter to ensure that the $S_k = \text{sp}(f_k)$ are pairwise disjoint. We have now that

$$\hat{f}_k(n) = \begin{cases} \varepsilon_k(n) \beta_{k,q}, & n \in S_k, \\ 0, & \text{otherwise.} \end{cases}$$

Put

$$c_n = \begin{cases} b_{k,q} \text{sgn } \hat{f}_k(n) |\hat{f}_k(n)|^{q-1}, & n \in S_k, \\ 0, & n \notin \bigcup_k S_k, \end{cases} \quad (3.3.4)$$

where $\{b_{k,q}\}$ is a sequence of positive numbers which will be chosen appropriately later. Then $\{c_n\} \in \ell^{p'}$ if and only if

$$\begin{aligned} \sum_{n \in \mathbb{Z}} |c_n|^{p'} &= \sum_{k=0}^{\infty} b_{k,q}^{p'} \sum_{n \in S_k} |\hat{f}_k(n)|^{(q-1)p'} \\ &= \sum_{k=0}^{\infty} b_{k,q}^{p'} \beta_{k,q}^{(q-1)p'} |S_k| < \infty . \end{aligned} \quad (3.3.5)$$

To ensure that ℓ is not continuous in the topology induced by A^q , we seek to arrange that

$$\sup_k \left(\frac{|\ell(f_k)|}{N_q(f_k)} \right) = \infty . \quad (3.3.6)$$

By (1.2.2), (1.2.10) and (3.3.3), $N_q(f_k) \sim \beta_{k,q} 2^{k/q}$. Also

$$\begin{aligned} |\ell(f_k)| &= \left| \sum_{n \in \mathbb{Z}} c_n \hat{f}_k(n) \right| \\ &= \sum_{n \in S_k} b_{k,q} |\hat{f}_k(n)|^q \\ &= b_{k,q} \beta_{k,q}^q |S_k| . \end{aligned}$$

Thus (3.3.6) can be replaced by the condition

$$\sup_k \left(\frac{b_{k,q} \beta_{k,q}^q |S_k|}{\beta_{k,q} 2^{k/q}} \right) = \infty ;$$

that is, by

$$\sup_k \left(b_{k,q} \beta_{k,q}^{(q-1)} 2^{-k/q} |S_k| \right) = \infty . \quad (3.3.7)$$

Now for f_k defined as in (3.3.3)

$$S_k = \{n \in \mathbb{Z} : v_k \leq n \leq v_k + 2^k - 1\} ,$$

so each S_k is a finite set with cardinality $|S_k| = 2^k$.

Hence (3.3.7) becomes

$$\sup_k \{b_{k,q} \beta_{k,q}^{(q-1)} 2^{-k/q} 2^k\} = \infty. \quad (3.3.8)$$

Choose

$$b_{k,q} = (k+1)^{-1/q'}, \quad k = 0, 1, 2, \dots, \quad (3.3.9)$$

and

$$\beta_{k,q} = (k+1)^{2/q} 2^{-k/q}, \quad k = 0, 1, 2, \dots. \quad (3.3.10)$$

Then (3.3.5) is satisfied, since

$$\begin{aligned} \|(c_n)\|_{p'}^{p'} &= \sum_{k=0}^{\infty} (k+1)^{-p'/q'} (k+1)^{2(q-1)p'/q} 2^{-k(q-1)p'/q} 2^k \\ &= \sum_{k=0}^{\infty} (k+1)^{p'/q'} 2^{-k(p'/q'-1)} \end{aligned}$$

where $\|(c_n)\|_{p'}^{p'} < \infty$ for $p \in (1, q)$.

Also (3.3.8) is satisfied since

$$\begin{aligned} \sup_k \{ &(k+1)^{-1/q'} (k+1)^{2(q-1)/q} 2^{-k(q-1)/q} 2^{-k/q} 2^k \} \\ &= \sup_k [(k+1)^{1/q'}] \end{aligned} \quad (3.3.11)$$

$$= \infty \quad \text{for } q \in (1, 2].$$

Finally, the choice

$$v_k = 2^{k+1}, \quad k = 0, 1, 2, \dots,$$

ensures that the S_k are pairwise disjoint.

For the case $q \in (1, 2]$ and $p = 1$, with $\{c_n\}$, $\{\beta_{k,q}\}$ and $\{b_{k,q}\}$ as in (3.3.4), (3.3.9) and (3.3.10), it follows from the above argument that $\ell \notin (A^q)'$. It remains for us to check that $\{c_n\} \in \ell^\infty$ to ensure that $\ell \in (A^1)'$. Now

$$\sup_{n \in \mathbb{Z}} |c_n| = \sup_k \left((k+1)^{1/q'} 2^{-k/q'} \right),$$

$$\| \{c_n\} \|_{\ell^\infty} \leq 1,$$

so that $\{c_n\} \in \ell^\infty$, and our construction is completed. //

3.3.2 CONSTRUCTION. *The strict inclusion (3.3.2).*

Consider a given $q \in (1, 2)$. We wish to construct a linear functional, ℓ say, on the space TP , such that

$$\ell(f) = \sum_{n \in \mathbb{Z}} c_n \hat{f}(n), \text{ for all } f \in TP,$$

where $\{c_n\}$ is chosen so that ℓ is continuous in the topology induced by A^q , but is not continuous in the topology induced by A^p for every $p \in (q, 2]$.

Write

$$f_k = (k+1)^{-1} 2^{-k/q'} p_k e_{v_k}, \quad k = 0, 1, 2, \dots \quad (3.3.11)$$

With f_k as in (3.3.11) the $S_k = \text{sp}\{f_k\}$ are finite sets with cardinality $|S_k| = 2^k$. For the choice

$$v_k = 2^{k+1}, \quad k = 0, 1, 2, \dots,$$

the S_k are pairwise disjoint. In the case $q \in (1, 2)$ the condition $\{c_n\} \in \ell^{q'}$ is sufficient to ensure the continuity of ℓ in the topology

induced by A^q . Put

$$c_n = \begin{cases} (k+1)^{-1/q'} \hat{f}_k(n) & , n \in S_k , \\ 0 & , n \notin \bigcup_k S_k . \end{cases} \quad (3.3.12)$$

With this choice,

$$\begin{aligned} \|(c_n)\|_{q'}^{q'} &= \sum_{n \in \mathbb{Z}} |c_n|^{q'} \\ &= \sum_{k=0}^{\infty} (k+1)^{-1} \sum_{n \in S_k} |\hat{f}_k(n)|^{q'} \\ &= \sum_{k=0}^{\infty} (k+1)^{-(1+q')} \\ &< \infty . \end{aligned}$$

Also, with the choices in (3.3.11) and (3.3.12) and using (1.2.2) and (1.2.10),

$$\begin{aligned} \sup_k \frac{|\ell(f_k)|}{N_p(f_k)} &= \sup_k \left[\frac{(k+1)^{-1/q'} (k+1)^{-2} 2^{-2k/q'} 2^k}{(k+1)^{-1} 2^{-k/q'} 2^{k/p}} \right] \\ &= \sup_k \left[(k+1)^{-(1+1/q')} 2^{k(1-1/q'-1/p)} \right] \\ &= \infty \text{ for } p \in (q, 2] . \end{aligned}$$

We now consider $q = 1$. In this case $\{c_n\} \in \ell^\infty$ is a sufficient condition for ℓ to be continuous in the topology induced by $A^1 = A$. Choose

$$c_n = \begin{cases} \operatorname{sgn} \hat{f}_k(n) & , n \in S_k , \\ 0 & , n \notin \bigcup_k S_k , \end{cases}$$

where

$$f_k = P_k e_{v_k}, \quad k = 0, 1, 2, \dots \quad (3.3.13)$$

Here too the $S_k = sp\{f_k\}$ are finite sets with cardinality $|S_k| = 2^k$, and can be made pairwise disjoint by the choice

$$v_k = 2^{k+1}, \quad k = 0, 1, 2, \dots$$

To prove that ℓ is not continuous in the topology induced by

$$A^p, \quad p \in (1, 2], \quad \text{we need to show that} \quad \sup_k \frac{|\ell(f_k)|}{N_p(f_k)} = \infty \quad \text{for all } p \in (1, 2].$$

Using (1.2.2), (1.2.10) and (3.3.13) we have

$$\begin{aligned} \sup_k \frac{|\ell(f_k)|}{N_p(f_k)} &= \sup_k \frac{\left| \sum_{n \in \mathbb{Z}} c_n \hat{f}_k(n) \right|}{N_p(f_k)} \\ &= \sup_k \left(2^{k(1-1/p)} \right) \\ &= \infty \quad \text{for all } p \in (1, 2]; \end{aligned}$$

and our construction is completed. //

3.3.3 THEOREM. *The following strict inclusions hold.*

$$(A^q, A^q) \subsetneq \bigcap_{p \in [1, q)} (A^p, A^p) \quad \text{if } q \in (1, 2], \quad (3.3.14)$$

and

$$\bigcup_{p \in (q, 2]} (A^p, A^p) \subsetneq (A^q, A^q) \quad \text{if } q \in [1, 2). \quad (3.3.15)$$

Proof. Suppose $q \in (1, 2]$. Then, by Construction 3.3.1, there exists a linear functional ℓ on TP satisfying

$$\ell \in \bigcap_{p \in [1, q)} (A^p)' \text{ and } \ell \notin (A^q)' . \quad (3.3.16)$$

Let ϕ correspond to ℓ as in Remark 3.2.2. Then (3.3.16) gives us that

$$\phi \in (A^p, C) , \text{ for all } p \in [1, q) , \text{ and } \phi \notin (A^q, C) . \quad (3.3.17)$$

Combination of (3.3.17) with Lemma 3.2.1 yields (3.3.14).

Similarly, we can combine Construction 3.3.2, Remark 3.2.2 and Lemma 3.2.1 to obtain (3.3.15). //

3.4 The multipliers (A^p, A^q) , $p \in [1, 2]$, $q \in [p, 2]$.

The following theorem provides an identification of these spaces.

3.4.1 THEOREM. Suppose $p \in [1, 2]$ and $q \in [p, 2]$. Then

$$(A^p, A^q) = (A^p, A^p) = M + F\ell^{p'} .$$

Proof. Consider $p \in [1, 2]$, $q \in [p, 2]$. Since $A^p \subseteq A^q$ with continuous injection,

$$(A^p, A^q) \supseteq (A^p, A^p) . \quad (3.4.1)$$

Conversely, since $A^q \subseteq C$ with continuous injection,

$$(A^p, A^q) \subseteq (A^p, C) . \quad (3.4.2)$$

Thus, use of Lemma 3.2.1 with (3.4.2) gives

$$(A^p, A^q) \subseteq (A^p, A^p) . \quad (3.4.3)$$

We can now combine (3.4.1) and (3.4.3) to obtain

$$(A^p, A^q) = (A^p, A^p) . \quad (3.4.4)$$

The identification

$$(A^p, A^q) = M + F\ell^{p'}$$

follows directly from (3.4.4) and Theorem 3.2.5. //

3.5 The multipliers (A^p, A^q) , $p \in (1, 2]$, $q \in [1, p)$.

We begin by giving a sufficient condition for membership of (A^p, A^q) , $p \in (1, 2]$, $q \in [1, p)$. From now on we shall use M^k , $k \in [1, \infty]$ to denote the space of those measures whose Fourier transforms belong to ℓ^k . Note that we may make the identifications $M^1 = A$ and $M^2 = L^2$.

3.5.1 THEOREM. Suppose $p \in (1, 2]$ and $q \in [1, p)$. Then

$$M^{pq/(p-q)} \subseteq (A^p, A^q) .$$

Proof. Suppose $p \in (1, 2]$, $q \in [1, p)$ and consider $\mu \in M^{pq/(p-q)}$. Then, since $\mu \in M$, for all $f \in TP$,

$$\|\mu * f\|_\infty \leq \|\mu\|_M \|f\|_\infty . \quad (3.5.1)$$

Also Hölder's inequality gives, for every $f \in TP$,

$$\sum_{n \in \mathbb{Z}} |\hat{\mu}(n) \hat{f}(n)|^q \leq \left(\sum_{n \in \mathbb{Z}} |\hat{f}(n)|^{qs} \right)^{1/s} \left(\sum_{n \in \mathbb{Z}} |\hat{\mu}(n)|^{qs'} \right)^{1/s'} .$$

For $s = p/q$, $s' = p/(p - q)$ this becomes

$$\sum_{n \in \mathbb{Z}} |\hat{\mu}(n) \hat{f}(n)|^q \leq \left(\sum_{n \in \mathbb{Z}} |\hat{f}(n)|^p \right)^{q/p} \left(\sum_{n \in \mathbb{Z}} |\hat{\mu}(n)|^{pq/(p-q)} \right)^{(p-q)/p} . \quad (3.5.2)$$

By (3.5.1) and (3.5.2) we have, for $f \in TP$,

$$\begin{aligned} N_q(\mu * f) &= \|\mu * f\|_\infty + \|(\mu * f)^\wedge\|_q \\ &\leq \|\mu\|_M \|f\|_\infty + \|\hat{\mu}\|_{pq/(p-q)} \|\hat{f}\|_p \\ &\leq \text{const. } N_p(f) . \end{aligned}$$

Appealing to (3.1.1) we can conclude that $\mu \in (A^p, A^q)$. //

We shall give now a necessary condition for membership of (A^2, A^q) , $q \in [1, 2)$. This condition, together with Theorem 3.5.1, will then be used in the identification of the space $(A^2, A^1) = (C, A)$.

3.5.2 THEOREM. Suppose $q \in [1, 2)$. Then

$$\phi \in (A^2, A^q) \Rightarrow \hat{\phi} \in \ell^{2q/(2-q)} .$$

Proof. If $\phi \in (A^2, A^q)$ then, since $A^2 = C$,

$$\begin{aligned} \|\hat{\phi} \cdot \hat{f}\|_q &\leq N_q(\phi * f) \\ &\leq \text{const. } \|f\|_\infty , \text{ for all } f \in C . \end{aligned} \quad (3.5.3)$$

From [8], Corollary 2.3, p.468 it follows that if $\hat{\phi} \cdot \hat{f} \in \ell^q$, $q \in [1, 2)$, for each $f \in C$ then $\hat{\phi} \in \ell^{2q/(2-q)}$. Combination of this with (3.5.3) gives our result. //

3.5.3 THEOREM. $(C, A) = L^2$.

Proof. By Theorem 3.5.1 for the case $p = 2$, $q = 1$,

$$L^2 = M^2 \subseteq (A^2, A^1) = (C, A) .$$

Conversely, suppose $\phi \in (C, A)$. Then, by Theorem 3.5.2, $\hat{\phi} \in \ell^2$, and so $\phi \in L^2$. Thus $(C, A) \subseteq L^2$; this completes our proof. //

We now establish preliminary results leading to necessary conditions for $\phi \in (A^p, A^q)$, $p \in (1, 2]$, $q \in [1, p)$.

For $p \in (1, 2]$, $\varepsilon > 0$, consider the space $S = S(p, \varepsilon)$ defined by

$$S = \{ (c_n)_{n \in \mathbb{Z}} \in \ell^p : \sum_{n \in \mathbb{Z} \setminus \{0\}} |c_n|^2 \log^{1+\varepsilon} |n| < \infty , \varepsilon > 0 \} . \quad (3.5.4)$$

Then, from [5], 14.3.6, p.205, for $(c_n) \in S$ almost all the series

$$\sum_{n \in \mathbb{Z}} r |n|^{(t)} c_n e_n$$

are the Fourier series of continuous functions (in fact, of functions in A^p). If $\phi \in (A^p, A^q)$ then, whenever $f \in A^p$, $\phi * f \in A^q$ and hence

$$\sum_{n \in \mathbb{Z}} |\hat{\phi}(n) c_n|^q < \infty , \text{ for all } (c_n) \in S . //$$

3.5.4 LEMMA. Let $p \in (1, 2]$. Then S , defined as in (3.5.4), is a Banach space under the norm

$$\| \cdot \|_S : (c_n) \mapsto \left(\sum_{n \in \mathbb{Z}} |c_n|^p \right)^{1/p} + \left(\sum_{n \in \mathbb{Z} \setminus \{0\}} |c_n|^2 \log^{1+\varepsilon} |n| \right)^{1/2} .$$

Proof. S is clearly a linear space. We need to show that under the above norm S is complete. To this end consider a sequence $(c^{(k)})_{k=1}^\infty$

which is Cauchy in S . Then, given $\delta > 0$, there exists $K > 0$ such that, for all $r, m > K$,

$$\left(\sum_{n \in \mathbb{Z}} |c_n^{(r)} - c_n^{(m)}|^p \right)^{1/p} < \delta \quad (3.5.5)$$

and

$$\left(\sum_{n \in \mathbb{Z} \setminus \{0\}} |c_n^{(r)} - c_n^{(m)}|^2 \log^{1+\epsilon} |n| \right)^{1/2} < \delta. \quad (3.5.6)$$

From (3.5.5) and the completeness of ℓ^p the sequence $(c^{(k)})_{k=1}^{\infty}$ has a limit, $c = (c_n)$ say in ℓ^p . Thus

$$\lim_{k \rightarrow \infty} \left| c_n^{(k)} \log^{\frac{1}{2}(1+\epsilon)} |n| - c_n \log^{\frac{1}{2}(1+\epsilon)} |n| \right| = 0, \text{ for all } n \in \mathbb{Z} \setminus \{0\}. \quad (3.5.7)$$

Now write $b^{(k)} = \left(c_n^{(k)} \log^{\frac{1}{2}(1+\epsilon)} |n| \right)_{n \in \mathbb{Z} \setminus \{0\}}$, $k = 1, 2, 3, \dots$. By (3.5.6)

and the completeness of ℓ^2 , the sequence $(b^{(k)})_{k=1}^{\infty}$ has a limit, $b = (b_n)$ say, in ℓ^2 . Hence

$$\lim_{k \rightarrow \infty} \left| c_n^{(k)} \log^{\frac{1}{2}(1+\epsilon)} |n| - b_n \right| = 0, \text{ for all } n \in \mathbb{Z} \setminus \{0\}. \quad (3.5.8)$$

Combination of (3.5.7) and (3.5.8) gives

$$b_n = c_n \log^{\frac{1}{2}(1+\epsilon)} |n|, \text{ for all } n \in \mathbb{Z} \setminus \{0\}. \quad (3.5.9)$$

Thus $c = (c_n)$ belongs to S and $\lim_{k \rightarrow \infty} \|c^{(k)} - c\|_S = 0$; therefore S is complete. //

3.5.5 LEMMA. Suppose $p \in (1, 2]$ and $q \in [1, p)$. Then for $\phi \in (A^p, A^q)$, the map $Q_\phi : S \rightarrow \ell^q$ defined by

$$Q_\phi : (c_n) \rightarrow \{\hat{\phi}(n)c_n\} \quad (3.5.10)$$

is continuous; hence for $K = K(\phi, \epsilon)$,

$$\left(\sum_{n \in \mathbb{Z}} |\hat{\phi}(n)c_n|^q \right)^{1/q} \leq K \left[\left(\sum_{n \in \mathbb{Z}} |c_n|^p \right)^{1/p} + \left(\sum_{n \in \mathbb{Z} \setminus \{0\}} |c_n|^2 \log^{1+\epsilon} |n| \right)^{1/2} \right]. \quad (3.5.11)$$

Proof. Suppose $\phi \in (A^p, A^q)$. The map Q_ϕ defined in (3.5.10) is clearly linear. Also S and ℓ^q are Banach spaces; we shall apply the closed graph theorem to show the continuity of Q_ϕ . To do this, consider a sequence $(c^{(k)})_{k=1}^\infty$ from S for which

$$\lim_{k \rightarrow \infty} \|c^{(k)} - c\|_S = 0 \quad (3.5.12)$$

and

$$\lim_{k \rightarrow \infty} \|Q_\phi c^{(k)} - b\|_q = 0, \quad (3.5.13)$$

where $c = (c_n)$ and $b = (b_n)$. From (3.5.12),

$$\lim_{k \rightarrow \infty} \|c^{(k)} - c\|_p = 0; \text{ hence}$$

$$\lim_{k \rightarrow \infty} |\hat{\phi}(n)c_n^{(k)} - \hat{\phi}(n)c_n| = 0, \text{ for all } n \in \mathbb{Z}. \quad (3.5.14)$$

From (3.5.13),

$$\lim_{k \rightarrow \infty} |\hat{\phi}(n)c_n^{(k)} - b_n| = 0, \text{ for all } n \in \mathbb{Z}; \quad (3.5.15)$$

and combination of (3.5.14) and (3.5.15) gives

$$b_n = \hat{\phi}(n) c_n, \text{ for all } n \in \mathbb{Z}.$$

Thus the graph of Q_ϕ is closed in $S \times \ell^2$ and, by the closed graph theorem, Q_ϕ is a continuous map from S to ℓ^q ; hence (3.5.11) holds. //

We are now in a position to establish the following result.

3.5.6 THEOREM. Suppose $p \in (1, 2]$ and $q \in [1, p)$. If $\phi \in (A^p, A^q)$ then, for every $\varepsilon > 0$ and every positive integer r ,

$$\sum_{|n| \leq r} |\hat{\phi}(n)|^{pq/(p-q)} \leq \max \left\{ 1, (K(1 + J_r))^{pq/(p-q)} \right\},$$

where $K = K(\phi, \varepsilon)$ is independent of r and

$$J_r^2 = \max_{0 < |n| \leq r} \left\{ |\hat{\phi}(n)|^{(2-p)q/(p-q)} \log^{1+\varepsilon} |n| \right\}. \quad (3.5.16)$$

Proof. Suppose $\phi \in (A^p, A^q)$. Then (3.5.11) holds. Choose $(c_n) \in S$ such that

$$c_n = \begin{cases} |\hat{\phi}(n)|^{q/(p-q)}, & |n| \leq r, \\ 0, & \text{otherwise} \end{cases}$$

and consider J_r as defined in (3.5.16). Then, on writing

$$\sigma_r = \sum_{|n| \leq r} |\hat{\phi}(n)|^{pq/(p-q)}, \quad (3.5.11) \text{ yields}$$

$$\sigma_r^{1/q} \leq K(\sigma_r^{1/p} + J_r \sigma_r^{1/2});$$

(3.5.16). Application of Theorem 3.5.6 now gives the desired result. //

that is,

$$\sigma_r^{1/q-1/p} \leq K(1 + J_r \sigma_r^{1/2-1/p}) . \quad (3.5.17)$$

Now (3.5.17) is equivalent to

$$\sigma_r \leq \left[K(1 + J_r \sigma_r^{-(2-p)/2p}) \right]^{pq/(p-q)}$$

which, for $\sigma_r \geq 1$, yields

$$\sigma_r \leq \left[K(1 + J_r) \right]^{pq/(p-q)} .$$

Hence it follows that

$$\sigma_r \leq \max \left[1, (K(1 + J_r))^{pq/(p-q)} \right] . \quad //$$

We shall now use Theorem 3.5.6 to establish the following two necessary conditions for membership of (A^p, A^q) , $p \in (1, 2]$, $q \in [1, p)$.

3.5.7 COROLLARY. Suppose $p \in (1, 2]$ and $q \in [1, p)$. If $\phi \in (A^p, A^q)$ then

$$\sum_{|n| \leq r} |\hat{\phi}(n)|^{pq/(p-q)} = o(\log^\Delta r) ,$$

where

$$\Delta = \frac{1}{2}(1 + \epsilon)pq/(p-q) , \text{ and } \epsilon > 0 .$$

Proof. Suppose $\phi \in (A^p, A^q)$. Then $\phi \in (A, C) = PM$, and it follows that $J_r = o((\log r)^{\frac{1}{2}(1+\epsilon)})$, where J_r is defined as in (3.5.16). Application of Theorem 3.5.6 now gives the desired result. //

POINTWISE MULTIPLIERS OF A^p , $p \in [1, 2]$

3.5.8 COROLLARY. Suppose $p \in (1, 2]$ and $q \in [1, p)$. If $\phi \in (A^p, A^q)$ and $J_r = o(1)$, where J_r is defined as in (3.5.16), then $\hat{\phi} \in \ell^{pq/(p-q)}$.

Proof. This result follows directly from Theorem 3.5.6. //

In this chapter we shall be concerned with pointwise multipliers from A^p to A^q ; that is, with integrable functions f for which $f \cdot A^p \subseteq A^q$. Following the notation of 1.3.7, we shall denote the space of such multipliers by $M(A^p, A^q)$ and, when $p = q$, abbreviate this to $M(A^p)$. It is not difficult to check that $M(A^p)$ is a Banach algebra under the norm $\|f\|_{M(p,p)} = \|f\|_{M(p,p)}$, where $\|f\|_{M(p,p)}$ is given by (1.3.2). Note here that we can make the identifications $M(A^1) = A$ and $M(A^2) = \mathbb{C}$.

4.2 Some inclusion results for pointwise multipliers

It is clear that $M(A^p, A^q) \subseteq A^q$, for all $p, q \in [1, 2]$; and that $A \subseteq M(A^p) \subseteq M(A^p, A^q)$ whenever $p \in [1, q]$. In particular, with continuous injections,

$$A \subseteq M(A^p) \subseteq A^p, \quad \text{for all } p \in (1, 2]. \quad (4.2.1)$$

We shall show that, with the appropriate restrictions on p , each of the inclusion results in (4.2.1) is strict. Actually, we shall prove slightly more than this (see Construction 4.2.3 and the following statement).

CHAPTER 4

POINTWISE MULTIPLIERS OF A^p , $p \in [1, 2]$

4.1 Introduction

In Chapter 3 we dealt with Fourier multipliers from A^p to A^q . These, as we saw there, are multipliers under *convolution*. In this chapter we shall be concerned with *pointwise* multipliers from A^p to A^q ; that is, with integrable functions f for which $f \cdot A^p \subseteq A^q$. Following the notation of 1.3.7, we shall denote the space of such multipliers by $M(A^p, A^q)$ and, when $p = q$, abbreviate this to $M(A^p)$. It is not difficult to check that $M(A^p)$ is a Banach algebra under the norm $f \mapsto \|f\|_{M(p,p)}$, where $\|f\|_{M(p,p)}$ is given by (1.3.2). Note here that we can make the identifications $M(A^1) = A$ and $M(A^2) = \mathbb{C}$.

4.2 Some inclusion results for pointwise multipliers

It is clear that $M(A^p, A^q) \subseteq A^q$, for all $p, q \in [1, 2]$; and that $A \subseteq M(A^p) \subseteq M(A^p, A^q)$ whenever $p \in [1, q]$. In particular, with continuous injections,

$$A \subseteq M(A^p) \subseteq A^p, \quad \text{for all } p \in [1, 2]. \quad (4.2.1)$$

We shall show that, with the appropriate restrictions on p , each of the inclusion results in (4.2.1) is strict. Actually, we shall prove slightly more than this (see Construction 4.2.3 and the following statement).

The strict inclusion $M(A^p) \subsetneq A^p$, $p \in (1,2)$, follows directly from 2.2. In fact, it was shown there that there exists $f \in A^\# \setminus M(A^p)$, for all $p \in (1,2)$; hence the strict inclusion $\bigcap_{1 < p < 2} M(A^p) \subsetneq A^\#$.

To show the strict inclusion $A \subsetneq M(A^p)$, $p \in (1,2]$, we shall make use of Corollary 1.3.5 for the case $G = \mathbb{Z}$ with the covering family indicated in Remark 1.3.6. This result, in the form we actually require, is contained in the following corollary.

4.2.1 COROLLARY. Suppose $p \in (1,2]$, $f \in C$ and that there exists a sequence $(k_v)_{v=1}^\infty$, with each $k_v \in \ell^1$, such that

$$\kappa = \sup_v \max \left[J(k_v), \|\hat{k}_v\|_\infty \right] < \infty$$

and

$$f = \lim_{v \rightarrow \infty} \hat{k}_v \text{ a.e. on } \mathbb{T},$$

where

$$J(k_v) = \sup_v \sup_{0 < |m| < 2^\beta} \sum_{|n| \geq 2^{\beta+1}} |k_v(n-m) - k_v(n)|.$$

Then $f \in M_p$.

4.2.2 CONSTRUCTION. The strict inclusion $A \subsetneq \bigcap_{p>1} M(A^p)$.

Consider any function $f \in C \setminus A$ satisfying the following conditions:

$$\hat{f}(n) \in \mathbb{R}, \text{ for all } n \in \mathbb{Z}. \quad (4.2.2)$$

$$\hat{f}(n) = -\hat{f}(-n), \text{ for all } n \in \mathbb{Z}. \quad (4.2.3)$$

$$\hat{f}(n) \rightarrow 0 \text{ as } n \rightarrow +\infty. \quad (4.2.4)$$

$$n\hat{f}(n) \rightarrow 0 \text{ as } n \rightarrow +\infty. \quad (4.2.5)$$

Note here that the set of all such f is non-empty; an example of such a function is $f = \hat{g}$, where

$$g(n) = \begin{cases} \frac{1}{n \log |n|} & , n \neq 0, \pm 1, \\ n & , n = 0, \pm 1. \end{cases}$$

By [5], 7.2.2, conditions (4.2.3) - (4.2.5) ensure that $f \in \mathcal{U}$.

For $v = 1, 2, \dots$, write

$$k_v(n) = \hat{f}(n) \xi_{[-v, v]}(n).$$

Then $k_v \in \ell^1$ and $\hat{k}_v = f * D_v = s_v f \rightarrow f$ pointwise on $\mathbb{T}$.

Now consider

$$J_\beta(k_v) = \sup_{0 < |m| < 2^\beta} \sum_{|n| \geq 2^{\beta+1}} |k_v(n-m) - k_v(n)|.$$

Fix v and note that the conditions (4.2.2), (4.2.3) and (4.2.4) ensure that $\sum_{|n| \geq 2^{\beta+1}} |k_v(n-m) - k_v(n)|$ is even in m . Thus for each positive integer β we need consider only integers $m \in (0, 2^\beta)$. Then

$$\begin{aligned} & \sum_{|n| \geq 2^{\beta+1}} |k_v(n-m) - k_v(n)| \\ &= \sum_{|n| \geq 2^{\beta+1}} |\hat{f}(n-m) \xi_{[-v, v]}(n-m) - \hat{f}(n) \xi_{[-v, v]}(n)| \end{aligned}$$

$$\begin{aligned}
&= \sum_{n \geq 2^{\beta+1}} (\hat{f}(n-m) \xi_{[-v, v]}(n-m) - \hat{f}(n) \xi_{[-v, v]}(n)) \\
&\quad + \sum_{n' \geq 2^{\beta+1}} |\hat{f}(-n'-m) \xi_{[-v, v]}(-n'-m) - \hat{f}(-n) \xi_{[-v, v]}(-n)| \\
&= \sum_{n \geq 2^{\beta+1}} (\hat{f}(n-m) \xi_{[-v, v]}(n-m) - \hat{f}(n) \xi_{[-v, v]}(n)) \\
&\quad + \sum_{n \geq 2^{\beta+1}} (-\hat{f}(n+m) \xi_{[-v, v]}(n+m) + \hat{f}(n) \xi_{[-v, v]}(n)) \\
&= \sum_{n \geq 2^{\beta+1}} (\hat{f}(n-m) \xi_{[-v, v]}(n-m) - \hat{f}(n+m) \xi_{[-v, v]}(n+m)) \\
&= \sum_{n=2^{\beta+1}-m}^v \hat{f}(n) - \sum_{n=2^{\beta+1}+m}^v \hat{f}(n) ,
\end{aligned}$$

where an empty sum is defined to be zero. Thus

$$\begin{aligned}
\sum_{|n| \geq 2^{\beta+1}} |k_v(n-m) - k_v(n)| &= \sum_{n=2^{\beta+1}-m}^{2^{\beta+1}+m-1} \hat{f}(n) \\
&= \sum_{j=0}^{2m-1} \hat{f}(2^{\beta+1}-m+j) .
\end{aligned}$$

By conditions (4.2.3) and (4.2.4), $\hat{f}(2^{\beta+1}-m+j)$ is positive and increases with m , so that

We are now in a position to apply Corollary 4.2.1 to conclude that

$$f * g \in \mathcal{P}^p, \text{ for all } g \in \mathcal{P}^p ;$$

and it follows that $f \in \mathcal{P}_1^{p, N(\mathcal{P})} \setminus \mathcal{A} . \quad \square$

$$\begin{aligned}
J_\beta(k_v) &\leq \sum_{j=0}^{2(2^\beta-1)-1} \hat{f}(2^{\beta+1} - 2^\beta + 1 + j) \\
&= \sum_{j=0}^{2^{\beta+1}-3} \hat{f}(2^\beta + 1 + j) \\
&\leq (2^{\beta+1} - 2) \hat{f}(2^\beta + 1) \quad (\text{using (4.2.4)}) \\
&< 2(2^\beta + 1) \hat{f}(2^\beta + 1).
\end{aligned}$$

Use of (4.2.5) now gives

$$J(k_v) = \sup_\beta J_\beta(k_v) \leq K < \infty,$$

where K is a constant which is independent of v ; and so

$$\sup_v J(k_v) < \infty. \quad (4.2.6)$$

In addition, for each v ,

$$\|\hat{k}_v\|_\infty = \|f * D_v\|_\infty = \|S_v f\|_\infty$$

and, since $f \in \mathcal{U}$,

$$\sup_v \|\hat{k}_v\|_\infty < \infty. \quad (4.2.7)$$

Combination of (4.2.6) and (4.2.7) gives

$$\sup_v \max \left\{ J(k_v), \|\hat{k}_v\|_\infty \right\} < \infty.$$

We are now in a position to apply Corollary 4.2.1 to conclude that

$$\hat{f} * g \in \ell^p, \text{ for all } g \in \ell^p;$$

and it follows that $f \in \bigcap_{p>1} M(A^p) \setminus A$. //

4.2.3 REMARKS . The functions constructed in 4.2.2 all belong to $A^\# \setminus A$.

²The question of whether or not $M(A^p) \subseteq M(A^q)$, for $1 < p < q < 2$, is still open. However we can give an example to show that any such inclusion must be strict.

4.2.4 THEOREM. Suppose $p \in [1, 2]$. Then

$$\bigcap_{q \in (p, 2]} M(A^q) \setminus M(A^p) \neq \emptyset .$$

Proof. Fix $p \in [1, 2]$ and put $\alpha = (2-p)/2p$. Now write

$$f = \sum_{n=1}^{\infty} e^{in \log n} n^{-\frac{1}{2}-\alpha} e_n . \quad (4.2.8)$$

The series defining f was first studied by Hardy and Littlewood in [14]. That $f \in \Lambda_\alpha$ is proved in [34], Chapter V, (4.9).

Application of a result of Hirschman (Theorem 2a of [18]) now gives $f \in M_q$ for all $q \in (p, 2]$; hence, since $f \in C$, it follows that $f \in M(A^q)$, for all $q \in (p, 2]$. However, using (4.2.8),

$$\sum_{n \in \mathbb{Z}} |\hat{f}(n)|^p = \sum_{n=1}^{\infty} |e^{in \log n} n^{-\frac{1}{2}-\alpha}|^{2/(1+2\alpha)}$$

$$= \sum_{n=1}^{\infty} n^{-1}$$

$$= \infty ;$$

thus $f \notin A^p$ and so, by (4.2.1), $f \notin M(A^p)$. //

4.2.5 We shall now indicate some inclusion results concerning $M(A^p, A^q)$, $1 \leq p < q \leq 2$. Application of Theorem 2.6.7 and Corollary 2.6.6 respectively give that

Now consider $A^p \subseteq M(A^p, A^q)$ only if $q \geq 3 - 2/p$

and

$$A^p \subseteq M(A^p, A^q) \text{ if } q \geq p^2/(2 + p^2 - 2p) .$$

$$4.3 \quad M(A^p, A^q) = \{0\} , \text{ if } 2 \geq p > q \geq 1 .$$

In order to make the above identification we first prove the following result. In so doing we shall make use of Lemma 1.3.8.

4.3.1 THEOREM. If $p, q \in [1, 2]$ then

$$L^1 * M(A^p, A^q) \subseteq M(A^p, A^q) ;$$

and, for $k \in L^1$, $f \in M(A^p, A^q)$,

$$\|k * f\|_{M(p, q)} \leq \|k\|_1 \|f\|_{M(p, q)} . \quad (4.3.1)$$

Proof. Consider $k \in L^1$, $f \in M(A^p, A^q)$ and $g \in TP$. Then

$$(k * f)g = \int_T k(y) \tau_y f \cdot g \, dy$$

and it follows from this that

$$\|(k * f)g\|_\infty \leq \int_T |k(y)| \|\tau_y f \cdot g\|_\infty \, dy . \quad (4.3.2)$$

Also, using the Fubini-Tonelli theorem,

$$\begin{aligned} \widehat{(k * f)g}(n) &= \int_T \left(\int_T k(y) \tau_y f(x) g(x) \, dy \right) e^{-inx} dx \\ &= \int_T k(y) \left(\int_T \tau_y f(x) g(x) e^{-inx} dx \right) dy \\ &= \int_T k(y) (\tau_y f \cdot g)^\wedge(n) \, dy . \end{aligned}$$

Thus (4.3.4) yields

Now consider $h \in \ell^{q'}$. Then

$$\begin{aligned}
 |((k * f)g)^\wedge * \check{h}(0)| &= \left| \sum_{n \in \mathbb{Z}} ((k * f)g)^\wedge(n) h(n) \right| \\
 &= \left| \sum_{n \in \mathbb{Z}} \left[\int_{\mathbb{T}} k(y) (\tau_y f \cdot g)^\wedge(n) dy \right] h(n) \right| \\
 &= \left| \int_{\mathbb{T}} k(y) \left[\sum_{n \in \mathbb{Z}} (\tau_y f \cdot g)^\wedge(n) h(n) \right] dy \right| \\
 &= \left| \int_{\mathbb{T}} k(y) (\tau_y f \cdot g)^\wedge * \check{h}(0) dy \right| \\
 &\leq \int_{\mathbb{T}} |k(y)| |(\tau_y f \cdot g)^\wedge * \check{h}(0)| dy ;
 \end{aligned}$$

and hence

$$\|((k * f)g)^\wedge\|_q \leq \int_{\mathbb{T}} |k(y)| \|(\tau_y f \cdot g)^\wedge\|_q dy . \quad (4.3.3)$$

From (4.3.2) and (4.3.3),

$$\begin{aligned}
 N_q((k * f)g) &= \|(k * f)g\|_\infty + \|((k * f)g)^\wedge\|_q \\
 &\leq \int_{\mathbb{T}} |k(y)| \|\tau_y f \cdot g\|_\infty dy + \int_{\mathbb{T}} |k(y)| \|(\tau_y f \cdot g)^\wedge\|_q dy \\
 &= \int_{\mathbb{T}} |k(y)| N_q(\tau_y f \cdot g) dy \\
 &\leq \int_{\mathbb{T}} |k(y)| \|\tau_y f\|_{M(p,q)} N_p(g) dy . \quad (4.3.4)
 \end{aligned}$$

However, if $f \in M(A^p, A^q)$ then $\tau_y f \in M(A^p, A^q)$ and

$$\|\tau_y f\|_{M(p,q)} = \|f\|_{M(p,q)} , \quad \text{for all } y \in \mathbb{T} .$$

Thus (4.3.4) yields

$$N_q((k * f)g) \leq \|k\|_1 \|f\|_{M(p,q)} N_p(g)$$

$< \infty$;

so $k * f \in M(A^p, A^q)$ and (4.3.1) follows immediately. //

4.3.2 THEOREM. $M(A^p, A^q) = \{0\}$, if $2 \geq p > q \geq 1$.

Proof. Let $2 \geq p > q \geq 1$ and suppose $f \in M(A^p, A^q)$.

From Theorem 4.3.1, $e_n * f \in M(A^p, A^q)$ for each $n \in \mathbb{Z}$,

and hence

$$N_q((e_n * f)g) < \infty, \text{ for all } g \in A^p. \quad (4.3.5)$$

It then follows that $f = 0$; since, if $f \neq 0$ then there is at least one $n \in \mathbb{Z}$ for which $\hat{f}(n) \neq 0$ and, for all $g \in A^p$,

$$\begin{aligned} N_q((e_n * f)g) &= N_q(\hat{f}(n)e_n g) \\ &= |\hat{f}(n)| N_q(e_n g) \\ &= |\hat{f}(n)| N_q(g), \text{ by (1.2.2)}. \end{aligned}$$

But this contradicts (4.3.5) since, by Construction 2.5.1 ,

$$A^q \subsetneq A^p. //$$

It is interesting to note that Theorem 4.3.1 can be strengthened to give the following result.

4.3.3 THEOREM. Suppose $p, q \in [1, 2]$. Then $L^1 * M(A^p, A^q) = M(A^p, A^q)$ if and only if $M(A^p, A^q)$ is translation-continuous.

Proof. (i) "if". We have already, from Theorem 4.3.1, that for $p, q \in [1, 2]$,

$$L^1 * M(A^p, A^q) \subseteq M(A^p, A^q)$$

and (4.3.1) holds. Suppose now that $M(A^p, A^q)$ is translation-continuous. It then satisfies all the conditions for a homogeneous Banach space in the sense of [23], Chapter I, Definition 2.10; thus [23], Chapter I, Theorem 2.11 ensures that, for $f \in M(A^p, A^q)$,

$$\lim_{N \rightarrow \infty} \|F_N * f - f\|_{M(p,q)} = 0.$$

From the argument given in [5], 7.5.1 we now have that

$$M(A^p, A^q) \subseteq L^1 * M(A^p, A^q).$$

(ii) "only if". Suppose that $L^1 * M(A^p, A^q) = M(A^p, A^q)$. Then for each $g \in M(A^p, A^q)$ there exist $k \in L^1$ and $f \in M(A^p, A^q)$ satisfying $k * f = g$. Thus, for $\alpha \in \mathbb{T}$,

$$\begin{aligned} \|\tau_\alpha g - g\|_{M(p,q)} &= \|\tau_\alpha(k * f) - k * f\|_{M(p,q)} \\ &= \|(\tau_\alpha k) * f - k * f\|_{M(p,q)} \\ &\leq \|\tau_\alpha k - k\|_1 \|f\|_{M(p,q)}, \end{aligned} \quad (4.3.6)$$

where the final step follows from (4.3.1). By (4.3.6) and the translation-continuity of L^1 we have

$$\lim_{\alpha \rightarrow 0} \|\tau_\alpha g - g\|_{M(p,q)} = 0,$$

as required. //

4.4 Some sufficient conditions for membership of $M(A^p)$.

We note here that the case $p = q$ in Theorem 4.3.3 gives us that $L^1 * M(A^p) = M(A^p)$ if and only if $M(A^p)$ is translation-continuous. ³We shall give some further results involving translation-continuity of functions in $M(A^p)$. We first prove the following lemma.

4.4.1 LEMMA. Suppose $p \in (1, 2)$. If $f \in M(A^p)$ and

$$\lim_{\alpha \rightarrow 0} \|\tau_\alpha f - f\|_{M(p,p)} = 0, \quad (4.4.1)$$

then $f \in [TP]_{M(p,p)}$, where $[TP]_{M(p,p)}$ denotes the closure in $M(A^p)$ of the space of trigonometric polynomials.

Proof. By (4.4.1), given $\varepsilon > 0$ there exists $\delta \in (0, \pi)$ such that whenever $|a| < \delta$,

$$\|\tau_a f - f\|_{M(p,p)} < \frac{1}{2}\varepsilon. \quad (4.4.2)$$

Now put $g = \frac{1}{2\delta} \chi_{[-\delta, \delta]}$. Then, using the continuity of $x \rightarrow \tau_x f$, we have

$$f * g - f = \int_T (\tau_y f - f) g(y) dy, \text{ an } M(A^p) \text{ - integral;}$$

and so

$$\begin{aligned} \|f * g - f\|_{M(p,p)} &\leq \int_T \|\tau_y f - f\|_{M(p,p)} g(y) dy \\ &< \frac{1}{2}\varepsilon. \end{aligned} \quad (4.4.3)$$

In addition we can find $t \in TP$ for which

$$\|f\|_{M(p,p)} \|g - t\|_1 < \frac{1}{2}\varepsilon.$$

Thus there exists $t \in TP$ satisfying

$$\begin{aligned} \|f - f * t\|_{M(p,p)} &\leq \|f - f * g\|_{M(p,p)} + \|f * g - f * t\|_{M(p,p)} \\ &< \frac{1}{2}\varepsilon + \|f\|_{M(p,p)} \|g - t\|_1 \\ &< \varepsilon. \end{aligned}$$

Since $f * t \in TP$ whenever $t \in TP$ it follows that

$$f \in [TP]_{M(p,p)}. //$$

4.4.2 THEOREM. Let $p \in [1, 2]$. If $f \in M(A^p)$ does not vanish on T and satisfies (4.4.1), then $f^{-1} \in M(A^p)$.

Proof. Note that $N_p(\frac{1}{2}e_n) = 1$ for all $n \in \mathbb{Z}$,
so by (1.2.2) and the definition of $\|\cdot\|_{M(p,p)}$,

$$\|e_n\|_{M(p,p)} = 1, \text{ for all } n \in \mathbb{Z}.$$

In particular,

$$\|1\|_{M(p,p)} = \|e_{-1}\|_{M(p,p)} = \|e_1\|_{M(p,p)} = 1$$

and so, arguing as in [23], Chapter VIII, 2.9, for any non-zero homomorphism ω of $M(A^p)$ into $\mathbb{C}$ there exists $x_0 \in T$ such that

$$\omega(t) = t(x_0), \text{ for all } t \in TP.$$

Continuity of ω (see [23], Chapter VIII, Lemma 2.8) then gives

$$\omega(f) = f(x_0), \text{ for all } f \in [TP]_{M(p,p)}. \quad (4.4.3)$$

Application of Lemma 4.4.1 shows that (4.4.3) holds for all the $f \in M(A^p)$ which satisfy (4.4.1). If, in addition, f does not vanish on T we can use [23], Chapter VIII, Corollary 2.8 to conclude that $f^{-1} \in M(A^p)$. //

4.4.3 THEOREM. Suppose $p \in (1,2)$, $f \in M(A^p)$ and that there exists a set $S \subseteq T$ with strictly positive interior measure such that $\{\tau_\alpha f : \alpha \in S\}$ is separable relative to $\|\cdot\|_{M(p,p)}$. Then the function $x \rightarrow \tau_x f$ is continuous from T to $M(A^p)$.

Proof. In view of the inner regularity of Lebesgue measure, we may assume that S is compact. Now $\{\tau_\alpha f : \alpha \in S\}$ is separable so that, given $\varepsilon > 0$, there exists $(x_n)_{n=1}^\infty$ from S such that

$$S = \bigcup_{n=1}^\infty \{x \in S : \|\tau_x f - \tau_{x_n} f\|_{M(p,p)} \leq \varepsilon\}.$$

Write

$$S_n = \{x \in S : \|\tau_x f - \tau_{x_n} f\|_{M(p,p)} \leq \varepsilon\}.$$

Now for each $\alpha \in T$, $g \in TP$, the function $x \rightarrow N_p((\tau_x f - \tau_\alpha f)g)$ is continuous; just use

$$\begin{aligned} |N_p((\tau_x f - \tau_\alpha f)g) - N_p((\tau_y f - \tau_\alpha f)g)| &\leq N_p((\tau_x f - \tau_y f)g) \\ &\leq N_p(\tau_x f - \tau_y f)N_1(g) \end{aligned}$$

and Theorem 2.6.9. It follows that $x \rightarrow \|\tau_x f - \tau_y f\|_{M(p,p)}$ is lower semicontinuous, so that each S_n is closed, and hence compact. Since S has strictly positive measure the same is true of S_n for at least one integer $n = n(\varepsilon)$. Now write

$$W_\varepsilon = \{x \in T : \|\tau_x f - \tau_{x_{n(\varepsilon)}} f\|_{M(p,p)} \leq \varepsilon\}.$$

As $W_\varepsilon \supseteq S_{n(\varepsilon)}$, W_ε has strictly positive interior measure. It then follows (see [16], (20.17)) that $W_\varepsilon - W_\varepsilon$ is a neighbourhood of 0. Also, for $x, y \in W_\varepsilon$,

$$\begin{aligned} \|\tau_{x-y} f - f\|_{M(p,p)} &= \|\tau_x f - \tau_y f\|_{M(p,p)} \\ &\leq \|\tau_x f - \tau_{x_{n(\varepsilon)}} f\|_{M(p,p)} + \|\tau_{x_{n(\varepsilon)}} f - \tau_y f\|_{M(p,p)} \\ &\leq 2\varepsilon, \end{aligned}$$

from which we can conclude that $x \rightarrow \tau_x f$ is continuous at zero; and hence, since $\|\cdot\|_{M(p,p)}$ is translation-invariant, continuous at each $x \in T$.

We shall adapt some results in [18] to give further sufficient conditions for membership of $M(A^p)$. Also, we shall derive these

conditions in a way which enables us to deduce that the

$f \in M(A^p)$ so obtained are actually translation-continuous.

Note here that $C \cap M_p \subseteq M(A^p)$ with a continuous injection, so that

$$\|\cdot\|_{M(p,p)} \leq \|\cdot\|_{\infty} + \|\cdot\|_{p,p}. \quad (4.4.4)$$

4.4.4 THEOREM. Suppose $p \in (1,2)$, $q \in (1,p)$, $\alpha \in \left[\frac{2-q}{q}, \frac{1}{2}\right]$ and $f \in \Lambda_{\alpha}$. Then $f \in M(A^p) \cap M(A^q)$ and

$$\lim_{\alpha \rightarrow 0} \|\tau_{\alpha} f - f\|_{M(p,p)} = 0.$$

Proof. Since $f \in \Lambda_{\alpha}$, Theorem 2a of [18] can be used to conclude that $f \in M_r$ for all $r \in \left[\frac{2}{1+2\alpha}, \frac{2}{1-2\alpha}\right]$.

Under the conditions of the theorem $p, q \in \left[\frac{2}{1+2\alpha}, \frac{2}{1-2\alpha}\right]$

and so $f \in M_p \cap M_q$; hence, since f is also continuous, $f \in M(A^p) \cap M(A^q)$. Furthermore, for all $\alpha \in \mathbb{T}$,

$\tau_{\alpha} f - f \in M_q \cap M_2$. By the Riesz-Thorin convexity theorem, for $\kappa \in (0,1)$ satisfying $\frac{1}{p} = \frac{1-\kappa}{q} + \frac{\kappa}{2}$,

$$\begin{aligned} \|\tau_{\alpha} f - f\|_{p,p} &\leq \|\tau_{\alpha} f - f\|_{q,q}^{1-\kappa} \|\tau_{\alpha} f - f\|_{2,2}^{\kappa} \\ &= \|\tau_{\alpha} f - f\|_{q,q}^{1-\kappa} \|\tau_{\alpha} f - f\|_{\infty}^{\kappa}. \end{aligned} \quad (4.4.5)$$

Thus, from (4.4.4) and (4.4.5),

$$\begin{aligned} \|\tau_{\alpha} f - f\|_{M(p,p)} &\leq \|\tau_{\alpha} f - f\|_{\infty} + \|\tau_{\alpha} f - f\|_{p,p} \\ &\leq \|\tau_{\alpha} f - f\|_{\infty} + \|\tau_{\alpha} f - f\|_{q,q}^{1-\kappa} \|\tau_{\alpha} f - f\|_{\infty}^{\kappa}, \end{aligned}$$

from which it follows that

$$\lim_{\alpha \rightarrow 0} \|\tau_{\alpha} f - f\|_{M(p,p)} = 0. //$$

In proving Theorems 4.4.6 and 4.4.8 below we follow the method of proof used by Hirschman in [18], theorems 2e and 2f. In addition we obtain estimates of the norm $\|\cdot\|_{M(p,p)}$ in these theorems; the one in Theorem 4.4.6 being by appeal to the results of the Appendix. Some of the above-mentioned proof is incorporated in Construction 4.4.5 which follows.

We denote the β -variation of f by $V_\beta[f]$; that is

$$V_\beta[f] = \sup \left\{ \sum_{k=0}^{r-1} |f(x_{k+1}) - f(x_k)|^\beta \right\}^{1/\beta},$$

the supremum being taken here with respect to all finite sequences $(x_k)_{k=0}^r$ of points in $[0, 2\pi]$. Note here that $V_1[f] = V[f]$.

4.4.5 CONSTRUCTION. Suppose $p \in (1, 2]$, $f \in C$, f is real-valued, $f(0) = 0$ and $V_\beta[f] < \infty$ for some $\beta \in [1, \frac{p}{2-p})$. Then, for each $m = 0, 1, 2, \dots$, we can construct piecewise linear functions f_m and functions g_m satisfying

$$V[f_m] \leq 2^{(\beta-1)m} (V_\beta[f])^\beta, \quad (4.4.6)$$

$$f = \sum_{m=0}^{\infty} g_m \text{ and } \|g_m\|_{\infty} \leq \nu 2^{-m}, \quad (4.4.7)$$

where $\nu = \max [6, \|f\|_{\infty}]$.

Proof. Fix m and then define a subset of $[0, 2\pi]$,

$$U_{N(m)} = \{x_0 = 0, x_1, \dots, x_{N(m)} = 2\pi\}$$

by recurrence as follows. If x_{k-1} has been defined then consider the set

$$S_k = \{x \in (x_{k-1}, 2\pi] : f(x) = f(x_{k-1}) \pm 2^{-m}\};$$

put $x_k = 2\pi$ if $S_k = \emptyset$ and $x_k = \inf S_k$ otherwise. In the latter case $x_k \in S_k$. This is clear when S_k is finite, and follows from the continuity of f when S_k is infinite. Since $V_\beta[f] < \infty$ there exists an integer (which we write as $N(m)$) such that $x_{N(m)} = 2\pi$. When this point is reached we end the above process. Now define f_m by putting $f_m(x_k) = f(x_k)$ for $x_k \in U_{N(m)}$ and by linear interpolation for $x \in (x_{k-1}, x_k)$, $k = 1, 2, \dots, N(m)$. The inequality (4.4.6) then follows by arguing as in [18] pp.227 and 228. For the functions g_m , $m = 0, 1, 2, \dots$, one can take

$$g_0 = f_0$$

$$g_m = f_m - f_{m-1}, \quad m = 1, 2, \dots$$

It is not difficult to see that $f = \sum_{m=0}^{\infty} g_m$, where the sum here is to be taken point-wise. Finally we prove that g_m satisfies the estimate in (4.4.7). Consider $x \in [0, 2\pi]$. Since $g(0) = 0$ we need consider only $x \in (0, 2\pi]$. Then there exists an integer $k \in [0, N(m))$ for which $x_k < x \leq x_{k+1}$ and, from the above construction,

$$\begin{aligned} |x - x_k| |f_m(x) - f(x_k)| &\leq |x_{k+1} - x_k| |f_m(x) - f(x_k)| \\ &= |x - x_k| |f(x_{k+1}) - f(x_k)|; \end{aligned}$$

it follows that

$$|f_m(x) - f(x_k)| \leq 2^{-m}. \quad (4.4.8)$$

Also from the construction

$$|f(x) - f(x_k)| \leq 2^{-m}$$

which, combined with (4.4.8), gives

$$|f_m(x) - f(x)| \leq 2^{1-m}.$$

Hence, for $m = 1, 2, \dots$,

$$\begin{aligned} |g_m(x)| &= |f_m(x) - f_{m-1}(x)| \\ &\leq |f_m(x) - f(x)| + |f(x) - f_{m-1}(x)| \\ &\leq 6 \cdot 2^{-m}, \end{aligned}$$

which gives $\|g_m\|_\infty \leq 6 \cdot 2^{-m}$. It is clear that $\|g_0\|_\infty \leq \|f\|_\infty$; thus we now have that $\|g_m\|_\infty \leq v \cdot 2^{-m}$ where $v = \max[6, \|f\|_\infty]$, as required for (4.4.7). //

4.4.6 THEOREM. Suppose $p \in (1, 2]$, $f \in C$ and $V_\beta[f] < \infty$ for some $\beta \in [1, \frac{p}{2-p}]$. Then $f \in M(A^p)$. Further, if $\varepsilon \in (0, p-1)$ then there exists $\omega_\varepsilon = \omega_\varepsilon(p) > \frac{\beta-1}{\beta}$ such that

$$\|f\|_{M(p,p)} \leq 3\sqrt{3}\|f\|_\infty + \sum_{m=0}^{\infty} K' 2^{[(\beta-1)(1-\omega_\varepsilon) - \omega_\varepsilon]m}, \quad (4.4.9)$$

where $K' = 2^{v\omega_\varepsilon} \left(2\|f\|_\infty + \frac{(2+\pi)}{2\pi} \left(\frac{36(1+\varepsilon)(9-7\varepsilon)}{\varepsilon(1-\varepsilon)} \right)^{\frac{1}{1+\varepsilon}} (1 + 2^{1-\beta})(V_\beta[f])^\beta \right)^{1-\omega_\varepsilon}$.

Proof. Suppose first that f is real-valued and that $f(0) = 0$. Then, as detailed in Construction 4.4.5, we can define sequences $\{f_m\}_{m=0}^\infty$ and $\{g_m\}_{m=0}^\infty$ satisfying (4.4.6) and (4.4.7). From (4.4.6) and the definition of g_m ,

$$V[g_m] \leq 2^{m(\beta-1)} (1 + 2^{1-\beta}) (V_\beta[f])^\beta,$$

so that we can now apply Corollary A.3.2 and (A.3.5) to conclude that, for any $q \in (1, \infty)$,

$$\begin{aligned}
\|g_m\|_{q,q} &\leq |\hat{g}_m(0)| + \frac{1}{2\pi} B_{[q,q']} V[g_m] \\
&\leq \|g_m\|_{\infty} + \frac{1}{2\pi} B_{[q,q']} V[g_m] \\
&\leq 2 \|f\|_{\infty} + \frac{1}{2\pi} B_{[q,q']} 2^{m(\beta-1)} (1 + 2^{1-\beta}) (V_{\beta}[f])^{\beta} \\
&\leq c_q 2^{m(\beta-1)}, \tag{4.4.10}
\end{aligned}$$

where $c_q = 2 \|f\|_{\infty} + \frac{1}{2\pi} B_{[q,q']} (1 + 2^{1-\beta}) (V_{\beta}[f])^{\beta}$. Put $\omega = 2(p-1)/p$.

Then $\omega > (\beta - 1)/\beta$ and we can choose $\varepsilon \in (0, p-1)$ for which

$\omega_{\varepsilon} = 2[(p-1) - \varepsilon]/p(1-\varepsilon)$ satisfies $\omega_{\varepsilon} \in (0, 1)$ and $\omega_{\varepsilon} > (\beta - 1)/\beta$;

and $\frac{1}{p} = \frac{1 - \omega_{\varepsilon}}{1 + \varepsilon} + \frac{\omega_{\varepsilon}}{2}$. Use of the Riesz-Thorin convexity theorem in conjunction with (4.4.7) and (4.4.10) now gives us that

$$\begin{aligned}
\|g_m\|_{p,p} &\leq \|g_m\|_{2,2}^{\omega_{\varepsilon}} \|g_m\|_{1+\varepsilon,1+\varepsilon}^{1-\omega_{\varepsilon}} \\
&= \|g_m\|_{\infty}^{\omega_{\varepsilon}} \|g_m\|_{1+\varepsilon,1+\varepsilon}^{1-\omega_{\varepsilon}} \\
&\leq v^{\omega_{\varepsilon}} c_{1+\varepsilon}^{1-\omega_{\varepsilon}} 2^{[(\beta-1)(1-\omega_{\varepsilon}) - \omega_{\varepsilon}]m}.
\end{aligned}$$

Since $\omega_{\varepsilon} > (\beta-1)/\beta$, $\sum_{m=0}^{\infty} \|g_m\|_{p,p} < \infty$. Thus, using (4.4.7) and

the estimates for $B_{[1+\varepsilon,(1+\varepsilon)']}$, obtained from Theorem A.2.1,

$$\begin{aligned}
\|f\|_{p,p} &\leq \sum_{m=0}^{\infty} \|g_m\|_{p,p} \\
&\leq \sum_{m=0}^{\infty} K 2^{[(\beta-1)(1-\omega_{\varepsilon}) - \omega_{\varepsilon}]m}, \tag{4.4.11}
\end{aligned}$$

where $K = v^{\omega_{\varepsilon}} \left[2 \|f\|_{\infty} + \frac{(2+\pi)}{2\pi} \left(\frac{36(1+\varepsilon)(9-7\varepsilon)}{\varepsilon(1-\varepsilon)} \right)^{\frac{1}{1+\varepsilon}} (1+2^{1-\beta}) (V_{\beta}[f])^{\beta} \right]^{1-\omega_{\varepsilon}}$.

Combination of (4.4.11) with (4.4.4) now gives us that

$$\|f\|_{M(p,p)} \leq \|f\|_{\infty} + \sum_{m=0}^{\infty} K 2^{[(\beta-1)(1-\omega_{\varepsilon}) - \omega_{\varepsilon}]m}. \quad (4.4.12)$$

If f is real-valued but $f(0) \neq 0$ write $f = g + f(0)$, where g is real-valued and $g(0) = 0$. Then $V_{\beta}[g] = V_{\beta}[f]$ and (4.4.12) holds for g . Hence

$$\begin{aligned} \|f\|_{M(p,p)} &\leq \|g\|_{M(p,p)} + \|f(0)\|_{M(p,p)} \\ &\leq \|g\|_{\infty} + \sum_{m=0}^{\infty} K 2^{[(\beta-1)(1-\omega_{\varepsilon}) - \omega_{\varepsilon}]m} + |f(0)| \\ &\leq 3 \|f\|_{\infty} + \sum_{m=0}^{\infty} K 2^{[(\beta-1)(1-\omega_{\varepsilon}) - \omega_{\varepsilon}]m}. \quad (4.4.13) \end{aligned}$$

Now, for arbitrary $f \in \mathcal{C}$ satisfying $V_{\beta}[f] < \infty$ we can write $f = f_1 + if_2$, where f_1 and f_2 are real-valued; they satisfy $V_{\beta}[f_j] \leq V_{\beta}[f]$, $j = 1, 2$. Thus (4.4.13) holds for both f_1 and f_2 and so

$$\begin{aligned} \|f\|_{M(p,p)} &\leq \|f_1\|_{M(p,p)} + \|f_2\|_{M(p,p)} \\ &\leq 3(\|f_1\|_{\infty} + \|f_2\|_{\infty}) + 2 \sum_{m=0}^{\infty} K 2^{[(\beta-1)(1-\omega_{\varepsilon}) - \omega_{\varepsilon}]m} \\ &\leq 3\sqrt{3} \|f\|_{\infty} + \sum_{m=0}^{\infty} K' 2^{[(\beta-1)(1-\omega_{\varepsilon}) - \omega_{\varepsilon}]m}, \end{aligned}$$

where $K' = 2K$; and this is the required estimate (4.4.9). //

4.4.7 COROLLARY. Suppose $p \in (1, 2]$, $f \in \mathcal{C}$ and $V_{\beta}[f] < \infty$ for some $\beta \in [1, \frac{p}{2-p})$. Then $f \in M(A^p)$ and

$$\lim_{\alpha \rightarrow 0} \|\tau_{\alpha} f - f\|_{M(p,p)} = 0.$$

Proof. This result follows directly from the estimate (4.4.9) obtained in Theorem 4.4.6. //

4.4.8 THEOREM. Suppose $p \in (1, 2]$, $f \in \Lambda_\alpha$ for some $\alpha \in (0, 1]$, and $V_\beta[f] < \infty$ for some $\beta \in [2, \frac{2p}{2-p})$. Then $f \in M(\Lambda^p)$. Further there exists $\varepsilon \in (0, 1)$ such that

$$\|f\|_{M(p,p)} \leq 3\sqrt{3} \|f\|_\infty + \sum_{m=0}^{\infty} K'' 2^{[\frac{1}{2}(\beta-2+\varepsilon)(1-\omega)-\omega]m}, \quad (4.4.14)$$

$$\text{for } K'' = 2v^\omega \left(2\|f\|_\infty + c(1 + 2^{2-\varepsilon-\beta}) (2\|f\|_{\Lambda_\alpha})^{\frac{1}{2}\varepsilon} (V_\beta[f])^{\frac{1}{2}\beta} \right)^{1-\omega}$$

where $\omega = 2(p-1)/p$, $c = c(\varepsilon, \alpha)$ depends only on ε and α , and v is as in (4.4.7).

Proof. Suppose first that f is real-valued and that $f(0) = 0$. We can then proceed as in Construction 4.4.5 to define sequences $(f_m)_{m=0}^\infty$ and $(g_m)_{m=0}^\infty$, the latter satisfying (4.4.7).

By the arguments given in [18], pp.227 and 228, $f_m \in \Lambda_\alpha$ with

$$\|f_m\|_{\Lambda_\alpha} \leq \|f\|_{\Lambda_\alpha} \text{ and, for any } \varepsilon \in (0, 1),$$

$$\left(V_{2-\varepsilon}[f_m] \right)^{2-\varepsilon} \leq 2^{(\beta-2+\varepsilon)m} \left(V_\beta[f] \right)^\beta. \quad (4.4.15)$$

Thus, from the method of construction of f_m and g_m , $\|g_m\|_{\Lambda_\alpha} \leq 2\|f\|_{\Lambda_\alpha}$

and

$$\left(V_{2-\varepsilon}[g_m] \right)^{2-\varepsilon} \leq 2^{(\beta-2+\varepsilon)m} \left(1 + 2^{2-\varepsilon-\beta} \right) \left(V_\beta[f] \right)^\beta. \quad (4.4.16)$$

Hence $V_{2-\varepsilon}[g_m] < \infty$ and $g_m \in \Lambda_\alpha$ so we can apply Lemma 2d of [18] to obtain (where the primed sum indicates omission of the case $n = 0$)

$$\begin{aligned} \|\hat{g}_m\|_1 &= |\hat{g}(0)| + \sum'_{n \in \mathbb{Z}} |\hat{g}_m(n)| \\ &\leq \|g_m\|_\infty + c \left(V_{2-\varepsilon}[g_m] \right)^{\frac{1}{2}(2-\varepsilon)} \|g_m\|_{\Lambda_\alpha}^{\frac{1}{2}\varepsilon}, \end{aligned} \quad (4.4.17)$$

where $c = c(\varepsilon, \alpha)$ depends only on ε and α . We can now

combine (4.4.17) with (4.4.15) to obtain

$$\begin{aligned} \|\hat{g}_m\|_1 &\leq 2 \|f\|_\infty + c 2^{\frac{1}{2}(\beta-2+\varepsilon)m} \left(1 + 2^{2-\varepsilon-\beta}\right)^{\frac{1}{2}} \left(2 \|f\|_{\Lambda_\alpha}\right)^{\frac{1}{2}\varepsilon} \left(V_\beta[f]\right)^{\frac{1}{2}\beta} \\ &\leq K'' 2^{\frac{1}{2}(\beta-2+\varepsilon)m}, \end{aligned} \quad (4.4.18)$$

where $K'' = 2 \|f\|_\infty + c(1 + 2^{2-\varepsilon-\beta})^{\frac{1}{2}} (2 \|f\|_{\Lambda_\alpha})^{\frac{1}{2}\varepsilon} (V_\beta[f])^{\frac{1}{2}\beta}$.

Now $\omega = 2(p-1)/p > (\beta - 2)/\beta$, whence it follows that for $\varepsilon > 0$

suitably small $\omega > (\beta - 2 + \varepsilon)/(\beta + \varepsilon)$. Also $\omega \in (0,1)$ and

$\frac{1}{p} = \frac{1-\omega}{1} + \frac{\omega}{2}$; application of the Riesz-Thorin convexity theorem

and use of (4.4.7) and (4.4.18) give

$$\|g_m\|_{p,p} \leq \|g_m\|_{1,1}^{1-\omega} \|g_m\|_{2,2}^\omega$$

$$= \|\hat{g}_m\|_1^{1-\omega} \|g_m\|_\infty^\omega$$

$$\leq K' 2^{[\frac{1}{2}(\beta-2+\varepsilon)(1-\omega) - \omega]m},$$

where $K' = K^{1-\omega} v^\omega$. Note that with this choice of ε

$\sum_{m=0}^{\infty} \|g_m\|_{p,p} < \infty$. We now use (4.4.4) and (4.4.7) to obtain

$$\|f\|_{M(p,p)} \leq \|f\|_\infty + \|f\|_{p,p}$$

$$\leq \|f\|_\infty + \sum_{m=0}^{\infty} \|g_m\|_{p,p}$$

$$\leq \|f\|_\infty + \sum_{m=0}^{\infty} K' 2^{[\frac{1}{2}(\beta-2+\varepsilon)(1-\omega) - \omega]m}.$$

Arguing now as in Theorem 4.4.6 we can deduce that, for arbitrary

$f \in \Lambda_\alpha$ for which $V_\beta[f] < \infty$,

$$\|f\|_{M(p,p)} \leq 3\sqrt{3} \|f\|_{\infty} + \sum_{m=0}^{\infty} K'' 2^{[\frac{1}{2}(\beta-2+\epsilon)(1-\omega) - \omega]m},$$

where $K'' = 2K'$; this is the required estimate (4.4.14). //

4.4.9 COROLLARY. Suppose $p \in (1,2]$, $f \in \Lambda_{\alpha}$ for some $\alpha \in (0,1]$, and $V_{\beta}[f] < \infty$ for some $\beta \in [2, \frac{2p}{2-p})$. Then $f \in M(A^p)$ and

$$\lim_{\alpha \rightarrow 0} \|\tau_{\alpha} f - f\|_{M(p,p)} = 0.$$

Proof. This result follows directly from the estimate (4.4.14) obtained in Theorem 4.4.8. //

4.5 An alternative approach.

We noted earlier that $M_p \cap C \subseteq M(A^p)$ for all $p \in [1,2]$; equality holds for the cases $p = 1$ and $p = 2$.⁴ The question whether or not equality holds for $p \in (1,2)$ remains unanswered. However, we shall give here an equivalent formulation of this question by obtaining conditions equivalent to membership of $M(A^p)$ and M_p respectively. To this end we shall define a multiplication for (some) distributions. It will be seen that this multiplication agrees with the usual one where such multiplication is already defined. We shall of course be considering all our spaces here as embedded in the space $\mathcal{D}$ of distributions on $\mathbb{T}$ in the usual way.

4.5.1 DEFINITION. For two distributions σ and ν , the product $\sigma\nu$ is the distributional limit (if it exists) of the sequence $\left\{ (F_N * \sigma)\nu \right\}_{N=0}^{\infty}$. //

Note here that if σ and ν are the distributions generated by

the continuous functions f and g respectively, then σv is the distribution generated by fg . We shall now show that for $p \in (1,2)$, σv is well-defined whenever $\sigma \in A^p$ and $v \in M + Fl^{p'}$.

4.5.2 LEMMA. Suppose $p \in (1,2)$, $f \in A^p$ and $\sigma \in M + Fl^{p'}$. Then $f\sigma$ exists and is given by

$$\langle f\sigma, u \rangle = \langle \sigma, fu \rangle, \text{ for all } u \in C^\infty.$$

Proof. As $f \in A^p$, so is the product $fu \in A^p$ for all $u \in C^\infty$; and, since

$$N_p \left[(F_N * f)u - fu \right] \leq \|u\|_{M(p,p)} N_p (F_N * f - f),$$

the sequence $\left[(F_N * f)u \right]_{N=0}^\infty$ is strongly, and hence weakly, convergent in A^p to fu . Thus, for all $u \in C^\infty$,

$$\begin{aligned} \lim_{N \rightarrow \infty} \langle (F_N * f)\sigma, u \rangle &= \lim_{N \rightarrow \infty} \langle \sigma, (F_N * f)u \rangle \\ &= \langle \sigma, fu \rangle. \quad // \end{aligned}$$

4.5.3 LEMMA. Suppose $p \in (1,2)$. Then $f \in M(A^p)$ if and only if $f \cdot (M + Fl^{p'}) \subseteq M + Fl^{p'}$.

Proof. (i) "only if". Suppose $f \in M(A^p)$. Since $M(A^p) \subseteq A^p$ it follows from Lemma 4.5.2 that $f\sigma$ is well-defined as a distribution for all $\sigma \in M + Fl^{p'}$. In order to conclude that $f\sigma \in M + Fl^{p'}$ we need to show that $f\sigma$ can be extended to a continuous linear functional on A^p . To do this consider $g \in A^p$ and any sequence $(g_k)_{k=1}^\infty$ from C^∞ for which $\lim_{k \rightarrow \infty} N_p(g - g_k) = 0$. Then use of Lemma 4.5.2 gives that

$$\begin{aligned}
| \langle f\sigma, g_k \rangle - \langle f\sigma, g_m \rangle | &= | \langle f\sigma, g_k - g_m \rangle | \\
&= | \langle \sigma, f(g_k - g_m) \rangle | \\
&\leq \text{const. } N_p \left[f(g_k - g_m) \right] \\
&\leq \text{const. } \|f\|_{M(p,p)} N_p(g_k - g_m); (4.5.1)
\end{aligned}$$

hence the sequence $\{\langle f\sigma, g_k \rangle\}_{k=1}^{\infty}$ is Cauchy in $\mathbb{C}$ and so $\lim_{k \rightarrow \infty} \langle f\sigma, g_k \rangle$ exists. It is not difficult to see that this limit is the same for all such sequences $(g_k)_{k=1}^{\infty}$. Now the required extension to A^p , written also as $f\sigma$, is obtained by writing

$$\langle f\sigma, g \rangle = \lim_{k \rightarrow \infty} \langle f\sigma, g_k \rangle,$$

where $g \in A^p$ and the sequence $(g_k)_{k=1}^{\infty}$ from C^{∞} is as described above. To see that the linear functional so defined is continuous we use (4.5.1) to obtain

$$| \langle f\sigma, g \rangle | = \lim_{k \rightarrow \infty} | \langle f\sigma, g_k \rangle |$$

$$\leq \text{const. } \lim_{k \rightarrow \infty} N_p(g_k)$$

$$= \text{const. } N_p(g).$$

(ii) "if". Suppose now that $f \cdot (M + Fl^{p'}) \subseteq M + Fl^{p'}$. Then, for $g \in A^p$ and a sequence $(g_k)_{k=1}^{\infty}$ from C^{∞} for which $\lim_{k \rightarrow \infty} N_p(g - g_k) = 0$, we have that for each $\sigma \in M + Fl^{p'}$,

$$\langle f\sigma, g \rangle = \lim_{k \rightarrow \infty} \langle f\sigma, g_k \rangle$$

$$= \lim_{k \rightarrow \infty} \langle \sigma, fg_k \rangle, \quad (4.5.2)$$

by Lemma 4.5.2. From (4.5.2) it then follows that $(fg_k)_{k=1}^{\infty}$ is weakly bounded; hence by [6], p.467, strongly bounded in A^p . Thus there exists a constant $K > 0$ for which

$$N_p(fg_k) \leq K, \text{ for all } k = 1, 2, \dots \quad (4.5.3)$$

Also, $((fg_k)^{\wedge})_{k=1}^{\infty}$ converges to $(fg)^{\wedge}$ pointwise on Z ; which fact, together with (4.5.3), implies that $(fg)^{\wedge} \in \ell^p$ (with $\|(fg)^{\wedge}\| \leq K$). It is clear that $fg \in C$ for all $g \in A^p$ so our proof is completed. //

4.5.4 LEMMA. Suppose $p \in (1, 2)$, $f \in \mathcal{D}$ and $f\sigma$ is defined for all $\sigma \in Fl^{p'}$. Then $f \in Fl^p$.

Proof. Consider $\sigma \in Fl^{p'}$. Then, from our hypothesis, $\lim_{N \rightarrow \infty} \langle (F_N * f)\sigma, u \rangle$ exists for all $u \in C^{\infty}$. In particular, since $1 \in C^{\infty}$, so we have that $\lim_{N \rightarrow \infty} \langle (F_N * f)\sigma, 1 \rangle = \lim_{N \rightarrow \infty} \langle \sigma, F_N * f \rangle$ exists for all $\sigma \in Fl^{p'}$. Thus $(F_N * f)_{N=0}^{\infty}$ is weakly, and hence strongly, bounded in Fl^p ; and so there exists a constant $K > 0$ for which

$$\|(F_N * f)^{\wedge}\|_p \leq K, \text{ for all } N = 0, 1, 2, \dots \quad (4.5.4)$$

From (4.5.4) and the fact that $(F_N)_{N=0}^{\infty}$ converges to 1 pointwise on Z it follows that $\hat{f} \in \ell^p$ and our proof is completed. //

4.5.5 LEMMA. Suppose $p \in (1, 2)$, $f \in \mathcal{D}$ and $f \cdot Fl^{p'} \subseteq M + Fl^{p'}$. Then $f \cdot A^p \subseteq Fl^p$.

Proof. Consider $\sigma \in Fl^{p'}$. Then from our assumptions, for $g \in A^p$, $\langle f\sigma, g \rangle = \lim_{k \rightarrow \infty} \langle f\sigma, g_k \rangle$, where $(g_k)_{k=1}^{\infty}$ is any sequence from C^{∞} for which $\lim_{k \rightarrow \infty} N_p(g - g_k) = 0$. From Lemma 4.5.4 it follows that $f \in Fl^p$ and so, for each $k = 1, 2, \dots$, $fg_k \in Fl^p$. Thus, since for each k ,

$$\| \left((F_N * f)g_k - fg_k \right)^\wedge \|_p \leq \| (F_N * f)^\wedge - \hat{f} \|_p \| \hat{g}_k \|_1 ,$$

the sequence $\left((F_N * f)g_k \right)_{N=0}^\infty$ converges strongly, and hence weakly, in Fl^p to fg_k . Thus

$$\begin{aligned} \langle f\sigma, g \rangle &= \lim_{k \rightarrow \infty} \lim_{N \rightarrow \infty} \langle \sigma, (F_N * f)g_k \rangle \\ &= \lim_{k \rightarrow \infty} \langle \sigma, fg_k \rangle, \text{ for all } \sigma \in Fl^{p'} . \end{aligned}$$

Arguing now as in the "if" proof of Lemma 4.5.3 it follows that $fg \in Fl^p$ and our proof is completed. //

We are now in a position to deduce the following (necessary and sufficient) condition for membership of $M(A^p)$.

4.5.6 THEOREM. Suppose $p \in (1,2)$ and $f \in C$. Then $f \in M(A^p)$ if and only if $f.Fl^{p'} \subseteq M + Fl^{p'}$.

Proof. (i) "only if". Suppose $f \in M(A^p)$. Then, since $Fl^{p'} \subseteq M + Fl^{p'}$ and $M(A^p) \subseteq A^p$, the required result follows directly from Lemma 4.5.3.

(ii) "if". Assume that $f.Fl^{p'} \subseteq M + Fl^{p'}$. Then, from Lemma 4.5.5, $f.A^p \subseteq Fl^p$ and, since $f \in C$, $f.A^p \subseteq Fl^p \cap C = A^p$, this completes our proof. //

Arguments similar to those in the preceding lemmas enable us to deduce a corresponding (necessary and sufficient) condition for membership of $M_p \cap C$.

4.5.7 THEOREM. Suppose $p \in (1,2)$ and $f \in \mathcal{D}$. Then $f \in M_p$ if and only if $f.Fl^{p'} \subseteq Fl^{p'}$.

Proof. (i) "only if". Note that $M_p \subseteq Fl^p$ and then argue as in the "only if" part of the proof of Lemma 4.5.3.

(ii) "if". Use Lemma 4.5.4 and then argue as in the proof of Lemma 4.5.5. //

4.5.8 COROLLARY. Suppose $p \in (1,2)$ and $f \in C$. Then $f \in M_p \cap C$ if and only if $f \cdot \text{Fl}^{p'} \subseteq \text{Fl}^{p'}$. //

A.1 Introduction.

Hilbert introduced two bilinear forms, known as his A and B forms. Details of these are available in [15], p.206 ff. Both of these forms have important applications in the theory of functions of a complex variable and have been studied by many people. We shall be concerned here with Hilbert's B form, and also with a generalization of it introduced by Stečkin in [32].

A.1.1 DEFINITION. For $p \in (1,\infty)$, $g \in \ell^p$, $h \in \ell^{p'}$, the bilinear form B defined by (where $'$ denotes conjugation of the case $m = n$)

$$B(g,h) = \sum_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} \frac{g(n) \bar{h}(m)}{m-n} \quad (\text{A.1.1})$$

is known as Hilbert's B form.

A.1.2 DEFINITION. The bilinear form B defined in A.1.1 is bounded in $[p,p']$ if and only if, for all g,h with finite support, there exists $c = c(p)$ depending only on p such that

$$\left| \sum_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} \frac{g(n) \bar{h}(m)}{m-n} \right| \leq c \|g\|_p \|h\|_{p'} \quad (\text{A.1.2})$$

APPENDIX A

BOUNDS FOR TWO SPECIAL BILINEAR FORMS

A.1 Introduction.

Hilbert introduced two bilinear forms, known as his A and B forms. Details of these are available in [15], p.206 ff. Both of these forms have important applications in the theory of functions of a complex variable and have been studied by many people. We shall be concerned here with Hilbert's B form, and also with a generalisation of it introduced by Stěckin in [32].

A.1.1 DEFINITION. For $p \in (1, \infty)$, $g \in \ell^p$, $h \in \ell^{p'}$, the bilinear form B defined by (where ' denotes omission of the case $m = n$)

$$B(g, h) = \sum_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}}' \frac{g(n) h(m)}{m - n} \quad (\text{A.1.1})$$

is known as Hilbert's B form.

A.1.2 DEFINITION. The bilinear form B defined in A.1.1 is bounded in $[p, p']$ if and only if, for all g, h with finite support, there exists $c = c(p)$ depending only on p such that

$$\left| \sum_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}}' \frac{g(n) h(m)}{m - n} \right| \leq c \|g\|_p \|h\|_{p'} . \quad (\text{A.1.2})$$

A.1.3 DEFINITION. For $p \in (1, \infty)$ we use $B_{[p, p']}$ to denote the infimum of all the $c = c(p)$ satisfying (A.1.2). We call $B_{[p, p]}$ the *bound* of B .

We shall prove here that B is bounded in $[p, p]$ for $p \in (1, \infty)$ and obtain estimates for $B_{[p, p]}$. In this context see the comments in [15], p.214. Note here that our Definition A.1.2 is equivalent to that given in [15], p.207.

A.2 Bounds for Hilbert's B form.

In our proof that Hilbert's B form is bounded in $[p, p]$ we shall use Corollary 1.3.5 for the case $G = \mathbb{Z}$ with the covering family indicated in Remark 1.3.6.

A.2.1 THEOREM. Suppose $p \in (1, \infty)$. Then Hilbert's B form is bounded in $[p, p]$; and

$$B_{[p, p]} \leq \begin{cases} (2 + \pi) \left(\frac{36p(16 - 7p)}{(p - 1)(2 - p)} \right)^{1/p}, & p \in (1, 2), \\ 3(2 + \pi), & p = 2, \\ B_{[p', p]}, & p \in (2, \infty). \end{cases} \quad (\text{A.2.1})$$

Proof. Consider $v = 1, 2, \dots$, and define $k_v \in \ell^1$ by

$$k_v(n) = \begin{cases} \frac{1}{n} \xi_{[-v, v]}(n), & n \neq 0, \\ 0, & \text{otherwise} \end{cases} \quad (\text{A.2.2})$$

Now $k_v(n)$ here is of the form

$$k_v(n) = \hat{f}(n) \xi_{[-v, v]}(n),$$

where the $\hat{f}(n)$ satisfy conditions (4.2.2), (4.2.3) and (4.2.4). Thus by the argument given in Construction 4.2.2 it follows that, for $\beta = 1, 2, \dots$,

$$\begin{aligned} J_\beta(k_v) &\leq 2(2^\beta - 1)/(2^\beta + 1) \\ &\leq 2; \end{aligned}$$

hence

$$J(k_v) = \sup_\beta J_\beta(k_v) \leq 2. \quad (\text{A.2.4})$$

In addition, for $x \in [0, 2\pi)$,

$$\begin{aligned} |\hat{k}_v(x)| &= \left| \sum_{|n| \leq v} n^{-1} e^{inx} \right| \\ &= \left| \sum_{1 \leq n \leq v} 2 n^{-1} \sin nx \right| \\ &\leq 2 + \pi; \end{aligned}$$

the final inequality follows from a result of D. Jackson in [20] (also to be found in [26], 3.5.5).

Hence

$$\|\hat{k}_v\|_\infty \leq 2 + \pi. \quad (\text{A.2.5})$$

From (A.2.4) and (A.2.5),

$$\max \left\{ J(k_v), \|\hat{k}_v\|_\infty \right\} \leq 2 + \pi$$

and so Corollary 1.3.5 gives us that $k_v * f \in \ell^p$ whenever $f \in \ell^p$, and

$$\|k_v * f\|_p \leq \begin{cases} (2 + \pi) \left(\frac{36p(16 - 7p)}{(p - 1)(2 - p)} \right)^{1/p} \|f\|_p, & p \in (1, 2), \\ 3(2 + \pi) \|f\|_2, & p = 2, \\ (2 + \pi) \left(\frac{36p'(16 - 7p')}{(p' - 1)(2 - p')} \right)^{1/p'} \|f\|_p, & p \in (2, \infty). \end{cases} \quad (\text{A.2.6})$$

We can now use (A.2.6) and Hölder's inequality to show that for all $v = 1, 2, \dots$,

$$\begin{aligned} \left| \sum_{m \in \mathbb{Z}} \sum_{\substack{n: |m-n| \leq v \\ m \neq n}} \frac{g(n) h(m)}{m - n} \right| &= \left| \sum_{m \in \mathbb{Z}} k_v * g(m) h(m) \right| \\ &\leq \|k_v * g\|_p \|h\|_{p'}, \\ &\leq c \|g\|_p \|h\|_{p'}, \end{aligned}$$

where $c = c(p)$ is given by

$$c = \begin{cases} (2 + \pi) \left(\frac{36p(16 - 7p)}{(p - 1)(2 - p)} \right)^{1/p}, & p \in (1, 2), \\ 3(2 + \pi), & p = 2, \\ (2 + \pi) \left(\frac{36p'(16 - 7p')}{(p' - 1)(2 - p')} \right)^{1/p'}, & p \in (2, \infty). \end{cases} \quad (\text{A.2.7})$$

Thus for $g \in \ell^p$, $h \in \ell^{p'}$, both with finite support, we can choose $v = v(g, h)$ to ensure that

$$\{(m, n) : 0 < |m - n| \leq v\} \supseteq \{(m, n) : m \in \text{supp } h, n \in \text{supp } g, m \neq n\};$$

hence

$$\left| \sum_{m \in \mathbb{Z}} \sum'_{n \in \mathbb{Z}} \frac{g(n) h(m)}{m - n} \right| = \left| \sum_{m \in \mathbb{Z}} k_v(g, h) * g(m) h(m) \right|$$

$$\leq \|g\|_p \|h\|_{p'}, \quad .$$

We thus have that B is bounded in $[p, p']$ with $B_{[p, p']}$ satisfying (A.2.1) as required. //

A.2.2 REMARKS. Hilbert himself proved the boundedness of B for the case $p = 2$. Schur in [31] considered the A and B forms together in order to obtain the estimate $B_{[2, 2]} \leq \pi$. Details of Schur's proof may also be found in [15]. In the general case Titchmarsh proved in [33] that for $p \in [2, \infty)$ there exist constants H and K independent of p such that $Hp \leq B_{[p, p']} \leq Kp^{4/3}$. In [27] M. Riesz improved this to $Hp \leq B_{[p, p']} \leq Kp$.

A.3 Bounds for Stěckin's bilinear form.

We shall give here a result of Stěckin ([32], pp.238 ff) concerning the boundedness in $[p, p']$ of the bilinear form S defined by (A.3.1) below. In the process we shall make use of the estimate (A.2.1) for $B_{[p, p']}$ to obtain an estimate for $S_{[p, p']}$.

A.3.1 THEOREM. Suppose $f \in BV$ and $p \in (1, \infty)$. Then for $g \in \ell^p$, $h \in \ell^{p'}$ the bilinear form S defined by

$$S(g, h) = \sum_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} \hat{f}(m - n) g(n) h(n) \quad (\text{A.3.1})$$

is bounded in $[p, p']$. Further,

$$S_{[p, p']} \leq \begin{cases} |\hat{f}(0)| + \frac{(2 + \pi)}{2\pi} \left[\frac{36p(16 - 7p)}{(p - 1)(2 - p)} \right]^{1/p} V[f], & p \in (1, 2), \\ |\hat{f}(0)| + \frac{3(2 + \pi)}{2\pi} V[f], & p = 2, \\ S_{[p', p]}, & p \in (2, \infty). \end{cases} \quad (\text{A.3.2})$$

Proof. Consider $g \in \ell^p$ and $h \in \ell^{p'}$, both with finite support.

Now write

$$S(g, h) = \hat{f}(0) \sum_{n \in \mathbb{Z}} g(n) h(n) + \sum_{m \in \mathbb{Z}} \sum'_{n \in \mathbb{Z}} \hat{f}(m - n) g(n) h(m) .$$

Then

$$|S(g, h)| \leq |\hat{f}(0)| \|g\|_p \|h\|_{p'} + |S'(g, h)| , \quad (\text{A.3.3})$$

where

$$\begin{aligned} S'(g, h) &= \frac{1}{2\pi} \int_0^{2\pi} f(t) \sum_{m \in \mathbb{Z}} \sum'_{n \in \mathbb{Z}} g(n) h(m) e^{-i(m-n)t} dt \\ &= \frac{1}{2\pi i} \int_0^{2\pi} \sum_{m \in \mathbb{Z}} \sum'_{n \in \mathbb{Z}} \frac{g(n) h(m) e^{-i(m-n)t}}{(m - n)} df(t) ; \end{aligned}$$

hence

$$|S'(g, h)| \leq \frac{1}{2\pi} \sup_{t \in [0, 2\pi]} \left| \sum_{m \in \mathbb{Z}} \sum'_{n \in \mathbb{Z}} \frac{g(n) e^{-imt} h(m) e^{int}}{(m - n)} \right| \int_0^{2\pi} |df(t)| .$$

From Theorem A.2.1,

$$|S'(g, h)| \leq \frac{1}{2\pi} B_{[p, p']} \|g\|_p \|h\|_{p'} \int_0^{2\pi} |df(t)| . \quad (\text{A.3.4})$$

Combination of (A.3.3) and (A.3.4) gives

$$|S(g, h)| \leq \left(|\hat{f}(0)| + \frac{1}{2\pi} B_{[p, p']} V[f] \right) \|g\|_p \|h\|_{p'} ;$$

hence S is bounded in $[p, p']$ and

$$S_{[p, p']} \leq |\hat{f}(0)| + \frac{1}{2\pi} B_{[p, p']} V[f] . \quad (\text{A.3.5})$$

The estimate (A.3.2) now follows from direct substitution in (A.3.5)

of the estimate (A.2.1). //

We shall now prove a corollary of Theorem A.3.1, the statement of which can be found in [32], p.240.

A.3.2 COROLLARY. (*Stěckin's multiplier theorem.*) Suppose $f \in BV$ and $p \in (1, 2]$. Then $\hat{f} * g \in \ell^p$ whenever $g \in \ell^p$ and

$$\|\hat{f} * g\|_p \leq S_{[p, p']} \|g\|_p ,$$

where S is the bilinear form defined in (A.3.1).

Proof. Consider $g \in \ell^p$ and $h \in \ell^{p'}$, both with finite support. By Theorem A.3.1,

$$\begin{aligned} |S(g, h)| &= \left| \sum_{m \in \mathbb{Z}} \sum_{n \in \mathbb{Z}} \hat{f}(m - n) g(n) h(m) \right| \\ &= \left| \sum_{m \in \mathbb{Z}} \hat{f} * g(m) h(m) \right| \\ &\leq S_{[p, p']} \|g\|_p \|h\|_{p'} , \end{aligned}$$

and hence, by the converse of Hölder's inequality ,

$$\|\hat{f} * g\|_p \leq S_{[p, p']} \|g\|_p \tag{A.3.6}$$

for all $g \in \ell^p$ with finite support. Now consider arbitrary $g \in \ell^p$ and define $(g_k)_{k=1}^\infty$ by

$$g_k(n) = \begin{cases} g(n) & , \quad |n| \leq k , \\ 0 & , \quad \text{otherwise} . \end{cases}$$

Clearly (A.3.6) holds for each g_k , $k = 1, 2, 3, \dots$. Since $\hat{f} \in \ell^2$ and $g \in \ell^p \subseteq \ell^2$, it follows that $\hat{f} * g(n)$ is defined a priori as the sum of the absolutely convergent series $\sum_{m \in \mathbb{Z}} \hat{f}(n - m) g(m)$ and it is

clear that $\hat{f} * g_k \rightarrow \hat{f} * g$ pointwise as $k \rightarrow \infty$. In addition, (A.3.6) shows that $\|\hat{f} * g_k\|_p$ is bounded with respect to k , so we can conclude that $\hat{f} * g \in \ell^p$ and (A.3.6) continues to hold for all $g \in \ell^p$. //

APPENDIX B

SOME ADDITIONAL RESULTS

B.1 Idempotent Fourier Multipliers from A^p to A^q , $p, q \in [1, 2]$.

In this section we turn our attention once more to the Fourier multipliers from A^p to A^q , $p, q \in [1, 2]$, which were previously considered in Chapter 3. Here we are interested in characterising the set of all such multipliers that are idempotent. By an idempotent multiplier from A^p to A^q we mean a distribution $\phi \in (A^p, A^q)$ for which $\phi * \phi = \phi$. We shall write $(A^p, A^q)_I$ for the set of all idempotent elements of (A^p, A^q) . In addition, we shall follow the notation of [39] and denote by J the set of all idempotent measures on T . Again ξ_E denotes the characteristic function of $E \subseteq Z$.

B.1.1 LEMMA. *To each $\phi \in (A^p, A^q)_I$ there corresponds (unique) $E \subseteq Z$ for which $\phi = \hat{\xi}_E$.*

Proof. Since $\phi * \phi = \phi$ it follows that $\hat{\phi}^2 = \hat{\phi}$ and so, for all $n \in Z$, $\hat{\phi}(n) \in \{0, 1\}$. Write

$$E = \{n \in Z : \hat{\phi}(n) = 1\} ;$$

then $\phi = \hat{\xi}_E$. //

B.1.2 THEOREM. *For every $E \subseteq Z$, $\hat{\xi}_E \in (A, A)_I$.*

Proof. This result follows directly from the fact that

$$A = F\ell^1 . //$$

In the proof of the following theorem we shall, for any positive integer k , denote by σ^{*k} the k -fold convolution of the distribution σ with itself.

B.1.3 THEOREM. $(A^p, A^p)_I = J$, $p \in (1, 2]$.

Proof. Let $p \in (1, 2]$. The inclusion $J \subseteq (A^p, A^p)_I$ is clear. We shall prove here that $(A^p, A^p)_I \subseteq J$. To this end suppose that $\phi \in (A^p, A^p)_I$. Then $\phi * \phi = \phi$ and, from Theorem 3.2.5, we can write $\phi = \mu - \sigma$, where $\mu \in M$ and $\hat{\sigma} \in \ell^{p'}$. Choose $n \in \mathbb{Z}$ to satisfy $2n \leq p' < 2(n+1)$. Then $\hat{\sigma}^{n+1} \in \ell^2$ and hence $\sigma^{*r} \in M$ for all $r \geq n+1$. Consideration of

$$\begin{aligned} \mu^{*s} &= (\phi + \sigma)^{*s} = \sum_{r=0}^s \binom{s}{r} \phi^{*s-r} * \sigma^{*r} \\ &= \sum_{r=0}^s \binom{s}{r} \phi * \sigma^{*r}, \quad s = n+1, n+2, \dots, 2n+1 \end{aligned}$$

gives us that

$$\begin{bmatrix} 1 & \binom{n+1}{1} & \dots & \binom{n+1}{n} \\ 1 & \binom{n+2}{1} & \dots & \binom{n+2}{n} \\ \cdot & \cdot & & \cdot \\ \cdot & \cdot & & \cdot \\ \cdot & \cdot & & \cdot \\ 1 & \binom{2n+1}{1} & \dots & \binom{2n+1}{n} \end{bmatrix} \begin{bmatrix} \phi \\ \phi * \sigma \\ \cdot \\ \cdot \\ \cdot \\ \phi * \sigma^{*n} \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \cdot \\ \cdot \\ \cdot \\ \mu_{n+1} \end{bmatrix} \quad (\text{B.1.1})$$

where $\mu_i \in M$, $i = 1, 2, \dots, n+1$.

Now write

$$w_{n(0)} = \begin{bmatrix} 1 & \binom{n+1}{1} & \binom{n+1}{2} & \cdot & \cdot & \cdot & \binom{n+1}{n} \\ 1 & \binom{n+2}{1} & \binom{n+2}{2} & \cdot & \cdot & \cdot & \binom{n+2}{n} \\ \cdot & \cdot & \cdot & & & & \cdot \\ \cdot & \cdot & \cdot & & & & \cdot \\ \cdot & \cdot & \cdot & & & & \cdot \\ 1 & \binom{2n+1}{1} & \binom{2n+1}{2} & \cdot & \cdot & \cdot & \binom{2n+1}{n} \end{bmatrix}$$

We shall show that $w_{n(0)}$ is non-singular and hence, from (B.1.1), that $\phi \in M$. The matrix

$$w_{n(1)} = \begin{bmatrix} 1 & \binom{n+1}{1} & \binom{n+1}{2} & \cdot & \cdot & \cdot & \binom{n+1}{n} \\ 0 & 1 & \binom{n+1}{1} & \cdot & \cdot & \cdot & \binom{n+1}{n-1} \\ 0 & 1 & \binom{n+2}{1} & \cdot & \cdot & \cdot & \binom{n+2}{n-1} \\ \cdot & \cdot & \cdot & & & & \cdot \\ \cdot & \cdot & \cdot & & & & \cdot \\ \cdot & \cdot & \cdot & & & & \cdot \\ 0 & 1 & \binom{2n}{1} & \cdot & \cdot & \cdot & \binom{2n}{n-1} \end{bmatrix}$$

is obtained from $w_{n(0)}$ by performing the following row operations:

row($k+1$) - row k , $k = n, n-1, \dots, 2, 1$; leave row 1 unchanged;

and using the result

$$\binom{t+1}{s} - \binom{t}{s} = \binom{t}{s-1}, \quad t \geq s-1. \quad (\text{B.1.2})$$

and our proof is completed. //

B.1.4 COROLLARY. If $p \in (1,2]$, $q \in [p,2]$, then

$$(A^p, A^q)_I = (A^p, A^p)_I = J .$$

Proof. This result follows directly by combination of Theorem B.1.3 with Theorem 3.4.1. //

B.1.5 THEOREM. Let $p \in (1,2]$, $q \in [1,p)$ be given. Then $\phi \in (A^p, A^q)_I$ if and only if $\phi = \hat{\xi}_E$ for some finite $E \subseteq \mathbb{Z}$.

Proof. (i) "if" ; trivial

(ii) "only if" . In the special case $p = 2$, $q \in [1,2)$ this result follows directly from Theorem 3.5.2. In general, suppose $\phi \in (A^p, A^q)_I$. Then, since $(A^p, A^q)_I \subseteq (A^p, A^p)_I$, it follows from Theorem B.1.3 that $\phi \in J$ and hence that $\phi = \hat{\xi}_E$ for some $E \subseteq \mathbb{Z}$ belonging to the coset ring of $\mathbb{Z}$ (see [39], p.61). Suppose now that E is an infinite set. By a theorem of Helson (see [35], [36]) E differs by a finite set from a periodic subset of $\mathbb{Z}$; thus there exist positive integers r, d , $a_1, a_2, \dots, a_r$ satisfying $a_1 < a_2 < \dots < a_r$ and $a_r - a_1 < d$, such that E differs in at most finitely many places from the set

$$\{a_1 + md , a_2 + md , \dots , a_r + md : m \in \mathbb{Z}\} .$$

We shall show that, for any such E , there exists $f \in A^p$ for which $\hat{\xi}_E f \notin \ell^q$; this is accomplished by slightly modifying Construction

2.5.2. Write $f = \sum_{k=0}^{\infty} f_k$, where

$$f_k = (k+1)^{-1/q} 2^{-k/q} p_k e_{v_k}, \quad k = 0, 1, 2, \dots,$$

and the $\{v_k\}$ will be chosen to ensure that the spectra S_k of the f_k are pairwise disjoint and that the number of elements in $S_k \cap E$ is sufficiently large. To do this first choose some positive integer m_0 for which $a_1 + md \in E$ for all $m \geq m_0$, and then choose an increasing sequence of positive integers $\{m_k\}$ satisfying $m_{k+1}d > m_kd + 2^k - 1$, $k = 0, 1, 2, \dots$. The sequence $\{v_k\}$ is then defined by

$$v_k = a_1 + m_k d, \quad k = 0, 1, 2, \dots$$

With this choice of $\{v_k\}$ we have that, for $p \in (1, 2]$, $q \in [1, p)$,

$$N_p(f) \leq \sum_{k=0}^{\infty} N_p(f_k) \sim \sum_{k=0}^{\infty} (k+1)^{-\frac{1}{q}} 2^{-\frac{k}{q}} 2^{\frac{k}{p}} < \infty$$

and so $f \in A^p$; but

$$\sum_{n \in \mathbb{Z}} |\xi_E \hat{f}(n)|^q = \sum_{k=0}^{\infty} \sum_{n \in S_k \cap E} |\hat{f}_k(n)|^q$$

$$= \sum_{k=0}^{\infty} (k+1)^{-1} 2^{-k} |E \cap S_k|$$

$$= \infty,$$

since $|E \cap S_k| \geq \frac{r}{d}(2^k - 1) \sim 2^k$. // Katelson (private conversation with the author). We first prove the following lemma.

B.1.6 REMARK. We could equally well have modified Construction 2.5.1 to show that the function $f = \sum_{k=0}^{\infty} f_k$, where

$$f_k = (k+1)^{-2} 2^{-k/p} P_k e_{v_k}, \quad k = 0, 1, 2, \dots,$$

with the $\{v_k\}$ chosen as above, also has the property that $f \in A^p$ but $\xi_E \hat{f} \notin \ell^q$. //

B.1.7 COROLLARY. If $p \in (1, 2]$, $q \in [1, p)$, then

$$(A^p, A^q)_I \subsetneq (A^p, A^p)_I.$$

Proof. As previously noted

$$(A^p, A^q)_I \subseteq (A^p, A^p)_I.$$

That this inclusion is strict follows from Theorem B.1.5 together with the fact that the identity map $\hat{\xi}_I$ belongs to $(A^p, A^p)_I$. //

B.2 $M_p \cap C = M(A^p)$, $p \in (1, 2)$.

In 4.5 we noted that, although it was clear that $M_p \cap C = M(A^p)$ for $p = 1$ and $p = 2$, the question of whether or not $M_p \cap C = M(A^p)$ for $p \in (1, 2)$ remained unanswered. We shall show here that equality holds in this case also. The proof of Theorem B.2.2 is essentially due to Professor Y. Katznelson (private conversation with the author). We first prove the following lemma.

B.2.1 LEMMA. Let $p \in (1,2)$ and suppose that

$$\|f\|_{p,p} \leq K \|f\|_{M(p,p)} \quad \text{for all } f \in TP, \quad (\text{B.2.1})$$

where K is a constant independent of f . Then

$$\|f\|_{p,p} \leq K \|f\|_{M(p,p)} \quad \text{for all } f \in M(A^p).$$

Proof. Assume (B.2.1) holds and consider $f \in M(A^p)$.

If $g \in TP$ and $\|\hat{g}\|_p = 1$ then

$$\begin{aligned} \left\| \left((F_N * f)g \right)^\wedge \right\|_p &\leq \|F_N * f\|_{p,p} \\ &\leq K \|F_N * f\|_{M(p,p)}, \text{ by (B.2.1)} \\ &\leq K \|f\|_{M(p,p)}, \text{ by Theorem 4.3.1;} \end{aligned}$$

thus

$$\begin{aligned} \|(fg)^\wedge\|_p &= \lim_{N \rightarrow \infty} \left\| \left((F_N * f)g \right)^\wedge \right\|_p \\ &\leq K \|f\|_{M(p,p)} \end{aligned}$$

and the result follows. //

We are now in a position to prove the following theorem.

B.2.2 THEOREM. $M_p \cap C = M(A^p)$, $p \in (1,2)$.

Proof. From (4.4.4) we have that

$$M_p \cap C \subseteq M(A^p), \quad p \in (1,2). \quad (\text{B.2.2})$$

We shall show here that, for all $f \in M(A^p)$,

$$\|f\|_{p,p} \leq 2\|f\|_{M(p,p)} \quad (\text{B.2.3})$$

and hence that

$$M(A^p) \subseteq M_p \cap C.$$

Because of Lemma B.2.1 we need show only that (B.2.3) holds for

all $f \in TP$. Consider $g \in TP$ such that $\|\hat{g}\|_p = 1$. By the estimates (1.2.10) it is possible, for $p \in (1,2)$, to choose a non-negative integer m so that the function h defined by $h = 2^{-m/p} P_m$ satisfies $\|g\|_\infty \|h\|_\infty \leq 1$. Note also that $\|\hat{h}\|_p = 1$. If s is a positive integer such that $s > \deg(fg)$ repeated use of Lemma 2.1.1 allows us to conclude that, for $f \in TP$,

$$\begin{aligned} \|(fg)^\wedge\|_p &= \|(fg)^\wedge\|_p \|\hat{h}\|_p \\ &= \|(fg)^\wedge\|_p \|\hat{h}_{(2s)}\|_p \\ &= \|(fgh_{(2s)})^\wedge\|_p \\ &\leq N_p(fgh_{(2s)}) \\ &\leq \|f\|_{M(p,p)} N_p(gh_{(2s)}) \\ &= \|f\|_{M(p,p)} [\|g\|_\infty \|h\|_\infty + \|\hat{g}\|_p \|\hat{h}\|_p] \\ &\leq 2\|f\|_{M(p,p)} ; \end{aligned}$$

from which it follows that

$$\|f\|_{p,p} \leq 2\|f\|_{M(p,p)}$$

and our proof is completed. //

B.2.3 COROLLARY. Let $p \in (1,2)$, $q \in (p,2)$. Then

$$M(A^p) \subsetneq M(A^q) .$$

Proof. The inclusion $M(A^p) \subseteq M(A^q)$ follows by combining Theorem B.2.2 with the result that $M_p \subseteq M_q$ (see [24] p.114). That the inclusion $M(A^p) \subseteq M(A^q)$ must be strict was the substance of Theorem 4.2.4. //

B.2.4 REMARKS. For $p \in (1,2)$ the multiplier spaces $M(A^p)$, unlike the spaces A^p themselves, are algebras; we shall consider here the maximal ideal space Δ_p of $M(A^p)$. It is clear that Δ_p contains all the maps of the form $\omega_x : f \mapsto f(x)$, for $x \in T$. It follows from a theorem of M. Zafran (see [40], Theorem 2) and a standard argument (for example see [39], p.140) that there exists a non-zero homomorphism from $M_p \cap \mathbb{C}$ to $\mathbb{C}$ which does not arise as an evaluation map. Thus, by Theorem B.2.2, Δ_p cannot be identified with T . It therefore follows that $[TP]_{M(p,p)} \neq M(A^p)$ and hence, by Lemma 4.4.1, that $M(A^p)$ is not translation-continuous. Recall that some sufficient conditions for $f \in M(A^p)$ to be translation-continuous were given earlier in Theorems 4.4.3 and 4.4.4 and Corollaries 4.4.7 and 4.4.9.

REFERENCES

- [1] L. Alpár. *Generalisation d'un théorème de Wiener et de Lévy.*
Acta.Math.Acad.Sci.Hung. 21(1970),11-19.
- [2] Lynette M. Bloom. *The Fourier multiplier problem for spaces of continuous functions with p -summable transforms.* J.Austral.Math.Soc. 17(1974),319-331.
- [3] Leon Brown. *Subalgebras of L_∞ of the circle group.*
Proc.Amer.Math.Soc. 25(1970),585-587.
- [4] Lynette M. Butler. *Certain non-algebras in harmonic analysis.* Bull.Austral.Math.Soc. 4(1971),247-254.
- [5] R. E. Edwards. *Fourier Series: a modern introduction.*
Vols. I,II. Holt, Rinehart and Winston, New York, Chicago, San Francisco, Toronto, London, 1967.
- [6] R. E. Edwards. *Functional analysis: theory and applications.* Holt, Rinehart and Winston, New York, Chicago, San Francisco, Toronto, London, 1965.
- [7] R. E. Edwards. *Some Fourier division problems.* J.Austral. Math.Soc. 12(1971),167-186.
- [8] R. E. Edwards. *Changing signs of Fourier coefficients.*
Pacific J.Math. 15(1965),463-475.
- [9] R. E. Edwards and G.I. Gaudry. *Littlewood-Paley and multiplier theory.* Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer-Verlag, Berlin, Heidelberg, New York, 1977.

- [10] R. E. Edwards and Kenneth A. Ross. *On p -Sidon sets.*
J. Functional Analysis 15(1974), 404-427.
- [11] M. Fekete. *Über die Faktorfolgen welche die "Klasse" einer Fourierschen Reihe unverändert lassen.*
Acta.Scient.Math.(Szeged) 1(1923), 148-166.
- [12] G. I. Gaudry. *Harmonic analysis on some special groups.*
Fourth year lecture notes, Flinders University of South Australia (1974).
- [13] Michael B. Gregory. *p -Helson sets, $1 < p < 2$.* Israel J.Math.12 (1972), 356-368.
- [14] G. H. Hardy and J. E. Littlewood. *Some problems of diophantine approximation : a remarkable trigonometrical series.* Proc.Nat.Acad.Sci. 2(1916), 583-586.
- [15] G. H. Hardy, J. E. Littlewood and G. Pólya. *Inequalities.*
2nd ed. Cambridge University Press, London, New York, 1952.
- [16] E. Hewitt and Kenneth A. Ross. *Abstract harmonic analysis*
Vols. I, II. Die Grundlehren der mathematischen Wissenschaften, Bände 115, 152. Academic Press, New York; Springer-Verlag, Berlin, Göttingen, Heidelberg, 1963, 1970.
- [17] Edwin Hewitt and Karl Stromberg. *Real and abstract analysis.*
Springer-Verlag, Berlin, Heidelberg, New York, 1969.

- [18] I. I. Hirschman, Jr. *On multiplier transformations*.
Duke Math. J. 26(1959), 221-242.
- [19] L. Hörmander. *Estimates for translation invariant operators in L^p spaces*. Acta Math. 104(1960), 93-140.
- [20] D. Jackson. *Über eine trigonometrische Summe*.
Rend Circ. Mat. Palermo. 32(1911), 257-262.
- [21] G. W. Johnson and Gordon S. Woodward. *On p -Sidon sets*. Indiana Univ. Math. J. 24(1974/1975), 161-167.
- [22] Jean-Pierre Kahane and Yitzhak Katznelson. *Sur les séries de Fourier uniformément convergentes*.
C.R.Acad.Sci.Paris, 261(1965), 3025-3028.
- [23] Yitzhak Katznelson. *An introduction to harmonic analysis*.
John Wiley and Sons, Inc., New York, London, Sydney, Toronto, 1968.
- [24] Ronald Larsen. *An introduction to the theory of multipliers*.
Die Grundlehren der mathematischen Wissenschaften.
Band 175. Springer-Verlag, Berlin, Heidelberg, New York, 1971.
- [25] J. R. McMullen and J. F. Price. *Rudin-Shapiro sequences for arbitrary compact groups*. J. Austral. Math. Soc. 17(Series A)(1976), 421-430.
- [26] D. S. Mitrinovic. *Analytic inequalities*. Die Grundlehren der mathematischen Wissenschaften. Band 165.
Springer-Verlag, Berlin, Heidelberg, New York, 1970.

- [27] Marcel Riesz. *Sur les maxima des formes bilinéaires et sur les fonctionnelles linéaires*. Acta.Math.49 (1927),465-497.
- [28] A. P. Robertson and W. J. Robertson. *Topological vector spaces*. 2nd ed. Cambridge University Press, London, New York, 1973.
- [29] Walter Rudin. *Some theorems on Fourier coefficients*. Proc. Amer. Math. Soc. 10(1959),855-859.
- [30] R. Salem. *A singularity of the Fourier series of continuous functions*. Duke Math. J. 10(1943), 711-716.
- [31] I. Schur. *Bemerkungen zur theorie der beschränkten bilinearformen mit unendlich vielen veränderlichen*. Journal f. Math. 140(1911),1-28.
- [32] S. B. Stečkin. *On bilinear forms*. C. R. (Doklady) de l'Academie des Sciences de l'URSS, N.S.71(1950),237-240.
- [33] E. C. Titchmarsh. *A contribution to the theory of Fourier transforms*. Proc. London Math. Soc. 23(1924),279-287.
- [34] A. Zygmund. *Trigonometric series*. Vols. I,II (combined). 2nd ed. Cambridge University Press, London, New York, 1968.

FURTHER REFERENCES

- [35] Henry Helson. *Note on harmonic functions.*
Proc.Amer.Math.Soc. 4(1953),686-691.
- [36] Henry Helson. *On a theorem of Szëgo.*
Proc.Amer.Math.Soc. 6(1955),235-242.
- [37] Edwin Hewitt and Gunter Ritter. *Über die Integrierbarkeit
von Fourier-Transformierten auf Gruppen. Teil I.
Stetige Funktionen mit kompaktem Träger und eine
Bemerkung über hyperbolische Differentialoperatoren.*
Math.Ann. 224(1976),77-96.
- [38] G. Ritter. *Konstruktion von Operatoren und Kernen mit
Hilfe von Absorptionsmengen.* Math.Ann. 216(1975),51-66.
- [39] W. Rudin. *Fourier Analysis on Groups.* Interscience
Publishers, New York, London, Sydney, 1962.
- [40] Misha Zafran. *The functions operating on multiplier algebras.*
J.Functional Analysis 26(1977),289-314.

INDEX SYMBOLS

NOTES

1. p.34 Consideration of the solutions of certain hyperbolic partial differential equations by Ritter in [38] led Hewitt and Ritter to prove in [37] an analogue, for non-discrete *LCA* groups, of the author's results (2.5.1) and (2.5.2).
2. p.71 The question posed here is answered in B.2.3.
3. p.75 That $M(A^p)$, $p \in (1,2)$, is not translation-continuous is contained in B.2.4.
4. p.87 The question posed here is answered in B.2.2.

INDEX OF SYMBOLS

$A, \ \cdot\ _A$	4	$M(E, F)$	15
A^p, N_p	5	$M\{A^p, A^q\}, \ \cdot\ _{M(p, q)}$	66, 16
$(A^p)'$	46	$M_p(G), M_p, \ \cdot\ _{p, p}$	13
$\{A^p, A^q\}$	45	PM	5
$A^\#$	5	P_m	11
BV	38	P_v^*	12
C	3	R	2
$C^\infty, \ \cdot\ _{(\infty)}$	3	$sp(f)$	19
$\mathbb{C}$	2	s_N^f, σ_N^f	9
$\mathcal{D}$	3	TP	5
D_N	9	T	2
e_k	4	$\tau_\alpha^f, \tau_\alpha^\sigma$	6, 7
(E, F)	13	$u, u_\circ, u^T, \ \cdot\ _u$	31
$\mathcal{F}\ell^p$	5	$V[f], V_\beta[f]$	38, 80
$\check{f}, \check{\sigma}$	6, 7	Z	2
F_N	9	* (convolution)	8
G, Γ	2	$\wedge$	4
$\Lambda_\alpha, \ \cdot\ _{\Lambda_\alpha}$	7	p'	4
Λ_α^1	7	ξ_E	6
$L^p(G), L^p, \ell^p$	4	//	end of proof
$M, \ \cdot\ _M$	5	Δ_p	109
M^k	48		