MICROTURBULENT MAGNETOACTIVE PLASMAS
AND POLARIZED RADIATION FROM THE SOLAR CORONA

by

W.N.-C. SY

A thesis submitted for examination for the degree of
Doctor of Philosophy at the Australian National University

Department of Theoretical Physics
Faculty of Science
Australian National University

Canberra, January 1973

STATEMENT OF ORIGINALITY

Except for references cited in the text, the work presented in this thesis is entirely my own.

ACKNOWLEDGEMENTS

My sincere thanks go to Dr. D.B. Melrose, my supervisor, for many valuable suggestions and numerous critical discussions throughout the course of this research. I express my gratitude to members of the Solar Group, Division of Radiophysics, C.S.I.R.O., Sydney, for discussions on solar radio bursts and for supplying preprints and unpublished materials. In particular, I thank K.V. Sheridan for supplying excellent prints of radio spectrographs, which are reproduced in this thesis. I also express my appreciation to Professor H.A. Buchdahl for his interest in this work and for his helpful comments.

The research scholarship awarded to me by the Australian National University is gratefully acknowledged.

W.N.-C. SY

ABSTRACT

A semiclassical theory of weak nonlinear interactions in a magnetoactive plasma is developed neglecting effects of binary collisions (chapters 2 and 4). Thomson scattering of magnetoionic waves by non-relativistic electrons and inverse Compton scattering of magnetoionic waves by relativistic electrons are discussed together with illustrative examples. Scattering by ions and by electrons in a Maxwellian plasma are treated and compared. Examples of nonlinear scattering of magnetoionic waves by thermal plasma ions are studied.

A theory of polarized radiation arising from nonlinear conversion of microturbulences in electron plasma waves and in electrostatic electron cyclotron waves is developed (chapter 5). On a simplifying assumption about the turbulence spectrum of electron plasma waves, polarized radiation arising from the "plasma emission processes" in a weak background magnetic field is studied in detail. Polarized radiation from nonlinear conversion of a microturbulence in electrostatic electron cyclotron waves is discussed for cases with either weak or strong background magnetic fields.

Type I solar radio bursts and associated noise storm phenomena are investigated in chapter 6. A model of type I solar radio bursts is constructed, based on the "plasma hypothesis" and including induced effects. A mechanism is proposed to explain the strong circular polarization of type I bursts from source regions where the background coronal magnetic field is expected to be weak.

CONTENTS

STATEMENT OF ORIGINALITY	ii
ACKNOWLEDGEMENTS	ii
ABSTRACT	iii

Chapter 1. INTRODUCTION

1. Scope of the Thesis	1
2. Assumptions	4
3. Formalism	6
4. Outline of the Thesis	8

PART I. THEORY

Chapter 2. ELECTRODYNAMIC PROPERTIES OF A MAGNETOACTIVE PLASMA

1. Fundamental Equations	11
2. Response of a Plasma	13
3. Reality, Causality and Symmetries	15
4. Response Tensors	19
5. Wave Properties	22
6. Waves in a Maxwellian Plasma	26
(a) Magnetoionic Waves	29
(b) Electron Plasma Waves	31
(c) Electron Cyclotron Waves	33
(d) Ion-acoustic Waves	36

Chapter 3. WAVE-PARTICLE INTERACTIONS AND PLASMA TURBULENCE

1. Introduction	38
2. Emission by a Single Charge	42
3. Quasilinear Equations	47
4. Wave Damping	49
5. Two-Stream Instabilities	53
(a) Hydrodynamic Instabilities	57
(b) Quasilinear Instabilities	59
6. Velocity-Anisotropy Instabilities	62
7. Drift Instabilities	64

Chapter 4. NONLINEAR INTERACTIONS IN A MAGNETOACTIVE PLASMA

1. Introduction	69
2. Evolution of Distribution Functions	71
3. Scattering of Waves	73
4. Thomson Scattering	76
(a) Scattering Cross-Sections	77
(b) Scattering by Non-relativistic Electrons	79
(c) Scattering by Relativistic Electrons	81
5. Nonlinear Scattering	87
(a) Inverse Maxwellian Tensor	89
(b) Scattering by Ions	90
(c) Scattering by Electrons	92
(d) Absorption Coefficients	94
6. Wave-Wave Interactions	95
7. Formation of Turbulence Spectra	96

Chapter 5. POLARIZED RADIATION FROM PLASMA TURBULENCE

1. Introduction	102
2. Electron Plasma Waves	103
(a) The Fundamental	106
(b) The Second Harmonic	108
(c) Conclusions	110
3. Electron Cyclotron Waves	110
(a) Case $\omega_p \gg \Omega_e$	111
(b) Case $\Omega_e \geq \omega_p$	116
(c) Conclusions	117
4. Amplification of Radiation	118

PART II. APPLICATIONS

Chapter 6. TYPE I SOLAR RADIO BURSTS AND ASSOCIATED NOISE STORM RADIATION

1. Introduction	120
2. Solar Corona	124
3. The Active Region Corona	125
(a) Electron Density	126
(b) Magnetic Field Strength	126
(c) Electron Temperature	131

4. Observations	131
(a) Intensity	135
(b) Polarization	135
(c) Duration	136
(d) Bandwidth	136
(e) Frequency Drift	137
(f) The Source	137
(g) Centre-Limb Variations	138
(h) Associated Phenomena	139
5. Review of Existing Theories	141
(a) Theory of Fung and Yip (1966)	145
(b) Theory of Takakura (1963)	148
(c) Theory of Trakhtengerts (1966)	152
6. Model of Type I Bursts	154
(a) Intensity	155
(b) Polarization	162
7. Discussion	165
 Chapter 7. CONCLUSIONS	
1. Review of Results	168
2. Suggestions for Further Research	170
 Appendix A. CALCULATION OF RESPONSE TENSOR	172
Appendix B. SUMMATION FORMULA (2.96)	175
Appendix C. CALCULATION OF CURRENT DENSITIES DUE TO A CHARGE SPIRALLING IN B_0	176
Appendix D. CALCULATION OF CURRENT DENSITIES IN THOMSON AND NONLINEAR SCATTERINGS	178
Appendix E. CONDITIONS FOR SIMPLIFICATION OF THE INVERSE MAXWELLIAN TENSOR	182
Appendix F. APPROXIMATE FORMS OF THE NONLINEAR RESPONSE TENSOR ϵ_{ijl}	186
 LIST OF SYMBOLS	188
 REFERENCES	191

CHAPTER 1

INTRODUCTION

1. SCOPE OF THE THESIS

In this thesis, a *plasma* is taken to be a fully ionized gas composed of electrons and positive ions. If these particles have speeds which are much less than that of light in vacuo, the plasma is *non-relativistic*. A plasma permeated by a background magnetic field is said to be *magnetoactive*.

Waves in *normal modes* in a plasma are called *plasma waves*. Each normal mode is defined by a dispersion relation, which is a solution or a suitable branch of a solution of the determinantal equation associated with the *homogeneous wave equation*. The homogeneous wave equation is the equation obtained from the *general inhomogeneous wave equation* (see equation (2.48) below) on setting the extraneous current density equal to zero and on neglecting all terms connected with non-linear responses, i.e., all terms quadratic, cubic, ..., in the electric field.

A plasma is *microturbulent* if plasma waves in it have been excited to a level higher than that corresponding to thermodynamic equilibrium. The prefix "micro" distinguishes the type of turbulence considered here from other types of turbulence which could be called "macroscopic". Macroscopic turbulence includes shock waves, bulk transport of gases, "hydromagnetic" turbulence, etc. Unless stated otherwise, the terms *turbulence* or *turbulent* shall henceforth refer to microscopic turbulence and all associated processes are assumed to be microscopic.

In laboratory plasmas and thermonuclear fusion devices, plasma turbulence is manifested by anomalously high electrical resistances, anomalous diffusion of particles, excessive radiation losses, etc. Astrophysically, the existence of plasma turbulence can be inferred from phenomena such as non-thermal radio emission from the solar corona, from the planetary magnetospheres, and from galactic and extragalactic radio sources. Our understanding of the physical processes in these diverse places is closely associated with our understanding of how plasma turbulence is generated and how plasma waves interact with particles and among themselves. The scattering of plasma waves by particles and coalescence and decay processes of plasma waves are examples of *nonlinear* processes, since at least some of the terms connected with nonlinear responses must now be retained in the general inhomogeneous wave equation (equation (2.48)) in treating these processes.

There are many important distinctions to be drawn between laboratory plasmas and astrophysical plasmas. A major difference is that laboratory plasmas generally have dimensions comparable with scale lengths or wavelengths of interest and they cannot be regarded as unconfined, whereas astrophysical plasmas often have dimensions much larger than scale lengths or wavelengths of interest and they can be regarded as unconfined. Because of this difference, laboratory plasmas generally have comparatively large gradients in density and other plasma parameters and the inhomogeneity due to such gradients usually cannot be ignored. By contrast, astrophysical plasmas have relatively small gradients and they can often be regarded as homogeneous. Other important differences are: (1) Laboratory plasmas are generally optically thin, i.e., radiation in emerging from the central regions of the plasma and propagating to the boundary of the plasma passes through

a comparatively short distance, in which no significant absorption takes place, whereas astrophysical plasmas are usually optically thick, particularly for radiation at certain frequencies. (2) Laboratory plasmas can emit radiation due to the presence of currents in the surface layers of the plasma, whereas radiation from astrophysical plasmas can rarely be attributed to such currents. In our theoretical studies, plasmas are assumed to be *homogeneous* and *unconfined*. In astrophysical applications, these assumptions sometimes need modifications, e.g., in treatment of absorption of radiofrequency waves propagating through a stellar atmosphere.

Because almost all plasmas, including laboratory plasmas and those of astrophysical interest, are magnetoactive, one should include the effects of a background magnetic field even if only to determine the conditions under which these effects are unimportant. The important rôle played by magnetic fields in astrophysics in general has been stressed by Alfvén and Fälthammar (1963), and Piddington (1969). Certain astrophysical phenomena may be explicable only by taking into account the effects of a background magnetic field. For example, the strong circular polarization of type I solar radio bursts may be one such case, a detailed discussion of which will be given in the second part of this thesis.

In summary, in the theoretical part of this thesis, we shall study problems associated with the generation and interactions of plasma microturbulence in a non-relativistic, unconfined, homogeneous, magnetoactive plasma, taking into account the effects of the background magnetic field in all our discussions. In the astrophysical part, we shall apply our theoretical results to the study of type I solar radio bursts.

2. ASSUMPTIONS

The following assumptions are made throughout the thesis:

- (a) Nonlinear interactions are *weak*.
- (b) Collisional effects can be neglected in treating the nonlinear interactions.
- (c) *The random phase approximation* for plasma waves is applicable to all nonlinear processes considered.

We discuss each of these assumptions in turn.

(a) Weak Nonlinearity

The Fourier transform of the phenomenological equation (see equation (2.11) below) describing the response of a plasma to a perturbing electric field $\tilde{E}(\underline{k}, \omega)$ can be written as (see equation (2.17) below)

$$D_i(\underline{k}, \omega) = \epsilon_{ij}(\underline{k}, \omega) E_j(\underline{k}, \omega) + \int \frac{d^3 \underline{k}' d\omega' d^3 \underline{k}'' d\omega''}{(2\pi)^4} \delta(\underline{k} - \underline{k}' - \underline{k}'') \times \delta(\omega - \omega' - \omega'') \epsilon_{ij\ell}(\underline{k}', \omega'; \underline{k}'', \omega'') E_j(\underline{k}', \omega') E_\ell(\underline{k}'', \omega'') + \dots, \quad (1.1)$$

where $D(\underline{k}, \omega)$ is the Fourier transform of the displacement current, $\epsilon_{ij}(\underline{k}, \omega)$ is the linear response (dielectric) tensor and $\epsilon_{ij\ell}(\underline{k}', \omega'; \underline{k}'', \omega'')$ is the first nonlinear response tensor. Nonlinear interactions in a plasma are considered *weak* if the series expansion in terms of $\tilde{E}(\underline{k}, \omega)$ on the right hand side of equation (1.1) converges sufficiently rapidly, i.e., only the first few terms need to be considered. In general, the nonlinear interaction can be regarded as weak for sufficiently small electric field amplitudes.

The weak nonlinearity condition can be expressed by the inequality (Tsytovich, 1970):

$$v_E/v_* \ll 1, \quad (1.2)$$

where v_E is the velocity amplitude of a plasma particle oscillating in the field of a wave with amplitude $|E|$ and v_* is the characteristic velocity of a nonlinear process (e.g., it might be a phase velocity in a wave process).

(b) The Neglect of Collisions

The mean time between two successive binary collisions τ_c may be defined by $\tau_c \equiv 1/\nu_c$ where ν_c is the effective collision frequency (see equation (3.44) below). If a process occurs in a time much shorter than τ_c , then binary collisions can be neglected in treating the process. In the literature a plasma is loosely referred to as *collisionless* if binary collisions can be neglected in treating all the processes discussed. For linear processes, a plasma can usually be regarded as collisionless under the condition $n_e \lambda_{De}^3 \gg 1$, where n_e is the electron number density and λ_{De} is the Debye length. For nonlinear processes, there is no such simple general condition for collisions to be negligible. The neglect of collisions usually has to be justified *a posteriori* for each individual nonlinear process.

The inclusion of distant binary collisions usually involves retaining a Fokker-Planck term on the right hand side of the Boltzmann equation (Montgomery and Tidman, 1964) and this complicates the problems considerably. We shall neglect collisions except in the treatments of damping of plasma waves. *Collisional damping* or damping by distant binary collisions (inverse-bremsstrahlung) is important in astrophysical plasmas for the following reasons. (1) Collisional damping is often the most efficient damping mechanism predominating over "collisionless" damping mechanisms, e.g., Landau and cyclotron dampings.

(2) The effect of collisional damping accumulates over the long distances through which the plasma waves propagate.

(c) The Random Phase Approximation

For waves having a width in the frequency spectrum $\delta\omega$, a *correlation time* or a time in which the relative phases of the waves shift significantly can be defined by $\tau_\phi \equiv 1/\delta\omega$. When these waves participate in a physical process which occurs in a time $\tau \gg \tau_\phi$, these waves can be regarded as *random-phase waves* for the purpose of treating the process. Furthermore, the ensemble averages of field quantities of the waves, which are assumed to be equal to the time averages of the corresponding quantities, all vanish over the time interval τ . Except for autocorrelations, all bilinear correlations of field quantities over the time interval τ vanish (see equation (3.1) below). The random phase approximation can be justified by comparing the correlation time τ_ϕ with the interaction time τ and showing $\tau \gg \tau_\phi$.

When the random phase approximation is not valid, we are dealing with processes involving fixed-phase waves whose mode of interaction is qualitatively different from those of random-phase waves.

3. FORMALISM

Among nonlinear processes, weak nonlinear processes have been studied in much more detail so far, because simple methods using series expansions of the form of equation (1.1) are available. Weak nonlinear processes in plasmas without a background magnetic field have been studied extensively (Davidson, 1972). Weak nonlinear processes in magnetoactive plasmas, however, have not been studied quite so extensively, due perhaps, in part, to the inherent mathematical complexities.

associated with spiralling motions of plasma particles in the presence of a background magnetic field. Nevertheless, a number of studies have been made (see references in chapter 4), and of these, *the semiclassical theory* of Tsytovich and Shvartsburg (1966, 1967) is perhaps the simplest and easiest to interpret physically.

When the random phase approximation is valid, the phases of the plasma waves are unnecessary in the description of the interactions and the plasma waves are described as phasmons (photons). This simplified description of plasma waves together with the quantum mechanical language of spontaneous and induced processes (Einstein, 1917) forms the basis of the semiclassical theory. The probabilities used in this formulation are all calculated classically by using the kinetic equations and Maxwell's equations. These probabilities together with the principle of detailed balance for microscopic processes are used to derive equations describing the evolution of distribution functions of plasmons and plasma particles. As the fields of the plasma waves have not really been quantized, the constant $\hbar = h/2\pi$, with $h \equiv$ Planck's constant, appearing in the equations are merely there for "book-keeping" purposes; equivalent equations can be written in which this constant does not appear. In brief, a quantum formalism is used to describe plasma physics in the classical regime.

In this thesis, we shall generalize and extend the semiclassical theory of Tsytovich and Shvartsburg (*ibid.*), and some aspects of the processes occurring in a magnetoactive plasma will be discussed more thoroughly here than in the existing literature. In symbols and notation, we follow a combination of those of Tsytovich (1970), Smith (1970), and Melrose and Sy (1972a,b), but with some alterations. Unless stated otherwise, c.g.s. units will be used throughout this study.

4. OUTLINE OF THE THESIS

In chapter 2, starting from the collisionless Boltzmann-Vlasov equation and Maxwell's equations, we define and derive response tensors for a magnetoactive plasma. We then obtain explicit formal expressions for the linear response (dielectric) tensor and the first nonlinear response tensor in terms of unperturbed one-particle distribution functions. Various properties, viz. those of reality, causality and symmetries of these tensors are discussed. The dielectric tensor for a bi-Maxwellian plasma is rederived. Wave properties are defined and summarized for wave-modes of interest in a Maxwellian magnetoactive plasma. Apart from the formal expression for the first nonlinear response tensor and the wave properties derived for electron cyclotron waves, this chapter is principally of a review nature.

In chapter 3, we review the relevant aspects of wave-particle interactions in a magnetoactive plasma within the framework of linear and quasilinear theory. Although it can be said that the basic physical principles behind wave-particle interactions are well understood, certain aspects of this theory, are either still under development or have been controversial in the literature. Included in the discussions are single-particle emission processes, quasilinear equations, and various classes of microinstabilities. Although new results are only incidental, the chapter has been found convenient (1) to provide a suitable basis for a comparison of linear with nonlinear theories, (2) to clarify the stance taken with respect to controversial issues on linear or quasilinear theory, and (3) to include instabilities, whose significance for astrophysics has remained incompletely explored.

In chapter 4, a semiclassical theory of weak nonlinear interactions in a magnetoactive plasma is developed. The principal contributions to the theory lie in the study of scattering of plasma waves.

After presentation of the semiclassical framework (sections 2 and 3), a scattering cross-section is defined and derived, and Thomson (and inverse Compton) scattering of magnetoionic waves by relativistic and non-relativistic particles is considered in section 4. In particular, Thomson scattering by non-relativistic electrons is treated generally, conditions under which the motion of relativistic electrons can be regarded as rectilinear in treating inverse Compton scattering are derived, and when such conditions are not fulfilled, we study a specific example in detail.

In section 5, after the general inverse Maxwellian tensor has been written down and the conditions under which this tensor simplifies have been derived (Appendix E), nonlinear scattering by thermal ions and electrons are treated and compared. Useful approximate forms of the first nonlinear response tensor are derived in Appendix F. In addition, the absorption coefficient (in time) and the scattering cross-section for spontaneous nonlinear scattering of magnetoionic waves by plasma ions are calculated for special cases of interest.

Formal probabilities for coalescence and decay processes are written down in section 6.

The final section of chapter 4 is devoted to a qualitative discussion of the effect of a background magnetic field on the formation of turbulence spectra.

In chapter 5, the general theory of weak nonlinear processes in a magnetoactive plasma is applied to the study of polarized radiation

arising from plasma turbulence. We consider in detail "plasma emission processes" where the microturbulence is in electron plasma waves. We also consider polarized radiation arising from a microturbulence in electron cyclotron waves, which do not seem to have been considered in this context in the literature. The amplification of radiation in magnetoionic waves, by induced effects, is discussed in the last section.

In chapter 6, we explore the possibilities of applying the results derived in the theoretical part of this thesis to polarized radiation from the solar corona. We have chosen to examine type I solar radio bursts in detail. After introductory discussions on radio emission from the sun (section 1), and on a morphological description of the solar corona (section 2), a simplified model of the solar corona above the active regions is adopted (section 3) in order to evaluate and develop quantitative theories of solar radio bursts. An updated review of the observational data on type I solar radio bursts is presented in section 4. Existing theories of type I solar radio bursts are reviewed and criticized in section 5. A model based on the "plasma hypothesis" is then constructed in an attempt to overcome some of the objections raised against the existing theories. The last section is devoted to a critical appraisal of the model proposed and to some concluding remarks.

In the final chapter, a synopsis of theoretical and astrophysical results found in our investigation is given. In conclusion, suggestions for other possible astrophysical applications of the theory and for further research are made.

PART I

THEORY

"There is a mask of theory over the whole face of nature"

— Whewell, 1831

CHAPTER 2

ELECTRODYNAMIC PROPERTIES OF A MAGNETOACTIVE PLASMA

1. FUNDAMENTAL EQUATIONS

Our theoretical study of a magnetoactive plasma is based on the collisionless Boltzmann-Vlasov equation

$$\frac{\partial f_\alpha}{\partial t} + \underline{v} \cdot \frac{\partial f_\alpha}{\partial \underline{r}} + q_\alpha \left(\underline{E} + \frac{\underline{v} \times \underline{B}}{c} + \frac{\underline{v} \times \underline{B}_0}{c} \right) \cdot \frac{\partial f_\alpha}{\partial \underline{p}} = 0, \quad (2.1)$$

where $f_\alpha \equiv f_\alpha(\underline{p}, \underline{r}, t)$ is the one-particle distribution of particles of species α , with charge q_α and rest mass m_α .

The coordinate axes are chosen so that the background magnetic field is given by $\underline{B}_0 = (0, 0, B_0)$. The electric and magnetic fields, $\underline{E} \equiv \underline{E}(\underline{r}, t)$ and $\underline{B} \equiv \underline{B}(\underline{r}, t)$, are caused by extraneous sources and by plasma particles themselves, so they are related consistently by Maxwell's equations,

$$\left. \begin{aligned} \text{curl } \underline{E} &= -\frac{1}{c} \frac{\partial \underline{B}}{\partial t}, & \text{div } \underline{E} &= 4\pi\rho, \\ \text{curl } \underline{B} &= \frac{1}{c} \frac{\partial \underline{E}}{\partial t} + \frac{4\pi}{c} \underline{j}, & \text{div } \underline{B} &= 0. \end{aligned} \right\} \quad (2.2)$$

The charge and current densities are separated by writing respectively,

$$\rho = \rho^{\text{ext}} + \rho^{\text{ind}}, \quad \underline{j} = \underline{j}^{\text{ext}} + \underline{j}^{\text{ind}}. \quad (2.3)$$

Extraneous charge and current densities are denoted by ρ^{ext} and $\underline{j}^{\text{ext}}$.

The induced charge and current densities are related to the particle distribution functions by

$$\begin{aligned}\rho^{\text{ind}} &= \sum_{\alpha} q_{\alpha} \int d^3 \underline{p} f_{\alpha}(\underline{p}, \underline{r}, t) , \\ \underline{j}^{\text{ind}} &= \sum_{\alpha} q_{\alpha} \int d^3 \underline{p} \underline{v} f_{\alpha}(\underline{p}, \underline{r}, t) .\end{aligned}\quad (2.4)$$

In the study of wave motion it is convenient to use Fourier transforms. The Fourier transform and its inverse for any quantity $F(\underline{r}, t)$ can be defined by

$$\begin{aligned}F(\underline{r}, t) &= \int \frac{d^3 \underline{k} d\omega}{(2\pi)^4} F(\underline{k}, \omega) e^{i(\underline{k} \cdot \underline{r} - \omega t)} , \\ F(\underline{k}, \omega) &= \int d^3 \underline{r} dt F(\underline{r}, t) e^{-i(\underline{k} \cdot \underline{r} - \omega t)} .\end{aligned}\quad (2.5)$$

To ensure the existence of the Fourier transform, the function $F(\underline{r}, t)$ is to be truncated outside a space-time volume whenever necessary, i.e.,

$$\begin{aligned}F(\underline{r}, t) &= 0 \quad \text{outside volume } V , \\ &= 0 \quad \text{outside time interval: } -T/2 < t < T/2 .\end{aligned}\quad (2.5)'$$

Fourier transforming (2.1) and (2.2), one has

$$\begin{aligned}-i\omega f_{\alpha}(\underline{p}, \underline{k}, \omega) + i\underline{k} \cdot \underline{v} f_{\alpha}(\underline{p}, \underline{k}, \omega) + q_{\alpha} \frac{(\underline{v} \times \underline{B}_0)}{c} \cdot \frac{\partial f_{\alpha}}{\partial \underline{p}}(\underline{p}, \underline{k}, \omega) \\ + \int d\lambda^{(2)} q_{\alpha} \left[\underline{E}(\underline{k}', \omega') + \frac{\underline{v} \times \underline{B}(\underline{k}', \omega')}{c} \right] \cdot \frac{\partial f_{\alpha}}{\partial \underline{p}}(\underline{p}, \underline{k}'', \omega'') = 0\end{aligned}\quad (2.6)$$

and

$$\left. \begin{aligned}\underline{k} \times \underline{E}(\underline{k}, \omega) &= \frac{\omega}{c} \underline{B}(\underline{k}, \omega) , & \underline{k} \cdot \underline{E}(\underline{k}, \omega) &= -4\pi i \rho(\underline{k}, \omega) , \\ -\underline{k} \times \underline{B}(\underline{k}, \omega) &= \frac{\omega}{c} \underline{E}(\underline{k}, \omega) + \frac{4\pi i}{c} \underline{j}(\underline{k}, \omega) , & \underline{k} \cdot \underline{B}(\underline{k}, \omega) &= 0 ,\end{aligned}\right\}\quad (2.7)$$

where

$$\int d\lambda^{(2)} \equiv \int \frac{d^3 \underline{k}' d\omega' d^3 \underline{k}'' d\omega''}{(2\pi)^4} \delta^3(\underline{k} - \underline{k}' - \underline{k}'') \delta(\omega - \omega' - \omega'') ,\quad (2.8)$$

and $\delta^3(\underline{k} - \underline{k}' - \underline{k}'')$ and $\delta(\omega - \omega' - \omega'')$ are Dirac delta functions. A quantity

on the left (right) hand side of $:\equiv (\equiv:)$ is defined by known quantities on the right (left) hand side.

2. RESPONSE OF A PLASMA

Electromagnetic fields in a plasma perturb particles and therefore alter their distribution functions. It is possible to make a perturbation expansion of the particle distribution functions in terms of the electric fields of small oscillations in a plasma:

$$f_{\alpha}(\underline{p}, \underline{k}, \omega) = f_{\alpha}^{(0)}(\underline{p}) + f_{\alpha}^{(1)}(\underline{p}, \underline{k}, \omega) + f_{\alpha}^{(2)}(\underline{p}, \underline{k}, \omega) + \dots \quad (2.9)$$

where $f_{\alpha}^{(0)}(\underline{p})$ is the unperturbed homogeneous time-independent distribution, normalized according to,

$$\int d^3p f_{\alpha}^{(0)}(\underline{p}) = 1. \quad (2.10)$$

Here, $f_{\alpha}^{(1)}(\underline{p}, \underline{k}, \omega)$ represents a correction to $f_{\alpha}^{(0)}(\underline{p})$ proportional to the electric field of the plasma waves, $f_{\alpha}^{(2)}(\underline{p}, \underline{k}, \omega)$ represents a higher order correction, proportional to the square of the electric field of the plasma waves, etc.

The phenomenological equation describing the response of a plasma to a perturbing electric field is given by (Landau and Lifshitz, 1960)

$$\underline{D}(\underline{r}, t) :\equiv \underline{E}(\underline{r}, t) + 4\pi \underline{P}(\underline{r}, t), \quad (2.11)$$

where the polarization $\underline{P}(\underline{r}, t)$ is defined by

$$\frac{\partial \underline{P}(\underline{r}, t)}{\partial t} :\equiv \underline{j}^{\text{ind}}(\underline{r}, t). \quad (2.12)$$

The Fourier transform of (2.11) on using (2.12), reads

$$\underline{D}(\underline{k}, \omega) = \underline{E}(\underline{k}, \omega) + \frac{4\pi i}{\omega} \underline{j}^{\text{ind}}(\underline{k}, \omega). \quad (2.13)$$

The corresponding expansion of the induced current density in terms of the electric field $E(\underline{k}, \omega)$ reads,

$$\underline{j}^{\text{ind}}(\underline{k}, \omega) = \underline{j}^{(1)}(\underline{k}, \omega) + \underline{j}^{(2)}(\underline{k}, \omega) + \dots \quad (2.14)$$

The n -th order current density can be written as

$$\begin{aligned} j_i^{(n)}(\underline{k}, \omega) &::= \int d\lambda^{(n)} \sigma_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n) E_{j_1}(\underline{k}_1, \omega_1) \dots E_{j_n}(\underline{k}_n, \omega_n) \\ &::= \sum_{\alpha} q_{\alpha} n_{\alpha} \int d^3 p \, v_i f_{\alpha}^{(n)}(\underline{p}, \underline{k}, \omega) \quad , \end{aligned} \quad (2.15)$$

where n_{α} denotes number density of particles of species α and

$\sigma_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n)$ denotes the n -th order conductivity tensor, with

$$\int d\lambda^{(n)} :: \int \frac{d^3 \underline{k}_1 d\omega_1}{(2\pi)^4} \dots \frac{d^3 \underline{k}_n d\omega_n}{(2\pi)^4} (2\pi)^4 \delta^3(\underline{k} - \underline{k}_1 - \dots - \underline{k}_n) \delta(\omega - \omega_1 - \dots - \omega_n) \quad . \quad (2.16)$$

Substitution of equations (2.14) and (2.15) in (2.13) yields the electric displacement in the form:

$$D_i(\underline{k}, \omega) = \sum_{n=1}^{\infty} \int d\lambda^{(n)} \epsilon_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n) E_{j_1}(\underline{k}_1, \omega_1) \dots E_{j_n}(\underline{k}_n, \omega_n) \quad , \quad (2.17)$$

where $\epsilon_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n)$ is the n -th order response tensor.

The first (linear) response tensor is just the dielectric tensor given by

$$\epsilon_{ij}(\underline{k}, \omega) :: \delta_{ij} + \frac{4\pi i}{\omega} \sigma_{ij}(\underline{k}, \omega) \quad . \quad (2.18)$$

The n -th order response tensor, for $n \geq 2$, is given by,

$$\epsilon_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n) :: \frac{4\pi i}{\omega} \sigma_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n) \quad . \quad (2.19)$$

3. REALITY, CAUSALITY AND SYMMETRIES

The expansion of the electric displacement $\underline{D}(\underline{r}, t)$, defined by (2.11), in terms of the electric field $\underline{E}(\underline{r}, t)$, reads

$$\begin{aligned} D_i(\underline{r}, t) = & \int d^3 \underline{r}_1 dt_1 \epsilon_{ij}(\underline{r}_1, t_1) E_j(\underline{r} - \underline{r}_1, t - t_1) + \int d^3 \underline{r}_1 dt_1 \int d^3 \underline{r}_2 dt_2 \times \\ & \times \epsilon_{ijl}(\underline{r}_1, t_1; \underline{r}_2, t_2) E_j(\underline{r} - \underline{r}_1, t - t_1) E_l(\underline{r} - \underline{r}_2, t - t_2) + \dots \end{aligned} \quad (2.20)$$

It is clear that the phenomenological coefficients (tensors), occurring in (2.20), whose Fourier transforms given by (2.18) and (2.19), are real quantities. Hence the response tensors (2.18) and (2.19) satisfy *reality conditions*:

$$\epsilon_{ij}^*(\underline{k}, \omega) = \epsilon_{ij}(-\underline{k}, -\omega), \quad (2.21)$$

$$\epsilon_{ijl}^*(\underline{k}_1, \omega_1; \underline{k}_2, \omega_2) = \epsilon_{ijl}(-\underline{k}_1, -\omega_1; -\underline{k}_2, -\omega_2), \quad (2.22)$$

and in general,

$$\epsilon_{ij_1 j_2 \dots j_n}^*(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n) = \epsilon_{ij_1 \dots j_n}(-\underline{k}_1, -\omega_1; \dots; -\underline{k}_n, -\omega_n). \quad (2.23)$$

Here, complex conjugation is denoted by '*'. The above conditions follow trivially from the Fourier transforms of real quantities.

The collisionless Boltzmann-Vlasov equation (2.1) is time-reversible, as may be seen by noting its invariance under the transformation: $\underline{p}, \underline{v}, \underline{B}, \underline{B}_0, t \rightarrow -\underline{p}, -\underline{v}, -\underline{B}, -\underline{B}_0, -t$. It cannot describe macroscopically irreversible processes such as relaxation to equilibrium without introducing additional hypotheses such as the stosszahlansatz in Boltzmann's theory and the initial condition in Bogoliubov's theory. It can, however, describe macroscopically irreversible processes such as small damping (or growth) of plasma waves, by imposing causal requirements on the response of the plasma to

small electromagnetic disturbances. That is, the requirement that the response be subsequent to the disturbances is imposed.

To impose causality, the Heaviside step function is introduced,

$$\theta(t) \equiv \begin{cases} 1 & (t > 0) \\ 0 & (t < 0) \end{cases} \quad (2.24)$$

whose Fourier transform can be written as

$$\theta(\underline{k}, \omega) = \frac{i}{\omega + i0} (2\pi)^3 \delta^3(\underline{k}) \quad (2.25)$$

where $i0$ is an infinitesimal positive imaginary constant. Explicit inclusion of causality alters (2.20) to read,

$$\begin{aligned} D_i(\underline{r}, t) = & \int d^3\underline{r}_1 dt_1 \epsilon_{ij}(\underline{r}_1, t_1) \theta(t_1) E_j(\underline{r} - \underline{r}_1, t - t_1) + \\ & + \int d^3\underline{r}_1 dt_1 \int d^3\underline{r}_2 dt_2 \epsilon_{ij\ell}(\underline{r}_1, t_1; \underline{r}_2, t_2) \theta(t_1) \theta(t_2) \times \\ & \times E_j(\underline{r} - \underline{r}_1, t - t_1) E_\ell(\underline{r} - \underline{r}_2, t - t_2) + \dots \end{aligned} \quad (2.26)$$

Fourier transforming (2.26) and equating this with (2.17), one derives the following *causality conditions*:

$$\epsilon_{ij}(\underline{k}, \omega) = \frac{i}{2\pi} \int \frac{d\omega'}{\omega - \omega' + i0} \epsilon_{ij}(\underline{k}, \omega') , \quad (2.27)$$

$$\epsilon_{ij\ell}(\underline{k}_1, \omega_1; \underline{k}_2, \omega_2) = \left(\frac{i}{2\pi}\right)^2 \int \frac{d\omega' d\omega''}{(\omega_1 - \omega' + i0)(\omega_2 - \omega'' + i0)} \epsilon_{ij\ell}(\underline{k}_1, \omega'; \underline{k}_2, \omega'') , \quad (2.28)$$

and in general,

$$\begin{aligned} \epsilon_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n) = & \left(\frac{i}{2}\right)^n \int \frac{d\omega' \dots d\omega^{(n)}}{(\omega_1 - \omega' + i0) \dots (\omega_n - \omega^{(n)} + i0)} \times \\ & \times \epsilon_{ij_1 \dots j_n}(\underline{k}_1, \omega'; \dots; \underline{k}_n, \omega^{(n)}) . \end{aligned} \quad (2.29)$$

The Fourier transforms of the phenomenological tensors $\epsilon_{ij}(\underline{r}, t)$, $\epsilon_{ij\ell}(\underline{r}_1, t_1; \underline{r}_2, t_2)$, etc., satisfy (2.27) - (2.29).

It is clear that the infinitesimal positive imaginary constants give rise to resonant contributions. However, only the non-resonant contributions to the nonlinear response tensors are of interest in this thesis. Hence, only the principal value parts of each of the singular integrals occurring in the nonlinear response tensors, is retained. Unless stated otherwise, the nonlinear response tensors are taken to mean the non-resonant parts of the tensors.

The symmetry properties of the linear response tensor (dielectric tensor) are given by the Onsager reciprocal relations, which are essentially consequences of the "time reversal invariance" of the equations of motion of individual particles (de Groot and Mazur, 1969). For a magnetoactive plasma, they are well-known and can be written in the form,

$$\epsilon_{ij}(\underline{k}, \omega, \underline{B}_0) = \epsilon_{ji}(-\underline{k}, \omega, -\underline{B}_0) = \epsilon_{ji}^*(\underline{k}, -\omega, -\underline{B}_0), \quad (2.30)$$

where reality condition (2.21) has been used.

Generalizations of the Onsager reciprocal relations to obtain symmetry relations for the nonlinear response tensors are not obvious. However, there is an obvious symmetry relation which arises from permuting dummy indices and their associated arguments in (2.16),

$$\epsilon_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n) = \epsilon_{i(j_1 \dots j_n)}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n) \quad (2.31)$$

where brackets (...) around the indices indicate that $1/n!$ times the sum of all tensors obtained by permutation of the indices and associated arguments is to be taken. This symmetry relation reflects the indistinguishability of the induced current density on spatially interchanging any pair of interacting electric fields. For the case $n=2$, equation (2.31) reads,

$$\epsilon_{ij\ell}(\underline{k}_1, \omega_1; \underline{k}_2, \omega_2) = \epsilon_{i\ell j}(\underline{k}_2, \omega_2; \underline{k}_1, \omega_1) . \quad (2.32)$$

The "time reversal invariance" symmetry, which arises from the invariance of the equations of motion of individual particles under the transformation: $\underline{p}, \underline{B}_0, t \rightarrow -\underline{p}, -\underline{B}_0, -t$, can be shown (Melrose, 1972a)[†] to be,

$$\epsilon_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n; \underline{B}_0) = \epsilon_{ij_1 \dots j_n}(\underline{k}_1, -\omega_1; \dots; \underline{k}_n, -\omega_n; -\underline{B}_0) . \quad (2.33)$$

[†] Melrose (1972a) expands the induced current density in terms of the vector potential $A(\underline{r}, t)$ and obtains a response tensor $\kappa_{ij_1 \dots j_n}$ related to $\epsilon_{ij_1 \dots j_n}$ by

$$\epsilon_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n) = 4\pi i \left(\frac{c}{i} \right)^n \frac{\kappa_{ij_1 \dots j_n}(\underline{k}, \omega; \underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n)}{\omega \omega_1 \omega_2 \dots \omega_n} . \quad (2.34)$$

There are also crossing symmetries, arising from the properties of the Poisson brackets in a dynamical formulation (Al'tshul' and Karpman, 1965; Melrose, 1972a). These symmetries have not been established for the general n -th order nonlinear response tensor. However for the cases $n=1$ and $n=2$, they can be expressed (see Melrose, 1972a) as

$$\epsilon_{ij}(\underline{k}, \omega) = \epsilon_{ji}(-\underline{k}, -\omega) , \quad (2.35)$$

$$\begin{aligned} \epsilon_{ij\ell}(\underline{k}_1, \omega_1; \underline{k}_2, \omega_2) &= \epsilon_{j\ell i}(-\underline{k}_1 - \underline{k}_2, -\omega_1 - \omega_2; \underline{k}_2, \omega_2) \\ &= \epsilon_{\ell j i}(\underline{k}_1, \omega_1; -\underline{k}_1 - \underline{k}_2, -\omega_1 - \omega_2) , \end{aligned} \quad (2.36)$$

where only non-resonant parts are retained.

4. RESPONSE TENSORS

The dielectric tensor and the nonlinear response tensors describe completely the electrodynamic properties of a plasma and they depend essentially on the unperturbed homogeneous time-independent distributions $f_{\alpha}^{(0)}(\underline{p})$. To write the tensors explicitly in terms of $f_{\alpha}^{(0)}(\underline{p})$, kinetic equations (2.6) and (2.9) are used to solve for $f_{\alpha}^{(n)}(\underline{p}, \underline{k}, \omega)$. Then application of equations (2.13), (2.14), (2.15) and (2.17) yields the required formal expressions for the tensors. The dielectric tensor and the first nonlinear response tensor read

$$\begin{aligned} \epsilon_{ij}(\underline{k}, \omega) &= \delta_{ij} + \frac{4\pi i}{\omega} \sigma_{ij}(\underline{k}, \omega) \\ &= \delta_{ij} + \frac{4\pi i}{\omega} \sum_{\alpha} \frac{|q_{\alpha}| n_{\alpha}}{\Omega_{\alpha}} \int d^3 \underline{p} \, v_i \int_{\epsilon_{\alpha}}^{\phi} d\phi' e^{P(\phi') - P(\phi)} g_{sj}(\underline{v}', \underline{k}, \omega) \frac{\partial f_{\alpha}^{(0)}}{\partial p'_s} \end{aligned} \quad (2.37)$$

and

$$\begin{aligned} \epsilon_{ijl}(\underline{k}_1, \omega_1; \underline{k}_2, \omega_2) &= \frac{4\pi i}{\omega} \sigma_{ijl}(\underline{k}_1, \omega_1; \underline{k}_2, \omega_2) \\ &= \frac{4\pi i}{\omega} \sum_{\alpha} \frac{q_{\alpha} n_{\alpha}}{\Omega_{\alpha}^2} \int d^3 \underline{p} \, v_i \int_{\epsilon_{\alpha}}^{\phi} d\phi' e^{P(\phi') - P(\phi)} g_{sj}(\underline{v}', \underline{k}_1, \omega_1) \frac{\partial f_{\alpha}^{(0)}}{\partial p'_s} \times \\ &\quad \times \left\{ \int_{\epsilon_{\alpha}}^{\phi'} d\phi'' e^{P_2(\phi'') - P_2(\phi')} g_{rl}(\underline{v}'', \underline{k}_2, \omega_2) \frac{\partial f_{\alpha}^{(0)}}{\partial p''_r} \right\}. \end{aligned} \quad (2.38)$$

The gyrofrequencies for non-relativistic plasma particles are denoted by $\Omega_{\alpha} = |q_{\alpha}| B_0 / m_{\alpha} c$, and also $\omega = \omega_1 + \omega_2$, $\underline{k} = \underline{k}_1 + \underline{k}_2$, in (2.38). The particle velocities are taken to be $\underline{v} = (v_{\perp} \cos \phi, v_{\perp} \sin \phi, v_{\parallel})$. Furthermore,

$$P(\phi) \equiv \frac{i}{\epsilon_{\alpha} \Omega_{\alpha}} \int_{\epsilon_{\alpha}}^{\phi} d\phi' (\omega - \underline{k} \cdot \underline{v}') , \quad (2.39)$$

$$P_2(\phi) \equiv \frac{i}{\epsilon_{\alpha} \Omega_{\alpha}} \int_{\epsilon_{\alpha}}^{\phi} d\phi' (\omega_2 - \underline{k}_2 \cdot \underline{v}') , \quad (2.40)$$

and

$$g_{ij}(\underline{v}, \underline{k}, \omega) \equiv \frac{q_\alpha}{\omega} \left[(\omega - \underline{k} \cdot \underline{v}) \delta_{ij} + k_i v_j \right]. \quad (2.41)$$

For purposes of calculation and further analysis, it is convenient to write equations (2.37) and (2.38) in forms involving infinite sums containing Bessel functions. After some tedious calculations (Appendix A), the equations read,

$$\begin{aligned} \epsilon_{ij}(\underline{k}, \omega) = & \delta_{ij} + \sum_{\alpha} \frac{8\pi^2 q_{\alpha}^2 n_{\alpha}}{\omega^2} \int_0^{\infty} p_{\perp} dp_{\perp} \int_{-\infty}^{\infty} dp_{\parallel} \\ & \left\{ \delta_{i3} \delta_{j3} \frac{v_{\parallel}}{v_{\perp}} \left[v_{\perp} \frac{\partial}{\partial p_{\parallel}} - v_{\parallel} \frac{\partial}{\partial p_{\perp}} \right] + \sum_{s=-\infty}^{\infty} \frac{V_i(s, \underline{k}, \underline{v}) V_j^*(s, \underline{k}, \underline{v})}{\omega - s\Omega_{\alpha} - k_{\parallel} v_{\parallel}} \right. \\ & \left. \left[\frac{\omega - k_{\parallel} v_{\parallel}}{v_{\perp}} \frac{\partial}{\partial p_{\perp}} + k_{\parallel} \frac{\partial}{\partial p_{\parallel}} \right] f_{\alpha}^{(0)}(p_{\perp}, p_{\parallel}) \right\}, \end{aligned} \quad (2.42)$$

and

$$\begin{aligned} \epsilon_{ij\ell}(\underline{k}_1, \omega_1; \underline{k}_2, \omega_2) = & \sum_{\alpha} \frac{8\pi^2 i q_{\alpha}^3 n_{\alpha}}{\omega \omega_1 \omega_2} \int_0^{\infty} p_{\perp} dp_{\perp} \int_{-\infty}^{\infty} dp_{\parallel} \sum_{s, \mu, \mu', v, v'} \\ & \delta_{s+\mu', \mu+v'+v} \exp\{i\epsilon_{\alpha}(\mu' \psi_1 - s\psi + (\mu - v + v')\psi_2)\} V_i(s, \underline{k}, \underline{v}) \\ & \frac{J_{\mu}(z_2)}{\omega - s\Omega_{\alpha} - k_{\parallel} v_{\parallel}} \left\{ \alpha_j^*(\mu', \underline{k}_1, \omega_1) \frac{\partial}{\partial p_{\perp}} + \beta_j^*(\mu', \underline{k}_1, \omega_1) \frac{i\epsilon_{\alpha}(v - v')}{p_{\perp}} \right. \\ & + \gamma_j^*(\mu', \underline{k}_1, \omega_1) \frac{\partial}{\partial p_{\parallel}} \left. \right\} \frac{J_v(z_2)}{\omega_2 - v'\Omega_{\alpha} - k_{2\parallel} v_{\parallel}} \left\{ \alpha_{\ell}^*(v', \underline{k}_2, \omega_2) \frac{\partial}{\partial p_{\perp}} \right. \\ & + \gamma_{\ell}^*(v', \underline{k}_2, \omega_2) \frac{\partial}{\partial p_{\parallel}} \left. \right\} f_{\alpha}^{(0)}(p_{\perp}, p_{\parallel}). \end{aligned} \quad (2.43)$$

The unperturbed distribution $f_{\alpha}^{(0)}(\underline{p})$ has been taken to be independent of azimuthal angles of particle velocities ϕ , i.e. $f_{\alpha}^{(0)}(\underline{p}) = f_{\alpha}^{(0)}(p_{\perp}, p_{\parallel})$.

The wavevectors are written as

$$\left. \begin{aligned} \underline{k} & \equiv (k_{\perp} \cos \psi, k_{\perp} \sin \psi, k_{\parallel}) \\ \underline{k}_1 & \equiv (k_{1\perp} \cos \psi_1, k_{1\perp} \sin \psi_1, k_{1\parallel}) \\ \underline{k}_2 & \equiv (k_{2\perp} \cos \psi_2, k_{2\perp} \sin \psi_2, k_{2\parallel}) \end{aligned} \right\} \quad (2.44)$$

In equation (2.42), $\delta_{s+v', \mu+\mu'+v}$ is the usual Kronecker delta and the quantities α_i^* , β_i^* , γ_i^* are defined as follows:

$$\begin{aligned}\alpha_1(v', \underline{k}_2, \omega_2) &::= \frac{\omega_2 - k_{2\parallel} v_{\parallel}}{v_{\perp}} V_1(v', \underline{k}_2, v) - \delta_{13} J_{v'}(z_2) \frac{v_{\parallel}}{v_{\perp}} (\omega_2 - v' \Omega_{\alpha} - k_{2\parallel} v_{\parallel}) , \\ \gamma_1(v', \underline{k}_2, \omega_2) &::= k_{2\parallel} V_1(v', \underline{k}_2, v) + \delta_{13} J_{v'}(z_2) (\omega_2 - v' \Omega_{\alpha} - k_{2\parallel} v_{\parallel}) , \\ \beta_1(v', \underline{k}_2, \omega_2) &::= \frac{-(\omega_2 - k_{2\parallel} v_{\parallel})}{v_{\perp}} V_2(v', \underline{k}_2, v) + k_{2\perp} v_{\perp} \sin \psi_2 J_{v'}(z_2) , \\ \beta_2(v', \underline{k}_2, \omega_2) &::= \frac{(\omega_2 - k_{2\parallel} v_{\parallel})}{v_{\perp}} V_1(v', \underline{k}_2, v) - k_{2\perp} v_{\perp} \cos \psi_2 J_{v'}(z_2) , \\ \beta_3(v', \underline{k}_2, \omega_2) &::= \frac{k_{\parallel} v_{\parallel}}{v_{\perp}} \{ \sin \psi_2 V_1(v', \underline{k}_2, v) - \cos \psi_2 V_2(v', \underline{k}_2, v) \} ,\end{aligned}\quad (2.45)$$

where

$$\begin{aligned}V_1(v', \underline{k}_2, v) &::= \left[\frac{v_{\perp}}{2} \{ \exp(i\varepsilon_{\alpha} \psi_2) J_{v',-1}(z_2) + \exp(-i\varepsilon_{\alpha} \psi_2) J_{v',+1}(z_2) \} , \right. \\ &\quad \left. -i\varepsilon_{\alpha} \frac{v_{\perp}}{2} \{ \exp(i\varepsilon_{\alpha} \psi_2) J_{v',-1}(z_2) - \exp(-i\varepsilon_{\alpha} \psi_2) J_{v',+1}(z_2) \} , v_{\parallel} J_{v'}(z_2) \right] .\end{aligned}\quad (2.46)$$

Occurring in the above expressions are the Bessel functions $J_{v'}(z_2)$, $J_{\mu'}(z_1)$, etc. where, as usual, the order is designated by v' , μ' , etc. and the kernels are defined by

$$z ::= \frac{k_{\perp} v_{\perp}}{\Omega_{\alpha}} , \quad z_1 ::= \frac{k_{1\perp} v_{\perp}}{\Omega_{\alpha}} , \quad z_2 ::= \frac{k_{2\perp} v_{\perp}}{\Omega_{\alpha}} . \quad (2.47)$$

Expressions for the dielectric tensor, equivalent to (2.42), have been written down elsewhere in the literature (Sitenko and Stepanov, 1957; Stepanov and Kitsenko, 1961; Akhiezer *et al.*, 1967; Melrose and Sy, 1972a). These provide the basis for the study of small amplitude oscillations in a magnetoactive plasma. A somewhat different, but rather unwieldy expression for the first nonlinear response tensor equivalent to (2.43) was written down by Tsytovich and Shvartsburg (1966). In its present form, it was reported by Melrose and Sy (1972a).

This tensor forms the basis for the study of nonlinear interaction of small amplitude oscillations in a magnetoactive plasma.

5. WAVE PROPERTIES

Waves are described by solutions of the relevant wave equation. The general inhomogeneous wave equation can be obtained from Maxwell's equations (2.7):

$$\underline{k} \times \{ \underline{k} \times \underline{E}(\underline{k}, \omega) \} + \frac{\omega^2}{c^2} \underline{D}(\underline{k}, \omega) = -\frac{4\pi i \omega}{c^2} \underline{j}^{\text{ext}}(\underline{k}, \omega), \quad (2.48)$$

where $\underline{D}(\underline{k}, \omega)$ is given by (2.12). This equation is essentially nonlinear, since $\underline{D}(\underline{k}, \omega)$ depends nonlinearly on the electric field $\underline{E}(\underline{k}, \omega)$. In the weak nonlinear approximation, it will be solved by a perturbation procedure. Application of equation (2.17) yields an expanded form of the above equation,

$$\left\{ \frac{k^2 c^2}{\omega^2} (\kappa_i \kappa_j - \delta_{ij}) + \epsilon_{ij}(\underline{k}, \omega) \right\} E_j(\underline{k}, \omega) = -\frac{4\pi i}{\omega} j_i^{\text{ext}}(\underline{k}, \omega) - \sum_{n=2}^{\infty} \int d\lambda^{(n)} \epsilon_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n) E_{j_1}(\underline{k}_1, \omega_1) \dots E_{j_n}(\underline{k}_n, \omega_n), \quad (2.49)$$

with $\underline{\kappa} \equiv \underline{k}/k$. Resonance contributions are retained only in the linear response tensor $\epsilon_{ij}(\underline{k}, \omega)$. By introducing hermitian (h) and antihermitian (a) parts,

$$\begin{aligned} \epsilon_{ij}^h(\underline{k}, \omega) &::= \frac{1}{2} \{ \epsilon_{ij}(\underline{k}, \omega) + \epsilon_{ji}^*(\underline{k}, \omega) \}, \\ \epsilon_{ij}^a(\underline{k}, \omega) &::= \frac{1}{2} \{ \epsilon_{ij}(\underline{k}, \omega) - \epsilon_{ji}^*(\underline{k}, \omega) \}, \end{aligned} \quad (2.50)$$

the causal condition (2.26) gives the Kramers-Kronig relations,

$$\begin{aligned} \epsilon_{ij}^h(\underline{k}, \omega) &= \frac{i}{\pi} \mathcal{P} \int_{-\infty}^{\infty} \frac{d\omega'}{\omega - \omega'} \epsilon_{ij}^a(\underline{k}, \omega'), \\ \epsilon_{ij}^a(\underline{k}, \omega) &= \frac{i}{\pi} \mathcal{P} \int_{-\infty}^{\infty} \frac{d\omega'}{\omega - \omega'} \epsilon_{ij}^h(\underline{k}, \omega'). \end{aligned} \quad (2.51)$$

The Onsager relations (2.30) can be used to establish,

$$\begin{aligned}\epsilon_{ij}^h(\underline{k}, \omega, B_0) &= \epsilon_{ij}^h(\underline{k}, -\omega, -B_0), \\ \epsilon_{ij}^a(\underline{k}, \omega, B_0) &= -\epsilon_{ij}^a(\underline{k}, -\omega, -B_0).\end{aligned}\quad (2.52)$$

It follows from the time-reversal operation: the simultaneous reversal of the signs of ω and B_0 , that ϵ_{ij}^h (ϵ_{ij}^a) describes the time-reversible (time-irreversible) dielectric properties of the plasma. The anti-hermitian part of the dielectric tensor describes such time-irreversible processes as the damping of waves in a plasma. Equation (2.49) can be rewritten as

$$\begin{aligned}\Lambda_{ij}(\underline{k}, \omega) E_j(\underline{k}, \omega) &= -\epsilon_{ij}^a(\underline{k}, \omega) E_j(\underline{k}, \omega) - \frac{4\pi i}{\omega} j_i^{\text{ext}}(\underline{k}, \omega) \\ &- \sum_{n=2}^{\infty} \int d\lambda^{(2)} \epsilon_{ij_1 \dots j_n}(\underline{k}_1, \omega_1; \dots; \underline{k}_n, \omega_n) E_{j_1}(\underline{k}_1, \omega_1) \dots E_{j_n}(\underline{k}_n, \omega_n),\end{aligned}\quad (2.53)$$

where

$$\Lambda_{ij}(\underline{k}, \omega) \equiv \frac{k^2 c^2}{\omega^2} (\kappa_i \kappa_j - \delta_{ij}) + \epsilon_{ij}^h(\underline{k}, \omega). \quad (2.54)$$

Regarding terms on the right hand side of equation (2.53) as source terms, one arrives at the homogeneous wave equation,

$$\Lambda_{ij}(\underline{k}, \omega) E_j(\underline{k}, \omega) = 0, \quad (2.55)$$

which describes free (undamped) oscillations in a plasma. The condition for a non-trivial solution is that the determinant of the coefficients vanishes:

$$\Lambda(\underline{k}, \omega) \equiv \det\{\Lambda_{ij}(\underline{k}, \omega)\} = 0. \quad (2.56)$$

A solution of (2.56) or a suitable branch of a solution $\omega = \omega^\sigma(\underline{k})$ is called the dispersion relation for waves in mode σ . On account of the reality condition (2.20) and the hermitian nature of $\Lambda_{ij}(\underline{k}, \omega)$ it is

seen that

$$\Lambda(\underline{k}, \omega) = \Lambda(-\underline{k}, -\omega) . \quad (2.57)$$

Hence there is an equivalent relation for positive and negative frequencies, viz.,

$$\omega = \omega^\sigma(\underline{k}) = -\omega^\sigma(-\underline{k}) . \quad (2.58)$$

To find the polarization vector $\underline{e}^\sigma(\underline{k})$ defined by,

$$\underline{e}^\sigma(\underline{k}) \equiv \frac{\underline{E}\{\underline{k}, \omega^\sigma(\underline{k})\}}{|\underline{E}\{\underline{k}, \omega^\sigma(\underline{k})\}|} , \quad (2.59)$$

the cofactor tensor $\lambda_{ij}(\underline{k}, \omega)$ of $\Lambda_{ij}(\underline{k}, \omega)$ is introduced (Sitenko and Kirochkin, 1966),

$$\Lambda_{ij}(\underline{k}, \omega) \lambda_{jl}(\underline{k}, \omega) \equiv \Lambda(\underline{k}, \omega) \delta_{il} . \quad (2.60)$$

The elements λ_{ij} expressed in terms of the elements Λ_{ij} read,

$$\lambda_{ij} = \frac{1}{2} \epsilon_{ikl} \epsilon_{jmn} \Lambda_{mk} \Lambda_{nl} , \quad (2.61)$$

where ϵ_{ijk} is the permutation symbol defined by,

$$\epsilon_{ijk} \equiv \begin{cases} 1 & (\text{ijk an even permutation of } 123) \\ -1 & (\text{ijk an odd permutation of } 123) \\ 0 & (\text{otherwise}) \end{cases} . \quad (2.62)$$

When $\Lambda = 0$, Λ_{ij} is of rank two (one less than its order) and so λ_{ij} is of rank one. Therefore any second order minor of λ_{ij} vanishes, i.e. for any $ijkl$,

$$\lambda_{ij} \lambda_{kl} - \lambda_{il} \lambda_{kj} = 0 , \quad \text{when } \Lambda = 0 . \quad (2.63)$$

From equations (2.55) and (2.60), it is easy to see that

$\lambda_{ij}\{\underline{k}, \omega^\sigma(\underline{k})\} \propto e_i^\sigma(\underline{k}) e_j^{\sigma*}(\underline{k})$. It then follows from equation (2.63) and the fact that λ_{ij} is hermitian

$$\lambda_{ij} \lambda_{ji}^* = \lambda_{ij} \lambda_{ij} = \lambda_{ii} \lambda_{jj} = (\lambda_{ss})^2, \quad (2.64)$$

where $\lambda_{ss} \equiv \lambda_{11} + \lambda_{22} + \lambda_{33}$ is the trace of λ_{ij} . By the normalization condition implicit in (2.59), it is thus proved that (Sitenko and Kirochkin, 1966)

$$\lambda_{ij}\{\underline{k}, \omega^\sigma(\underline{k})\} = \lambda_{ss}\{\underline{k}, \omega^\sigma(\underline{k})\} e_i^\sigma(\underline{k}) e_j^{\sigma*}(\underline{k}). \quad (2.65)$$

From this, it is straightforward to show,

$$e_i^\sigma(\underline{k}) = \frac{\lambda_{ij}^\sigma a_j}{\left(a_i^* \lambda_{ij}^\sigma a_j \lambda_{ss}^\sigma \right)^{1/2}}, \quad (2.66)$$

where $\lambda_{ij}^\sigma \equiv \lambda_{ij}\{\underline{k}, \omega^\sigma(\underline{k})\}$, λ_{ss}^σ is the trace of λ_{ij}^σ and a_i is an arbitrary vector, whose phase can be fixed arbitrarily. The phase of e_i^σ can be arbitrarily fixed by a suitable choice of $\underline{a}$.

Apart from the dispersion relation and the polarization vector, the other quantity useful in characterizing waves in mode σ is the ratio of the electrical energy density in the range $d^3\underline{k}/(2\pi)^3$ at $\underline{k}$ to the total energy density in the same range (Melrose, 1968a):

$$\left(\frac{W_E}{W_T} \right)^\sigma \equiv \frac{W_E^\sigma(\underline{k})}{W_T^\sigma(\underline{k})} = \left(\frac{\lambda_{ss}(\underline{k}, \omega)}{\omega \frac{\partial}{\partial \omega} \Lambda(\underline{k}, \omega)} \right)_{\omega=\omega^\sigma(\underline{k})}. \quad (2.67)$$

For longitudinal waves, this expression is replaced by

$$\left(\frac{W_E}{W_T} \right)^\ell = 1 / \left(\omega \frac{\partial}{\partial \omega} \epsilon^\ell(\underline{k}, \omega) \right)_{\epsilon^\ell(\underline{k}, \omega)=0}, \quad (2.67)'$$

where $\epsilon^\ell(\underline{k}, \omega) \equiv \kappa_i \kappa_j \epsilon_{ij}^h(\underline{k}, \omega)$.

6. WAVES IN A MAXWELLIAN PLASMA

A detailed study of free oscillations in a plasma requires assumptions to be made concerning the actual form of the particle functions. From a theoretical point of view, a reasonably general form of the particle distribution functions for which the integrals in (2.42) remain easily tractable is the bi-Maxwellian distribution:

$$f_{\alpha}(p_{\perp}, p_{\parallel}) = \frac{1}{(2\pi)^{3/2} m_{\alpha}^3 V_{\alpha\perp}^2 V_{\alpha\parallel}} \exp \left\{ -\frac{v_{\perp}^2}{2V_{\alpha\perp}^2} - \frac{(v_{\parallel} - u_{\alpha})^2}{2V_{\alpha\parallel}^2} \right\}, \quad (2.68)$$

where

$$V_{\alpha\perp} \equiv \left(\frac{\kappa T_{\alpha\perp}}{m_{\alpha}} \right)^{1/2}, \quad V_{\alpha\parallel} \equiv \left(\frac{\kappa T_{\alpha\parallel}}{m_{\alpha}} \right)^{1/2}. \quad (2.69)$$

The particles are assumed to be non-relativistic, i.e. $V_{\alpha\perp}, V_{\alpha\parallel} \ll c$. The temperatures $T_{\alpha\perp}, T_{\alpha\parallel}$ are measured in °K and κ is a Boltzmann's constant. The bi-Maxwellian streaming distribution includes temperature anisotropies ($V_{\alpha\perp} \neq V_{\alpha\parallel}$) and anisotropies due to streaming motion ($u_{\alpha} \neq 0$).

Insertion of (2.68) into (2.42) yields integrals in $p_{\perp}$ over products of Bessel functions. These integrals can be evaluated from the standard integral (Watson, 1944):

$$\int_0^{\infty} x dx e^{-\rho^2 x^2} J_s(\alpha x) J_s(\beta x) = \frac{1}{2\rho^2} \exp \left\{ -\frac{\alpha^2 + \beta^2}{4\rho^2} \right\} I_s \left(\frac{\alpha\beta}{2\rho^2} \right), \quad (2.70)$$

where I_s is a modified Bessel function of the first kind.

Integrals over $p_{\parallel}$ give (see e.g., Stix, 1962; Akhiezer *et al.*, 1967; Sitenko, 1967)

$$\int_{-c}^c dv_{\parallel} \frac{\exp \left\{ -\frac{(v_{\parallel} - u_{\alpha})^2}{2V_{\alpha\parallel}^2} \right\}}{\omega - s\Omega_{\alpha} - k_{\parallel} v_{\parallel} + i0} \approx \int_{-\infty}^{\infty} dv_{\parallel} \frac{\exp \left\{ -\frac{(v_{\parallel} - u_{\alpha})^2}{2V_{\alpha\parallel}^2} \right\}}{\omega - s\Omega_{\alpha} - k_{\parallel} v_{\parallel} + i0} = -\frac{i\sqrt{\pi}}{k_{\parallel}} F^{(0)}(y_{\alpha s}),$$

where

$$y_{\alpha s} \equiv \frac{\omega - s\Omega_\alpha - k_{\parallel} v_\alpha}{\sqrt{2} k_{\parallel} v_{\alpha\parallel}} , \quad (2.71)$$

$$F^{(0)}(y) \equiv \sqrt{\pi} \frac{k_{\parallel}}{|k_{\parallel}|} e^{-y^2} + \frac{1}{y} \phi(y) ,$$

and

$$\begin{aligned} \phi(y) &\equiv \frac{y}{\sqrt{\pi}} \bigcirc \int_{-\infty}^{\infty} dt \frac{e^{-t^2}}{y-t} = 2ye^{-y^2} \int_0^y e^{t^2} dt \\ &\approx \begin{cases} 2y^2 + \dots & (|y| \ll 1) \\ 1 + \frac{1}{2y^2} + \frac{3}{4y^4} + \dots & (|y| \gg 1) . \end{cases} \end{aligned}$$

Fried and Conte (1961) has published a table of $Z(y) \equiv -iF^{(0)}(-y)$, with $k_{\parallel} = |k_{\parallel}|$.

After some straightforward but lengthy calculations, the dielectric tensor for a bi-Maxwellian-streaming particle distribution reads (Melrose, 1972b)

$$\epsilon_{ij}(\underline{k}, \omega) = \delta_{ij} + \sum_{\alpha} \{C_{ij}^{\alpha}(\underline{k}, \omega) + D_{ij}^{\alpha}(\underline{k}, \omega)\} , \quad (2.72)$$

where

$$C_{ij}^{\alpha}(\underline{k}, \omega) = -\frac{\pi_{\alpha}^2}{\omega^2} \begin{pmatrix} \tau_{\alpha} & 0 & -\frac{k_{\perp}}{k_{\parallel}} \tau_{\alpha} \\ 0 & \tau_{\alpha} & 0 \\ -\frac{k_{\perp}}{k_{\parallel}} \tau_{\alpha} & 0 & \frac{k_{\perp}^2}{k_{\parallel}^2} \tau_{\alpha} - \frac{\omega^2}{k_{\parallel}^2 v_{\alpha\parallel}^2} \end{pmatrix}$$

$$D_{ij}^{\alpha}(\underline{k}, \omega) \equiv \sum_{s=-\infty}^{\infty} \frac{\pi_{\alpha}^2}{\omega^2} \frac{ie^{-\lambda_{\alpha}}}{\sqrt{2} k_{\parallel} v_{\alpha\parallel}} F^{(0)}(y_{\alpha s}) \pi_{ij}^{\alpha}(\underline{k}, \omega; s)$$

$$\{s\Omega_{\alpha} + (\omega - s\Omega_{\alpha} - k_{\parallel} v_{\alpha})(1 - \tau_{\alpha})\} , \quad (2.73)$$

where

$$\pi_{ij}^{\alpha}(\underline{k}, \omega; s) \equiv$$

$$\begin{pmatrix} \frac{s^2}{\lambda_{\alpha}} I_s & i\epsilon_{\alpha} s(I'_s - I_s) & \frac{s}{\lambda_{\alpha}} \frac{k_{\perp}}{k_{\parallel}} \left(\frac{\omega - s\Omega_{\alpha}}{\Omega_{\alpha}} \right) I_s \\ -i\epsilon_{\alpha} s(I'_s - I_s) & \left(\frac{s^2}{\lambda_{\alpha}} + 2\lambda_{\alpha} \right) I_s - 2\lambda_{\alpha} I'_s & -i\epsilon_{\alpha} \frac{k_{\perp}}{k_{\parallel}} \left(\frac{\omega - s\Omega_{\alpha}}{\Omega_{\alpha}} \right) (I'_s - I_s) \\ \frac{s}{\lambda_{\alpha}} \frac{k_{\perp}}{k_{\parallel}} \left(\frac{\omega - s\Omega_{\alpha}}{\Omega_{\alpha}} \right) I_s & i\epsilon_{\alpha} \frac{k_{\perp}}{k_{\parallel}} \left(\frac{\omega - s\Omega_{\alpha}}{\Omega_{\alpha}} \right) (I'_s - I_s) & \frac{1}{\lambda_{\alpha}} \frac{k_{\perp}}{k_{\parallel}} \left(\frac{\omega - s\Omega_{\alpha}}{\Omega_{\alpha}} \right) I_s \end{pmatrix}. \quad (2.74)$$

The plasma frequency of particles of species α is denoted by π_{α} . The argument of the modified Bessel functions I_s is λ_{α} , and I'_s is the derivative of I_s with respect to λ_{α} . These quantities are defined as follows:

$$\pi_{\alpha}^2 \equiv \frac{4\pi q_{\alpha}^2 m_{\alpha}}{m_{\alpha}}, \quad \tau_{\alpha} \equiv 1 - \frac{V_{\alpha\perp}^2}{V_{\alpha\parallel}^2}, \quad \lambda_{\alpha} \equiv \left(\frac{k_{\perp} V_{\alpha\perp}}{\Omega_{\alpha}} \right)^2. \quad (2.75)$$

Of special interest are the waves in a stationary equilibrium plasma ($u_{\alpha} = 0$, $\tau_{\alpha} = 0$). In this case, on writing $V_{\alpha\perp} = V_{\alpha\parallel} = V_{\alpha}$, the dielectric tensor reads (Sitenko and Stepanov, 1957; Stix, 1962; Akhiezer *et al.*, 1967; Sitenko, 1967)

$$\begin{aligned} \epsilon_{ij}(\underline{k}, \omega) &= \delta_{ij} - \sum_{\alpha} \frac{\pi_{\alpha}^2}{\omega^2} \left\{ e^{-\lambda_{\alpha}} \sum_{s=-\infty}^{\infty} \frac{y_{\alpha 0}}{y_{\alpha s}} \pi_{ij}^{\alpha}(\underline{k}, \omega; s) \times \right. \\ &\quad \left. \times \left[\phi(y_{\alpha s}) - i\sqrt{\pi} \frac{k_{\parallel}}{|k_{\parallel}|} e^{-y_{\alpha s}^2} \right] - 2y_{\alpha 0}^2 \delta_{i3} \delta_{j3} \right\}, \end{aligned} \quad (2.76)$$

where

$$y_{s\alpha} \equiv \frac{\omega - s\Omega_{\alpha}}{\sqrt{2} k_{\parallel} V_{\alpha}} \quad \text{and} \quad \lambda_{\alpha} \equiv \left(\frac{k_{\perp} V_{\alpha}}{\Omega_{\alpha}} \right)^2. \quad (2.77)$$

From equation (2.76), since $\pi_{ij}^{\alpha}(\underline{k}, \omega; s)$ is hermitian, the electric tensor can be separated easily into hermitian and antihermitian parts:

$$\epsilon_{ij}^h(\underline{k}, \omega) = \delta_{ij} - \sum_{\alpha} \frac{\pi_{\alpha}^2}{\omega^2} \left\{ e^{-\lambda_{\alpha}} \sum_{s=-\infty}^{\infty} \frac{y_{\alpha 0}}{y_{\alpha s}} \pi_{ij}^{\alpha}(\underline{k}, \omega; s) \phi(y_{\alpha s}) - 2y_{\alpha 0}^2 \delta_{i3} \delta_{j3} \right\}, \quad (2.78)$$

and

$$\epsilon_{ij}^a(\underline{k}, \omega) = i\sqrt{\pi} \frac{k_{||}}{|k_{||}|} \sum_{\alpha} \frac{\pi_{\alpha}^2}{\omega^2} e^{-\lambda_{\alpha}} \sum_{s=-\infty}^{\infty} \frac{y_{\alpha 0}}{y_{\alpha s}} \pi_{ij}^{\alpha}(\underline{k}, \omega; s) e^{-y_{\alpha s}^2}. \quad (2.79)$$

The free oscillations, described by the homogeneous wave equation (2.55) with $\Lambda_{ij}(\underline{k}, \omega)$ given by (2.54) and (2.78), have been fairly thoroughly studied to various degrees of approximation (Sitenko and Stepanov, 1957; Ginzburg, 1964; Stix, 1962; Sitenko, 1967; Shafranov, 1967; Bernstein, 1958; among others). The general procedure for obtaining wave properties of interest has been outlined in the previous section. A summary of these wave properties for various wave modes is given below.

(a) Magnetoionic Waves

In a magnetoactive plasma, magnetoionic waves in the ordinary (O) and extraordinary (X) modes can exist. Above their cutoff frequencies, they can escape from the medium to infinity. In this sense, at least, they are the counterpart of transverse electromagnetic waves in an isotropic (unmagnetized) plasma. Their importance astrophysically lies simply in the fact that radiofrequency waves from many astrophysical sources are just magnetoionic waves escaping from magnetoactive plasmas.

Spatial dispersion has only a slight effect on the properties of magnetoionic waves in a non-relativistic plasma (Silin, 1960; Sitenko, 1967). The cold plasma approximation: $\lambda_{\alpha} \ll 1$, $|y_{\alpha 0}|$, $|y_{\alpha \pm}| \gg 1$, is adequate in describing magnetoionic waves. For convenience, the two natural frequencies of a magnetoactive plasma, ω_p and Ω_e , are

incorporated into the formulae by introducing two magnetoionic parameters:

$$X \equiv \frac{\omega_p^2}{\omega^2}, \quad Y \equiv \frac{\Omega_e}{\omega}. \quad (2.80)$$

The wavevector is taken to be $\underline{k} = k (\sin\theta \cos\psi, \sin\theta \sin\psi, \cos\theta)$ and the wave modes are denoted by $\sigma = 0, X$ for the ordinary and extraordinary modes respectively. The wave properties are given by (Melrose and Sy, 1972a)

$$\eta_\sigma^2 = 1 - \frac{XT_\sigma}{T_\sigma - Y\cos\theta}, \quad (2.81)$$

$$\underline{e}^\sigma = (1 + K_\sigma^2 + T_\sigma^2)^{-1/2} (a_\sigma \cos\psi - i\sin\psi, a_\sigma \sin\psi + i\cos\psi, b_\sigma), \quad (2.82)$$

and

$$\left(\frac{W_E}{W_T} \right)^\sigma = \frac{1 + K_\sigma^2 + T_\sigma^2}{2(1 + T_\sigma^2) \eta_\sigma \frac{\partial}{\partial \omega} (\omega \eta_\sigma)}, \quad (2.83)$$

where

$$\begin{aligned} a_\sigma &\equiv K_\sigma \sin\theta + T_\sigma \cos\theta, & b_\sigma &\equiv K_\sigma \cos\theta - T_\sigma \sin\theta, \\ K_\sigma &\equiv \frac{XYT_\sigma \sin\theta}{(1-X)(T_\sigma - Y\cos\theta)} = \frac{XY\sin\theta(1 + T_\sigma Y\cos\theta)}{(1-X)(1 - Y^2) - XY^2\sin^2\theta}, \end{aligned} \quad (2.84)$$

$$T_\sigma \equiv \frac{Y(1-X)\cos\theta}{\frac{1}{2}Y^2\sin^2\theta \mp \Delta} = \frac{-\frac{1}{2}Y^2\sin^2\theta \mp \Delta}{Y(1-X)\cos\theta},$$

and

$$\Delta^2 = \frac{1}{4} Y^4 \sin^4\theta + (1-X)^2 Y^2 \cos^2\theta.$$

The upper (lower) signs refer to $\sigma = 0$ ($\sigma = X$). Equation (2.81) is sometimes known as the Appleton-Hartree dispersion relation (Stix, 1962). It can be seen from (2.82) and (2.84) that T_σ and K_σ describe respectively the transverse and longitudinal parts of the polarization. Furthermore, T_σ is the axial ratio of the polarization ellipse, whilst the sign of T_σ determines the handedness (RH for $T_\sigma > 0$).

(b) Electron Plasma Waves

In a magnetoactive plasma, the longitudinal (electrostatic) oscillations at high frequencies (above ω_p) are the electron plasma waves and the electrostatic electron cyclotron waves. These waves in general have much lower phase velocities than the magnetoionic waves (Furutani and Kalman, 1965). Consequently, they couple more readily to non-relativistic or mildly relativistic charges, or in the language of modern electrodynamics, the emission and absorption of these waves by non-relativistic or mildly relativistic charges are preferred. Suprathermal non-relativistic charges in a magnetoactive plasma could generate a turbulence of these waves by coherent or incoherent emission.

The dispersion equation for high-frequency longitudinal waves may be obtained from equations (2.54), (2.55) and (2.78). It reads (Stix, 1962; Sitenko, 1967),

$$\epsilon^l(\underline{k}, \omega) = 1 + \frac{\omega_p^2}{k^2 V_e^2} \left\{ 1 - \sum_{s=-\infty}^{\infty} \frac{y_{eo}}{y_{es}} \phi(y_{es}) \zeta_s(\lambda_e) \right\} = 0, \quad (2.85)$$

where ionic contributions have been justifiably neglected,

$$\zeta_s(\lambda_e) := e^{-\lambda_e} I_s(\lambda_e) \quad \text{and} \quad \lambda_e := (k_{\perp} V_e / \Omega_e)^2. \quad (2.86)$$

On the assumption that

$$|y_{es}| = \left| \frac{\omega - s\Omega_e}{\sqrt{2} k_{\parallel} V_e} \right| \gg 1, \quad \text{for all } s, \quad (2.87)$$

and by the equations (2.71), the dispersion equation (2.85) can be written approximately as

$$\epsilon^l(\underline{k}, \omega) \approx 1 - \frac{\omega_p^2}{\omega^2} \cos^2 \theta \zeta_0(\lambda_e) - \frac{2\omega_p^2 \sin^2 \theta}{\omega^2 - \Omega_e^2} \frac{\zeta_1(\lambda_e)}{\lambda_e} + \frac{\omega_p^2 \sin^2 \theta}{\Omega_e^2} \frac{1 - \zeta_0(\lambda_e) - 2\zeta_1(\lambda_e)}{\lambda_e} - \frac{\omega_p^2 \sin^2 \theta}{\Omega_e^2} \sum_{s=2}^{\infty} \frac{2\omega^2}{\omega^2 - s^2 \Omega_e^2} \frac{\zeta_s(\lambda_e)}{\lambda_e} = 0$$

To the zeroth order in λ_e , equation (2.88) reads

$$\epsilon^{\ell}(\underline{k}, \omega) = 1 - \frac{\omega_p^2}{\omega^2} \cos^2 \theta - \frac{\omega_p^2 \sin^2 \theta}{\omega^2 - \Omega_e^2} = 0, \quad (2.89)$$

which gives rise to the well-known cold-plasma dispersion relations for electron plasma waves:

$$\omega_{\pm}^2 = \frac{1}{2} (\omega_p^2 + \Omega_e^2) \pm \frac{1}{2} \{ (\omega_p^2 + \Omega_e^2)^2 - 4\omega_p^2 \Omega_e^2 \cos^2 \theta \}^{1/2}. \quad (2.90)$$

Small corrections to (2.90), to allow for the effects of thermal electron motion, can be estimated by taking equation (2.88) to first order in λ_e . Including thermal corrections, the dispersion relations for electron plasma waves read (Sitenko, 1967)

$$(\omega^{el})^2 = \omega_{\pm}^2 + 3k^2 v_e^2 F(\omega_{\pm}, \theta), \quad (2.91)$$

where

$$F(\omega, \theta) := \frac{\omega^4}{1 + \omega_p^2 \Omega_e^2 \sin^2 \theta / (\omega^2 - \Omega_e^2)^2} \times \\ \times \left\{ \frac{\sin^4 \theta}{(\omega^2 - \Omega_e^2)(\omega^2 - 4\Omega_e^2)} + \frac{\sin^2 \theta \cos^2 \theta (2\omega^4 - \omega^2 \Omega_e^2 + \frac{1}{3} \Omega_e^4)}{\omega^2 (\omega^2 - \Omega_e^2)^3} + \frac{\cos^4 \theta}{\omega^4} \right\}. \quad (2.92)$$

In a plasma where $\omega_p > \Omega_e$ the upper branch (+ve sign) of the electron plasma waves corresponds directly to the Langmuir waves in an isotropic plasma, as might be seen by allowing $\Omega_e \rightarrow 0$.

The electron plasma waves have polarization vector $\underline{e} = \underline{\kappa} \equiv \underline{k}/k$ and their ratios of electrical to total energy can be approximated from (2.89) to give

$$\left(\frac{W_E}{W_T} \right)^{el} = \frac{1}{2} \left\{ 1 + \frac{\omega_p^2 \Omega_e^2 \sin^2 \theta}{(\omega^2 - \Omega_e^2)^2} \right\}^{-1}. \quad (2.93)$$

One must bear in mind in equations (2.91) - (2.93) ω and θ are not independent variables: ω depends explicitly on θ .

(c) Electron Cyclotron Waves

The electron cyclotron waves are longitudinal waves at harmonics of the electron gyrofrequency. When $\Omega_e > \omega_p$, the fundamental coincides with the upper branch of the electron plasma waves. When the $\omega_p > \Omega_e$, the fundamental is strongly damped.

The electron cyclotron waves were first correctly treated by Bernstein (1958); for this reason, they are sometimes known as Bernstein waves.

Equation (2.88), which is valid under the assumption (2.87), can be written as

$$\begin{aligned} \epsilon^l(\underline{k}, \omega) = 1 - \frac{\omega_p^2 \cos^2 \theta}{\omega^2} \zeta_0(\lambda_e) - \frac{\omega_p^2 \sin^2 \theta}{\omega^2 - \Omega_e^2} \frac{2\zeta_1(\lambda_e)}{\lambda_e} \\ - \frac{1}{k^2 \lambda_{De}^2} \sum_{s=2}^{\infty} \frac{2s^2 \Omega_e^2}{\omega^2 - s^2 \Omega_e^2} \zeta_s(\lambda_e) = 0, \end{aligned} \quad (2.94)$$

where $\lambda_{De} \equiv V_e / \omega_p$ is the Debye length for electrons. For wave propagation perpendicular to the background magnetic field

$$\epsilon^l(\underline{k}, \omega) = 1 - \frac{1}{k^2 \lambda_{De}^2} \sum_{s=1}^{\infty} \frac{2s^2 \Omega_e^2}{\omega^2 - s^2 \Omega_e^2} \zeta_s(\lambda_e) = 0, \quad (2.95)$$

a result derived by Bernstein (1958). It is shown in Appendix B that

$$\sum_{s=1}^{\infty} \frac{2s^2 \zeta_s(\lambda_e)}{s^2 - q^2} \equiv 1 - \pi q \cot \pi q \zeta_q(\lambda_e). \quad (2.96)$$

Hence, the dispersion equation (2.95) can be expressed as

$$k^2 \lambda_{De}^2 = \alpha(q, \lambda_e) \equiv 1 - \pi q \cot \pi q \zeta_q(\lambda_e), \quad (2.97)$$

with $q \equiv \omega / \Omega_e$. Dispersion curves for longitudinal waves propagating perpendicularly to the background magnetic field for various thermal plasmas, have been obtained by Stone and Auer (1965), see Fig. 2.1.

Fig. 2.1: Dispersion curves for longitudinal waves propagating at right angles to the background magnetic field in a thermal plasma. For this figure only, v_0 is the thermal velocity of electrons and ω_b is the electron gyrofrequency (Stone and Auer, 1965).

At arbitrary angles of propagation, the more general dispersion equation (2.94) must be used. On account of the smallness of $\zeta_s(\lambda_e)$ for $s \geq 2$, all terms in the sum of equation (2.94) are negligible except in the neighbourhood of resonances $\omega \approx s\Omega_e$. The dispersion equation reads

$$\begin{aligned} \epsilon_o^l(\underline{k}, \omega) = 1 - \frac{\omega_p^2 \cos^2 \theta}{\omega^2} \zeta_o(\lambda_e) - \frac{\omega_p^2 \sin^2 \theta}{\omega^2 - \Omega_e^2} \frac{\zeta_1(\lambda_e)}{\lambda_e} \\ - \frac{\zeta_s(\lambda_e)}{k^2 \lambda_{De}^2} \frac{2s^2 \Omega_e^2}{\omega^2 - s^2 \Omega_e^2} = 0. \end{aligned} \quad (2.98)$$

In general, a cubic equation in ω^2 must be solved to give a dispersion relation for cyclotron waves at $\omega \approx s\Omega_e$. However, if

$$\frac{\zeta_s(\lambda_e)}{k^2 \lambda_{De}^2} \ll \epsilon_o^l(\underline{k}, s\Omega_e), \quad (2.99)$$

where

$$\epsilon_o^l(\underline{k}, \omega) \equiv 1 - \frac{\omega_p^2 \cos^2 \theta}{\omega^2} \zeta_o(\lambda_e) - \frac{\omega_p^2 \sin^2 \theta}{\omega^2 - \Omega_e^2} \frac{\zeta_1(\lambda_e)}{\lambda_e}, \quad (2.100)$$

an approximate dispersion relation can be found:

$$\omega^2 = s^2 \Omega_e^2 + 2s^2 \Omega_e^2 \frac{\zeta_s(\lambda_e)}{k^2 \lambda_{De}^2 \epsilon_o^l(\underline{k}, s\Omega_e)} \quad (2.101)$$

In this case, the ratio of the electrical to total energy density is given by

$$\left(\frac{W_E}{W_T} \right)^c = \frac{\zeta_s(\lambda_e)}{k^2 \lambda_{De}^2 \{ \epsilon_o^l(\underline{k}, s\Omega_e) \}^2}. \quad (2.102)$$

Contrary to Sugihara's approximations (1963), $\epsilon_o^l(\underline{k}, s\Omega_e) < 1$ generally, and $\epsilon_o^l(\underline{k}, s\Omega_e) = 1$, only when $\omega^2 \approx s^2 \Omega_e^2 \gg \omega_p^2 + \Omega_e^2$.

The assumption that wave damping is exponentially small (2.87) and equation (2.101) together determine a cone of propagation, outside

which the theory presented is no longer valid. This cone is small and is in a direction perpendicular to the background magnetic field, with

$$|\cot\theta| \ll \frac{s}{\sqrt{2}\lambda_e} \frac{\zeta_s(\lambda_e)}{k^2 \lambda_{De}^2 \epsilon_o^l(\underline{k}, s\Omega_e)}, \quad (2.103)$$

for cyclotron waves at $\omega \approx s\Omega_e$.

(d) Ion-Acoustic Waves

Ion-acoustic waves are longitudinal waves, which exist only when the electron temperature is at least a few times the ion temperature, and whose frequencies are much lower than the ion-cyclotron frequency Ω_i . Specifically, under the following conditions:

$$\left. \begin{aligned} v_s \ll v_A \ll v_e, \\ \lambda_i, \lambda_e \ll 1, \quad \text{and} \quad |k_{\parallel}| v_i \ll \omega \ll |k_{\parallel}| v_e, \end{aligned} \right\} \quad (2.104)$$

where the sound speed and the Alfvén speed are defined respectively by,

$$v_s^2 \equiv \frac{z_i k T_e}{m_i} = \pi_i^2 \lambda_{De}^2, \quad v_A^2 \equiv \frac{B_0^2}{4\pi m_i n_i} = \frac{\Omega_i^2}{\pi_i^2} c^2, \quad (2.105)$$

a direct application of the equations (2.71) and (2.78) gives a simplified dielectric tensor,

$$\epsilon_{ij}^h(\underline{k}, \omega) = \frac{c^2}{v_A^2} \begin{pmatrix} 1 & \frac{i\omega}{\Omega_i} & \cdot \\ -\frac{i\omega}{\Omega_i} & 1 & -\frac{i\Omega_i}{\omega} \tan\theta \\ \cdot & \frac{i\Omega_i}{\omega} \tan\theta & -\frac{\Omega_i^2}{\omega^2} \left(1 - \frac{\omega^2}{k^2 v_s^2 \cos^2\theta} \right) \end{pmatrix}. \quad (2.106)$$

From this expression it is straightforward to use equations (2.54), (2.55) and (2.67)' to obtain the wave properties for the ion-acoustic

waves,

$$\omega^2 = k^2 v_s^2 \cos^2 \theta, \quad (2.107)$$

and

$$\begin{aligned} \left(\frac{W_E}{W_T} \right)^S &= \frac{k^2 \lambda_{De}^2}{2} \left(1 + \frac{v_s^4}{v_A^4} \frac{\Omega_i^2}{\omega^2} \sin^2 \theta \cos^2 \theta \right) / \left(1 + \frac{c^2}{v_A^2} k^2 \lambda_{De}^2 \right) \\ &\approx \frac{k^2 \lambda_{De}^2}{2}. \end{aligned} \quad (2.108)$$

The ion-acoustic waves can be identified as slow magnetoacoustic waves (Sitenko, 1967) when $\omega \ll \Omega_i$, as assumed above. However, ion-acoustic waves can exist up to frequency $\pi_i = (m_e/m_i)^{1/2} \omega_p$. In the range $\Omega_i \ll \omega \leq \pi_i$, their wave properties are essentially those of ion-acoustic waves in an equilibrium plasma without a background magnetic field:

$$\omega^2 = k^2 v_s^2 \left(\frac{1}{1+k^2 \lambda_{De}^2} + \frac{3T_i}{T_e} \right), \quad (2.109)$$

and

$$\left(\frac{W_E}{W_T} \right)^S = \frac{k^2 \lambda_{De}^2}{2} \left(\frac{1}{1+k^2 \lambda_{De}^2} - \frac{6T_i}{T_e} \right). \quad (2.110)$$

These relations can be derived in a straightforward way, from equations (2.67)' and (2.78) by allowing $\Omega_\alpha \rightarrow 0$.

CHAPTER 3

WAVE-PARTICLE INTERACTIONS AND PLASMA TURBULENCE

1. INTRODUCTION

Amongst the topics of current interest in plasma physics, which could fall within the general category of wave-particle interactions in a magnetoactive plasma, are the emission of plasma waves by single charges, statistical acceleration of charges by plasma turbulence, wave damping and certain classes of microinstabilities. From the point of view of plasma turbulization, microinstabilities are particularly efficient mechanisms, as they can generate a suprathermal level of plasma waves in an avalanche fashion. This chapter is devoted principally to the study of wave-particle interactions in the random phase approximation, with particular emphasis on the causes of plasma turbulence.

Interactions involving random phase waves (plasmons) is qualitatively different from interactions involving fixed phase waves (Tsytovich, 1970). For this reason, the range of validity of the random phase approximation should be clarified. When the interaction time of a physical process τ , say the growth time of plasma waves in a microinstability, is much longer than the correlation (phase-shift) time τ_ϕ of the waves, i.e. $\tau \gg \tau_\phi$, the phase of a wave must change rapidly and repeatedly during the time of interaction. Hence, the ensemble averages of field quantities of the wave all vanish over the time interval τ . Except for autocorrelations, all bilinear correlations of field quantities vanish (Sitenko, 1967; Landau and Lifshitz,

1969):

$$\langle E_i^\sigma(\underline{k}, \omega) E_j^{\sigma*}(\underline{k}', \omega') \rangle = \langle E_i^\sigma E_j^{\sigma*} \rangle(\underline{k}, \omega) (2\pi)^4 \delta(\underline{k} - \underline{k}') \delta(\omega - \omega') , \quad (3.1)$$

where the brackets $\langle \dots \rangle$ denote the averaging operation and all other symbols have their usual meanings.

Thus when the random phase approximation is applicable, the essentially wave-like characteristics of the plasma waves, viz., their phases, are unnecessary in the description of the interaction. This means a purely "quantum" description of the plasma waves, the plasmons, is sufficient. This simplification together with Einstein's (1917) concept of induced emission and absorption forms the basis for a relatively simple semiclassical approach (Tsytovich, 1967; Kovrizhnykh, 1968; Melrose, 1968a), which has the advantage of easy physical interpretation.

Plasmons with energy $\hbar\omega^\sigma(\underline{k})$ and momentum $\hbar\underline{k}$, where $\hbar \equiv h/2\pi$ and h is Planck's constant, are described by a distribution function $N^\sigma(\underline{k})$ which is the number of plasmons in the range $d^3\underline{k}/(2\pi)^3$ at $\underline{k}$. The total energy density in the waves of mode σ is then given by

$$W^\sigma = \int \frac{d^3\underline{k}}{(2\pi)^3} \hbar\omega^\sigma(\underline{k}) N^\sigma(\underline{k}) , \quad (3.2)$$

where the quantum formalism has been used for convenience, and the fields of the waves have not really been quantized. In fact, Fourier transforming the electrical energy density by (2.5), one obtains

$$W_E^\sigma(\underline{k}) = \lim_{V \rightarrow \infty} \frac{|\underline{E}^\sigma(\underline{k})|^2}{8\pi V} , \quad (3.3)$$

where $\underline{E}\{\underline{k}, \omega^\sigma(\underline{k})\} \equiv: \underline{E}^\sigma(\underline{k}) 2\pi\delta(\omega - \omega^\sigma(\underline{k}))$ and V is the spatial volume containing waves of mode σ . The quantities in (3.2) are related to the classical electric field amplitude $\underline{E}^\sigma(\underline{k})$ by equations (3.3) and (2.67),

viz.,

$$\hbar\omega^\sigma(\underline{k}) N^\sigma(\underline{k}) = \lim_{V \rightarrow \infty} \frac{|\underline{E}^\sigma(\underline{k})|^2}{8\pi V} \left(\frac{\omega}{\lambda_{ss}} \frac{\partial \Lambda}{\partial \omega} \right)_{\omega=\omega^\sigma(\underline{k})}. \quad (3.4)$$

To describe the particles quantum mechanically, a set of quantum numbers $\{p_{||}, n\}$ is introduced, where the continuous quantum number $p_{||}$ describes motion along the background magnetic field B_0 and the simple harmonic oscillator quantum number n defined by (Tsytovich, 1962; Melrose, 1968a)

$$p_{\perp}^2 = m_{\alpha} \Omega_{\alpha} \hbar (2n+1), \quad n = 0, 1, 2, \dots \quad (3.5)$$

$$\Omega_{\alpha} = |q_{\alpha}| B_0 / m_{\alpha} c,$$

describes motion perpendicular to B_0 .

Finally, the semiclassical theory of wave-particle interactions requires the introduction of a probability of spontaneous emission $w_{\alpha}^{\sigma}(s, \underline{p}, \underline{k})$, which is the probability per unit time that a particle of species α , with momentum $\underline{p}$, emits a plasmon of mode σ , with momentum $\hbar \underline{k}$ and at the same time decreases its harmonic quantum number by s . The probability is defined by

$$\bar{P}^{\sigma} \equiv: \sum_{s=-\infty}^{\infty} \int \frac{d^3 \underline{k}}{(2\pi)^3} \hbar \omega^{\sigma}(\underline{k}) w^{\sigma}(s, \underline{p}, \underline{k}), \quad (3.6)$$

where subscript α has been dropped for convenience. The probability $w^{\sigma}(s, \underline{p}, \underline{k})$ is calculated classically.

From equations (2.53) and (2.54), the electric field generated by an extraneous current $\underline{j}^{\text{ext}}(\underline{k}, \omega)$ is given by

$$\underline{E}_i(\underline{k}, \omega) = -\frac{4\pi i}{\omega} \frac{\lambda_{ij}(\underline{k}, \omega)}{\Lambda(\underline{k}, \omega)} j_j^{\text{ext}}(\underline{k}, \omega), \quad (3.7)$$

where $\lambda_{ij}(\underline{k}, \omega)$ is defined by (2.61). The electric field in mode σ is found by setting $\omega \approx \omega^\sigma(\underline{k})$, whereupon,

$$\frac{1}{\Lambda(\underline{k}, \omega)} \approx \frac{1}{(\omega - \omega^\sigma(\underline{k}) + i0) \left(\frac{\partial \Lambda}{\partial \omega} \right)_{\omega = \omega^\sigma(\underline{k})}}. \quad (3.8)$$

From equations (2.65) and (2.67),

$$\left(\frac{\lambda_{ij}(\underline{k}, \omega)}{\frac{\partial}{\partial \omega} \Lambda(\underline{k}, \omega)} \right)_{\omega = \omega^\sigma(\underline{k})} = \omega^\sigma(\underline{k}) \left(\frac{W_E}{W_T} \right)^\sigma e_i^{\sigma*}(\underline{k}) e_j^{\sigma*}(\underline{k}). \quad (3.9)$$

Classically, the time-averaged power radiated by a charge into waves in mode σ is given by (see e.g. McKenzie, 1967)

$$\begin{aligned} \bar{P}^\sigma &= - \lim_{T \rightarrow \infty} \frac{1}{T} \int_{-T/2}^{T/2} dt \int d^3 \underline{r} \underline{E}^\sigma(\underline{r}, t) \cdot \underline{j}^{\text{ext}}(\underline{r}, t) \\ &= - \lim_{T \rightarrow \infty} \frac{1}{T} \int \frac{d\omega}{2\pi} \int \frac{d^3 \underline{k}}{(2\pi)^3} \text{Re} \left(\underline{E}^\sigma(\underline{k}, \omega) \cdot \underline{j}^{\text{ext}}(\underline{k}, \omega) \right), \end{aligned} \quad (3.10)$$

where a Fourier transform has been taken and 'Re' denotes the real part. Application of (3.7) - (3.10) together with the Plemelj formula,

$$\frac{1}{\omega - \omega' \pm i0} = \mathcal{P} \frac{1}{\omega - \omega'} \mp i\pi \delta(\omega - \omega'), \quad (3.11)$$

yields

$$\bar{P}^\sigma = - \lim_{T \rightarrow \infty} \frac{4\pi}{T} \int \frac{d^3 \underline{k}}{(2\pi)^3} \left(\frac{W_E}{W_T} \right)^\sigma |\underline{e}^\sigma \cdot \underline{j}^{\text{ext}}(\underline{k}, \omega^\sigma(\underline{k}))|^2. \quad (3.12)$$

For a charge spiralling in B_0 , it can be shown (Appendix C) that

$$\underline{j}^{\text{ext}}(\underline{k}, \omega) = \sum_{s=-\infty}^{\infty} 2\pi q e^{-is(\phi + \epsilon\psi)} \underline{V}(s, \underline{k}, \underline{v}) \delta(\omega - k_{\parallel} v_{\parallel} - s\Omega^*), \quad (3.13)$$

where $\Omega^* \equiv \Omega mc^2/E$, $E^2 = (p_{\perp}^2 c^2 + p_{\parallel}^2 c^2 + m^2 c^4)$, $\underline{k}$ is given by (2.44) and $\underline{V}(s, \underline{k}, \underline{v})$ is defined by (2.46) with $z \equiv k_{\perp} v_{\perp} / \Omega^*$. On substitution of (3.13) into (3.12) and noting that from (2.5)',

$$\sum_{s'=-\infty}^{\infty} \delta(\omega - s\Omega^* - k_{\parallel} v_{\parallel}) \delta(\omega - s'\Omega^* - k_{\parallel} v_{\parallel}) = \frac{T}{2\pi} \delta(\omega - s\Omega^* - k_{\parallel} v_{\parallel}) , \quad (3.14)$$

the time-averaged power radiated reads

$$\bar{P}^{\sigma} = \sum_{s=-\infty}^{\infty} \int \frac{d^3 \underline{k}}{(2\pi)^3} 8\pi^2 q^2 \left(\frac{W_E}{W_T} \right)^{\sigma} |e^{\sigma*} \cdot \underline{V}(s, \underline{k}, v_{\parallel})|^2 \delta(\omega^{\sigma}(\underline{k}) - s\Omega^* - k_{\parallel} v_{\parallel}) . \quad (3.15)$$

It is clear that from geometrical symmetry, $\bar{P}^{\sigma}$ is independent of the azimuthal angle ψ of the wavevector $\underline{k}$. Hence for the present discussion $\underline{V}(s, \underline{k}, v_{\parallel})$ can be set equal to

$$\underline{V}(s, z, v_{\parallel}) \equiv \left(v_{\perp} \frac{s}{z} J_s(z), -i\epsilon v_{\perp} J_s'(z), v_{\parallel} J_s(z) \right) , \quad (3.16)$$

which corresponds to the case $\psi = 0$ in $\underline{V}(s, \underline{k}, v_{\parallel})$ with $z \equiv k_{\perp} v_{\perp} / \Omega^*$. From (3.6), (3.15) and (3.16), the probability of spontaneous emission

$w_{\alpha}^{\sigma}(s, \underline{p}, \underline{k})$ is identified to be

$$w_{\alpha}^{\sigma}(s, \underline{p}, \underline{k}) = \frac{8\pi^2 q_{\alpha}^2}{\hbar \omega^{\sigma}(\underline{k})} \left(\frac{W_E}{W_T} \right)^{\sigma} |e^{\sigma*} \cdot \underline{V}(s, z_{\alpha}, v_{\parallel})|^2 \delta(\omega^{\sigma}(\underline{k}) - s\Omega_{\alpha}^* - k_{\parallel} v_{\parallel}) . \quad (3.17)$$

A probability essentially the same as (3.17) was derived by Melrose (1968a). Earlier calculations for low frequency waves, based on the same approach were made by Tsytovich (1963). Still earlier calculations for the probability based on the Hamiltonian formalism were made by Ginzburg and Eidman (1959), though they did not take spatial dispersion into account.

2. EMISSION BY A SINGLE CHARGE

The problem of radiation from an electron in a magnetoactive plasma is of considerable interest in various aspects of astrophysics, magnetospheric physics and thermonuclear research. For example, it is well known that the radio and optical spectra of the Crab Nebula can be

attributed to synchrotron radiation from relativistic electrons. Type IV solar radiation at radiofrequencies has a similar origin (Kundu, 1965; Zheleznyakov, 1970). The decametric radiation from Jupiter has been attributed (Ellis, 1962) to cyclotron (gyromagnetic) radiation from non-relativistic electrons in the Jovian magnetosphere. The same physical idea has been used to explain certain features of the very low frequency (VLF) and low frequency (LF) emissions (3 - 300 kHz) from the earth's magnetosphere (McKenzie, 1963; Liemohn, 1965). Attempts have been made (Sugihara, 1963; Canobbio and Croci, 1963; Stone and Auer, 1965) to explain the observed noise radiation at electron cyclotron harmonics from laboratory plasmas (Crawford, 1965) on the basis of electron-cyclotron wave emission from non-relativistic electrons. In all the above theories, the electrons emit waves incoherently, i.e., the total emission is taken to be the arithmetic sum of emission by single electrons.

Radiation from a charge in a magnetoactive plasma has been studied by Eidman (1958, 1959, 1962), Liemohn (1965), Mansfield (1967), McKenzie (1963, 1967), Melrose (1968a) and Trulsen and Fejer (1970). In the last mentioned paper, many errors and inconsistencies of previous derivations are pointed out. Our time-averaged power radiated as given by equation (3.15) agrees with those derived by the last three authors. Some numerical examples are given by McKenzie (1967) and Trulsen and Fejer (1970).

As examples, consider time-averaged power radiated by a charge into magnetoionic waves and longitudinal waves.

For magnetoionic waves, with spatial dispersion neglected, direct application of equations (2.82) - (2.84) and (3.16) gives the power radiated per unit frequency range per unit solid angle as

$$\frac{d^2 \bar{P}^\sigma}{d\omega d\Omega} = \sum_{s=-\infty}^{\infty} \frac{q^2 \omega^2 \eta_\sigma \beta_\perp^2}{2\pi c(1+T_\sigma^2)} \left\{ \frac{a_\sigma - \beta_\parallel \eta_\sigma T_\sigma}{\beta_\perp \eta_\sigma \sin\theta} J_s(z) - \epsilon J_s'(z) \right\}^2 \delta\{\omega(1-\eta_\sigma \beta_\parallel \cos\theta) - s\Omega^*\}, \quad (3.18)$$

where $z = \omega \eta_\sigma \beta_\perp \sin\theta / \Omega^*$ and the right hand side is often known as the emissivity. In the limit of ultrarelativistic electrons radiating in vacuo, equation (3.18) coincides with the well known results for synchrotron radiation. Because of its importance for the study of cyclotron and synchrotron radiation in a plasma, equation (3.18) has been derived many times in the literature (Eidman, 1958, 1959; Melrose, 1968b, 1972b; Ramaty, 1969; Trulsen and Fejer, 1970; Sakurai, 1972).

For longitudinal waves, equations (3.15) and (3.16) gives

$$\bar{P}^\sigma = \sum_{s=-\infty}^{\infty} \int \frac{d^3 k}{(2\pi)^3} 8\pi^2 q^2 \left(\frac{W_E}{W_T} \right)^\sigma \frac{(\omega^\sigma(k))^2}{k^2} J_s^2(z) \delta(\omega^\sigma(k) - s\Omega^* - k_\parallel v_\parallel), \quad (3.19)$$

where σ denotes any longitudinal wave mode.

The frequencies of electron plasma waves have a strong angular dependence, but a weak wavenumber dependence, as may easily be seen from equation (2.91). Consequently it is not very useful to write down equations in the form of (3.18) for electron plasma waves. Instead, the delta function in (3.19) is integrated away with respect to the wavenumber k to obtain the time-averaged power radiated per unit solid angle,

$$\frac{dP}{d\Omega} = \sum_{s=-\infty}^{\infty} \frac{q^2 \omega^2}{2\pi |v_\parallel \cos\theta|} J_s^2 \left(\frac{\omega - s\Omega^*}{\Omega^*} \frac{v_\perp \sin\theta}{v_\parallel \cos\theta} \right) / \left(1 + \frac{\omega_p^2 \Omega_e^2 \sin^2\theta}{(\omega^2 - \Omega_e^2)^2} \right), \quad (3.20)$$

where the frequencies are given by

$$\omega^2 = \omega_\pm^2 + 3 \left(\frac{\omega_\pm - s\Omega^*}{v_\parallel \cos\theta} \right)^2 v_e^2 F(\omega_\pm, \theta), \quad (3.21)$$

subjected to the usual proviso that the last term in (3.21) is a small

correction. Consequently, $(v_{\parallel} \cos \theta)^2 \gg 3F(\omega_{\pm}, \theta) V_e^2 (\omega_{\pm} - s\Omega_e^*)^2 / \omega_{\pm}^2$, and the apparent divergence in (3.20) as $|v_{\parallel} \cos \theta| \rightarrow 0$ is not permitted in our approximation. Yip (1970) derived a more complicated expression than (3.20) by including the effects of spatial dispersion in the delta function of (3.19). However, as his numerical results show, effects of spatial dispersion are slight; the simpler expression (3.20) gives essentially the same results.

Consider the emission of electron-cyclotron waves at frequency $\omega \approx n\Omega_e$. The weak damping condition (2.87) together with the Doppler condition in (3.19) implies

$$\frac{v_{\parallel}^2}{2V_e^2} \left(\frac{\omega - n\Omega_e}{\omega - s\Omega_e^*} \right)^2 \gg 1, \quad (3.22)$$

where, for non-relativistic electrons, $\Omega_e^* = \Omega_e$ and (3.22) can only be satisfied in general for $n=s$ and $v_{\parallel}^2 \gg V_e^2$. It is possible to use (3.19) directly to obtain the average power radiated. However, because of the complicated k dependence of $(W_E/W_T)^c$, it is more illuminating to obtain an alternative expression for time-averaged power radiated by using equations (3.7), (3.10), (3.13) and (3.14) only:

$$\bar{P}^L = \sum_{s=-\infty}^{\infty} \int \frac{d^3 k}{(2\pi)^4} d\omega \, 8\pi^2 q^2 \, \text{Im} \left(\frac{1}{\epsilon_L} \right) \frac{\omega}{k^2} J_s^2(z) \delta(\omega - s\Omega_e^* - k_{\parallel} v_{\parallel}). \quad (3.23)$$

This equation is valid for longitudinal waves and is equivalent to equation (3.19). For electron cyclotron waves in the vicinity of $\omega \approx n\Omega_e$, an emissivity analogous to (3.18) can be written down,

$$\frac{d^2 \bar{P}^c}{d\omega d\Omega} = \frac{q^2}{\pi^2} \int dk \, \omega J_n^2(z) \, \text{Im} \left(\frac{1}{\epsilon_L} \right) \delta(\omega - n\Omega_e - k_{\parallel} v_{\parallel}). \quad (3.24)$$

It can be deduced from the dispersion relation (2.98) that (Stone and Auer, 1965; Bekefi, 1966)

$$\text{Im}\left(\frac{1}{\epsilon_l}\right) \approx \frac{\pi}{2} k_0 \delta(k-k_0), \quad (3.25)$$

where k_0 is given, for any ω and θ , by $\epsilon^l(\omega, \theta, k_0) = 0$. Substitution of (3.25) in (3.24) gives

$$\frac{d^2 \bar{P}^c}{d\omega d\Omega} = \frac{q^2 \omega k_0}{2\pi} J_n^2 \left(\frac{k_0 v_{\perp} \sin \theta}{\Omega_e} \right) \delta(\omega - n\Omega_e - k_0 v_{\parallel} \cos \theta). \quad (3.24)'$$

Integration over solid angles, bearing in mind $\sin \theta \approx 1$, gives

$$\frac{d\bar{P}^c}{d\omega} = \frac{q^2 \omega}{|v_{\parallel}|} J_n^2 \left(\frac{k_0 v_{\perp}}{\Omega_e} \right), \quad (3.26)$$

which differs from Stone and Auer's (1965, eqn. (37)) result by a factor $8\pi^2$.[†] Numerical examples have been given by these authors. Significant emission occurs only for $(k_0 v_{\perp})^2 \geq n^2 \Omega_e^2$ and waves are not strongly damped only if $v_{\parallel}^2 \gg v_e^2$.

It is straightforward to use equations (2.107) and (2.108) to obtain the emissivity of ion-acoustic waves by charges in a magnetoactive plasma

$$\frac{d\bar{P}^s}{d\omega d\Omega} = \frac{q^2 \omega^4}{2\pi \Omega_i^2 v_s |\cos^3 \theta|} \sum_{n=-\infty}^{\infty} J_n^2 \left(\frac{\omega v_{\perp}}{\Omega_i^* v_s} \tan \theta \right) \delta \left\{ \omega \left(1 - \frac{v_{\parallel}}{v_s} \right) - n\Omega_i^* \right\}. \quad (3.27)$$

It is clear from the above equation that Cerenkov emission is in general not possible. Cyclotron emission by electrons is highly unlikely, because one requires $|v_{\parallel}| \approx (n\Omega_e/\omega) v_s \gg (m_i/m_e)^{1/2} v_e$, at which speed, other wave modes are preferentially emitted. Cyclotron emission by ions is possible provided $v_e > |v_{\parallel}| = |n\Omega_i v_s/\omega| \gg v_s$, and it is via normal Doppler emission ($n > 0$) when $v_{\parallel} < 0$ and via anomalous Doppler emission ($n < 0$) when $v_{\parallel} > 0$. In either case,

$$\frac{d\bar{P}^s}{d\Omega} = \frac{q^2 \Omega_i^2 v_s^4}{2\pi |v_{\parallel}|^5 |\cos^3 \theta|} \sum_{n=1}^{\infty} n^4 J_n^2 \left(\frac{n v_{\perp} \tan \theta}{v_{\parallel}} \right), \quad (3.28)$$

[†] Stone and Auer have made a numerical slip in their equation (36).

where $|\cos\theta| > n\Omega_i/kV_e$.

3. QUASILINEAR EQUATIONS

Due to various deviations from thermodynamic equilibrium, such as those caused by the presence of drifting or streaming particles or those caused by temperature anisotropies, a plasma is frequently found to be unstable with respect to the generation of plasma waves. The general criteria for determining whether a plasma with given particle distribution functions, is unstable or not, are well-known (see e.g., Akhiezer *et al.*, 1967, p.86). To follow the development of such micro-instabilities, the first logical attempt is to consider the changes in the unstable particle distribution functions resulting from the "feedback" effect of the waves.

From such considerations, a set of quasilinear equations was derived by Vedenov *et al.* (1961, 1962) and Drummond and Pines (1962), for an isotropic electron plasma. The basic assumptions, the range of validity and the consequences of the quasilinear theory have been subjects of controversy, see for example Klozenberg and Bernstein (1970), Montgomery and Bodner (1971), Fukai and Harris (1972) and references therein. Much of the confusion disappears if one bears clearly in mind that the time-irreversible processes, such as damping of waves, described by the quasilinear equations, must arise from the resonant parts of the equations, involving delta functions. The resonant parts are direct consequences of imposing causality on the time-reversible Boltzmann-Vlasov equation (2.1), as done in the previous chapter. The non-resonant parts, which appear most naturally from a classical derivation, describe reversible adiabatic diffusion (Kadomtsev, 1965; Sagdeev and Galeev, 1969), which apparently heats the non-resonant particles, on account of increased kinetic energy associated with

increased wave amplitudes. However, it is the resonant interactions, which describe the alteration to the particle distribution functions.

In almost all derivations of the quasilinear equations, except that of Al'tshul' and Karpman (1966), the random phase approximation has been used. In some derivations, emphasis has been placed on the non-resonant parts (Vedenov *et al.*, 1962; Drummond and Pines, 1962; Vahala and Montgomery, 1970) and in others, on the resonant parts (Pines and Schrieffer, 1962; Tsytovich, 1967; Harris, 1969; Fukai and Harris, 1972). A semiclassical or quantum mechanical derivation makes the random phase approximation and takes into account only resonant interactions, but it introduces the effects of spontaneous emission easily and naturally into the equations.

In a semiclassical derivation for a magnetoactive plasma (Tsytovich and Shvartsburg, 1966; Melrose, 1968a) the total (spontaneous and induced) emission probability and the absorption probability is given respectively by (Einstein, 1917) $\{1 + N^\sigma(\underline{k})\} w_\alpha^\sigma$ and $N^\sigma(\underline{k}) w_\alpha^\sigma$, where

$$w_\alpha^\sigma \equiv w_\alpha^\sigma(\underline{p}, \underline{k}) \equiv \sum_{s=-\infty}^{\infty} w_\alpha^\sigma(s, \underline{p}, \underline{k}) \quad (3.29)$$

and the wave mode and particle species are denoted as usual by σ and α respectively. The change in particle momenta on emission or absorption of a plasmon of momentum $\hbar \underline{k}$ is given by

$$\begin{aligned} \Delta p_{\parallel} &= \hbar k_{\parallel} \ll p_{\parallel}, \\ \Delta p_{\perp}^2 &= \hbar m_{\alpha} \gamma s \Omega_{\sigma} \ll p_{\perp}^2, \quad (s = 0, 1, 2, \dots). \end{aligned} \quad (3.30)$$

By a Taylor series expansion, the change in particle distribution function in an emission or absorption process is given by

$$\Delta f_{\alpha}(\underline{p}) = \pm (D_s \pm \frac{\hbar}{2} D_s^2 + \dots) f_{\alpha}(\underline{p}), \quad (3.31)$$

where

$$D_s \equiv \frac{s\Omega_\alpha}{v_\perp} \frac{\partial}{\partial p_\perp} + k_\parallel \frac{\partial}{\partial p_\parallel}, \quad (3.32)$$

and the + (-) sign applies to absorption (emission) processes. From the above considerations, it is straightforward to obtain the following set of quasilinear equations (see e.g., Melrose, 1968a),

$$\left(\frac{\partial}{\partial t} + \mathbf{v}_g^\sigma \cdot \frac{\partial}{\partial \mathbf{r}} \right) N^\sigma(\mathbf{k}) = \sum_\alpha \sum_{s=-\infty}^{\infty} n_\alpha \int d^3p w_\alpha^\sigma(s, \mathbf{p}, \mathbf{k}) \{1 + N^\sigma(\mathbf{k}) \hbar D_s\} f_\alpha(\mathbf{p}), \quad (3.33)$$

$$\frac{\partial f_\alpha(\mathbf{p})}{\partial t} = \sum_\sigma \sum_{s=-\infty}^{\infty} \int \frac{d^3k}{(2\pi)^3} \hbar D_s \{w_\alpha^\sigma(s, \mathbf{p}, \mathbf{k}) \{1 + N^\sigma(\mathbf{k}) \hbar D_s\} f_\alpha(\mathbf{p})\}, \quad (3.34)$$

where $\mathbf{v}_g^\sigma \equiv \partial \omega^\sigma(\mathbf{k}) / \partial \mathbf{k}$ is the group velocity of waves in mode σ . It is frequently convenient to define a quasilinear growth (or damping) rate in (3.33) by

$$\gamma_\alpha^\sigma(\mathbf{k}) \equiv - \sum_{s=-\infty}^{\infty} n_\alpha \int d^3p w_\alpha^\sigma(s, \mathbf{p}, \mathbf{k}) \hbar D_s f_\alpha(\mathbf{p}), \quad (3.35)$$

where induced growth (damping) of plasma waves in mode σ , by particles of species α , occurs when $\gamma_\alpha^\sigma(\mathbf{k}) < 0$ ($\gamma_\alpha^\sigma(\mathbf{k}) > 0$). The quasilinear process is exactly analogous to collisionless Landau ($s=0$) and cyclotron damping ($s \neq 0$), only that a microinstability has been made possible by the presence of non-Maxwellian distributions of particles.

4. WAVE DAMPING

Waves in an equilibrium magnetoactive plasma may be damped by at least two processes: a collisionless wave-particle interaction process, which has been called Landau-cyclotron damping or gyromagnetic resonance absorption, and a collisional, inverse-bremsstrahlung process, which arises from binary collisions between plasma particles.

The collisionless damping decrements can be calculated by retaining only the first term on the right hand side of the inhomogeneous wave equation (2.53) and solving the resulting equation approximately. Or equivalently, they can be calculated from the quasilinear theory by taking an appropriate Maxwellian particle distribution function in equation (3.35). The latter procedure, together with equations (2.70), (2.80) - (2.84) and (3.17), gives, after some straightforward calculations, the collisionless damping decrement for magnetoionic waves by Maxwellian plasma electrons,

$$\gamma^\sigma(\omega, \theta) = \sum_{s=-\infty}^{\infty} \left(\frac{\pi}{2} \right)^{1/2} \frac{\omega_p^2}{\omega \eta_\sigma \beta_e |\cos \theta|} \frac{e^{-\lambda_e} A(\omega, \theta, s)}{(1+T_\sigma^2) \eta_\sigma \frac{\partial}{\partial \omega} (\omega \eta_\sigma)} \exp \left\{ - \frac{(\omega - s \Omega_e)^2}{2 \omega^2 \eta_\sigma^2 \beta_e^2 \cos^2 \theta} \right\}, \quad (3.36)$$

where

$$A(\omega, \theta, s) := \frac{(\mu_s^2 + s^2)}{\lambda_e} I_s(\lambda_e) + 2(\mu_s - \lambda_e) \{ I_s'(\lambda_e) - I_s(\lambda_e) \}, \quad (3.37)$$

with

$$\mu_s := \mu(\omega, \theta, s) := (\omega b_\sigma \sin \theta / \Omega_e + s T_\sigma) / \cos \theta,$$

$$\lambda_e = \eta_\sigma^2 \omega^2 \beta_e^2 \sin^2 \theta / \Omega_e, \quad \text{and} \quad \beta_e := v_e / c.$$

All other symbols have their usual meanings. Collisionless damping has been treated by many authors, see for example Landau (1946), Sitenko and Stepanov (1957), Gershman (1960), Stepanov (1960) and Zheleznyakov (1970). However, the collisionless damping decrement (3.36) is more general than previous analyses, since fewer approximations have been made here. In the various limits, previous results can be recovered. For example, when $\omega \approx n \Omega_e$ and $\lambda_e \approx n^2 \eta_\sigma^2 \beta_e^2 \sin^2 \theta \ll 1$, $A(\omega, \theta) \approx (a_\sigma + 1)^2 n^2 \lambda_e^{n-1} / 2^n n!$ and

$$\gamma^\sigma(\omega, \theta) \approx \left(\frac{\pi}{2}\right)^{\frac{1}{2}} \frac{\omega_p^2}{\omega} \frac{n^{2n}}{2^n n!} \frac{(a_\sigma + 1)^2}{(1 + T_\sigma^2)} \frac{\eta_\sigma^{2n-4} \beta_e^{2n-3}}{\partial(\omega \eta_\sigma) / \partial \omega} \frac{\sin^{2n-3} \theta}{|\cos \theta|} \exp \left\{ - \frac{(\omega - n\Omega_e)^2}{2\omega^2 \eta_\sigma^2 \beta_e^2 \cos^2 \theta} \right\}. \quad (3.38)$$

In the limit $\omega \gg \omega_p$, (3.38) becomes particularly simple, since $\eta_\sigma \approx 1$ and therefore, close to the vacuum limit,

$$\gamma^\sigma(\omega, \theta) \approx \left(\frac{\pi}{8}\right)^{\frac{1}{2}} \frac{\omega_p^2}{\omega} \frac{n^{2n}}{2^n n!} \frac{(1 \mp \cos \theta)^2}{|\cos \theta|} \frac{\sin^{2n-3} \theta}{|\cos \theta|} \beta_e^{2n-3} \exp \left\{ - \frac{(\omega - n\Omega_e)^2}{2\omega^2 \beta_e^2 \cos^2 \theta} \right\}, \quad (3.39)$$

where the upper (lower) sign refers to ordinary (extraordinary) mode waves. Similar application of equation (3.35) gives the collisionless damping decrements for longitudinal waves by Maxwellian plasma electrons,

$$\gamma^\sigma(\underline{k}) = \sum_s \left(\frac{\pi}{2}\right)^{\frac{1}{2}} \frac{\omega_p^2 \omega^2}{k^3 v_e^3 |\cos \theta|} \left(\frac{W_E}{W_T}\right)^l \zeta_s(\lambda_e) \exp \left\{ - \frac{(\omega - s\Omega_e)^2}{2k^2 v_e^2 \cos^2 \theta} \right\}, \quad (3.40)$$

where 'l' denotes any longitudinal wave mode and $\zeta_s(\lambda_e)$ has been defined by equation (2.86). The well-known decrements for electron plasma waves and ion acoustic waves (Sitenko, 1967) can easily be recovered from (3.40):

$$\gamma^{el}(\underline{k}) = \left(\frac{\pi}{8}\right)^{\frac{1}{2}} \frac{\omega^2}{k^3 \lambda_{De}^3 \omega_p |\cos \theta|} \left\{ 1 + \frac{\omega_p^2 \Omega_e^2 \sin^2 \theta}{(\omega^2 - \Omega_e^2)^2} \right\}^{-1} \left\{ \exp \left(- \frac{\omega^2}{2k^2 v_e^2 \cos^2 \theta} \right) + \frac{k^2 v_e^2 \sin^2 \theta}{2\Omega_e^2} \left[\exp \left(- \frac{(\omega - \Omega_e)^2}{2k^2 v_e^2 \cos^2 \theta} \right) + \exp \left(- \frac{(\omega + \Omega_e)^2}{2k^2 v_e^2 \cos^2 \theta} \right) \right] + \dots \right\},$$

($\lambda_e \ll 1$), (3.41)

and

$$\gamma^s(\underline{k}) = \left(\frac{\pi}{8}\right)^{\frac{1}{2}} \left(\frac{m_e}{m_i}\right)^{\frac{1}{2}} \omega. \quad (3.42)$$

In addition, the collisionless, cyclotron damping decrements can easily be written down for electron-cyclotron waves. For example, equations (2.98) - (2.102) give, for the case $\omega \approx n\Omega_e \gg \omega_p^2 + \Omega_e^2$,

$$\gamma^c(\underline{k}) = \left(\frac{\pi}{2}\right)^{\frac{1}{2}} \frac{\omega^2}{\omega_p} \frac{\zeta_n^2(\lambda_e)}{k^5 \lambda_{De}^5 |\cos\theta|} \exp \left\{ -\frac{(\omega - n\Omega_e)^2}{2k^2 V_e^2 \cos^2\theta} \right\}. \quad (3.43)$$

Classically, collisional damping lies outside the scope of the Boltzmann-Vlasov equation and it can only be properly treated, in general, by including a Fokker-Planck term on the right hand side of (2.1), see for example Chandrasekhar (1943), Rosenbluth *et al.* (1957), McBride (1969a,b) and references contained in Krishan and Selim (1971). However, for an equilibrium plasma, it is possible to use Kirchhoff's law together with an approximate bremsstrahlung emissivity to estimate the collisional damping decrements (Bekefi, 1966; Zheleznyakov, 1970).

For high frequency waves ($\omega \geq \omega_p$), collisional damping is determined essentially by electron-ion interactions whose damping decrements are generally $O(\omega^2/k^2 V_e^2)$ greater than those arising from electron-electron interactions (Buti and Jain, 1965; Buti, 1967; McBride, 1969a,b). When $\omega_p \gg \Omega_e$, the magnetic field B_0 does not affect the nature of the collisions, and to the lowest order, the damping decrement is independent of the effects of B_0 (*loc. cit.*),

$$\gamma_c \approx \frac{1}{3} \left(\frac{2}{\pi}\right)^{\frac{1}{2}} \frac{\omega_p^2}{\omega^2} \nu_c, \quad (3.44)$$

where the effective collision frequency is given by

$$\nu_c \equiv 3\omega_p \frac{\ln \Lambda_e}{\Lambda_e}, \quad \Lambda_e \equiv 12\pi n_e \lambda_{De}^3. \quad (3.45)$$

5. TWO-STREAM INSTABILITIES

Depending on the relative amount of bunching in physical and velocity space, it is possible theoretically to discern three distinct types of coherent mechanisms whereby streams of charges can generate plasma turbulence and give rise to non-thermal radiation.

The first mechanism operates when a group of, say, N monoenergetic particles of charge q , is spatially bunched to such a degree that

$$kd \ll 1, \quad (3.46)$$

where d is the dimension of bunch measured in the direction of emitted waves of wavenumber k . The bunch then radiates like a single particle of charge Nq . The total power radiated is N^2 times the power radiated by a single charge q or N times the total power radiated incoherently by all the charges. This mechanism, which requires bunches of charges to be postulated, has been invoked in theories of pulsar radiation by Caroff and Scargle (1970) and Komesaroff (1970) (among others); it has been criticized by Ginzburg and Zheleznyakov (1970) and defended by Goldreich *et al.* (1972). However, it is unlikely to be applicable to non-thermal radiation by non-relativistic or mildly relativistic particles under cosmic conditions.

The other two mechanisms are the well known two stream instabilities in plasma physics. They operate when there is no appreciable spatial bunching and in fact when the opposite limit of (3.46) is satisfied. Two classes of two stream instabilities are distinguished here: one is hydrodynamic or reactive and the other is quasilinear, resistive or maser-like. The hydrodynamic instability can occur only when the velocity spread of the stream is sufficiently small, so that the hydrodynamic approximation is applicable, i.e. when

$$\left| \frac{\omega - s\Omega - k_{\parallel} u_s}{\sqrt{2} k_{\parallel} V_{s\parallel}} \right| \sim \frac{\gamma_h^{\sigma}(k, u_s, s)}{\sqrt{2} k_{\parallel} V_{s\parallel}} \gg 1, \quad (s = 0, 1, \dots), \quad (3.47)$$

where $V_{s\parallel}$ is the parallel thermal spread of the stream and $\gamma_h^{\sigma}(k, u_s, s)$ is the hydrodynamic growth rate when the beam is in resonance, at the s -th harmonic, with the unstable waves in mode σ . A quantum mechanical or semiclassical approach based on Einstein's coefficients is not valid in this case, because a random phase description of the waves is not admissible, i.e., beam particles are not in resonance with several waves (Twiss, 1963; Ginzburg *et al.*, 1962; Ginzburg and Zheleznyakov, 1965).

By arguments exactly analogous to those given by Shapiro (1963) and Tsytovich (1970, p.183), it can be shown that the hydrodynamic instability is accompanied by a significant increase in the thermal spread of the beam. If nonlinear interaction is unimportant, it will lead eventually to the situation where

$$\left| \frac{\omega - s\Omega - k_{\parallel} u_s}{\sqrt{2} k_{\parallel} V_{s\parallel}} \right| \lesssim 1, \quad (s = 0, 1, \dots). \quad (3.48)$$

At this stage, the hydrodynamic instability passes over to the quasilinear instability. The energy density of the unstable waves resulted from a hydrodynamic instability is less than $(n_s/n_e)^{1/3}$ times the energy density of the initial stream. On account of the applicability of the random phase approximation to the quasilinear instability, classical considerations based on the kinetic equation coincide with quantum mechanical or semiclassical considerations based on Einstein's coefficients. In the absence of significant nonlinear interactions, the quasilinear equations (3.33) - (3.35) describe the relaxation or flattening of the beam particle distribution function and the decrease in quasilinear growth rates of unstable waves (Harris, 1967; Register

and Oberman, 1968).

Analogous to the derivation of (2.53), the wave equation, including effects of a stream but neglecting damping by the background plasma and nonlinear effects, reads

$$\{\Lambda_{ij}(\underline{k}, \omega) + \Delta\epsilon_{ij}(\underline{k}, \omega)\} E_j(\underline{k}, \omega) = 0, \quad (3.49)$$

where $\Lambda_{ij}(\underline{k}, \omega)$ is defined by (2.54),

$$\Delta\epsilon_{ij}(\underline{k}, \omega) \equiv \epsilon_{ij}'(\underline{k}, \omega) - \delta_{ij}, \quad (3.50)$$

and $\epsilon_{ij}'(\underline{k}, \omega)$ is given by (2.42), with primed plasma parameters referring to the stream. In the hydrodynamic limit,

$$\Delta\epsilon_{ij}(\underline{k}, \omega) = \epsilon_{ij}^{h'}(\underline{k}, \omega) - \delta_{ij}, \quad (3.51)$$

and the instabilities for this case have been extensively studied, see e.g. Fainberg (1962), Neufeld and Wright (1963) and Briggs (1964).

The hydrodynamic instability has been divided into two types: convective and absolute instabilities. In the convective instabilities, plasma waves are amplified as they move away from the origin of the instability. In the absolute instabilities, plasma waves grow at the origin of the instability. Sturrock (1958) showed that the dispersion relations are sufficient to determine whether an instability is convective or absolute. In general, a necessary and sufficient condition for hydrodynamic instability is that real $\underline{k}$ should give rise to complex ω solutions of the dispersion equation. Furthermore, the instability is convective (absolute) if real ω give rise to complex (real) $\underline{k}$ solutions of the dispersion equation (Polovin, 1962; Akhiezer *et al.*, 1967; Akhiezer and Polovin, 1971). In the quasilinear limit,

$$\Delta\epsilon_{ij}(\underline{k}, \omega) = \epsilon_{ij}^{a'}(\underline{k}, \omega), \quad (3.52)$$

and the instabilities for this case can be described by the quasilinear equations (3.33) - (3.35), provided the density of the stream is sufficiently low so that wave properties are only slightly perturbed by the presence of the stream.

For such low density streams, the dispersion equation, viz., the determinantal equation of (3.49), can be solved generally to find the growth rates of various plasma wave modes. A perturbation expansion of

$$\det\{\Lambda_{ij}(\underline{k}, \omega) + \Delta\epsilon_{ij}(\underline{k}, \omega)\} = 0, \quad (3.53)$$

with respect to ω for a given real $\underline{k}$ gives to first order,

$$\lambda_{ij}(\underline{k}, \omega^\sigma(\underline{k})) \Delta\epsilon_{ji}(\underline{k}, \omega) + \Delta\omega \left. \frac{\partial \Lambda}{\partial \omega} \right|_{\omega=\omega^\sigma} = 0, \quad (3.54)$$

where $\omega = \omega^\sigma(\underline{k}) = -\omega^\sigma(-\underline{k})$ is the dispersion relation for waves in mode σ , obtained from the zeroth order equation $\Lambda(\underline{k}, \omega) = 0$, and $\Delta\omega \equiv \omega - \omega^\delta(\underline{k}) = \delta \exp(i\chi)$, say, is the frequency perturbation caused by the presence of the stream. Application of (3.9) yields

$$\frac{\Delta\omega}{\omega^\sigma} = - \left(\frac{W_E}{W_T} \right)^\sigma e_i^{\sigma*}(\underline{k}) e_j^\sigma(\underline{k}) \Delta\epsilon_{ij}(\underline{k}, \omega^\sigma + \Delta\omega), \quad (3.55)$$

where $\Delta\epsilon_{ij}(\underline{k}, \omega^\sigma + \Delta\omega) \approx \epsilon_{ij}'(\underline{k}, \omega^\sigma)$ gives essentially the same growth rates as (3.35), in the quasilinear limit. The growth rates from a hydrodynamic instability can only be obtained in general by solving equation (3.55) for $\Delta\omega$. Since wave amplitudes change in time according to $\exp(-i\omega t)$, the temporal growth rate is given by $\text{Im}(\Delta\omega) = \delta \sin\chi$.

Treatments of the two-stream instabilities, involving a relativistic stream have been given by Blanken *et al.*, (1969) and Fainberg *et al.*, (1970).

(a) Hydrodynamic Instabilities

To determine the growth rates of hydrodynamic instabilities, equation (2.42) can be used to write

$$\Delta \epsilon_{ij}(\underline{k}, \omega) = \epsilon_{ij}^{h'}(\underline{k}, \omega) - \delta_{ij} = -\frac{\omega_p'^2}{\omega^2} \left\{ \delta_{i3} \delta_{j3} + \sum_{n=-\infty}^{\infty} \int d^3 p f_s(\underline{p}) \times \left[\frac{\omega - k_{\parallel} v_{\parallel}}{(\omega - n\Omega' - k_{\parallel} v_{\parallel})} \frac{1}{v_{\perp}} \frac{\partial}{\partial v_{\perp}} (v_{\perp} v_j^*) + \frac{k_{\parallel}^2 v_{\perp} v_j^*}{(\omega - n\Omega' - k_{\parallel} v_{\parallel})^2} \right] \right\}, \quad (3.56)$$

where $V_i = V_i(n, \underline{k}, \underline{v})$, $f_s(\underline{p})$ is the distribution function of the stream particles and dashed quantities refer to the stream. For a stream with

$$f_s(\underline{p}) = C f_{s\perp}(p_{\perp}) \delta(v_{\parallel} - u_s), \quad (3.57)$$

where C is a normalization constant, it is possible to choose values of n and $k_{\parallel}$ such that $\omega^{\sigma}(\underline{k}) - n\Omega' - k_{\parallel} u_s \ll \Delta\omega$. Under these conditions, a comparison of imaginary parts in (3.55), using equations (3.56) and (3.57) gives a cubic equation for δ , for a specific value of n ,

$$\delta_n^3 + a_n \delta_n + b_n = 0, \quad (3.58)$$

where

$$a_n \equiv \frac{2\pi m' C \omega_p'^2}{\omega^{\sigma}} \left(\frac{W_E}{W_T} \right)^{\sigma} n\Omega' \int p_{\perp} dp_{\perp} f_{s\perp}(p_{\perp}) \frac{1}{v_{\perp}} \frac{\partial}{\partial v_{\perp}} |\underline{e}^{\sigma} \cdot \underline{v}|^2, \quad (3.59)$$

and

$$b_n \equiv \frac{2\pi m' C \omega_p'^2}{\omega^{\sigma}} \left(\frac{W_E}{W_T} \right)^{\sigma} 2k_{\parallel}^2 \cos \chi \int p_{\perp} dp_{\perp} f_{s\perp}(p_{\perp}) |\underline{e}^{\sigma} \cdot \underline{v}|^2, \quad (3.60)$$

with $\underline{v} = \underline{v}(n, \underline{k}, v_{\perp}, u_s)$. Substituting the appropriate wave properties, equations (3.58) - (3.60) recover the growth rates for hydrodynamic instabilities of whistler mode waves (Bell and Buneman, 1964) and of electron plasma waves (Yip, 1970, $\beta_T' = 0$).

Consider a stream with a bi-Maxwellian distribution function of a form given by equation (2.68). The above procedure together with

equations (2.72) - (2.75) and condition (3.47) gives again an equation of the form (3.58) with

$$a_n := \frac{\omega_p'^2}{\omega^\sigma} \left(\frac{W_E}{W_T} \right)^\sigma e_i^{\sigma*} e_j^\sigma \pi_{ij}'(\underline{k}, \omega^\sigma, n) e^{-\lambda'} n \Omega', \quad (3.61)$$

and

$$b_n := \frac{\omega_p'^2}{\omega^\sigma} \left(\frac{W_E}{W_T} \right)^\sigma e_i^{\sigma*} e_j^\sigma \pi_{ij}'(\underline{k}, \omega^\sigma, n) e^{-\lambda'} 2(1-\tau') k_{||}^2 v_{s||}^2 \cos \chi, \quad (3.62)$$

where $\pi_{ij}'(\underline{k}, \omega^\sigma, n)$, with stream parameters, is given by (2.74) and the temperature anisotropy parameter is $\tau' := 1 - v_{s||}^2 / v_{s\perp}^2$. Coefficients (3.61) and (3.62) can also be obtained from (3.59) and (3.61) by taking $f_{s\perp}(p_\perp)$ to be Maxwellian. For small gyroradii, i.e., $\lambda' \ll 1$, the coefficients (3.61) and (3.62) are very small except for small harmonic numbers n .

As an example, consider longitudinal waves, then the coefficients simplify considerably to

$$a_n = \omega_p'^2 \omega^\ell \left(\frac{W_E}{W_T} \right)^\ell \zeta_n(\lambda') \frac{n \Omega'}{k^2 v_{s\perp}^2}, \quad (3.63)$$

and

$$b_n = \omega_p'^2 \omega^\ell \left(\frac{W_E}{W_T} \right)^\ell \zeta_n(\lambda') 2 \cos^2 \theta \cos \chi, \quad (3.64)$$

where $\zeta_n(\lambda')$, with $\lambda' = k_\perp^2 v_{s\perp}^2 / \Omega'^2$, is given by (2.86). As might be obvious from a physical argument, equations (3.63) and (3.64) show clearly that temperature anisotropies are not important in hydrodynamic instabilities. For the case $n=0$, equation (3.58) gives the hydrodynamic growth rate,

$$\gamma_h^\sigma(\underline{k}, n=0) = (\delta_0 \sin \chi)_{\max} = \frac{\sqrt{3}}{2} \left\{ \frac{\omega_p'^2}{\omega^\sigma} \left(\frac{W_E}{W_T} \right)^\sigma e_i^{\sigma*} e_j^\sigma \pi_{ij}'(\underline{k}, \omega^\sigma, n) \times e^{-\lambda'} (1-\tau') k_{||}^2 v_{s||}^2 \right\}^{1/3}, \quad (3.65)$$

and in particular,

$$\gamma_h^{\ell}(\underline{k}, n=0) = \frac{\sqrt{3}}{2} \left\{ \omega_p'^2 \omega^{\ell} \left(\frac{W_E}{W_T} \right)^{\ell} \zeta_0(\lambda') \cos^2 \theta \right\}^{1/3}. \quad (3.66)$$

For the case $n \neq 0$, in general, the cubic equation (3.58) has to be solved. However, when $a_n^3 \gg b_n^2$,

$$\gamma_h^{\sigma}(\underline{k}, n) = (\delta_n \sin \chi)_{\max} = a_n^{1/2}. \quad (3.67)$$

For example, the case $n = -1$ for longitudinal waves in the limit $\lambda' \ll 1$ could satisfy the inequality, $a_{-1}^3 \gg b_{-1}^2$ or

$$\frac{\omega_p'^2 \omega^{\ell}}{\Omega'^3} \left(\frac{W_E}{W_T} \right)^{\ell} \frac{\sin^6 \theta}{2\lambda'^2 \cos^4 \theta} \gg 1, \quad (3.68)$$

provided θ is not too small. Thus,

$$\gamma_h^{\ell}(\underline{k}, n=-1) = \omega_p' \sin \theta \left\{ \frac{\omega^{\ell}}{\Omega'} \left(\frac{W_E}{W_T} \right)^{\ell} \right\}^{1/2}, \quad (3.69)$$

which gives a growth rate for electron plasma waves, which coincides with the one found by Stepanov and Kitsenko (1961). Note that when $a_n^3 \gg b_n^2$, hydrodynamic instability can occur only when $n < 0$.

(b) Quasilinear Instabilities

The quasilinear growth rates can be found either by the application of equation (3.52) and (3.55) or by the direct use of (3.35). For a bi-Maxwellian streaming distribution function of the form (2.68), the former procedure is simpler, as the integrals have already been performed in (2.72) and (2.73). It gives

$$\begin{aligned} \gamma^{\sigma}(\underline{k}) = & - \sum_{n=-\infty}^{\infty} \frac{\omega_p'^2}{\omega^{\sigma}} \left(\frac{W_E}{W_T} \right)^{\sigma} \sqrt{\pi} \exp(-y_n'^2) e_i^{\sigma*} e_j^{\sigma} \pi_{ij}'(\underline{k}, \omega^{\sigma}; n) \\ & \times e^{-\lambda'} \frac{k_{\parallel}}{|k_{\parallel}|} \{y_0' - \tau' y_n'\}, \end{aligned} \quad (3.70)$$

where the symbols have their usual meanings and $y_n' := (\omega^\sigma - n\Omega' - k_{||} u_s) / \sqrt{2} k_{||} V_{s||}$, with streaming velocity u_s and velocity spread parallel to the background magnetic field $V_{s||}$. The condition for growth of unstable waves is

$$n\Omega' + \frac{V_{s||}^2}{V_{s||}^2} (\omega^\sigma - n\Omega' - k_{||} u_s) < 0, \quad (3.71)$$

which reduces to $\omega^\sigma - k_{||} u_s < 0$, in the absence of temperature anisotropies. Equation (3.70) can be rewritten in the form

$$\gamma^\sigma(k) = - \sum_{n=-\infty}^{\infty} \sqrt{\pi} \frac{\omega_p'^2}{\omega^\sigma} \left(\frac{W_E}{W_T} \right)^\sigma e_i^{\sigma*} e_j^{\sigma} \pi_{ij}'(k, \omega^\sigma; n) e^{-\lambda'} f(X_n), \quad (3.72)$$

where

$$f(X_n) := (C_n + dX_n) e^{-X_n^2}, \quad (3.73)$$

$$C_n := \frac{n\Omega'}{\sqrt{2} |k_{||}| V_{s||}}, \quad d := 1 - \tau' = \frac{V_{s||}^2}{V_{s||}^2}, \quad X_n := \frac{\omega^\sigma - n\Omega' - k_{||} u_s}{\sqrt{2} |k_{||}| V_{s||}}.$$

For each n , the growth rate (3.72) can be maximized by finding the largest negative value of $f(X_n)$. Simple calculus gives

$$(X_n)_{\max} = -\frac{c_n}{2d} - \frac{1}{2} \left\{ \left(\frac{c_n}{d} \right)^2 + 2 \right\}^{\frac{1}{2}}; \quad (3.74)$$

at this value $f(X_n)$ is not exponentially small only if $(c_n/2d)^2 \leq 1$ or

$$\frac{|n\Omega'|}{\sqrt{2} |k_{||}| V_{s||}} \leq \frac{2V_{s||}^2}{V_{s||}^2}. \quad (3.75)$$

Note that this necessary condition derived from (3.74) is independent of the streaming velocity u_s and it is trivially satisfied for $n=0$ as may be seen from (3.74).

For magnetoionic waves, it is convenient to regard the wavenumber k as dependent on ω and θ . It is easily shown that for $\underline{e}^\sigma = (a_\sigma, i, b_\sigma)$,

$$e_i^{\sigma*} e_j^{\sigma} \pi_{ij}'(\omega, \theta; n) = \frac{A(\omega, \theta, n)}{1 + K_{\sigma}^2 + T_{\sigma}^2}, \quad (3.76)$$

where $A(\omega, \theta, n)$ has the form of (3.37), except the plasma parameters now refer to the stream. Substitution of (3.76) and wave property (2.83) into equation (3.72) gives

$$\gamma^{\sigma}(\omega, \theta) = - \sum_{n=-\infty}^{\infty} \frac{\sqrt{\pi}}{2} \frac{\omega_p'^2}{\omega} \frac{e^{-\lambda'}}{(1 + T_{\sigma}^2) \eta_{\sigma} \frac{\partial}{\partial \omega} (\omega \eta_{\sigma})} f(X_n) . \quad (3.77)$$

For longitudinal waves, equation (3.72) yields

$$\gamma^{\ell}(\underline{k}) = - \sum_{n=-\infty}^{\infty} \sqrt{\pi} \frac{\omega_p'^2 \omega^{\ell}}{k^2 V_{s||}^2} \left(\frac{W_E}{W_T} \right)^{\ell} \zeta_n(\lambda') f(X_n) , \quad (3.78)$$

which, for the special case of $V_{s\perp} = V_{s||} = V_s$, $\lambda' \ll 1$ and for electron plasma waves, recovers the result found first by Stepanov and Kitsenko (1961),

$$\frac{\gamma^{e\ell}(\underline{k})}{\omega} = - \sum_{n=-\infty}^{\infty} \frac{\sqrt{\pi} \lambda' |n|}{2 |n| + 1 |n|!} \frac{\omega_p'^2}{k^2 V_s^2} X_0 \exp(-X_n^2) \left\{ 1 + \frac{\omega_p'^2 \Omega_e^2 \sin^2 \theta}{(\omega^2 - \Omega_e^2)^2} \right\}^{-1} . \quad (3.79)$$

For the novel case of electron-cyclotron waves generated by electron streams at a frequency $\omega \approx r \Omega_e$ ($r > 0$), the condition that growth is not exponentially small, together with the weak damping condition (2.87) requires, for $u_s \gg V_{s||}$,

$$\frac{u_s^2}{V_e^2} \left(\frac{\omega - r \Omega_e}{\omega - n \Omega_e} \right)^2 \gg 1 , \quad (3.80)$$

which can be satisfied in general only for $n = r$ and $u_s \gg V_e$. Under these circumstances,

$$\frac{\gamma^c(\underline{k})}{\omega} = - \frac{\sqrt{\pi}}{k^2 V_{s||}^2} \frac{\omega_p'^2}{k^2 \lambda_{De}^2} \frac{\zeta_r(\lambda_e) \zeta_r(\lambda')}{\{\epsilon_0^{\ell}(\underline{k}, r \Omega_e)\}^2} f(X_r) , \quad (3.81)$$

where $\varepsilon_0^{\ell}(\underline{k}, r\Omega_e)$ is defined by (2.100). For optically thin laboratory plasmas, condition (3.80) is not necessary and Cerenkov excitation of the electron-cyclotron waves is possible. Mizuno *et al.* (1971) have observed the Cerenkov and cyclotron excitation of these waves by electron beams in a laboratory plasma.

6. VELOCITY-ANISOTROPY INSTABILITIES

Harris (1959, 1961) was the first to show that a plasma with anisotropic velocity distributions could give rise to micro-instabilities. For this reason, such instabilities are often known as Harris instabilities. Sometimes, they are also known as instabilities due to temperature or pressure anisotropies. Velocity-anisotropy instabilities can be studied by substituting various anisotropic velocity distribution functions into equation (2.42) and examining dispersion equation (2.56) for complex frequency solutions. Such investigations have been carried out quite extensively, particularly in connection with cyclotron harmonic wave phenomena, see for example Crawford (1968), Tataronis and Crawford (1970a,b) and Brambilla (1970) and the references therein. Although the instabilities themselves have been well investigated, the properties of the waves which exist in such plasmas with anisotropic velocity distributions have not been studied to sufficient analytic detail. However, from the point of view of astrophysical applications and some laboratory experiments (Perkins and Barr, 1966) it seems that a relevant type of velocity-anisotropy instability is the double-distribution instability, where a group of particles with anisotropic velocity distribution exists in an equilibrium plasma (with isotropic velocity distributions). Provided the density of the "extraneous" particles is sufficiently low, such instabilities can be treated by equation (3.55) and the wave properties

can be taken to be those of the background equilibrium plasma.

Consider an anisotropic velocity distribution of particles of the form,

$$f_s(\underline{p}) = \frac{1}{(2\pi)^3 m'^3 v_{s\perp}^2} \exp \left[-\frac{v_{\perp}^2}{2v_{s\perp}^2} \right] \delta(v_{\parallel}) , \quad (3.82)$$

where dashed quantities refer to the "extraneous" particles. Again, provided values n exist such that $\omega^\sigma(\underline{k}) - n\Omega' \ll \Delta\omega$, equations (3.55) and (3.82) gives a cubic equation (3.58), with coefficients for magneto-ionic waves:

$$a_n = \frac{\omega_p'^2}{2\omega} \frac{n\Omega' e^{-\lambda'}}{(1+T_\sigma^2) \eta_\sigma \frac{\partial}{\partial\omega} (\omega\eta_\sigma)} \left\{ \frac{(a_\sigma^2+1) n^2 I_n}{\lambda'} + 2(na_\sigma - \lambda') (I_n' - I_n) \right\} , \quad (3.83)$$

$$b_n = \frac{\omega_p'^2 \omega v_{s\perp}^2}{c^2} \frac{\eta_\sigma \cos^2\theta \cos\chi}{(1+T_\sigma^2) \frac{\partial}{\partial\omega} (\omega\eta_\sigma)} \left\{ \frac{(a_\sigma^2+1) n^2 I_n}{\lambda'} + 2(na_\sigma - \lambda') (I_n' - I_n) \right\} ,$$

with $\lambda' = \eta_\sigma^2 \omega^2 v_{s\perp}^2 \sin^2\theta / c^2 \Omega'^2$ and for longitudinal waves:

$$a_n = \frac{\omega_p'^2}{\omega} \left(\frac{W_E}{W_T} \right)^\lambda \frac{n^3 \Omega' \Omega_e^2}{k^2 v_{s\perp}^2} \zeta_n(\lambda') , \quad (3.84)$$

$$b_n = \frac{\omega_p'^2}{\omega} \left(\frac{W_E}{W_T} \right)^\lambda 2n^2 \Omega_e^2 \cos^2\theta \cos\chi \zeta_n(\lambda') ,$$

with $\lambda' = k_\perp^2 v_{s\perp}^2 / \Omega'^2$. Hence groups of non-streaming charges with anisotropic velocity distributions could give rise to hydrodynamic instabilities of magnetoionic and cyclotron waves at frequencies $\omega \approx n\Omega'$, in an equilibrium plasma.

Similarly, a non-streaming bi-Maxwellian distribution of "extraneous" charges gives rise to a quasilinear instability, with growth rates given by (3.72), (3.77), and (3.78) with $X_n = (\omega^\sigma - n\Omega')/\sqrt{2} |k_\parallel| v_{s\parallel}$. Instabilities in a bi-Maxwellian plasma (not

double-distribution instabilities) have received considerable attention in the literature, see for example Hall and Heckrotte (1964), Shima and Hall (1965) and references in Crawford (1968). Zayed and Kitsenko (1968) pointed out, as shown here, velocity-anisotropy instabilities are also possible for magnetoionic waves, which can be radiated from the plasma directly.

Instabilities of magnetoionic waves in plasmas with anisotropic temperatures have been studied by Landau and Cuperman, 1970; Gaffey *et al.*, 1972; and Davidson *et al.*, 1972). Bursts of radiation have been observed from such plasmas adiabatically compressed by a pulsed magnetic field (Schwartz and Lichtenberg, 1972). The quasilinear relaxation of a weak temperature anisotropy in a rarefied and hot plasma without a background magnetic field has been theoretically studied by Montes *et al.* (1970) and Montes and Peyraud (1972).

7. DRIFT INSTABILITIES

Microinstabilities can arise in a current-carrying plasma. Such currents, which are just relative motions of electrons and ions, may be driven by $\mathbf{E} \times \mathbf{B}$ drifts or by drifts due to gradients in density, temperature or magnetic field and they are present in certain collisionless magnetosonic shocks (see e.g., Tidman and Krall, 1971) and in some theta-pinch experiments (Davis *et al.*, 1972). The microturbulence generated by the drift instabilities, which is usually ion-acoustic turbulence, can heat the plasma particles (see e.g. Keilhacker and Steuer, 1971; Paul, 1971) and it can also produce radiation at electron cyclotron harmonic frequencies (Livshitz and Tsytovich, 1968).

To study instabilities from drifts parallel to the background magnetic field, the dielectric tensor (2.42) can still be used, since the

cylindrical symmetry assumed in its derivation is preserved. However, for drifts perpendicular or at an oblique angle to the background magnetic field, cylindrical symmetry is broken and the dielectric tensor (2.42), as it stands, is no longer directly applicable. Under these circumstances, there are two equivalent ways of proceeding: the first is to derive the dispersion equations *ab initio*, not assuming cylindrical symmetry (Krall and Rosenbluth, 1963) and the second is to transform dielectric tensor (2.42) in the rest frame of drifting electrons back to the laboratory frame by a Galilean transformation. Galilean transformations of wave properties and the dielectric tensor have been treated by Melrose (1972c), from which it is easy to show

$$\begin{aligned} \omega^2 \{ \epsilon_{ij}(\underline{k}, \omega) - \delta_{ij} \} &= (\omega - \underline{k} \cdot \underline{v})^2 \{ \epsilon'_{ij}(\underline{k}, \omega - \underline{k} \cdot \underline{v}) - \delta_{ij} \} + (\omega - \underline{k} \cdot \underline{v}) \\ &\times k_s \{ v_i \{ \epsilon'_{sj}(\underline{k}, \omega - \underline{k} \cdot \underline{v}) - \delta_{sj} \} + v_j \{ \epsilon'_{is}(\underline{k}, \omega - \underline{k} \cdot \underline{v}) - \delta_{is} \} \} \\ &+ v_i v_j k_r k_s \{ \epsilon'_{rs}(\underline{k}, \omega - \underline{k} \cdot \underline{v}) - \delta_{rs} \} , \end{aligned} \quad (3.85)$$

where the prime refers to the rest frame of the electrons with drift $\underline{v}$. From (3.85), it is easily deduced that

$$\epsilon^{\ell}(\underline{k}, \omega) = \epsilon^{\ell'}(\underline{k}, \omega - \underline{k} \cdot \underline{v}) . \quad (3.86)$$

Consider the drift instability of ion-acoustic waves at (angular) frequency range $\Omega_i \ll \omega \leq \tau_i$, caused by electrons drifting relative to ions with velocity $(0, u_d, u_s)$. For such ion-acoustic waves, the wave properties are determined principally by unmagnetized ions and the instability may be regarded as resulting from the perturbing effects of the drifting electrons.

Straightforward application of (2.42) and (3.86) gives

$$\epsilon^{\ell(e)}(\underline{k}, \omega) = \frac{1}{k^2 \lambda_{De}^2} \left\{ 1 - \sum_{n=-\infty}^{\infty} \int d^3 \underline{v} f_e J_n^3 \left(\frac{k_{\perp} v_{\perp}}{\Omega_e} \right) \times \frac{\omega - k_y u_D - k_{\parallel} u_s}{\omega - k_y u_D - k_{\parallel} u_s - n \Omega_e - k_{\parallel} v_{\parallel}} \right\}, \quad (3.87)$$

where f_e is a Maxwellian electron distribution and u_D , the perpendicular component of drift, might be dependent on $v_{\perp}$, e.g., $u_D = \epsilon v_{\perp}^2 / 2 \Omega_e$, with $\epsilon = (dB/dx)/B$. On integration with respect to $v_{\parallel}$, (3.87) becomes

$$\epsilon^{\ell(e)}(\underline{k}, \omega) = \frac{1}{k^2 \lambda_{De}^2} \left\{ 1 + \sum_{n=-\infty}^{\infty} \int v_{\perp} dv_{\perp} \frac{i y_{en} F^{(0)}(y_{en})}{v_e^2} \times \exp \left(- \frac{v_{\perp}^2}{2 v_e^2} \right) J_n^2 \left(\frac{k_{\perp} v_{\perp}}{\Omega_e} \right) \right\}, \quad (3.88)$$

where $F^{(0)}(y)$ is given by (2.71) and

$$y_{en} \equiv \frac{\omega - k_y u_D - k_{\parallel} u_s - n \Omega_e}{\sqrt{2} k_{\parallel} v_e}. \quad (3.89)$$

When $|y_{en}| \gg 1$, which is the case, for example when $\theta \rightarrow \pi/2$, for all values of n

$$\epsilon^{\ell(e)}(\underline{k}, \omega) = - \frac{1}{k^2 \lambda_{De}^2} \sum_{n=-\infty}^{\infty} \int \frac{v_{\perp} dv_{\perp}}{v_e^2} \exp \left(- \frac{v_{\perp}^2}{2 v_e^2} \right) J_n^2 \left(\frac{k_{\perp} v_{\perp}}{\Omega_e} \right) \times \frac{n \Omega_e}{\omega - k_y u_D - k_{\parallel} u_s - n \Omega_e}. \quad (3.90)$$

For drifts which are independent of $v_{\perp}$, for example $u_D = c |\underline{E} \times \underline{B}| / B^2$, equation (3.90) can be integrated;

$$\epsilon^{\ell(e)}(\underline{k}, \omega) = \frac{1}{k^2 \lambda_{De}^2} \sum_{n=1}^{\infty} \frac{2 n^2 \Omega_e^2 \zeta_n(\lambda_e)}{n^2 \Omega_e^2 - (\omega - k_y u_D - k_{\parallel} u_s)^2},$$

which, by equation (2.95), can be summed to give

$$\epsilon^{\ell(e)}(\underline{k}, \omega) = \frac{1}{k^2 \lambda_{De}^2} \{ 1 - \pi q \cot \pi q \zeta_q(\lambda_e) \} \quad (3.91)$$

with $q = |\omega - k_y u_D - k_{\parallel} u_s| / \Omega_e$. Equation (3.91) is valid, in general, only for almost perpendicular propagation of the waves, so that

$q = |\omega - k_y u_D| / \Omega_e$, $\lambda_e = k^2 v_e^2 / \Omega_e^2$. Now, $k \sim \lambda_{De}^{-1}$ implies $\lambda_e \sim \omega_p^2 / \Omega_e^2$, which is much greater than unity when $\omega_p \gg \Omega_e$. Hence, $\zeta_q(\lambda_e) \approx (2\pi\lambda_e)^{-1/2}$.

Also, when $\omega - k_y u_D - n\Omega_e \approx \Delta\omega$, $q \approx n + \Delta\omega / \Omega_e$ and $\pi q \cot \pi q \approx n\Omega_e / \Delta\omega$, so that

$$\epsilon^{(e)}(k, \omega) \approx -\frac{1}{k^2 \lambda_{De}^2} \cdot \frac{n\Omega_e}{\Delta\omega} \cdot \frac{\Omega_e}{\sqrt{2\pi} k v_e}. \quad (3.92)$$

Substitution of (2.110) and (3.92) into equation (3.55) gives a growth rate of perpendicular propagating ion-acoustic waves, in direction of the drift u_D ,

$$\frac{\gamma_s}{\Omega_e} = \left\{ \frac{m_e}{2\pi m_i} \left(\frac{1}{1+k^2 \lambda_{De}^2} + \frac{3T_i}{T_e} \right) \right\}^{1/4} \left\{ \frac{\omega_p}{2\Omega_e} \left(\frac{u_D - \omega/k}{v_e} \right) \left(\frac{1}{1+k^2 \lambda_{De}^2} + \frac{6T_i}{T_e} \right) \right\}^{1/2}, \quad (3.93)$$

which reduces, under the condition $T_e \gg T_i$, to

$$\frac{\gamma_s}{\Omega_e} = \left\{ \frac{m_e}{8\pi m_i} \frac{\omega_p^2 / \Omega_e^2}{(1+k^2 \lambda_{De}^2)^3} \right\}^{1/4} \left(\frac{u_D - \omega/k}{v_e} \right)^{1/2}, \quad (3.94)$$

which agrees essentially with the results of Gary and Sanderson (1970), who used a different method of calculation. For ion-acoustic waves propagating parallel to the background magnetic field, there is no drift instability of the hydrodynamic type, as may be readily verified by the above equations. However, there is a quasilinear type drift instability and its growth rate can be found from equation (3.78).

Under the conditions $T_e \gg T_i$ and $(u_s - \omega/k)/v_e \ll 1$, it reads

$$\frac{\gamma_s}{\Omega_e} = \frac{1}{1+k^2 \lambda_{De}^2} \left(\frac{\pi}{8} \frac{m_e}{m_i} \right)^{1/2} \left(\frac{u_s - \omega/k}{v_e} \right). \quad (3.95)$$

It is clear from a comparison of equations (3.94) and (3.95) that when $\omega_p > (m_e/m_i)^{1/2} \Omega_e \approx 43 \Omega_e$, the growth of ion-acoustic turbulence is faster in the perpendicular direction than in the parallel direction, for the

same drift speed.

The importance of ion-acoustic waves in turbulent heating processes at fronts of perpendicularly propagating MHD shocks is well-known (Lampe *et al.*, 1972). Ion-acoustic turbulence may give rise to suprathermal particles which could then be exciting agencies of non-thermal radiation (Lacombe and Mangeney, 1969; Smith, 1971).

CHAPTER 4

NONLINEAR INTERACTIONS IN A MAGNETOACTIVE PLASMA

1. INTRODUCTION

Nonlinear effects manifest themselves frequently in plasma physics. For example, they are important in studies on development of micro-instabilities, determinations of quasistationary spectra of turbulence, anomalous diffusion in confined plasmas and turbulent heating (see references below). Of particular relevance to this thesis are scattering of magnetoionic waves and nonlinear conversion of plasma turbulence into magnetoionic waves which are easily radiated from the plasma.

Many cases of interest fall within the weak turbulence regime where

$$W_{th} \ll W \ll W_p. \quad (4.1)$$

That is, energy density in plasma turbulence W is much greater than that in thermal equilibrium fluctuation W_{th} , but much less than kinetic energy density of the plasma particles W_p . When the frequency of turbulent plasma waves ω is of the order of, or greater than the electron plasma frequency ω_p , condition (1.2) is usually satisfied and the above cases also fall within the weak nonlinearity regime, where perturbation techniques are available.

Especially in the weak nonlinear approximation, nonlinear interactions have been intensively studied both for fixed-phase (coherent) waves and for random-phase (turbulent) waves. See for example, Kadomtsev (1965), Vedenov (1968), Kalman and Feix (1969), Sagdeev and

Galeev (1969), Tsytovich (1970), Stenflo (1972), Davidson (1972) and references contained therein. However, all these studies concentrate principally on plasmas without a background magnetic field, in contrast to the present study which includes a background magnetic field.

In addition to the random-phase approximation, our study of nonlinear interactions in a turbulent magnetoactive plasma also makes the collisionless assumption, i.e.,

$$\tau_{\phi} \ll \tau \ll \tau_c, \quad (4.2)$$

where the characteristic time of nonlinear interaction τ is short compared to mean time between particle collisions τ_c , but long compared to characteristic correlation time τ_{ϕ} .

Studies of weak nonlinear interactions in a turbulent magnetoactive plasma, making similar assumptions, have been undertaken by Kadomtsev and Petviashvili (1963), Karpman (1963), Tintsadze (1965), Kropotkin and Pustovalov (1966, 1967), Tsytovich and Shvartsburg (1966, 1967) and Gratzl (1971). The semiclassical theory of Tsytovich and Shvartsburg (1966, 1967) has been generalized and extended by Melrose and Sy (1972a,b, hereafter referred to as I,II), who include finite gyroradii effects and allow particles to have arbitrary energies. In particular they discuss Thomson scattering of magnetoionic waves by non-relativistic electrons in detail (I) and they study polarized radiation from nonlinear conversion of turbulent electron plasma waves in a plasma where $\omega_p \gg \Omega_e$ (II).

In this chapter, some nonlinear interactions in a magnetoactive plasma are investigated following the formalism presented in I and II. Thomson scattering of magnetoionic waves by both relativistic and non-relativistic particles, nonlinear scattering, coalescence and decay

processes of plasma waves are discussed. In addition, nonlinear effects on the formation of turbulence spectra are qualitatively considered.

2. EVOLUTION OF DISTRIBUTION FUNCTIONS

In the semiclassical theory, scattering of plasma waves by particles, which is a nonlinear wave-particle process, is described by a spontaneous scattering probability $w_{\alpha}^{\sigma\sigma'}(s, \underline{p}, \underline{k}, \underline{k}')$, which is the probability per unit time that a particle of species α , with momentum $\underline{p}$, scatters a plasmon of mode σ' in the range $d^3\underline{k}'/(2\pi)^3$ at wavevector $\underline{k}'$ into a plasmon of mode σ in the range $d^3\underline{k}/(2\pi)^3$ at $\underline{k}$ and at the same time decreases its harmonic quantum number by s . Standard procedures based on the use of Einstein coefficients give equations describing the evolution of distribution functions $N^{\sigma}(\underline{k})$, $N^{\sigma'}(\underline{k}')$ of plasmons and $f_{\alpha}(\underline{p})$ of scattering particles, see I for example. The evolution is due to the combined effects of spontaneous scattering $\sigma' \rightarrow \sigma$, $\sigma \rightarrow \sigma'$ and induced scattering $\sigma' \leftrightarrow \sigma$:

$$\left(\frac{\partial}{\partial t} + \underline{v}_g^{\sigma} \cdot \frac{\partial}{\partial \underline{r}} \right) N^{\sigma}(\underline{k}) = \sum_{\alpha} \sum_{s=-\infty}^{\infty} n_{\alpha} \int d^3\underline{p} \int \frac{d^3\underline{k}'}{(2\pi)^3} w_{\alpha}^{\sigma\sigma'}(s, \underline{p}, \underline{k}, \underline{k}') \{N^{\sigma'}(\underline{k}') - N^{\sigma}(\underline{k}) + N^{\sigma'}(\underline{k}') N^{\sigma}(\underline{k}) \hbar \hat{D}\} f_{\alpha}(\underline{p}), \quad (4.3)$$

$$\left(\frac{\partial}{\partial t} + \underline{v}_g^{\sigma'} \cdot \frac{\partial}{\partial \underline{r}} \right) N^{\sigma'}(\underline{k}') = - \sum_{\alpha} \sum_{s=-\infty}^{\infty} n_{\alpha} \int d^3\underline{p} \int \frac{d^3\underline{k}}{(2\pi)^3} w_{\alpha}^{\sigma\sigma'}(s, \underline{p}, \underline{k}, \underline{k}') \{N^{\sigma'}(\underline{k}') - N^{\sigma}(\underline{k}) + N^{\sigma'}(\underline{k}') N^{\sigma}(\underline{k}) \hbar \hat{D}\} f_{\alpha}(\underline{p}), \quad (4.4)$$

and

$$\frac{\partial f_{\alpha}(\underline{p})}{\partial t} = \sum_{s=-\infty}^{\infty} \int \frac{d^3\underline{k}}{(2\pi)^3} \int \frac{d^3\underline{k}'}{(2\pi)^3} \hbar \hat{D} \{w_{\alpha}^{\sigma\sigma'}(s, \underline{p}, \underline{k}, \underline{k}') \{N^{\sigma'}(\underline{k}') - N^{\sigma}(\underline{k}) + N^{\sigma'}(\underline{k}') N^{\sigma}(\underline{k}) \hbar \hat{D}\} f_{\alpha}(\underline{p})\}, \quad (4.5)$$

where n_α is the number density of scattering particles of species α , $f_\alpha(\underline{p})$ is normalized by (2.10), $\underline{v}_g^\sigma := \partial\omega^\sigma(\underline{k})/\partial\underline{k}$, $\underline{v}_g^{\sigma'} := \partial\omega^{\sigma'}(\underline{k}')/\partial\underline{k}'$ are group velocities and $\hat{D}$ is the differential operator,

$$\hat{D} = \frac{s\Omega_\alpha}{v_\perp} \frac{\partial}{\partial p_\perp} + (\underline{k}_\parallel - \underline{k}'_\parallel) \frac{\partial}{\partial p_\parallel}, \quad (4.6)$$

$$s\Omega_\alpha = \omega^\sigma(\underline{k}) - \omega^{\sigma'}(\underline{k}') - (\underline{k}_\parallel - \underline{k}'_\parallel)v_\parallel.$$

The terms linear in $N^{\sigma'}$ in (4.3) to (4.5) describe the effects of spontaneous scattering $\sigma' \rightarrow \sigma$, those linear in N^σ the effects of spontaneous scattering $\sigma \rightarrow \sigma'$, while the remaining terms involving $N^{\sigma'}N^\sigma$ describe the effects of induced scattering. If the waves are in the same mode ($\sigma = \sigma'$) (4.3) and (4.4) are the same while (4.5) includes the effect of each scattering twice and so only one half of the right hand side of (4.5) is to be taken. It is apparent from (4.3) and (4.4) that scattering conserves the total number of plasmons.

Similarly, in the semiclassical theory, nonlinear wave-wave interaction is described by a spontaneous probability $u^{\sigma\sigma'\sigma''}(\underline{k}, \underline{k}', \underline{k}'')$ which is the probability per unit time that a plasmon of mode σ' in the range $d^3\underline{k}'/(2\pi)^3$ at wavevector $\underline{k}'$ coalescences with a plasmon of mode σ'' in the range $d^3\underline{k}''/(2\pi)^3$ at $\underline{k}''$ to form a plasmon of mode σ in the range $d^3\underline{k}/(2\pi)^3$ at $\underline{k}$. The equations describing the evolution of the distribution functions of plasmons due to the decay and coalescence processes $\sigma' + \sigma'' \leftrightarrow \sigma$ are unchanged from the case of a plasma without a background magnetic field:

$$\left(\frac{\partial}{\partial t} + \underline{v}_g^\sigma \cdot \frac{\partial}{\partial \underline{r}} \right) N^\sigma(\underline{k}) = \int \frac{d^3\underline{k}'}{(2\pi)^3} \int \frac{d^3\underline{k}''}{(2\pi)^3} u^{\sigma\sigma'\sigma''}(\underline{k}, \underline{k}', \underline{k}'') \{ N^{\sigma'}(\underline{k}') N^{\sigma''}(\underline{k}'') - N^\sigma(\underline{k}) N^{\sigma'}(\underline{k}') - N^\sigma(\underline{k}) N^{\sigma''}(\underline{k}'') \}, \quad (4.7)$$

$$\left(\frac{\partial}{\partial t} + \mathbf{v}_g^{\sigma'} \cdot \frac{\partial}{\partial \mathbf{r}} \right) N^{\sigma'}(\mathbf{k}') = - \int \frac{d^3 \mathbf{k}}{(2\pi)^3} \int \frac{d^3 \mathbf{k}''}{(2\pi)^3} u^{\sigma\sigma'\sigma''}(\mathbf{k}, \mathbf{k}', \mathbf{k}'') \{ N^{\sigma'}(\mathbf{k}') N^{\sigma''}(\mathbf{k}'') - N^{\sigma}(\mathbf{k}) N^{\sigma'}(\mathbf{k}') - N^{\sigma}(\mathbf{k}) N^{\sigma''}(\mathbf{k}'') \}, \quad (4.8)$$

together with a further equation obtained by interchanging primed and double primed quantities in (4.8).

From these equations, total energy density of interacting waves can be shown to be conserved, i.e., $W^{\sigma} + W^{\sigma'} + W^{\sigma''} = \text{const.}$, where W^{σ} is defined by equation (3.2), etc.

3. SCATTERING OF WAVES

In this section, a general theory of scattering of waves in a magnetoactive plasma is presented ignoring intrinsically quantum mechanical effects (important for particles or plasmons of very high energies) but otherwise allowing arbitrary particle energies and taking into account effects of the background magnetic field on wave properties, on unperturbed (by waves) motions of scattering particles and on shielding fields which depend on dielectric properties of the plasma.

Scattering of waves in a plasma arises from contributions due to (a) Thomson (Compton) scattering, which can also occur in vacuo, and (b) nonlinear scattering, which can only occur in a medium. Thomson scattering (Fig. 4.1a) is connected with perturbed motions of a particle in the field of unscattered waves, while nonlinear scattering (Fig. 4.1b) is connected with nonlinear interactions of shielding fields of a particle with the field of unscattered waves.

In Thomson scattering of a wave in mode σ' , the current density which gives rise to scattered waves reads (Appendix D)

Fig. 4.1: Schematic diagrams of (a) Thomson scattering, (b) nonlinear scattering, and (c) a coalescence process. The circle represents a nonlinear interaction, the horizontal line a particle, the solid line a plasma wave and the dotted line a virtual plasma wave.

$$j_1^q(\underline{k}, \omega) = \frac{2\pi i q^2}{m} \sum_{s=-\infty}^{\infty} \int \frac{d^3 \underline{k}'}{(2\pi)^4} \frac{d\omega'}{\omega'} A_{ij}^q(s; \omega, \underline{k}; \omega', \underline{k}'; \underline{v}) E_j^{\sigma'}(\underline{k}', \omega') \delta\{\omega - \omega' - (\underline{k}_{\parallel} - \underline{k}'_{\parallel})v_{\parallel} - s\Omega\}, \quad (4.9)$$

where subscripts for particles have been dropped,

$$A_{ij}^q(s; \omega, \underline{k}; \omega', \underline{k}'; \underline{v}) = \sum_{\nu=-\infty}^{\infty} A_{ij}^q(s-\nu, \underline{k}, \omega; \nu, \underline{k}', \omega'; \underline{v}) \quad (4.10)$$

and

$$\begin{aligned} A_{ij}^q(s, \underline{k}, \omega; s', \underline{k}', \omega'; \underline{v}) &= \frac{e^{-i\epsilon(s'\psi' + s\psi)}}{\gamma \omega_s \omega_{s'}} \left\{ \omega_s \omega_{s'} J_s(z) J_{s'}(z') \tau_{ij} \right. \\ &+ \omega_s J_s(z) \tau_{im} k_m v_j^*(s', \underline{k}', \underline{v}) + \omega_{s'} J_{s'}(z') v_i(s, \underline{k}, \underline{v}) k'_m \tau_{mj} \\ &\left. + \left(k_\ell \tau_{\ell m} k'_m - \frac{\omega \omega'}{c^2} \right) v_i(s, \underline{k}, \underline{v}) v_j^*(s', \underline{k}', \underline{v}) \right\}, \quad (4.11) \end{aligned}$$

with symbols having their usual meanings, see equations (2.46), (2.47) for example,

$$\omega_s = \omega - s\Omega - k_{\parallel} v_{\parallel} , \quad \omega'_s = \omega' - s'\Omega - k'_{\parallel} v_{\parallel} , \quad (4.12)$$

$\tau_{ij} = \tau_{ij}(\omega_s)$ and

$$\tau_{ij}(\omega) \equiv \begin{pmatrix} \frac{\omega^2}{\omega^2 - \Omega^2} & \frac{i\epsilon\omega\Omega}{\omega^2 - \Omega^2} & 0 \\ \frac{-i\epsilon\omega\Omega}{\omega^2 - \Omega^2} & \frac{\omega^2}{\omega^2 - \Omega^2} & 0 \\ 0 & 0 & 1 \end{pmatrix} . \quad (4.13)$$

The current density which gives rise to scattered waves in non-linear scattering of a wave in mode σ' reads (Appendix D)

$$j_i^{n.s.}(\underline{k}, \omega) = \frac{2\pi i q^2}{m} \sum_{s=-\infty}^{\infty} \int \frac{d^3 \underline{k}'}{(2\pi)^4} \frac{d\omega'}{\omega'} \Gamma_{ijm}(\omega, \underline{k}; \omega', \underline{k}') V_m(s, \underline{k} - \underline{k}', \underline{v}) E_j^{\sigma'}(\underline{k}', \omega') \delta\{\omega - \omega' - (k_{\parallel} - k'_{\parallel}) v_{\parallel} - s\Omega\} , \quad (4.14)$$

where $V_m(s, \underline{k} - \underline{k}', \underline{v})$ is given by (2.46) and

$$\Gamma_{ijm}(\omega, \underline{k}; \omega', \underline{k}') \equiv \frac{2im\omega\omega'}{q(\omega - \omega')} \epsilon_{ij\ell}(\underline{k}', \omega'; \underline{k} - \underline{k}', \omega - \omega') \frac{\lambda_{\ell m}(\underline{k} - \underline{k}', \omega - \omega')}{\Lambda(\underline{k} - \underline{k}', \omega - \omega')} . \quad (4.15)$$

The total current density associated with the scattering is given by the sum of (4.9) and (4.14) and it is given by equation (4.9), with A_{ij}^q replaced by

$$A_{ij}(s; \underline{k}, \omega; \underline{k}', \omega'; \underline{v}) = A_{ij}^q(s; \underline{k}, \omega; \underline{k}', \omega'; \underline{v}) + \Gamma_{ijm}(\underline{k}, \omega; \underline{k}', \omega') V_m(s, \underline{k} - \underline{k}', \underline{v}) . \quad (4.16)$$

By a procedure exactly analogous to that outlined in the introduction of the last chapter, the scattering probability is derived from

the total current density associated with the scattering:

$$w_{\alpha}^{\sigma\sigma'}(s, \underline{p}, \underline{k}, \underline{k}') = \frac{4(2\pi)^3 q_{\alpha}^4}{m_{\alpha}^2} \left(\frac{W_E}{W_T} \right)^{\sigma} \left(\frac{W_E}{W_T} \right)^{\sigma'} \frac{|A^{\sigma\sigma'}(s, \underline{k}, \underline{k}', \underline{v})|^2}{\omega_{\sigma} \omega_{\sigma'}} \delta\{\omega^{\sigma} - \omega^{\sigma'} - (\underline{k}_{\parallel} - \underline{k}'_{\parallel}) v_{\parallel} - s\Omega_{\alpha}\}, \quad (4.17)$$

with

$$A^{\sigma\sigma'}(s, \underline{k}, \underline{k}', \underline{v}) \equiv e_i^{\sigma*}(\underline{k}) e_j^{\sigma'}(\underline{k}') A_{ij}(s; \underline{k}, \omega^{\sigma}; \underline{k}', \omega^{\sigma'}; \underline{v}). \quad (4.18)$$

On account of the symmetry properties of A_{ij}^q , ϵ_{ij} and ϵ_{ijl} , the probability (4.17), which is derived for scattering $\sigma' \rightarrow \sigma$, is equal to $w_{\alpha}^{\sigma'\sigma}(-s, \underline{p}, \underline{k}', \underline{k})$, a probability of the inverse scattering $\sigma \rightarrow \sigma'$. *A fortiori*, $w_{\alpha}^{\sigma\sigma'}(\underline{p}, \underline{k}, \underline{k}') = w_{\alpha}^{\sigma'\sigma}(\underline{p}, \underline{k}', \underline{k})$, with

$$w_{\alpha}^{\sigma\sigma'}(\underline{p}, \underline{k}, \underline{k}') \equiv \sum_{s=-\infty}^{\infty} w_{\alpha}^{\sigma\sigma'}(s, \underline{p}, \underline{k}, \underline{k}'). \quad (4.19)$$

4. THOMSON SCATTERING

When $|\Delta \underline{k}| \gg \lambda_{De}^{-1}$, with $|\Delta \underline{k}| \equiv |\underline{k} - \underline{k}'|$, shielding effects are negligible and the scattering process can be regarded as Thomson scattering. Astrophysical applications of Thomson scattering in a magnetoactive plasma have been considered by Canuto *et al.* (1971) and Goldstein and Lenchek (1971). Canuto *et al.* considered Thomson scattering of magnetoionic waves by non-relativistic electrons in the atmosphere of white dwarf stars with strong magnetic fields. Goldstein and Lenchek considered inverse Compton scattering, which is Thomson scattering by relativistic electrons, but they neglected the spiralling motion of the scattering electrons. In this section, Thomson scattering is studied more exactly, more generally, and in greater detail than in previous studies.

(a) Scattering Cross-Sections (I)

A quantity useful for comparing the different processes which determine a plasmon's mean free path is the differential scattering cross-section $\Sigma^{\sigma\sigma'}(\theta_r, \psi; \theta'_r, \psi')$ which relates the power generated in scattered plasmons propagating in the range of solid angle $d\cos\theta_r d\psi$ to the energy density flux in unscattered plasmons in the range of solid angle $d\cos\theta'_r d\psi'$. The plasmons propagate with group velocity

$$\underline{v}_g^\sigma(\underline{k}) \equiv \frac{\partial \omega^\sigma(\underline{k})}{\partial \underline{k}} \equiv v_g^\sigma(\sin\theta_r \cos\psi, \sin\theta_r \sin\psi, \cos\theta_r), \quad (4.20)$$

which is not parallel to $\underline{k} = |\underline{k}| (\sin\theta \cos\psi, \sin\theta \sin\psi, \cos\theta)$ in general. The energy density flux in the range $d^3\underline{k}/(2\pi)^3$ at $\underline{k}$, $F^\sigma(\underline{k})$, which is the Poynting vector only for non-spatially-dispersive media, may be identified as

$$\underline{F}^\sigma(\underline{k}) = \underline{v}_g^\sigma(\underline{k}) \hbar \omega^\sigma(\underline{k}) N^\sigma(\underline{k}). \quad (4.21)$$

Retaining only the effects of spontaneous scattering $\sigma' \rightarrow \sigma$ due to a single scattering particle in (4.3), the power generated in scattered plasmons is given by

$$P^\sigma = \int \frac{d^3\underline{k}}{(2\pi)^3} \int \frac{d^3\underline{k}'}{(2\pi)^3} \hbar \omega^\sigma(\underline{k}) w_{\alpha}^{\sigma\sigma'}(\underline{p}, \underline{k}, \underline{k}') N^{\sigma'}(\underline{k}'). \quad (4.22)$$

On the other hand,

$$P^\sigma \equiv \int_0^\infty d\Omega \int_0^{2\pi} d\psi \int_{-1}^{+1} d\cos\theta_r \int_0^\infty d\Omega' \int_0^{2\pi} d\psi' \int_{-1}^{+1} d\cos\theta'_r \Sigma^{\sigma\sigma'}(\theta_r, \psi; \theta'_r, \psi') I^{\sigma'}(\omega', \theta'_r, \psi'), \quad (4.23)$$

where $I^{\sigma'}(\omega', \theta'_r, \psi')$ is the intensity of a monochromatic flux of unscattered plasmons (power per unit surface area per unit solid angle about the ray direction per unit frequency range) and

$$\int_0^\infty d\omega' \int_0^{2\pi} d\psi' \int_{-1}^{+1} d\cos\theta'_r I^{\sigma'}(\omega', \theta'_r, \psi') := \int \frac{d^3\mathbf{k}'}{(2\pi)^3} |F^{\sigma'}(\mathbf{k}')| . \quad (4.24)$$

From (4.22) and (4.23), the cross-section reduces to

$$\Sigma^{\sigma\sigma'}(\theta_r, \psi; \theta'_r, \psi') = \int_0^\infty d\omega \frac{\omega^3 \eta_\sigma^2}{(2\pi c)^3 \omega'} \frac{\partial \cos\theta}{\partial \cos\theta_r} \frac{\partial \cos\theta'}{\partial \cos\theta'_r} \frac{1}{v_g^{\sigma'}} \frac{\partial}{\partial \omega} (\omega \eta_\sigma) w^{\sigma\sigma'}(\mathbf{p}, \mathbf{k}, \mathbf{k}') , \quad (4.25)$$

where $\eta_\sigma := |\mathbf{k}|c/\omega^\sigma$ is the refractive index.

For the scattering of magnetoionic waves the cross-section (4.25) reduces to

$$\Sigma^{\sigma\sigma'}(\theta_r, \psi; \theta'_r, \psi') = \sum_{s=-\infty}^{\infty} \left(\frac{q^2}{mc^2} \right) \frac{\eta_\sigma \omega^2}{\eta_{\sigma'} \omega'^2} \frac{(1+K_\sigma^2+T_\sigma^2)}{(1+T_\sigma^2)} \frac{(1+K_{\sigma'}^2+T_{\sigma'}^2)}{(1+T_{\sigma'}^2)} \frac{\frac{\partial \cos\theta}{\partial \cos\theta_r} \frac{\partial \cos\theta'}{\partial \cos\theta'_r} \frac{|A^{\sigma\sigma'}|^2}{(1+\delta_{\sigma'}^2)^{1/2} \left| 1 - \frac{\partial}{\partial \omega} (\omega \eta_\sigma) \frac{v_{||} \cos\theta}{c} \right|}} , \quad (4.26)$$

where (see I)

$$v_g^\sigma = \frac{c(1+\delta_\sigma^2)^{1/2}}{\partial(\omega \eta_\sigma)/\partial \omega} , \quad \cos\theta_r = \frac{\cos\theta + \delta_\sigma \sin\theta}{\{1+\delta_\sigma^2\}^{1/2}} , \quad (4.27)$$

$$\sin\theta_r = \frac{\sin\theta - \delta_\sigma \cos\theta}{\{1+\delta_\sigma^2\}^{1/2}} , \quad \delta_\sigma := \frac{1}{\eta_\sigma} \frac{\partial \eta_\sigma}{\partial \theta} = \frac{K_\sigma T_\sigma}{1+T_\sigma^2} ,$$

and the frequency of scattered wave ω is given by the solution of

$$\omega \left(1 - \eta_\sigma \frac{v_{||}}{c} \cos\theta \right) = \omega' \left(1 - \eta_{\sigma'} \frac{v_{||}}{c} \cos\theta' \right) + s\Omega \quad (4.28)$$

for each value of s , with $\eta_\sigma \equiv \eta_\sigma(\omega)$ and $\eta_{\sigma'} \equiv \eta_{\sigma'}(\omega')$. The result (4.26) disagrees with those of Canuto *et al.* (1971), because they have neglected spatial dispersion.

(b) Scattering by Non-relativistic Electrons (I)

If scattering electrons are non-relativistic, the small gyroradii conditions $z, z' \ll 1$ are sometimes satisfied. For magnetoionic waves, we write $z = \omega \eta_0 \beta \sin \theta \sin \alpha / \Omega_e$ and $z' = \omega' \eta_{\sigma'} \beta \sin \theta' \sin \alpha / \Omega_e$, where the pitch angle α is defined by $v_{\perp} \equiv |\underline{v}| |\sin \alpha|$. The conditions $z, z' \ll 1$ are satisfied provided ω, ω' are not much greater than Ω_e . Under these circumstances, significant scattering occurs only for small harmonic numbers s in (4.26), since the Bessel functions for given small arguments decrease rapidly with increases in s .

The scattering probability and cross-section in general depend on the azimuthal angle between the planes defined by $\underline{k}', \underline{B}_0$ and $\underline{k}, \underline{B}_0$. However, in practice, only waves distributed with axial symmetry about $\underline{B}_0$ are considered; therefore it is convenient to average over this azimuthal angle from the outset and denote the average by $\langle \dots \rangle$.

For zeroth harmonic scattering by non-relativistic electrons equation (4.28) gives $\omega \approx \omega'$ and (4.11) and (4.18) give $|A^{\sigma\sigma'}|^2 = |\tau_e^{\sigma\sigma'}|^2 \equiv |e_i^{\sigma*} e_j^{\sigma'} \tau_{ij}(\omega, \Omega_e)|^2$. Equation (2.82), together with the averaging process yields

$$\langle |\tau_e^{\sigma\sigma'}|^2 \rangle = \left\{ \frac{1+Y^2}{2(1-Y^2)^2} \left[(1+a_{\sigma}a_{\sigma'})^2 + (a_{\sigma}+a_{\sigma'})^2 \right] + \frac{2Y}{(1-Y^2)^2} \right. \\ \left. (1+a_{\sigma}a_{\sigma'})(a_{\sigma}+a_{\sigma'}) + (b_{\sigma}b_{\sigma'})^2 \right\} / (1+K_{\sigma}^2+T_{\sigma}^2)(1+K_{\sigma'}^2+T_{\sigma'}^2), \quad (4.29)$$

where a_{σ}, b_{σ} , etc., are given by (2.84) and $Y \equiv \Omega_e/\omega$. The corresponding scattering cross-section can be obtained by substituting (4.29) into (4.26):

$$\Sigma^{\sigma\sigma'}(\theta_r, \theta_r') = \left(\frac{e^2}{mc^2} \right) \frac{\eta_{\sigma}}{\eta_{\sigma'}} \frac{\partial \cos \theta}{\partial \cos \theta_r} \frac{\partial \cos \theta'}{\partial \cos \theta_r'} \left\{ \frac{1+Y^2}{2(1-Y^2)^2} \left[(1+a_{\sigma}a_{\sigma'})^2 + (a_{\sigma}+a_{\sigma'})^2 \right] \right. \\ \left. + \frac{2Y}{(1-Y^2)^2} (1+a_{\sigma}a_{\sigma'})(a_{\sigma}+a_{\sigma'}) + (b_{\sigma}b_{\sigma'})^2 \right\} / (1+T_{\sigma}^2)(1+T_{\sigma'}^2)(1+\delta_{\sigma'}^2)^{\frac{1}{2}}. \quad (4.30)$$

Example

When $\omega_p \ll \omega \ll \Omega_e$, a situation which is presumably possible for the magnetospheres of white dwarf stars, pulsars and perhaps Jupiter, the cross-section reduces to

$$\Sigma^{\sigma\sigma'}(\theta_r, \theta_r') = \left(\frac{e^2}{mc^2}\right)^2 \left\{ \frac{1+Y^2}{2(1-Y^2)^2} \left[(1+a_\sigma a_{\sigma'})^2 + (a_\sigma + a_{\sigma'})^2 \right. \right. \\ \left. \left. + \frac{2Y}{(1-Y^2)^2} (1+a_\sigma a_{\sigma'}) (a_\sigma + a_{\sigma'}) + (b_\sigma b_{\sigma'})^2 \right] / (1+T_\sigma^2)(1+T_{\sigma'}^2) \right\}. \quad (4.31)$$

From an analysis of (4.31), using (2.84), it can be shown that except for ordinary mode waves propagating perpendicularly to the background magnetic field, all other cross-sections are smaller than the classical Thomson cross-section $\sigma_T \equiv (8\pi/3)(e^2/mc^2)^2$ by approximately a factor $\omega^2/\Omega_e^2 \ll 1$, a result in qualitative agreement with Canuto *et al.* (1971), who considered only zeroth harmonic scattering ($\omega = \omega'$). However, shielding effects are important when $|\Delta k| \lesssim \lambda_{De}^{-1}$ and $|\Delta\omega| \ll \omega_p$, both conditions of which are satisfied for the zeroth harmonic Thomson scattering considered here.

For the first harmonic scattering however,

$\Omega_e/|k_{||}-k'_{||}|V_e \gtrsim \Omega_e c/|\omega-\omega'|V_e \approx c/V_e \gg 1$ and shielding effects are unimportant. When magnetoionic waves are scattered by non-relativistic electrons under the conditions $z, z' \ll 1$, equation (4.28) gives $\omega \approx \omega' + \Omega_e$ and equations (4.11) and (4.18) give

$$\langle |A^{\sigma\sigma'}|^2 \rangle = \{J_1^2(z) + J_1^2(z')\} \langle |\tau_e^{\sigma\sigma'}|^2 \rangle + 2J_1(z) J_1(z') b_\sigma b_{\sigma'}, \\ (a_\sigma a_{\sigma'} + a_\sigma Y - a_{\sigma'} Y - 1)/(1-Y^2)(1+K_\sigma^2+T_\sigma^2)(1+K_{\sigma'}^2+T_{\sigma'}^2), \quad (4.32)$$

where $\langle |\tau_e^{\sigma\sigma'}|^2 \rangle$ is given by (4.29). From the above expression and equation (4.26), it is seen that first harmonic scattering is smaller than zeroth harmonic scattering by a factor of the order of

$$J_1^2(z) \approx z^2/4 \ll 1.$$

In retrospect, the following conclusions have been reached with regard to Thomson scattering of magnetoionic waves by non-relativistic electrons:

(1) When the conditions $z, z' \ll 1$ are satisfied, Thomson scattering by non-relativistic electrons weakens as the harmonic number of scattering s is increased.

(2) In the presence of a strong background magnetic field, such that $\omega_p \ll \omega \ll \Omega_e$, Thomson scattering cross-section is about ω^2/Ω_e^2 times the classical Thomson cross-section (save in a special case).

However, we point out that shielding effects cannot be neglected for zeroth harmonic scattering by non-relativistic electrons and that scattering at higher harmonic numbers becomes important for mildly relativistic electrons and predominates for ultra-relativistic electrons, when $z, z' \gg 1$.

(c) Scattering by Relativistic Electrons

Thomson scattering by highly relativistic electrons, usually called inverse Compton scattering, leads to scattered waves of much greater frequencies than the unscattered waves. In this limit, it is possible, under certain conditions, to regard motion of the scattering electrons as rectilinear in treating the scattering. Such conditions are derived here for electrons with $\gamma \gg 1$.

The spiralling motion, neglected by Goldstein and Lenchek (1971), enters through the Bessel function in (4.11) and through the fact that τ_{ij} defined by (4.13) differs from the unit tensor δ_{ij} . The spiralling motion can be regarded as approximately rectilinear, for the purposes of treating the scattering, when τ_{ij} can be approximated by δ_{ij} and

when the arguments of the Bessel functions are much greater than unity, i.e. $z, z' \gg 1$. For high frequency magnetoionic waves ($\omega, \omega' \gg \omega_p$), the last requirement leads to

$$\gamma \gg \frac{\Omega_e}{\omega' \sin \theta' \sin \alpha} \quad \text{and} \quad \gamma \gg \frac{\Omega_e}{\omega \sin \theta \sin \alpha}. \quad (4.33)$$

Furthermore, the requirement $\tau_{ij} \approx \delta_{ij}$ leads to $\omega_s \approx \omega' (1 - \beta) \approx \omega' / \gamma^2 \gg \Omega_e / \gamma$ or

$$\frac{\omega'}{\Omega_e} \gg \gamma. \quad (4.34)$$

Combining the inequalities (4.33) and (4.34), it is found that spiraling motion for electrons with $\gamma \gg 1$ can be neglected provided

$$\frac{\omega'}{\Omega_e} \gg \gamma \gg \max \left(\frac{\Omega_e}{\omega' \sin \theta' \sin \alpha}, \frac{\Omega_e}{\omega \sin \theta \sin \alpha} \right). \quad (4.35)$$

These conditions necessarily entail $\omega' \gg \Omega_e$ and they are trivially satisfied as $\Omega_e \rightarrow 0$.

When the condition (4.35) is not satisfied, the anisotropy introduced by the background magnetic field cannot be ignored. Such cases do not appear to have been discussed in the literature. An example is presented in detail below.

Example

Consider inverse Compton scattering by a relativistic electron moving in the direction of a background magnetic field B_0 . Conditions (4.33) are violated since $\sin \alpha = 0$ implies $z = z' = 0$. But when condition (4.34) holds, i.e. $\omega' \gg \gamma \Omega_e$, τ_{ij} can be approximated by δ_{ij} . Equations (4.10) and (4.11) then give

$$A_{ij}^q(s=0, \underline{k}, \omega; \underline{k}', \omega'; \underline{v}) = \frac{1}{\gamma} \left\{ \delta_{ij} + \frac{k_i \delta_{j3} v}{\omega' - k_{\parallel}' v} + \frac{\delta_{i3} k_j' v}{\omega - k_{\parallel} v} \right. \\ \left. + \frac{\delta_{i3} \delta_{j3} v^2}{(\omega' - k_{\parallel}' v)(\omega - k_{\parallel} v)} \left[\underline{k} \cdot \underline{k}' - \frac{\omega \omega'}{c^2} \right] \right\}, \quad (4.36)$$

with $\omega' - k_{\parallel}' v = \omega - k_{\parallel} v$. For magnetoionic waves this can be rewritten in the form:

$$A_{ij}^q(s=0; \omega, \theta; \omega', \theta'; \underline{v}) = \frac{1}{\gamma} \left\{ \delta_{ij} + \frac{\kappa_i \delta_{j3} \eta_{\sigma} \beta}{(1 - \eta_{\sigma} \beta \cos \theta)} + \frac{\delta_{i3} \kappa_j' \eta_{\sigma'} \beta}{(1 - \eta_{\sigma'} \beta \cos \theta')} + \frac{\delta_{i3} \delta_{j3} \beta^2 (\eta_{\sigma} \eta_{\sigma'} \kappa \cdot \kappa' - 1)}{(1 - \eta_{\sigma} \beta \cos \theta)(1 - \eta_{\sigma'} \beta \cos \theta')} \right\}, \quad (4.37)$$

where $\kappa = k/|k|$, $\beta = v/c$, etc., as usual. From this and equations (2.82) and (4.18), it is simple to show

$$\begin{aligned} \langle |A^{\sigma\sigma'}|^2 \rangle &= \left\{ \frac{1}{2} \left[a_{\sigma} a_{\sigma'} + 1 + \frac{b_{\sigma} b_{\sigma'} \eta_{\sigma} \eta_{\sigma'} \beta^2 \sin \theta \sin \theta'}{(1 - \eta_{\sigma} \beta \cos \theta)(1 - \eta_{\sigma'} \beta \cos \theta')} \right]^2 \right. \\ &\quad + \frac{1}{2} (a_{\sigma} + a_{\sigma'})^2 + \left[b_{\sigma} b_{\sigma'} + \frac{\kappa_{\sigma} b_{\sigma'} \eta_{\sigma} \beta}{(1 - \eta_{\sigma} \beta \cos \theta)} + \frac{\kappa_{\sigma'} b_{\sigma} \eta_{\sigma'} \beta}{(1 - \eta_{\sigma'} \beta \cos \theta')} \right. \\ &\quad \left. \left. + \frac{b_{\sigma} b_{\sigma'} (\eta_{\sigma} \eta_{\sigma'} \cos \theta \cos \theta' - 1)}{(1 - \eta_{\sigma} \beta \cos \theta)(1 - \eta_{\sigma'} \beta \cos \theta')} \right]^2 \right\} / \gamma^2 (1 + \kappa_{\sigma}^2 + T_{\sigma}^2) (1 + \kappa_{\sigma'}^2 + T_{\sigma'}^2). \quad (4.38) \end{aligned}$$

This rather complicated expression simplifies considerably under the condition $\gamma \Omega_e \geq \omega_p$, in which case equations (2.81) and (2.84) give $\eta_{\sigma} \approx \eta_{\sigma'} \approx 1$, $T_{\sigma} \gg \kappa_{\sigma} \approx 0$, $T_{\sigma'} \gg \kappa_{\sigma'} \approx 0$, $a_{\sigma} \approx T_{\sigma} \cos \theta$, $b_{\sigma} \approx -T_{\sigma} \sin \theta$, etc. Equation (4.38) then reduces to

$$\begin{aligned} \langle |A^{\sigma\sigma'}|^2 \rangle &= \left\{ \frac{1}{2} \left[T_{\sigma} T_{\sigma'} \cos \theta \cos \theta' + 1 + \frac{T_{\sigma} T_{\sigma'} \beta^2 \sin^2 \theta \sin^2 \theta'}{(1 - \beta \cos \theta)(1 - \beta \cos \theta')} \right]^2 \right. \\ &\quad + \frac{1}{2} (T_{\sigma} \cos \theta + T_{\sigma'} \cos \theta')^2 + (T_{\sigma} T_{\sigma'} \sin \theta \sin \theta')^2 \\ &\quad \left. \left[1 + \frac{\beta^2 (\cos \theta \cos \theta' - 1)}{(1 - \beta \cos \theta)(1 - \beta \cos \theta')} \right]^2 \right\} / \gamma^2 (1 + T_{\sigma}^2) (1 + T_{\sigma'}^2). \quad (4.39) \end{aligned}$$

Case (i)

Consider an unscattered wave propagating along the lines of the background magnetic field, i.e. $\cos \theta' = \pm 1$. On application of (2.84), equation (4.39) reads

$$\langle |A^{\sigma\sigma'}|^2 \rangle = \frac{(1 \pm T_{\sigma} \cos \theta)}{2\gamma^2 (1 + T_{\sigma}^2)}, \quad (4.40)$$

where the upper sign refers to unscattered waves in 0 (X) mode propagating antiparallel (parallel) to $\underline{y}$ or $\underline{B}_0$, and the lower sign refers to unscattered waves in X (0) mode propagating antiparallel (parallel) to $\underline{y}$ or $\underline{B}_0$. Under quite general conditions the scattered waves are quasi-longitudinal ($\cos^2\theta \gg Y^2 \sin^4\theta/4$). This is the case for example under the conditions $Y \approx 10^{-2}$, $\gamma \approx 10$ for $\theta \leq 86^\circ$ and $\theta \geq 94^\circ$. Granted this, (4.40) reduces further to

$$\langle |A^{\sigma\sigma'}|^2 \rangle = \frac{(1 \pm \cos\theta)^2}{4\gamma^2}, \quad (4.41)$$

where the upper sign refers to $X \rightarrow 0$ or $0 \rightarrow X$ ($0 \rightarrow 0$ or $X \rightarrow X$) scattering with the unscattered waves propagating antiparallel (parallel) to $\underline{y}$ or $\underline{B}_0$ and the lower sign refers to $0 \rightarrow 0$ or $X \rightarrow X$ ($X \rightarrow 0$ or $0 \rightarrow X$) scattering with the unscattering waves propagating antiparallel (parallel) to $\underline{y}$ or $\underline{B}_0$.

The total scattering cross-section for a plasmon of mode σ' , propagating initially in direction $(\sin\theta'_r \cos\psi', \sin\theta'_r \sin\psi', \cos\theta'_r)$ and then scattering into a plasmon of mode σ can be defined by

$$\Sigma^{\sigma\sigma'}(\theta'_r, \psi') \equiv \int_0^{2\pi} d\psi \int_{-1}^{+1} d\cos\theta_r \Sigma^{\sigma\sigma'}(\theta_r, \psi; \theta'_r, \psi'), \quad (4.42)$$

where $\Sigma^{\sigma\sigma'}(\theta_r, \psi; \theta'_r, \psi')$ is determined by equation (4.25). When the scattering process is axisymmetric, the integrals can be recasted to read

$$\begin{aligned} \Sigma^{\sigma\sigma'}(\theta'_r) &= \int_{-1}^{+1} d\cos\theta_r \int_0^\omega d\omega \frac{2\pi\omega^3 \eta_\sigma^2}{(2\pi c)^3 \omega'} \frac{\partial \cos\theta}{\partial \cos\theta'_r} \frac{\partial \cos\theta'}{\partial \cos\theta'_r} \\ &\quad \frac{1}{v_g} \frac{\partial}{\partial \omega} (\omega \eta_\sigma) \langle w_e^{\sigma\sigma'}(\underline{p}, \underline{k}, \underline{k}') \rangle, \end{aligned} \quad (4.43)$$

where $\langle \dots \rangle$ again denotes an average over azimuthal angles.

For the present problem $\omega' \gg \gamma \Omega_e \geq \omega_p$, application of this expression together with an integration with respect to ω over the delta

function in the probability gives

$$\Sigma^{\sigma\sigma'}(\cos\theta' = \mp 1) = 2\pi \left(\frac{e^2}{mc^2}\right)^2 (1 \pm \beta)^2 \int_{-1}^{+1} d\cos\theta \frac{\langle |A^{\sigma\sigma'}|^2 \rangle}{(1 - \beta \cos\theta)^3}, \quad (4.44)$$

where the approximations $\theta' = \theta'_r$ and $\theta = \theta_r$ have been used ($\omega, \omega' \gg \omega_p$). The upper (lower) sign refers to antiparallel (parallel) propagation of unscattered waves. $\langle |A^{\sigma\sigma'}|^2 \rangle$ is given by equation (4.41). The integration over θ includes a small cone about the perpendicular direction where the quasilongitudinal approximation fails for scattered waves. But it can be shown that this introduces only a negligible error.

Evaluation of the integral in (4.44), by using standard integrals of the form

$$\int_{-1}^{+1} \frac{d\cos\theta}{(1 - \beta \cos\theta)^n} = \frac{1}{\beta(n-1)} \left\{ \frac{(1+\beta)^{n+1} - (1-\beta)^{n+1}}{(1-\beta^2)^{n-1}} \right\}, \quad (n \geq 2) \quad (4.45)$$

gives the total scattering cross-section,

$$\Sigma^{\sigma\sigma'}(\cos\theta' = \pm 1) = 2\pi \left(\frac{e^2}{mc^2}\right)^2 g(\gamma) \quad (4.46)$$

where $g(\gamma)$ is given by the following table of approximate values:

Table 4.1

Mode of Scattering $\cos\theta'$	$0 \rightarrow 0$ $X \rightarrow X$	$0 \rightarrow X$ $X \rightarrow 0$
-1	$16\gamma^2$	$\ln(4\gamma^2)/\gamma^2$
+1	$\ln(4\gamma^2)/16\gamma^4$	1

These results can be compared with those for the case of an isotropic distribution of unscattered waves without a background magnetic field where $g(\gamma) = 16\gamma^2/9$ (Melrose, 1971).

Case (ii)

Consider an unscattered wave propagating perpendicular to the background magnetic field. The total scattering cross-section in this case reads

$$\Sigma^{\sigma\sigma'}(\cos\theta'=0) = 2\pi \left(\frac{e^2}{mc^2} \right)^2 \int_{-1}^{+1} d\cos\theta \frac{\langle |A^{\sigma\sigma'}|^2 \rangle}{(1-\beta\cos\theta)^3}, \quad (4.47)$$

where we have from equation (4.39),

$$\begin{aligned} \langle |A^{\sigma\sigma'}|^2 \rangle = & \left\{ \frac{1}{2} (1 + T_{\sigma}^2 \cos^2\theta + T_{\sigma'}^2 \sin^2\theta) + \frac{T_{\sigma} T_{\sigma'} \beta^2 \sin^2\theta (1 - 2T_{\sigma} T_{\sigma'})}{(1 - \beta\cos\theta)} \right. \\ & \left. + \frac{1}{2} \frac{T_{\sigma}^2 T_{\sigma'}^2 \beta^2 \sin^2\theta (2 + \sin^2\theta)}{(1 - \beta\cos\theta)^2} \right\} / \gamma^2 (1 + T_{\sigma}^2) (1 + T_{\sigma'}^2). \end{aligned} \quad (4.48)$$

On account of the large momentum transmitted by the electron in process of scattering, the scattered waves can be regarded as satisfying the quasilongitudinal approximation for the purpose of estimating the total scattering cross-section. This will be justified *a posteriori*. Equation (4.48) then reads

$$\begin{aligned} \langle |A^{\sigma\sigma'}|^2 \rangle = & \left\{ \frac{1}{2} (1 + \cos^2\theta + T_{\sigma'}^2 \sin^2\theta) - \frac{T_{\sigma'}^2 \beta^2 \sin^2\theta (2T_{\sigma'} \pm 1)}{(1 - \beta\cos\theta)} \right. \\ & \left. + \frac{1}{2} \frac{T_{\sigma'}^2 \beta^2 \sin^2\theta (2 + \sin^2\theta)}{(1 - \beta\cos\theta)^2} \right\} / 2\gamma^2 (1 + T_{\sigma'}^2), \end{aligned} \quad (4.49)$$

where the upper (lower) sign refers to scattered waves in the 0 (X) mode. Again the integral in (4.47), with $\langle |A^{\sigma\sigma'}|^2 \rangle$ given by (4.49), can be evaluated using integral formulae (4.45) to give

$$\begin{aligned} \Sigma^{\sigma\sigma'}(\cos\theta'=0) = & \left(\frac{e^2}{mc^2} \right)^2 \frac{\pi}{\gamma^2 (1 + T_{\sigma'}^2)} \left\{ \frac{1}{2\beta^3} \left[\frac{4\beta^3}{(1 - \beta^2)^2} + (1 + T_{\sigma'}^2) \ln \left(\frac{1 + \beta}{1 - \beta} \right) \right] \right. \\ & \left. - \frac{4T_{\sigma'} (2T_{\sigma'} \pm 1) (3 - 2\beta^2)}{3(1 - \beta^2)^2} + \frac{T_{\sigma'}^2}{2\beta^3} \left[\frac{2\beta(3 + 8\beta^2 - 7\beta^4)}{3(1 - \beta^2)^2} + \ln \left(\frac{1 + \beta}{1 - \beta} \right) \right] \right\}, \end{aligned} \quad (4.50)$$

which, for highly relativistic electrons $\gamma \gg 1$, is approximated by

$$\Sigma^{\sigma\sigma'}(\cos\theta'=0) = \left(\frac{e^2}{mc^2}\right)^2 \frac{\pi}{\gamma^2(1+T_{\sigma'}^2)} \left\{ \frac{1}{2} (4\gamma^4 - T_{\sigma'}^2 \ln(4\gamma^2)) - \frac{4}{3} \gamma^4 T_{\sigma'} (2T_{\sigma'} \pm 1) + \frac{4}{3} T_{\sigma'} (2T_{\sigma'} \pm 1) + \frac{4}{3} \gamma^6 T_{\sigma'}^2 \right\} \quad (4.51)$$

For unscattered waves propagating perpendicularly to the background magnetic field, $T_{\sigma'}^2 \rightarrow \infty$ for O-mode waves and $T_{\sigma'}^2 \rightarrow 0$ for X-mode waves. Hence the total scattering cross-section for scattered waves in either mode is given again by (4.46) with

$$g(\gamma) = \begin{cases} 2\gamma^4/3 & \text{(O-mode unscattered waves)} \\ \gamma^2 & \text{(X-mode unscattered waves)} \end{cases} \quad (4.52)$$

The negligible effect of scattered waves in a small cone about $\theta = \pi/2$ on the estimates of the total scattering cross-sections can be shown as follows. For $\theta = \pi/2$, $\langle |A^{\sigma\sigma'}|^2 \rangle$ and also the integrand in (4.47) can be approximated by (see equation (4.48))

$$\left(\frac{1}{2} + T_{\sigma} T_{\sigma'} + \frac{T_{\sigma}^2 T_{\sigma'}^2}{2\gamma^2} \right) / \gamma^2 (1+T_{\sigma}^2) (1+T_{\sigma'}^2), \quad (4.53)$$

which is smaller than the corresponding integrands for scattered waves in the quasilongitudinal approximation by at least a factor $\gamma^4 \ll 1$. Hence the assumption made in expression (4.49) is justified *a posteriori*.

In conclusion, we note that relativistic electrons moving in the direction of a background magnetic field are particularly effective in scattering O-mode magnetoionic waves propagating perpendicularly to the background magnetic field under the conditions $\omega' \gg \gamma \Omega_e \geq \omega_p$.

5. NONLINEAR SCATTERING

The effects of shielding become important when the changes in wavevector $|\Delta \mathbf{k}| = |\mathbf{k} - \mathbf{k}'|$ and in frequency $|\Delta \omega| = |\omega - \omega'|$ on scattering

satisfy $|\Delta k| \leq \lambda_{De}^{-1}$ and $|\Delta \omega| \ll \omega_p$ (Bekefi, 1966).

Assuming the nonlinear response tensor ϵ_{ijl} is determined principally by electrons, one can infer that the two terms in (4.16), one due to Thomson scattering, the other due to nonlinear scattering, always have opposite signs for electrons (as scatterers) and have the same sign for ions. In this sense, the two scattering processes interfere destructively for electrons and constructively for ions. For relativistic particles, Thomson or inverse Compton scattering always predominates over nonlinear scattering. For non-relativistic electrons, there are large cancellations between the two terms in (4.16). Although the cancellation is not as complete as for the case without a background magnetic field, the resultant scattering is nevertheless at least a factor $z^2 = (k_{\perp} v_{\perp} / \Omega_e)^2 \ll 1$ smaller than either Thomson scattering or nonlinear scattering taken alone. For non-relativistic ions however, nonlinear scattering predominates over Thomson scattering by a factor $(m_i / m_e)^2 \gg 1$.

In a plasma, where $T_e \geq T_i$, when

$$\Delta \omega \ll |\Delta k_{\parallel}| v_i \ll |\Delta k_{\parallel}| v_e, \quad (4.54)$$

nonlinear scattering by ions is the dominant process. When

$$|\Delta k_{\parallel}| v_i \ll \Delta \omega \ll |\Delta k_{\parallel}| v_e, \quad (4.55)$$

scattering by ions is exponentially small and scattering by electrons is dominant. Finally when

$$|\Delta k_{\parallel}| v_e \ll \Delta \omega, \quad (4.56)$$

scattering by either ions or electrons is exponentially small.

(a) Inverse Maxwellian Tensor

The tensor $\lambda_{ij}(\underline{k}, \omega)/\Lambda(\underline{k}, \omega)$, occurring in (3.7) and (4.15) for example, is often called the inverse Maxwellian tensor. For arbitrary directions $\underline{k} = (\sin\theta\cos\psi, \sin\theta\sin\psi, \cos\theta)$ and arbitrary dielectric tensor $\epsilon_{ij} = \epsilon_{ij}(\underline{k}, \omega)$, standard matrix theory gives (see equations (2.54) and (2.56))

$$\Lambda(\underline{k}, \omega) = A\eta^4 - B\eta^2 + C, \quad (4.57)$$

with

$$\left. \begin{aligned} A &= \epsilon_{11} \sin^2\theta + 2\epsilon_{13} \sin\theta\cos\theta + \epsilon_{33} \cos^2\theta, \\ B &= \epsilon_{22} A + \epsilon_{11}\epsilon_{33} - \epsilon_{13}^2 + (\epsilon_{23} \cos\theta - \epsilon_{12} \sin\theta)^2, \\ C &= \det(\epsilon_{ij}), \end{aligned} \right\} \quad (4.58)$$

and

$$\left. \begin{aligned} \lambda_{11} &= \eta^4 \sin^2\theta \cos^2\psi - \eta^2 \{ 2\text{Re}\epsilon_{23} \cos\theta \sin\theta \sin\psi + \epsilon_{22} \sin^2\theta \\ &\quad + \epsilon_{33} (1 - \sin^2\theta \sin^2\psi) \} + (\epsilon_{22}\epsilon_{33} - \epsilon_{23}^2), \\ \lambda_{22} &= \eta^4 \sin^2\theta \sin^2\psi - \eta^2 \{ 2\text{Re}\epsilon_{13} \cos\theta \sin\theta \cos\psi + \epsilon_{11} \sin^2\theta \\ &\quad + \epsilon_{33} (1 - \sin^2\theta \cos^2\psi) \} + (\epsilon_{11}\epsilon_{33} - \epsilon_{13}^2), \\ \lambda_{33} &= \eta^4 \cos^2\theta - \eta^2 \{ 2\text{Re} \sin^2\theta \cos\psi \sin\psi + \epsilon_{11} (1 - \sin^2\theta \sin^2\psi) \\ &\quad + \epsilon_{22} (1 - \sin^2\theta \cos^2\psi) \} + (\epsilon_{11}\epsilon_{22} - \epsilon_{12}^2), \\ \lambda_{12} &= \lambda_{21}^* = \eta^4 \sin^2\theta \cos\psi \sin\psi + \eta^2 \{ (\epsilon_{23}^* \cos\psi + \epsilon_{13} \sin\psi) \cos\theta \sin\theta \\ &\quad + (\epsilon_{12} + \epsilon_{33} \cos\psi \sin\psi) \sin^2\theta \} + (\epsilon_{23}\epsilon_{13} - \epsilon_{12}\epsilon_{33}), \\ \lambda_{23} &= \lambda_{32}^* = \eta^4 \cos\theta \sin\theta \sin\psi + \eta^2 \{ (\epsilon_{12}^* \cos\theta + \epsilon_{13} \sin\theta \sin\psi) \sin\theta \cos\psi \\ &\quad + \epsilon_{23} (1 - \sin^2\theta \cos^2\psi) - \epsilon_{11} \cos\theta \sin\theta \sin\psi \} + (\epsilon_{12}\epsilon_{13} - \epsilon_{11}\epsilon_{23}), \\ \lambda_{13} &= \lambda_{31}^* = \eta^4 \cos\theta \sin\theta \cos\psi + \eta^2 \{ (\epsilon_{12} \cos\theta + \epsilon_{23} \sin\theta \cos\psi) \sin\theta \sin\psi \\ &\quad + \epsilon_{13} (1 - \sin^2\theta \sin^2\psi) - \epsilon_{22} \cos\theta \sin\theta \sin\psi \} + (\epsilon_{12}\epsilon_{23} - \epsilon_{22}\epsilon_{13}), \end{aligned} \right\} \quad (4.59)$$

with $\eta^2 \equiv |\underline{k}|^2 c^2 / \omega^2$ and '*' denoting complex conjugation. Tsytovich

and Shvartsburg (1967) have written down an inverse Maxwellian tensor, which is valid only when $\sin\theta = 0$. No such assumption has been made here.

Under quite general conditions, discussed in detail in Appendix E for various cases,

$$\eta^2 = |\Delta \underline{k}|^2 c^2 / |\Delta \omega|^2 \gg |\epsilon_{ij}(\Delta \underline{k}, \Delta \omega)|, \quad (4.60)$$

and the inverse Maxwellian tensor reduces to

$$\frac{\lambda_{ij}(\Delta \underline{k}, \Delta \omega)}{\Lambda(\Delta \underline{k}, \Delta \omega)} = \frac{\kappa_i \kappa_j}{\epsilon^{\ell}(\Delta \underline{k}, \Delta \omega)}, \quad (4.61)$$

where $\underline{\kappa} = \Delta \underline{k} / |\Delta \underline{k}|$ and $\epsilon^{\ell}(\Delta \underline{k}, \Delta \omega) = A$ (see (4.58)). When (4.60) is not satisfied, the inverse Maxwellian tensor has to be obtained from equations (4.57) to (4.59). It is assumed, in what follows, that the inverse Maxwellian tensor can be approximated by (4.61).

(b) Scattering by Ions (II)

In the scattering of high frequency waves, the conditions:

$$\frac{\omega}{|\underline{k}|} \gg V_e, \quad \text{and} \quad \frac{\omega'}{|\underline{k}'|} \gg V_e \gg \frac{|\Delta \omega|}{|\Delta \underline{k}|}, \quad (4.62)$$

are often fulfilled. The nonlinear response tensor then reads (see (F.6), Appendix F)

$$\begin{aligned} \epsilon_{ij\ell}(\underline{k}', \omega'; \Delta \underline{k}, \Delta \omega) = & -\frac{ie}{2m_e \omega^2} \left\{ \tau_{ij}^{(e)}(\omega) \Delta k_s \{ \epsilon_{s\ell}^{(e)}(\Delta \underline{k}, \Delta \omega) - \delta_{s\ell} \} \right. \\ & \left. + \frac{\omega_p^2}{\omega'} \frac{\Delta k_s}{\Delta \omega} \{ \tau_{i\ell}^{(e)}(\omega) \tau_{sj}^{(e)}(\omega') - \tau_{is}^{(e)}(\omega) \tau_{\ell j}^{(e)}(\omega') \} \right\} \quad (4.63) \end{aligned}$$

where $\tau_{ij}^{(e)}(\omega)$ and $\tau_{ij}^{(e)}(\omega')$ are given by (4.13) with $\Omega = \Omega_e$ and $\epsilon = -1$, and $\epsilon_{s\ell}^{(e)}$ is the electronic contribution to the dielectric tensor.

Substitution of (4.61) and (4.63) into (4.15) gives

$$\Gamma_{ijm}(\underline{k}, \omega; \underline{k}', \omega') = \left(\frac{e}{q} \right) \left(\frac{m}{m_e} \right) \left(\frac{\omega'}{\omega} \right) \tau_{ij}^{(e)}(\omega) \frac{\Delta k_m}{\Delta \omega} \left\{ \frac{\epsilon^{\ell(e)}(\Delta \underline{k}, \Delta \omega) - 1}{\epsilon^{\ell}(\Delta \underline{k}, \Delta \omega)} \right\}. \quad (4.64)$$

For the case of scattering by ions with $q = Z_i e$ and $m = m_i$, equation (E.10) can be used to reduce (4.64) to

$$\Gamma_{ijm}(\underline{k}, \omega; \underline{k}', \omega') = \left(\frac{m_i}{Z_i m_e} \right) \left(\frac{\omega'}{\omega} \right) \frac{\Delta k_m}{\Delta \omega} \frac{\tau_{ij}^{(e)}(\omega)}{(1 + Z_i T_e / T_i)}. \quad (4.65)$$

It is convenient to define a probability averaged over the distribution of scattering particles $f_\alpha(\underline{v})$:

$$w_\alpha^{\sigma\sigma'}(s, \underline{k}, \underline{k}') \equiv \int d^3 \underline{v} f_\alpha(\underline{v}) w_\alpha^{\sigma\sigma'}(s, \underline{p}, \underline{k}, \underline{k}'), \quad (4.66)$$

where $f_\alpha(\underline{v})$ is normalized as usual and $w_\alpha^{\sigma\sigma'}(s, \underline{p}, \underline{k}, \underline{k}')$ is given by (4.17).

Consider nonlinear scattering by a Maxwellian distribution of ions. After some calculations using (4.16), (4.17) and (4.65), the averaged probability (4.66) simplifies to

$$w_i^{\sigma\sigma'}(\underline{k}, \underline{k}') = \frac{4(2\pi)^{5/2} Z_i^2 e^4}{m_e^2 \omega^\sigma \omega^{\sigma'}} \left(\frac{W_E}{W_T} \right)^\sigma \left(\frac{W_E}{W_T} \right)^{\sigma'} \frac{|e_i^{\sigma*} \tau_{ij}^{(e)}(\omega) e_j^{\sigma'}|^2}{(1 + Z_i T_e / T_i)^2} \\ \times \sum_{s=-\infty}^{\infty} \frac{\zeta_s(\lambda_i)}{|\Delta k_{\parallel}| V_i} \exp \left\{ -\frac{(\omega^\sigma - \omega^{\sigma'} - s \Omega_i)^2}{2 |\Delta k_{\parallel}|^2 V_i^2} \right\}, \quad (4.67)$$

where $\lambda_i = (\Delta k_{\perp} V_i / \Omega_i)^2$ and $\zeta_s(\lambda_i)$ is given by (2.86).

In the limiting case of negligible small ion gyroradii, i.e., $\lambda_i \ll 1$, the contribution from different values of s decreases rapidly with increase in $|s|$. The contribution from $s = 0$, with $\zeta_0(\lambda_i) \approx 1$, dominates.

In the opposite limiting case of arbitrarily large ion gyroradii, i.e., $\lambda_i \gg 1$, the motion of the ions can be regarded as rectilinear for the purpose of treating scattering. The δ -function in (4.17) can be

replaced by $\delta\{(\omega^\sigma - \omega^{\sigma'}) - (\underline{k} - \underline{k}') \cdot \underline{v}\}$ and $V_m(s, \underline{k} - \underline{k}', \underline{v})$ can be replaced by v_m . With this approximation, the probability (4.67) is replaced by

$$w_i^{\sigma\sigma'}(\underline{k}, \underline{k}') = \frac{4(2\pi)^{5/2} Z_i^2 e^4}{m_e^2 \omega^\sigma \omega^{\sigma'}} \left(\frac{W_E}{W_T} \right)^\sigma \left(\frac{W_E}{W_T} \right)^{\sigma'} \frac{|e_i^{\sigma*} \tau_{ij}^{(e)}(\omega) e_j^{\sigma'}|^2}{(1 + Z_i T_e / T_i)^2} \times \frac{1}{|\Delta \underline{k}| v_i} \exp \left\{ -\frac{(\omega^\sigma - \omega^{\sigma'})^2}{2 |\Delta \underline{k}|^2 v_i^2} \right\}. \quad (4.68)$$

Except for cases of little interest, the conditions under which the contributions from $|s| = 1$ could be comparable to that from $s = 0$, are just those for (4.68) to be a reasonable approximation.

(c) Scattering by Electrons

In the regime (4.55), it is evident from the results of the previous section that scattering by ions is an exponentially small effect and scattering by electrons is dominant.

In the limit of arbitrarily large gyroradii of electrons, i.e., $k_\perp v_\perp / \Omega_e \gg 1$ for all relevant waves, the motion of the electrons can be regarded as rectilinear for the purpose of treating scattering and the results would reduce to those for the case of a plasma without a background magnetic field. It is interesting to consider the opposite limiting case of negligibly small electron gyroradii, i.e. $k_\perp v_\perp / \Omega_e \ll 1$. For this purpose, it is convenient to use the current density due to nonlinear scattering in the form given by (D.15) with (D.16) and (D.17), since this facilitates calculations of (4.16).

For the zeroth harmonic scattering, which is dominant in this context, Thomson scattering by non-relativistic electrons contributes a term (see (4.11)):

$$A_{ij}^q(\underline{k}, \omega; \underline{k}', \omega'; \underline{v}) = J_0(z) J_0(z') \tau_{ij}^{(e)}(\omega), \quad (4.69)$$

whilst nonlinear scattering contributes, on calculation using (D.15), (D.16), (D.17), (2.46), (E.17) and (4.63), the term,

$$A_{ij}^{n.s.}(\underline{k}, \omega; \underline{k}', \omega'; \underline{v}) = -J_0(z) J_0(z') \tau_{ij}^{(e)}(\omega) + \frac{i\Delta k_{\perp} v_{\perp}}{\Delta\omega} J_0'(z) J_0(z') \\ \times \sin(\psi'' + \psi) \tau_{ij}^{(e)}(\omega), \quad (4.70)$$

where

$$\tan\psi'' \equiv \frac{k_{\perp} \sin\psi - k'_{\perp} \sin\psi'}{k_{\perp} \cos\psi - k'_{\perp} \cos\psi'}. \quad (4.70)'$$

Substituting the sum of terms (4.69) and (4.70) into (4.17), the appropriate scattering probability is obtained. On averaging over a Maxwellian distribution of electrons, by (4.66), the probability reads

$$w_e^{\sigma\sigma'}(\underline{k}, \underline{k}') = \frac{4(2\pi)^{5/2} e^4}{m_e^2 \omega^{\sigma} \omega'^{\sigma'}} \left(\frac{W_E}{W_T} \right)^{\sigma} \left(\frac{W_E}{W_T} \right)^{\sigma'} |e^{\sigma*} \tau_{ij}^{(e)}(\omega) e_j^{\sigma'}|^2 \\ \times \left(\frac{k_{\perp} v_e}{\Delta\omega} \right)^2 \frac{2\lambda_e \sin^2(\psi'' + \psi)}{|\Delta k_{\parallel}| v_e} \exp \left\{ -\frac{(\omega^{\sigma} - \omega'^{\sigma'})}{2|\Delta k_{\parallel}|^2 v_e^2} \right\}, \quad (4.71)$$

where the approximations $J_0(z') \approx 1$ and $\exp(-\lambda_e) \{I_0(\lambda_e) - I_0'(\lambda_e)\} \approx 1$ have been made and here, $\lambda_e \equiv k_{\perp}^2 v_e^2 / \Omega_e^2 \ll 1$.

Comparison of (4.67) and (4.71) shows that scattering by electrons is more effective or comparable to scattering by ions provided

$$\frac{|\Delta\omega|^2}{2|\Delta k_{\parallel}|^2 v_i^2} \gtrsim \ln \left\{ \frac{Z_i^2 v_e / 2v_i}{(1 + Z_i T_e / T_i)^2} \left(\frac{\Delta\omega}{k_{\perp} v_e} \right)^2 \left(\frac{\Omega_e}{k_{\perp} v_e} \right)^2 \right\}, \quad (4.72)$$

where it is assumed that $v_e \gg v_i$.

It is a simple matter to write down scattering cross-sections for scattering by ions or electrons, by merely substituting the appropriate probability into (4.25).

Example

Consider nonlinear scattering of magnetoionic waves by plasma ions under the conditions $\omega_p \ll \omega \ll \Omega_e$ (cf. example in section 4b). Insertion of either (4.67) or (4.68) into (4.25) gives the cross-section,

$$\begin{aligned} \Sigma^{\sigma\sigma'}(\theta, \theta') = & \sqrt{\left(\frac{\pi}{2}\right)} \frac{Z_i^2}{(1+Z_i T_e/T_i)^2} \frac{(e^2/m_e c^2)^2}{\xi(1+T_\sigma^2)(1+T_{\sigma'}^2)} \left\{ \frac{1+Y^2}{2(1-Y^2)^2} ((1+a_\sigma a_{\sigma'})^2 \right. \\ & + (a_\sigma + a_{\sigma'})^2) + \frac{2Y}{(1-Y^2)^2} (1+a_\sigma a_{\sigma'})(a_\sigma + a_{\sigma'}) + (b_\sigma b_{\sigma'}) \\ & \left. \int_0^\infty \frac{d\omega}{\omega'} \exp \left\{ -\frac{(1-\omega/\omega')^2}{2\xi^2} \right\} \right\}, \quad (4.73) \end{aligned}$$

where $\xi = |\cos\theta - \cos\theta'| V_i/c$ when $\lambda_i \ll 1$, $\xi = \sqrt{2(1-\kappa\kappa')} V_i/c$ when $\lambda_i \gg 1$, and approximations $\omega \approx \omega'$, $\theta = \theta_r$ and $\theta' = \theta'_r$ have been made. The integral in (4.73) can be accurately approximated ($\xi \ll 1$) to give

$$\begin{aligned} \Sigma^{\sigma\sigma'}(\theta, \theta') = & \frac{Z_i^2}{(1+Z_i T_e/T_i)^2} \frac{\pi(e^2/m_e c^2)^2}{(1+T_\sigma^2)(1+T_{\sigma'}^2)} \left\{ \frac{1+Y^2}{2(1-Y^2)^2} ((1+a_\sigma a_{\sigma'}) \right. \\ & + (a_\sigma + a_{\sigma'})^2) + \frac{2Y}{(1-Y^2)^2} (1+a_\sigma a_{\sigma'})(a_\sigma + a_{\sigma'}) + (b_\sigma b_{\sigma'})^2 \left. \right\}. \quad (4.73)' \end{aligned}$$

From this, it is seen that for certain directions of propagation (e.g., $\theta' = 0$) the scattering cross-section is reduced by a factor $\omega^2/\Omega_e^2 \ll 1$.

(d) Absorption Coefficients

The Landau-cyclotron damping decrements and the collisional damping decrements, written down in section 4 of chapter 3, are also known as true absorption coefficients, since the wave amplitudes are truly attenuated by the corresponding interactions with plasma particles. There is also an absorption coefficient due to scattering by plasma particles and this is generally scattering by ions, provided the electron temperature T_e does not greatly exceed the ion temperature T_i .

(Bekefi, 1966).

From (4.3), such absorption coefficients due to spontaneous scattering reads

$$\gamma_{sc}^{\sigma}(\underline{k}) = n_i \int d^3p \int \frac{d^3\underline{k}'}{(2\pi)^3} w_i^{\sigma\sigma'}(p, \underline{k}, \underline{k}') f_i(p). \quad (4.74)$$

As an example, consider the nonlinear scattering of high frequency magnetoionic waves by plasma ions. The conditions $\omega' \geq 2\omega_p \gg \Omega_e$ imply $T_{\sigma}^2 \approx K_{\sigma} \approx \Omega_e/\omega$, so that $a_{\sigma} \approx T_{\sigma} \cos\theta$, $b_{\sigma} \approx -T_{\sigma} \sin\theta$, $T_{\sigma}^2 \approx 1$, etc. After a little manipulation, the equations (4.66), (4.67) and (4.29) give

$$\begin{aligned} \gamma_{sc}^{\sigma}(\omega, \theta) = & \left(\frac{\pi}{2}\right)^{1/2} \frac{Z_i^2 e^4}{2m_e^2 c^3 \omega} \frac{n_i}{(1 + Z_i T_e/T_i)^2} \int_{-1}^{+1} \frac{d\cos\theta'}{\xi} \{ (1 + \cos^2\theta)(1 + \cos^2\theta') \\ & + 2(1 - \cos^2\theta)(1 - \cos^2\theta') \} \int_0^{\infty} d\omega' \exp\left\{-\frac{(1 - \omega'/\omega)^2}{2\xi^2}\right\}, \quad (4.75) \end{aligned}$$

where $\xi = |\cos\theta - \cos\theta'| V_i/c$ when $\lambda_i \ll 1$ and $\xi = \sqrt{2(1 - \cos\theta \cos\theta')} V_i/c$ when $\lambda_i \gg 1$. Integration gives finally

$$\gamma_{sc}^{\sigma} = \frac{8\pi}{3} \left(\frac{e^2}{m_e c^2}\right)^2 c \frac{n_i Z_i^2}{(1 + Z_i T_e/T_i)^2}, \quad (4.76)$$

where $(8\pi/3)(e^2/m_e c^2)^2 = \sigma_T$ = Thomson cross-section. Such absorption processes have been considered in connection with ionospheric research (see Bekefi, 1966).

In conclusion, we have discussed nonlinear scattering by ions and electrons in a magnetoactive plasma and have shown that nonlinear scattering of magnetoionic waves by plasma ions can be strongly suppressed under the conditions $\omega_p \ll \omega \ll \Omega_e$.

6. WAVE-WAVE INTERACTIONS

The evolution of distribution functions due to resonant interaction of three plasma waves (Fig. 4.1c) is described by equations

(4.7), (4.8), etc. Like these wave-kinetic equations, the spontaneous probability $u^{\sigma\sigma'\sigma''}(\underline{k}, \underline{k}', \underline{k}'')$ of the coalescence and decay processes is formally identical to that for the case of a plasma in the absence of a background magnetic field. The effect of the background magnetic field on the probability enters through the nonlinear response tensor and the wave properties.

Standard calculations, for example, substitution of the appropriate form of nonlinear current density (see (2.49)) into (3.12), yield a probability which reads (see (14) of II)

$$u^{\sigma\sigma'\sigma''}(\underline{k}, \underline{k}', \underline{k}'') = 2(2\pi)^5 \hbar \omega^{\sigma} \omega^{\sigma'} \omega^{\sigma''} \left(\frac{W_E}{W_T} \right)^{\sigma} \left(\frac{W_E}{W_T} \right)^{\sigma'} \left(\frac{W_E}{W_T} \right)^{\sigma''} \\ \times |\epsilon^{\sigma\sigma'\sigma''}(\underline{k}', \underline{k}'')|^2 \delta^3(\underline{k} - \underline{k}' - \underline{k}'') \delta(\omega^{\sigma} - \omega^{\sigma'} - \omega^{\sigma''}), \quad (4.77)$$

with

$$\epsilon^{\sigma\sigma'\sigma''}(\underline{k}', \underline{k}'') \equiv e_i^{\sigma*} e_j^{\sigma'} e_l^{\sigma''} \epsilon_{ijl}(\underline{k}', \omega^{\sigma'}; \underline{k}'', \omega^{\sigma''}). \quad (4.78)$$

When there is only a single distribution of waves in mode $\sigma' = \sigma''$, the factor 2 must be dropped from (4.77).

7. FORMATION OF TURBULENCE SPECTRA

Turbulence spectra of plasma waves are formed in many experimental and astrophysical situations. For example, ion-acoustic turbulence has been observed directly in perpendicular shock experiments (Paul, 1971) and it may play an important rôle in collisionless shock waves associated with type II solar radio bursts (Smith, 1972). The suppression of quasilinear relaxation of charged particle streams associated with type III solar radio bursts has been attributed to the formation of a turbulence spectrum of electron plasma waves (Kaplan and Tsytovich, 1968; Smith, 1970).

Plasma waves excited by sources are usually confined to narrow regions of wavevector $\mathbf{k}$ space. When their dissipation is not rapid, energy will be transferred by nonlinear processes to other waves, which are not directly excited by the sources. The balance between the nonlinear inflow and the outflow of energy due to dissipative processes, determines the quasistationary spectra of turbulence (Kropotkin, 1968). Quasistationary spectra of ion-acoustic turbulence driven by a current have been discussed by Kadomtsev (1965, p.68) and Tsytovich (1971), whilst the quasistationary spectra of turbulence in electron plasma waves, arising from the two-stream instability in the context of type III solar radio bursts, have been discussed by Kaplan and Tsytovich (1968) and Smith and Fung (1970). However, in these studies the effects of a background magnetic field have been largely ignored. Although it is out of the scope of this thesis to do the computations required to solve the corresponding problems for a magnetoactive plasma, it is nevertheless desirable to discuss the qualitative effects of a background magnetic field on the nonlinear transfer of energy across the turbulence spectra.

In the references cited, nonlinear scattering by plasma ions determines the spectral transfer of Langmuir waves, which are upper-branch electron plasma waves (see equations (2.90) - (2.93)) in the limit $\Omega_e = 0$. In the limit of a weak background magnetic field such that

$$\omega_p \gg \Omega_e, \quad \text{and} \quad \lambda_i > \lambda_e = (\Delta k_{\perp} V_e / \Omega_e)^2 \gg 1, \quad (4.79)$$

the modified dispersion relation reads

$$\omega^2 = \omega_p^2 + \Omega_e^2 \sin^2 \theta + 3k_{\perp}^2 V_e^2, \quad (4.80)$$

and the averaged probability reads approximately (4.68),

$$w_i^{\ell\ell'}(\underline{k}, \underline{k}') = \frac{(2\pi)^{5/2} Z_i^2 e^4}{m_e^2 \omega_p^2} \frac{|\underline{k} \cdot \underline{k}'|^2}{(1 + Z_i T_e / T_i)^2} \frac{1}{|\Delta \underline{k}| V_i} \exp \left\{ - \left(\frac{3 V_e^2 (k^2 - k'^2)}{2 \sqrt{2} \omega_p |\Delta \underline{k}| V_i} \right)^2 \right\}, \quad (4.81)$$

which is identical to the averaged probability for the case of a plasma without a background magnetic field. All the assumptions made in deriving (4.81), e.g. (4.61) and (4.62), can be justified *a posteriori*. Under the conditions specified by (4.79), the background magnetic field can be neglected in treating spectral transfer of electron plasma waves.

On the other hand, under the conditions:

$$\omega_p \geq \Omega_e, \quad \text{and} \quad (\Delta k_{\perp} V_e / \Omega_e)^2 = \lambda_e < \lambda_i \ll 1, \quad (4.82)$$

the background magnetic field has significant effects on spectral transfer and turbulence formation.

Induced scattering of longitudinal waves by a Maxwellian distribution of ions is described by a wave-kinetic equation (4.3):

$$\frac{dN^{\ell}(\underline{k})}{dt} = -\mu^{\ell}(\underline{k}) N^{\ell}(\underline{k}), \quad (4.83)$$

where

$$\mu^{\ell}(\underline{k}) = \frac{\hbar n_i}{m_i V_i^2} \int \frac{d^3 \underline{k}'}{(2\pi)^3} N^{\ell'}(\underline{k}') (\omega^{\ell} - \omega^{\ell'}) w_i^{\ell\ell'}(\underline{k}, \underline{k}'), \quad (4.84)$$

with $d/dt = \partial/\partial t + \underline{v}_g^{\ell} \cdot (\partial/\partial \underline{r})$ and $w_i^{\ell\ell'}(\underline{k}, \underline{k}')$ determined by (4.67). Under the conditions specified by (4.82), zeroth harmonic scattering dominates and from maximization of (4.84), the condition for most efficient induced scattering reads

$$\omega^{\ell} = \omega^{\ell'} - |\Delta k_{\parallel}| V_i, \quad (4.85)$$

which implies that in the induced scattering $\ell' \rightarrow \ell$, the scattered wave has a decreased frequency. In the case of electron plasma waves in a relatively strong background magnetic field, this means induced

scattering has a tendency to produce anisotropic turbulence spectra along the direction of B_0 .

When (4.85) is satisfied, bearing in mind, $\omega^\ell \approx \omega^{\ell'} \gg |\Delta k_\parallel| V_i$ in a strong magnetic field, equation (4.84) reduces to

$$\mu^{el}(\underline{k}) = -\frac{3\sqrt{2\pi}}{5(1+Z_i T_e/T_i)^2} \frac{\omega_p^4}{\omega^3} \left(\frac{W_E}{W_T}\right)^{el} \times \int \frac{d^3 \underline{k}'}{(2\pi)^3} \frac{W^{el}(\underline{k}')}{n_i m_i V_i^2} \left(\frac{W_E}{W_T}\right)^{el'} \langle |\kappa_i \tau_{ij}^{(e)}(\omega) \kappa_j'|^2 \rangle, \quad (4.86)$$

where $\langle \dots \rangle$ denotes again an average over the unimportant azimuthal angles.

From (4.86), it is seen that induced scattering of electron plasma waves by ions has a characteristic time given in order of magnitude by

$$\tau \sim (W_i/W^{el}) \omega_p^{-1} \gg \omega_p^{-1}, \quad (4.87)$$

where W_i is the kinetic energy density of the plasma ions and W^{el} is the turbulence energy density of the electron plasma waves

$$W^{el} = \int \frac{d^3 \underline{k}}{(2\pi)^3} W^{el}(\underline{k}). \quad (4.88)$$

Comparison of (4.76) and (4.86) shows that the neglect of spontaneous scattering is justified provided

$$W^{el}/W_i \gg \sigma_T c n_e / \omega_p, \quad (4.89)$$

where σ_T is the Thomson cross-section, numerically equal to $6.65 \times 10^{-25} \text{ cm}^2$. Since the factor on the right hand side of (4.89) is of the order of $\sqrt{n_e} \times 10^{-20}$, condition (4.89), together with the weak turbulence assumption $W^{el}/W_i \ll 1$, can be satisfied in most situations of interest.

Bound up intimately with the problem of formation of turbulence spectra of electron plasma waves arising from the two-stream instability

is the question of stabilization of the stream itself. Within certain ranges of plasma parameters, nonlinear spectral transfer of growing waves from the resonance region into the non-resonance region can be so rapid that the instability is suppressed and the stream stabilized (Tsytovich and Shapiro, 1965). Analysis so far has pertained only to plasmas without a background magnetic field.

In the presence of a background magnetic field, a one-dimensional treatment of the problem becomes possible. The conditions under which such a treatment is valid, are discussed here. Analysis of the quasi-linear instability due to an electron stream of velocity u_s along B_0 and thermal spread V_s (see section 5(b) of chapter 3) shows that in both limiting cases $\lambda' \ll 1$ and $\lambda' \gg 1$, with $\lambda' = (k_{\perp} V_s / \Omega_e)^2$, the electron plasma waves grow fastest along B_0 and the maximum growth rates can be approximated by (see (3.78))

$$\gamma^{el} = \frac{3}{10} \left(\frac{\pi}{2} \right)^{\frac{1}{2}} \left(\frac{u_s}{V_s} \right)^2 \frac{n'}{n_e} \omega_p, \quad (4.90)$$

where n' is the number density of the streaming electrons.

From equations (4.81) and (4.86), it is clear that induced nonlinear scattering by ions tends to maintain the anisotropy along B_0 . Hence a one-dimensional treatment for these cases is qualitatively justified. The corresponding absorption coefficient due to induced nonlinear scattering by ions reads

$$\mu^{el} = \frac{3}{10} \left(\frac{\pi}{2} \right)^{\frac{1}{2}} \frac{W^{el}/W_i}{(1 + Z_i T_e / T_i)^2} \omega_p. \quad (4.91)$$

When $\epsilon \equiv \gamma^{el} / \mu^{el} \ll 1$, the speed of spectral transfer of waves from the resonance region can be sufficiently fast to quench the instability.

Taking $W^{el} \approx n' m_e u_s V_s$ and $W_i = W_e = n_e m_e V_e^2$, one has

$$\epsilon = \left(1 + \frac{Z_i T_e}{T_i} \right)^2 \left(\frac{u_s}{V_e} \right) \left(\frac{V_e}{V_s} \right)^3. \quad (4.92)$$

If in addition,

$$\epsilon \ln(\epsilon W^{el}/W_0) \ll 1, \quad (4.93)$$

where $W_0 \approx m_e V_e^2 / \lambda_{De}^3$, the distribution function of the stream changes little for the instability (Tsytovich and Shapiro, 1965).

Summing up, we have written down conditions under which a one-dimensional treatment of the problems of turbulence spectra formation, can be justified. But we note that quantitative solutions are needed particularly for a plasma with a strong background magnetic field.

CHAPTER 5

POLARIZED RADIATION FROM PLASMA TURBULENCE

1. INTRODUCTION

Nonlinear conversion of plasma turbulence into observable radiation is now firmly established in laboratory and astrophysical plasmas; see for example, the monographs cited at the beginning of the last chapter, and also Kaplan and Tsytovich (1969). However, little attention, so far, has been devoted explicitly to the consideration of polarization of radiation arising from such mechanisms. Since polarization yields important and sometimes crucial information concerning the sources of radiation, it is worthwhile considering such questions here.

Polarized radiation can appear from a turbulent plasma even in the absence of a background magnetic field (Tsytovich, 1969). However, such cases must be regarded as marginal, since *ad hoc* assumptions, such as the existence of anisotropic turbulence, etc., have to be made concerning the plasma. In general, we assumed polarized radiation arises from a plasma with a background magnetic field.

To study polarization of radiation in a magnetoactive plasma, the magnetoionic character of the escaping radiation must be taken into account. For the purpose of discussion the name "magnetoactive" will henceforth be restricted to refer to those waves of magnetoionic theory which can escape directly to infinity. These waves are waves in the ordinary mode (O-mode) at angular frequencies $\omega > \omega_p$ and waves in the extraordinary mode (X-mode) at $\omega > \omega_X$, where the cutoff frequency is given by

$$\omega_X = \frac{1}{2} \Omega_e + \frac{1}{2} (4\omega_p^2 + \Omega_e^2)^{1/2}. \quad (5.1)$$

The generation of polarized radiation from microturbulence in a magnetoactive plasma will be studied in this chapter.

2. ELECTRON PLASMA WAVES (II)

For the purposes of this chapter, "electron plasma waves" is restricted to refer to the longitudinal waves (2.90) in the upper frequency range $\max(\omega_p, \Omega_e) \leq \omega \leq (\omega_p^2 + \Omega_e^2)^{1/2}$. To obtain approximate probabilities for the nonlinear conversion of electron plasma waves into magnetoionic waves, the mode σ in probabilities (4.67), (4.68) or (4.77) is identified with one or the other of the magnetoionic waves and the modes σ' and σ'' are identified with the "el" mode, designating electron plasma waves.

Since the phase velocities of electron plasma waves are much less than the velocity of light, we may approximate $|\Delta k_{\parallel}| \approx k' |\cos \theta'|$ in (4.67) and $|\Delta k| \approx k'$ in (4.68). In the limiting of negligibly small ion gyroradii, i.e., $\lambda_i = (\Delta k_{\perp} V_i / \Omega_i)^2 \ll 1$, zeroth harmonic scattering is dominant in (4.67) and the simplified probability for scattering of electron plasma waves into magnetoionic waves by thermal ions, after averaging over the azimuthal angles, reduces to (see II)

$$\begin{aligned} \langle w_i^{\sigma el}(\underline{k}, \underline{k}') \rangle &= \frac{2(2\pi)^{5/2} Z_i^2 e^4}{m_e^2 \omega \omega' (1 + Z_i T_e / T_i)^2} \left(\frac{W_E}{W_T} \right)^{el} \frac{F^{\sigma el}(\omega, \theta, \theta')}{V_i k' |\cos \theta'|} \\ &\times \exp \left\{ - \frac{(\omega - \omega')^2}{2k'^2 V_i^2 |\cos \theta'|^2} \right\}, \quad (5.2) \end{aligned}$$

where ω' and $(W_E/W_T)^{el}$ are functions of θ' , given by (2.90), (2.91) and (2.93), and

$$F^{\sigma\ell}(\omega, \theta, \theta') = \frac{1}{(1+T_\sigma^2) \eta_\sigma} \frac{\partial(\omega \eta_\sigma)}{\partial \omega} \left\{ \frac{1+Y^2}{2(1-Y^2)^2} (1+a_\sigma^2) \sin^2 \theta' + \frac{2Y}{(1-Y^2)^2} a_\sigma \sin^2 \theta' + b_\sigma^2 \cos^2 \theta' \right\}, \quad (5.3)$$

with a_σ and b_σ given by (2.84), and $Y = \Omega_e/\omega$ as usual. Expression (5.3) can be rewritten in terms of T_σ , X , Y and θ by using

$$a_\sigma = \frac{\cos \theta - Y T_\sigma}{T_\sigma - Y \cos \theta}, \quad b_\sigma = \frac{\cos^2 \theta - T_\sigma^2}{\sin \theta (T_\sigma - Y \cos \theta)} \quad (5.4)$$

and

$$\eta_\sigma \frac{\partial}{\partial \omega} (\omega \eta_\sigma) = 1 + \frac{XY \cos \theta T_\sigma}{2(T_\sigma - Y \cos \theta)^2} \left\{ 1 + \frac{(1+X)(1-T_\sigma^2)}{(1-X)(1+T_\sigma^2)} \right\},$$

with $X = \omega_p^2/\omega^2$. In the opposite limit of $\lambda_1 \gg 1$, the probability (4.68) must be used and this leads to a simplified probability of the form of (5.2), with $k' |\cos \theta'|$ replaced by k' .

The probability for the coalescence process $e\ell + e\ell \rightarrow \sigma$ can be found by inserting the appropriate wave properties in (4.77) with $\epsilon_{ij\ell}$ given by equation (F.8). Since $e\ell$ -waves have phase velocities much less than the velocity of light, $|\underline{k}'|, |\underline{k}''| \gg |\underline{k}|$; the requirement $\underline{k} = \underline{k}' + \underline{k}''$ leads to $\underline{k}'' \approx -\underline{k}'$, which reduces the number of independent directions in the problem and therefore results in considerable simplification. The average probability reduces to (see II)

$$\langle u^{\sigma e\ell e\ell}(\underline{k}, \underline{k}', \underline{k}'') \rangle = \frac{(2\pi)^5 \hbar e^2 \omega_p^2}{m_e^2 c^2 \omega'} \left\{ \left(\frac{W_E}{W_T} \right)^{e\ell} \right\}^2 G^{\sigma\ell}(\omega', \theta, \theta') \times \delta^3(\underline{k} - \underline{k}' - \underline{k}'') \delta(\omega - 2\omega'), \quad (5.5)$$

where ω' and $(W_E/W_T)^{e\ell}$ are again given by (2.90) - (2.93), and

$$\begin{aligned}
G^{\sigma\ell}(\omega', \theta, \theta') = & \frac{\eta_\sigma}{(1+T_\sigma^2) \partial(\omega\eta_\sigma)/\partial\omega} \left\{ \frac{\sin^2\theta \sin^4\theta'}{8(1-Y'^2)^4} \{ (3+Y'^2)(a_\sigma+Y')^2 \right. \\
& + (1+3Y'^2)(a_\sigma Y'+1)^2 \} + \frac{\cos^2\theta \sin^2\theta' \cos^2\theta'}{2(1-Y'^2)^2} \{ (a_\sigma+Y')^2 + (1+a_\sigma Y')^2 \} \\
& + \frac{2\sin\theta \cos\theta \sin^2\theta' \cos^2\theta'}{(1-Y'^2)^2} (a_\sigma+Y') b_\sigma + \frac{(1+Y'^2)}{2(1-Y'^2)^2} \\
& \left. \sin^2\theta \sin^2\theta' \cos^2\theta' b_\sigma^2 + \cos^2\theta \cos^4\theta' b_\sigma^2 \right\}, \quad (5.6)
\end{aligned}$$

with $Y' = \Omega_e/\omega'$.

Similar considerations give an average probability describing the coalescence of an electron plasma wave and an ion-acoustic wave into a magnetoionic wave $el + s \rightarrow \sigma$ (see II):

$$\begin{aligned}
\langle u^{\sigma els}(\underline{k}, \underline{k}', \underline{k}'') \rangle = & \frac{(2\pi)^5 \hbar e^2}{2m_e^2} \left(\frac{m_i}{Z_i m_e} \right) \frac{(\omega'')^3}{k'^2 V_e^4} \left(\frac{W_E}{W_T} \right)^{el} \\
& \times F^{\sigma\ell}(\omega', \theta, \theta') \delta^3(\underline{k}-\underline{k}'-\underline{k}'') \delta(\omega-\omega'-\omega''), \quad (5.7)
\end{aligned}$$

where the assumptions $\omega \approx \omega' \gg \omega''$ and $\underline{k}' \approx -\underline{k}''$ have been made and $F^{\sigma\ell}$ is given by (5.3).

Particularly in the context of solar radio astronomy, "plasma emission processes" refers to the nonlinear plasma processes whereby electron plasma waves are converted to escaping radiation at about the plasma frequency (the fundamental) and twice this frequency (the second harmonic). In a magnetoactive plasma, on account of preferential emission of one or other of the magnetoionic modes, the escaping radiation can be expected, in general, to be polarized.

It is possible at this stage to estimate the degrees of polarization for both the fundamental and the second harmonic in the region of emission. The estimated degrees of polarization will coincide with the degrees of polarization actually observed provided the polarization

limiting region is at the source. However, if the polarization limiting region is far removed from the source, the estimated degrees of circular polarization will coincide with those actually observed (even in the presence of Faraday rotation) provided mode-mode coupling and preferential absorption of one mode or the other can be neglected.

(a) The Fundamental

Emission at the fundamental can be 100% polarized in the sense of the O-mode, when emission is at frequencies $\omega < \omega_X$ below the cutoff of the X-mode. This occurs in a plasma with a strong background magnetic field $\Omega_e \approx \omega_p$, provided the wavenumbers k' of the electron plasma waves satisfy the condition $k^2 \lambda_{De}^2 \lesssim 0.6$. This condition is usually satisfied.

In the weak field limit $\omega_p \gg \Omega_e$, equation (5.1) reduces to

$$\omega_X \approx \omega_p + \frac{1}{2} \Omega_e, \quad (5.8)$$

while the frequency of el-waves is given by

$$\omega'(\theta') \approx \omega_p + (\Omega_e^2 / 2\omega_p) \sin^2 \theta' + 3(V_e^2 / 2v_\phi^2) \omega_p, \quad (5.9)$$

where $v_\phi = \omega' / k' \approx \omega_p / k'$ is the phase velocity. From the probability (5.2), emission occurs at frequencies $\omega < \omega_X$ when

$$\omega' \{1 + (V_i / v_\phi) |\cos \theta'|\} \lesssim \omega_X, \quad (5.10)$$

which requires

$$v_\phi \geq \max\{V_e (3\omega_p / \Omega_e)^{1/2}, V_i (2\omega_p / \Omega_e)\}. \quad (5.11)$$

When the condition (5.11) is satisfied the radiation in the region of emission is 100% polarized in the sense of the O-mode. When the opposite of the inequality (5.11) is satisfied, the emission is in general expected to be a mixture of the two modes.

The case of most practical interest is for $e\ell$ -waves propagating nearly along the field lines, i.e., $\sin\theta' \approx 0$. Consider the following limiting cases for emission at a given angle θ , with $Y < 1-X$ (the reverse of inequality (5.11)).

$$(i) \quad \text{For} \quad \frac{1}{4} Y^2 \sin^4 \theta \ll (1-X)^2 \cos^2 \theta ,$$

the degree of circular polarization in the sense of the O-mode can be estimated by (see II)

$$r_c = \{F^{O\ell}(\omega, \theta, \theta') - F^{X\ell}(\omega, \theta, \theta')\} / \{F^{O\ell}(\omega, \theta, \theta') + F^{X\ell}(\omega, \theta, \theta')\} , \quad (5.12)$$

where $F^{\sigma\ell}$ is given by (5.3), with $\sin\theta' = 0$. Since

$$T_O \approx -\cos\theta / |\cos\theta| , \quad K_O \approx XY \sin\theta (1-Y|\cos\theta|) / (1-X) , \quad (5.13)$$

$$T_X \approx \cos\theta / |\cos\theta| , \quad K_X \approx XY \sin\theta (1+Y|\cos\theta|) / (1-X) ,$$

it follows that $F^{\sigma\ell} \approx \frac{1}{2} \sin^2 \theta$ and

$$r_c = 2Y(1-X) |\cos\theta| / \{Y^2 + (1-X)^2 \cos^2 \theta\} , \quad (5.14)$$

where $1-X \ll 1$ and $Y \ll 1$ have been assumed. It is noted here that at $|\cos\theta| = Y/(1-X)$ the radiation is fully polarized in the sense of the O-mode.

$$(ii) \quad \text{For} \quad \frac{1}{4} Y^2 \sin^4 \theta \gg (1-X)^2 \cos^2 \theta ,$$

$$T_O \approx -Y \sin^2 \theta / (1-X) \cos \theta , \quad K_O \approx XY \sin \theta / (1-X) , \quad (5.15)$$

$$T_X \approx (1-X) \cos \theta / Y \sin^2 \theta , \quad K_X \approx XY / (1-X-Y^2) ,$$

and it follows that

$$F^{O\ell}(\omega, \theta \approx \frac{1}{2}\pi, \theta' \approx 0) \approx 1 \gg F^{X\ell}(\omega, \theta \approx \frac{1}{2}\pi, \theta' \approx 0) . \quad (5.16)$$

Thus for emission nearly perpendicular to the field lines the power is

almost all in the O-mode even for $Y \ll 1-X$, and the radiation at the source is almost 100% linearly polarized in the sense of O-mode waves.

To illustrate the range of the limiting cases, the angle defined by $\frac{1}{4} Y^2 \sin^4 \theta = (1-X)^2 \cos^2 \theta$ is $\theta = 65^\circ$ for $Y = 1-X$ and is $\theta = 77^\circ$ for $Y = \frac{1}{2}(1-X)$.

(b) The Second Harmonic

For the second harmonic emission, $X \approx \frac{1}{4}$ and the condition $\frac{1}{4} Y^2 \sin^4 \theta \ll (1-X)^2 \cos^2 \theta$, which is equivalent to $Y' \sin^2 \theta \ll 3|\cos \theta|$, with $Y' = \Omega_e / \omega_p$, is satisfied for a wide range of angles θ . For example, the angle defined by $Y' \sin^2 \theta = 3|\cos \theta|$ is $\theta = 72.5^\circ$ for $Y' = 1$, and it is $\theta = 81^\circ$ for $Y' = 2$. It is reasonable to approximate the properties of the magnetoionic waves by

$$T_O = -\frac{\cos \theta}{|\cos \theta|} - \frac{Y' \sin^2 \theta}{3 \cos \theta}, \quad T_X = \frac{\cos \theta}{|\cos \theta|} - \frac{Y' \sin^2 \theta}{3 \cos \theta}, \quad (5.17)$$

which have been expanded to first order in $Y' \sin^2 \theta / 3|\cos \theta| \ll 1$.

The degree of circular polarization of the second harmonic, in the sense of the O-mode, can be estimated by (see II)

$$r_c = \{G^{O\ell}(\omega', \theta, \theta') - G^{X\ell}(\omega', \theta, \theta')\} / \{G^{O\ell}(\omega', \theta, \theta') + G^{X\ell}(\omega', \theta, \theta')\}, \quad (5.18)$$

where $G^{\sigma\ell}$ is given by (5.6).

For the case $\sin \theta' = 0$,

$$G^{\sigma\ell}(\omega', \theta, \theta') = \eta_\sigma^2 b_\sigma^2 \cos^2 \theta / (1 + T_\sigma^2) \eta_\sigma \partial(\omega \eta_\sigma) / \partial \omega. \quad (5.19)$$

Application of (5.17) gives an expression of $G^{\sigma\ell}$ to first order in $Y' \sin^2 \theta / 3|\cos \theta| \ll 1$:

$$G^{\sigma\ell}(\omega', \theta, \theta' = 0) = \frac{1}{2} \sin^2 \theta \cos^2 \theta (1+a) \left(\frac{1 \pm 2a/\sin^2 \theta}{1 \pm b} \right)^2$$

$$\left(1 - \frac{1 \pm b}{4(1 \pm b)} \right) / \left(1 \mp \frac{Y' |\cos \theta| (1 + 5a/3)}{16(1 \pm b)^2} \right), \quad (5.20)$$

where upper (lower) signs refer to 0 (X) mode waves and

$$a \equiv a(Y', \theta) \equiv \frac{Y' \sin^2 \theta}{3 |\cos \theta|}, \quad b \equiv b(Y', \theta) \equiv \frac{Y' (2 + \cos^2 \theta)}{6 |\cos \theta|}. \quad (5.21)$$

On the explicit assumption that $Y' = \Omega_e / \omega_p \ll 1$, (5.20) reduces to

$$G^{\sigma\ell}(\omega', \theta, \theta' = 0) = \frac{3}{8} \sin^2 \theta \cos^2 \theta \left\{ 1 \pm \frac{(16 + 11 \cos^2 \theta)}{48 |\cos \theta|} \frac{\Omega_e}{\omega_p} \right\}, \quad (5.22)$$

from which it is found, by (5.18), that

$$r_c \approx \frac{(16 + 11 \cos^2 \theta)}{48 |\cos \theta|} \frac{\Omega_e}{\omega_p} \quad (\sin \theta' = 0). \quad (5.23)$$

At the maximum power, $\cos^2 \theta = \sin^2 \theta = \frac{1}{2}$, and

$$r_c \approx 0.63 \frac{\Omega_e}{\omega_p} \quad (\theta' = 0, \theta = \frac{\pi}{4}). \quad (5.24)$$

At a quarter of the maximum power, $\theta = 75^\circ$, and

$$r_c = 1.35 \frac{\Omega_e}{\omega_p} \quad (\theta' = 0, \theta = \frac{5\pi}{12}). \quad (5.25)$$

Thus the second harmonic is weakly polarized in the sense of the 0-mode, in a weak background magnetic field $\Omega_e \ll \omega_p$.

However, in a relatively strong background magnetic field, the second harmonic can be significantly polarized. Take for example, $Y' = 1/5$, $\cos \theta = 1/2$, which corresponds roughly to three quarters maximum power, one finds $Y' \sin^2 \theta / 3 \cos \theta = 0.1 \ll 1$ and direct calculation from (5.20) and (5.18) gives $r_c \approx 26\%$. Again $Y' = 3\sqrt{2}/10 \approx 0.42$ and $\cos^2 \theta = 0.5$ imply $Y' \sin^2 \theta / 3 \cos \theta = 0.1 \ll 1$ and $r_c \approx 33\%$.

(c) Conclusions

In conclusion we summarize the results obtained in this section.

General formulae have been derived for estimating the degrees of circular polarization at the source for the fundamental and the second harmonic radiation arising from the "plasma emission processes".

In a strong background magnetic field ($\Omega_e \approx \omega_p$), the fundamental is 100% polarized in the sense of the 0-mode provided the wavenumbers k' of the electron plasma waves satisfy the condition $k^2 \lambda_{De}^2 \leq 0.6$.

For the case of electron plasma waves propagating nearly along the lines of the background magnetic field, the radiation at the fundamental and the radiation at the second harmonic have net degrees of circular polarization at the source in the sense of the 0-mode. In a weak background magnetic field ($\omega_p \gg \Omega_e$) the second harmonic is always weakly polarized, whilst the fundamental can be strongly polarized in the sense of the 0-mode. Under the condition (5.11), viz.

$v_\phi \geq \max\{V_e (3\omega_p/\Omega_e)^{1/2}, V_i (2\omega_p/\Omega_e)\}$, the fundamental radiation is 100% polarized in the sense of the 0-mode.

3. ELECTRON CYCLOTRON WAVES

Despite numerous observations of radiation at electron cyclotron harmonics from laboratory plasmas (see Crawford, 1968), a detailed theory of radiation from electrostatic electron cyclotron waves does not seem to have appeared in the literature. One reason for this might be that in this case, it is important to take into account the effects of the background magnetic field on the dielectric properties of the medium, on the waves as well as on the conversion processes. Such effects are taken into account in a theory presented here.

In a plasma with a strong background magnetic field ($\Omega_e > \omega_p$), radiation can arise in principle from parallel electron cyclotron waves at Ω_e and $2\Omega_e$, and from perpendicular electron cyclotron waves at higher harmonics. Radiation at the fundamental in this case can be identified as radiation arising from electron plasma waves in the frequency range $\Omega_e \leq \omega \leq (\Omega_e^2 + \omega_p^2)^{1/2}$.

In a plasma with a weak background magnetic field ($\omega_p \gg \Omega_e$) only perpendicularly propagating electron cyclotron waves at high harmonic numbers $n > \omega_p / \Omega_e \gg 1$ can lead to observable radiation. For the purposes of discussion, we shall refer to electron cyclotron waves propagating perpendicularly to the background magnetic field as Bernstein waves. From the results of chapter 3, it can be shown that for given plasma parameters, the power radiated into Bernstein waves by non-relativistic electrons decreases with increases in harmonic number n . Therefore, when the microturbulence in Bernstein waves is generated coherently, e.g., by a microinstability, one expects the energy density to be highest for the generated Bernstein waves in the lowest harmonic number. Thus most intense radiation from astrophysical plasmas containing such microturbulence is expected to arise from Bernstein waves at the lowest harmonic n satisfying $n > \omega_p / \Omega_e$ at the source.

(a) Case $\omega_p \gg \Omega_e$

Consider Bernstein waves at frequencies $\omega \approx s\Omega_e$ with

$$s \geq \omega_p / \Omega_e \gg s-1 \gg 1. \quad (5.26)$$

From equation (2.102), it can be shown that the ratio of the electrical to total energy density $(W_E/W_T)^c$ for Bernstein waves at frequencies $\omega \approx s\Omega_e$ would be very small unless the condition $\Omega_e \geq s \gg 1$ is satisfied.

Assuming that this condition is fulfilled we find

$\varepsilon_0^{\ell}(\underline{k}, s\Omega_e) = 1 - \{\omega_p^2 / (s^2 - 1)\Omega_e^2\} \{\zeta_1(\lambda_e) / \lambda_e\} \approx 1 - \zeta_1(\lambda_e) / \lambda_e \approx 1$. Hence the dispersion relation (2.101) can be written as

$$\omega = s\Omega_e + \{s\zeta_s(\lambda_e) / \lambda_e\} (\omega_p^2 / \Omega_e^2) \Omega_e, \quad (5.27)$$

where the last term determines the natural bandwidth of Bernstein waves at the s -th harmonic. From the theory of Bessel functions

(Watson, 1944) it can be established (i) $s\zeta_s(\lambda_e) / \lambda_e = \zeta_{s-1}(\lambda_e) + \zeta_{s+1}(\lambda_e)$, (ii) $\zeta_{s-1}(\lambda_e) > \zeta_{s+1}(\lambda_e)$, (iii) $\zeta_s(\lambda_e)$ maximizes at $\lambda_e \approx s^2$ for fixed s and (iv) $\zeta_s(\lambda_e) \approx (2\pi\lambda_e)^{-1/2}$ for $\lambda_e \gg s \gg 1$. From these relationships, the maximum natural bandwidth of the waves can be estimated to be

$$(\Delta\omega)_{\max} \approx \{s/\sqrt{2} (s-1)^3\} (\omega_p^2 / \Omega_e^2) \Omega_e. \quad (5.28)$$

However in general the range of values of λ_e , which determines the actual bandwidth of the Bernstein waves, is itself determined by the excitation processes in question and the actual bandwidth could be much narrower than $(\Delta\omega)_{\max}$.

Since $\lambda_i = (\Delta k_{\perp} V_i / \Omega_i)^2 \approx (k' V_i / V_i)^2 \gg s(m_i / m_e) \gg 1$, where k' is the wavenumber of Bernstein waves in the s -th harmonic, the probability for nonlinear scattering of the Bernstein waves by thermal ions is given by (4.68). On noting $k' \gg k$, where k is the wavenumber of magnetoionic waves, the probability averaged over azimuthal angles reads

$$\begin{aligned} \langle w_i^{\sigma c}(\underline{k}, \underline{k}') \rangle &= \frac{2(2\pi)^{5/2} Z_i^2 e^4}{m_e^2 \omega \omega' (1 + Z_i T_e / T_i)^2} \left(\frac{W_E}{W_T} \right)^c \frac{F^{\sigma \ell}(\omega, \theta, \theta' = \pi/2)}{k' V_i} \\ &\quad \times \exp \left\{ - \frac{(\omega - \omega')^2}{2k'^2 V_i^2} \right\}, \end{aligned} \quad (5.29)$$

where primed (unprimed) quantities refer to Bernstein (magnetoionic) waves, and $F^{\sigma \ell}$ is determined by (5.3).

From the above probability, it is seen that emission occurs at frequencies $\omega < \omega_X$, when

$$\omega' + k'V_i \leq \omega_X, \quad (5.30)$$

which, on application of equation (5.8), reads

$$s - \omega_p/\Omega_e + (\Delta\omega + k'V_i)/\Omega_e \leq \frac{1}{2}, \quad (5.31)$$

where $\Delta\omega$ is the bandwidth of the Bernstein waves determined by the excitation processes, and $k' \geq \sqrt{s} \Omega_e/V_e$. For example, in the case where $\omega_p \approx s\Omega_e$ and $k' \approx \sqrt{s} \Omega_e/V_e$, equation (5.31) can be satisfied in an isothermal plasma ($T_e = T_i$) provided $\Delta\omega \ll \Omega_e$ and $\sqrt{s} \leq 20$. When equation (5.31) is satisfied, emission is purely in 0-mode waves.

On the other hand, when the plasma parameters are such that ω_p lies midway between $(s-1)\Omega_e$ and $s\Omega_e$, i.e., $s - \omega_p/\Omega_e \approx \frac{1}{2}$, equation (5.31) is always violated, and the emission consists of a mixture of 0-mode and X-mode waves. In this case, and in others when (5.31) is violated, the degree of circular polarization of the radiation at the source in the sense of one mode or the other, can again be estimated by (5.12) or an analogous expression.

From (5.3), one finds

$$\begin{aligned} F^{\sigma\ell}(\omega, \theta, \theta' = \pi/2) &= \frac{\{(1+Y^2)(1+a_\sigma^2) + 4a_\sigma Y\}}{2(1-Y^2)^2(1+T_\sigma^2) \eta_\sigma \partial(\omega\eta_\sigma)/\partial\omega} \\ &= \frac{\cos^2\theta + T_\sigma^2}{2(T_\sigma - Y\cos\theta)^2(1+T_\sigma^2) \eta_\sigma \partial(\omega\eta_\sigma)/\partial\omega}, \end{aligned} \quad (5.32)$$

where $Y = \Omega_e/\omega \approx s^{-1}$ and the quantities a_σ , T_σ , etc., are given by (2.84) and (5.4).

Consider the following limiting cases for emission at a given angle θ , when the reverse of inequality (5.31) obtains.

(i) For $\frac{1}{4} Y^2 \sin^4 \theta \ll (1-X)^2 \cos^2 \theta$,

$$T_0 \approx -\frac{\cos \theta}{|\cos \theta|} - \frac{Y \sin^2 \theta}{2(1-X) \cos \theta}, \quad T_X \approx \frac{\cos \theta}{|\cos \theta|} - \frac{Y \sin^2 \theta}{2(1-X) \cos \theta}, \quad (5.33)$$

and it follows that

$$F^{\sigma\ell}(\omega, \theta, \theta' = \pi/2) = \frac{(1+\cos^2 \theta)}{4} \left\{ 1 - \frac{Y \sin^2 \theta}{2(1-X)^2} \mp \frac{Y}{2} \left[\frac{\sin^2 \theta (1+3\cos^2 \theta)}{|\cos \theta| (1-X) (1+\cos^2 \theta)} + 3|\cos \theta| \right] \right\}. \quad (5.34)$$

The degree of circular polarization in the sense of the X-mode reads

$$r_c = \frac{Y}{2} \left[\frac{\sin^2 \theta (1+3\cos^2 \theta)}{|\cos \theta| (1-X) (1+\cos^2 \theta)} + 3|\cos \theta| \right] / \left[1 - \frac{Y \sin^2 \theta}{2(1-X)^2} \right]. \quad (5.35)$$

The assumptions $1-X \ll 1$ and $Y \ll 1$ have been made in deriving (5.34) and (5.35). At the maximum power, $\cos^2 \theta = 1$, and $r_c = 3Y/2$, in the sense of the X-mode.

(ii) For $\frac{1}{4} Y^2 \sin^4 \theta \gg (1-X)^2 \cos^2 \theta$,

$$T_0 \approx -Y \sin^2 \theta / (1-X) \cos \theta, \quad T_X \approx (1-X) \cos \theta / Y \sin^2 \theta, \quad (5.36)$$

and it follows that

$$F^{X\ell}(\omega, \theta \approx \pi/2, \theta' = \pi/2) \approx \frac{1}{2} \gg F^{O\ell}(\omega, \theta \approx \pi/2, \theta' = \pi/2). \quad (5.37)$$

Thus for emission nearly perpendicular to the direction of the background magnetic field, the power is almost all in the X-mode provided the reverse of inequality (5.31) is satisfied. The radiation at the source is almost 100% linearly polarized in the sense of the X-mode waves.

Consider next, radiation from Bernstein waves at frequencies $\omega \approx n\Omega_e$ with $n > s$, under the conditions specified by equation (5.26). In this case, the radiation from the nonlinear scattering of these Bernstein waves by thermal ions always consists of a mixture of O-mode

and X-mode waves, and the degrees of polarization can be estimated by the formulae presented above.

Coalescence of two Bernstein waves at $\omega \approx s\Omega_e$ can give rise to radiation at $\omega \approx 2s\Omega_e$. The degree of circular polarization at the sources can again be estimated by (5.18) or by analogous expressions. In this case, we find

$$G^{\sigma\ell}(\omega', \theta, \theta'=\pi/2) = \frac{\eta_\sigma^2 \sin^2 \theta \{ (3+Y'^2)(a_\sigma+Y')^2 + (1+3Y'^2)(a_\sigma Y'+1)^2 \}}{8(1-Y'^2)^4(1+T_\sigma^2) \eta_\sigma \partial(\omega\eta_\sigma)/\partial\omega}, \quad (5.38)$$

from which it is seen that at maximum power $\sin^2 \theta = 1$,

$$G^{X\ell}(\omega', \theta=\pi/2, \theta'=\pi/2) \approx \frac{3}{32} >> G^{0\ell}(\omega', \theta=\pi/2, \theta'=\pi/2), \quad (5.39)$$

on the assumption $Y' = \Omega_e/s\Omega_e = s^{-1} \ll 1$.

However, since we have $X \sim 1/4$ and $Y \approx 1/2s \ll 1$, emission is also possible for a wide range of angles θ satisfying the condition $\frac{1}{4} Y^2 \sin^4 \theta \ll (1-X)^2 \cos^2 \theta$. Under this condition, equation (5.38) together with (5.4) and various wave properties gives, to first order in $Y' \ll 1$,

$$G^{\sigma\ell}(\omega', \theta, \theta'=\pi/2) = \frac{\sin^2 \theta}{16} (1-X)(1+3\cos^2 \theta) \{1 \mp Y' \chi(\theta, X)\}, \quad (5.40)$$

where the upper (lower) sign refers to O (X) mode waves and

$$\chi(\theta, X) \equiv \frac{1 + \cos^2 \theta (28 - 23X + X^2) + \cos^4 \theta (3 - 21X + 3X^2)}{4 |\cos \theta| (1-X)(1+3\cos^2 \theta)}. \quad (5.41)$$

The degree of circular polarization in the sense of the X-mode is then given by

$$r_c \approx s^{-1} \chi(\theta, X) \quad (\theta' \approx \pi/2). \quad (5.42)$$

As a rough estimate, take $X = 1/4$, we find

$$\chi(\theta, X=1/4) = \frac{1 + 357 \cos^2 \theta - 33 \cos^4 \theta}{48 |\cos \theta| (1+3\cos^2 \theta)}. \quad (5.43)$$

Thus for $\theta \approx \pi/3$, one finds $r_c \approx 2.1/s$ and for $\theta \approx \pi/6$, $r_c \approx 1.85/s$, with radiation polarized in the sense of the X-mode.

(b) Case $\Omega_e \geq \omega_p$

In the case of a plasma with $\Omega_e > \omega_p$, one must be cautioned that microinstabilities at electron cyclotron harmonics in general have reduced growth rates and in some cases, particularly when $\Omega_e \gg \omega_p$, the instabilities are suppressed altogether. For example, from (3.78) it can be shown that in the two-stream instability, parallel electron cyclotron waves at $\omega \approx \Omega_e$ have a growth rate reduced by a factor ω_p/Ω_e compared with the corresponding growth rate of electron plasma waves in a plasma with $\omega_p > \Omega_e$. Brambilla (1970), for example, examined density thresholds for the onset of anisotropy and loss-cone instabilities in hot electron plasmas and found that some instabilities are suppressed in hot electron plasmas.

Suppose parallel electron cyclotron waves at $\omega \approx \Omega_e$ have been excited in a plasma with $\Omega_e > \omega_p$ and $\Omega_e \approx \omega_p$. Since the cutoff frequency X-mode waves is given by $\omega_X = 1.6 \Omega_e$, from an inequality exactly analogous to (5.11), we find that radiation at $\omega \approx \Omega_e$ from nonlinear scattering of these waves by thermal ions consists purely of 0-mode waves provided $k'^2 v_e^2 / \Omega_e^2 \approx k'^2 \lambda_{De}^2 \leq 0.4$, i.e., the phase velocities v_ϕ of the electron cyclotron waves satisfy the condition $v_\phi > 1.6 V_e$. Similarly, in a plasma with a strong background magnetic field ($\Omega_e \gg \omega_p$), the cutoff frequency of X-mode waves reads $\omega_X \approx \Omega_e + \omega_p^2/\Omega_e$, and the radiation at $\omega \approx \Omega_e$ is purely in 0-mode waves provided $k'^2 \lambda_{De}^2 \leq 0.67$ or the phase velocities v_ϕ of the electron cyclotron waves satisfy the condition $v_\phi \geq 1.24 V_e (\Omega_e/\omega_p)$. When the radiation consists of a mixture of 0-mode and X-mode waves, the degree of circular polarization can again be estimated by using equations (5.12), (5.32), etc.

Radiation at the second or higher harmonics can occur by the coalescence of electron cyclotron waves at frequencies $\omega \approx \Omega_e$ or by non-linear (Rayleigh) scattering of electron cyclotron waves at higher harmonics. In any case, the radiation in general consists of a mixture of O-mode and X-mode waves. Procedures for estimating the degrees of polarization follow exactly the method presented above.

(c) Conclusions

On recapitulation, a theory of radiation arising from the non-linear conversion of electrostatic electron cyclotron waves has been presented and general formulae for estimating the degrees of polarization of the radiation at the source, at harmonics of the electron gyrofrequency, i.e., at $s\Omega_e$, with $s = 1, 2, \dots$ have been derived.

In general, when the radiation at $\omega = s\Omega_e$ ($s \geq 2$) satisfies the condition $s\Omega_e > \omega_X$, where ω_X is the cutoff frequency of the X-mode, it has a net degree of polarization at the source in the sense of the X-mode. The actual degree of polarization depends on the ratio ω_p/Ω_e , the harmonic number s and the angle of emission θ .

Exceptions to this rule occurs when the condition $s(s-1) < \omega_p^2/\Omega_e^2 < s^2$ is satisfied for the radiation at $\omega \approx s\Omega_e$. In this case the radiation is completely polarized in the sense of the O-mode. This could happen under the following situations:

(i) $\omega_p \geq \Omega_e$: for at most one harmonic $s \geq 2$ (sometimes none, e.g. $\omega_p/\Omega_e = \sqrt{5}$).

(ii) $\Omega_e \geq \omega_p$: from parallel electron cyclotron waves provided $k'^2 \lambda_{De}^2 \leq 0.4$ (i.e. $v_\phi \geq 1.6 V_e$).

(iii) $\Omega_e \gg \omega_p$: from parallel electron cyclotron waves provided $k'^2 \lambda_{De}^2 \leq 0.67$ (i.e. $v_\phi \geq 1.24 V_e \Omega_e/\omega_p$).

The wavenumber and phase velocity of the electron cyclotron waves have been denoted by k' and v_ϕ respectively.

4. AMPLIFICATION OF RADIATION

Under certain circumstances, radiation from nonlinear Rayleigh scattering of turbulent plasma waves can be amplified by induced processes. In these cases, the absorption coefficient is negative and it is appropriate to speak of the "maser" effect. Such effect is required (Smith, 1970) to explain the brightness temperatures of some radiation at the fundamental from the solar corona.

Radiation into magnetoionic waves by induced Rayleigh scattering of longitudinal plasma waves is described by a wave-kinetic equation (4.3):

$$\frac{dN^\sigma(\underline{k})}{dt} = -\mu^\sigma(\underline{k}) N^\sigma(\underline{k}) , \quad (5.44)$$

with

$$\mu^\sigma(\underline{k}) = \frac{\hbar n_i}{m_i v_i^2} \int \frac{d^3 \underline{k}'}{(2\pi)^3} N^{\ell'}(\underline{k}') (\omega^\sigma - \omega^{\ell'}) w_i^{\sigma \ell'}(\underline{k}, \underline{k}') , \quad (5.45)$$

where $w_i^{\sigma \ell'}(\underline{k}, \underline{k}')$ is determined by either (4.67) or (4.68). When $\lambda_i = (\Delta k_\perp v_i / \Omega_i)^2 \ll 1$, $w_i^{\sigma \ell'}(\underline{k}, \underline{k}')$ is given by (4.67) with contribution from $s=0$ dominating, and the amplification is most effective under the condition:

$$\omega^\sigma = \omega^{\ell'} - |\Delta k_\parallel| v_i . \quad (5.46)$$

On the other hand, when $\lambda_i \gg 1$, $w_i^{\sigma \sigma'}(\underline{k}, \underline{k}')$ is given by (4.68) and the amplification is most effective when

$$\omega^\sigma = \omega^{\ell'} - |\Delta \underline{k}| v_i . \quad (5.47)$$

In either case, when the amplification process is most efficient, the absorption coefficients read

$$\mu^\sigma(\underline{k}) = -\frac{3\sqrt{2\pi}}{5(1+Z_i T_e/T_i)^{3/2}} \frac{\omega_p^4}{\omega^3} \int \frac{d^2 k'}{(2\pi)^3} \frac{W^{\ell'}(\underline{k}')}{n_i m_i V_i^2} \left(\frac{W_E}{W_T}\right)^{\ell'} F^{\sigma\ell'}(\omega, \theta, \theta') , \quad (5.48)$$

where $F^{\sigma\ell'}(\omega, \theta, \theta')$ is given by (5.3). Note that if O-mode and X-mode waves are uncoupled in the amplification region, the differences in the absorption coefficients can give rise to a high degree of polarization in the sense of one mode or the other.

PART II

APPLICATIONS

"Sic itur ad astra"

— Virgil, 30 B.C.

CHAPTER 6

TYPE I SOLAR RADIO BURSTS AND
ASSOCIATED NOISE STORM RADIATION1. INTRODUCTION

Radio emission from the sun has been classified into three distinct components (Kundu, 1965): (1) Quiet sun component, which is simply thermal emission by bremsstrahlung from the solar atmosphere; (2) Slowly varying component, which is also thermal emission but from high density regions called condensations; (3) Non-thermal component, which can consist of radio bursts of numerous spectral types. Studies of the thermal components contributed usefully to our understanding of the large-scale structure of the chromosphere and corona. But it is the non-thermal component which has stimulated new developments in our understanding of generation, propagation and transformation of waves in plasmas.

Radio bursts are generally associated with solar flares, though not always (see e.g., Martres *et al.*, 1972), and they originate between the lower chromosphere (millimetre and centimetre waves) and the outer corona to heights of several solar radii (metre and decametre waves). Their observed source brightness temperature can reach as high as 10^{12} °K, and their suggested emission mechanisms are enhanced bremsstrahlung (Tidman and Dupree, 1965), cyclotron-synchrotron radiation and plasma emission processes. Burst radiation is characterized by its intensity and polarization as functions of position, time, and frequency. On the basis of these spectral and temporal characteristics,

the bursts have been classified into various types:

Bursts on centimetre wavelengths have been classified (Kundu, 1965) into three distinct types: (1) Impulsive bursts; (2) Post-burst; and (3) Bursts exhibiting gradual rise and fall of intensity. Bursts on decimetre wavelengths consist of a variety of fast drifting elements superimposed on a background continuum. There is a great variety of bursts on metre and decametre wavelengths. Apart from rarer events such as drift pair bursts, Hook bursts, U bursts and spike bursts, bursts on metre and decametre wavelengths have been classified (Wild and McCready, 1950; Wild *et al.*, 1959) as types I, II, III, IV, and V, on the basis of their spectral characteristics. Examples of the principal spectral types of solar radio bursts on metre wavelengths, taken with the Dapto Solar Radio Spectrograph, are shown in Fig. 6.1. Their principal characteristics are summarized in Table 6.1.

Whilst considerable progress has been made in our understanding of type III, type II and moving type IV bursts during the last decade (see Wild and Smerd, 1972), the same cannot be said for type I bursts. In view of the fact that type I bursts and associated noise storms are the most common events on metre wave bands, being present about one tenth of the time during periods of the disturbed sun (Kundu, 1965), the noise storm phenomenon presents to us a theoretical challenge. The complexity of the phenomenon can perhaps be attributed to the significant rôle played by the coronal magnetic field, since type I bursts are strongly associated with the appearance of sunspots and with optical activity in the photosphere and the chromosphere. Evidences for this are the strong circular polarization and the high degree of directivity of type I bursts (see section 4). The emission mechanisms in a magnetoactive plasma, considered in the first part of this thesis,

The main figure shows records taken with the Dapto Solar Radio Spectrograph (near Sydney, Australia). The records chosen show typical examples of the six main types of events observable at meter wavelengths. Frequency is measured vertically and time horizontally, the intensity is given by the shade: intense bursts appear white, weak ones grey, and the background is black. The bright lines which are observed at a constant frequency are interference from radio transmitters.

The bottom figure shows in the same form, an idealization of the most complete sequence of events which can follow a major flare. Other details of the particular events are as follows:

Type I (22-12-59 record starts 0210 U.T.). This is a typical "noise storm" which may last for hours. This particular event occurred simultaneously with a large number of type III bursts (see below) which should not be confused with the type I.

Type II (30-11-59 record starts 0251 U.T.) This is also called a slow-drift burst. The direction of drift is always the same and the rate of drift does not vary greatly. Note, particularly near the end, the two pairs of bands, one at twice the frequency of the other.

Type III (7-10-60 record starts 0547 U.T.) These are also called fast-drift bursts. Each of the individual, almost vertical lines is a separate burst. The direction of drift is the same as for type II bursts.

Type IV (11-11-60 record starts 0408 U.T.) The example shows a form of continuum radiation (so called for its slow variation with frequency and time) which commonly follows type II bursts. It is sometimes called "type IV," but the term has been loosely applied to various forms of continuum.

Type V (4-2-60 record starts 0422 U.T.) This is a continuum event, of much shorter duration, which follows immediately after a group of type III bursts.

Drifting pairs (D.P.) (12-2-60 record starts 0422 U.T.) A *drifting pair* is a pair of short-duration, narrow-bandwidth bursts, very similar to one another, which occur separated in time by about two seconds. Both parts drift rapidly in frequency, either from high to low frequencies ("forward-drift pairs") or from low to high ("reverse-drift pairs"). This record shows a portion of a particularly dense form of drifting pairs and includes drifts of both senses.

(Recorded with the Dapto Radio Spectrograph of the Division of Radiophysics C. S. I. R. O., University Grounds, Sydney, Australia. Courtesy, Paul Wild.)

Fig. 6.1: Examples of principal spectral types of solar radio bursts.
(After L.H. Aller, 1963: The Atmosphere of the Sun and Stars,
New York, Ronald Press.)

Type	IV			I Noise Storm		III	V	II
	mA	mB		Continuum	Burst			
Max. Flux $10^{-22} \text{ Wm}^{-2} \text{ Hz}^{-1}$	$< 10^5$			$\leq 10^4$	$\geq 10^4$		$< 10^5$	
Circular Polarization	weak			strong		partial		very weak
Linear Polarization	$< 10\%$			?	very weak	observed?		very weak
Mode	extraordinary			ordinary		mixed		
Bandwidth MHz	~ 200			≤ 200	~ 4	50-100		~ 5
Duration	10 min-hours		hours-days		1 sec (chains 5 sec)	10 sec (groups 1 min)	1 min	5-30 min
Frequency Drift MHz sec^{-1}	sometimes similar Type II		none		1-20 (chains 0.16-0.25)	~ 20	(following Type III)	< 1
Height of Source $R_\odot$	0.5-5.0		0.2-1.0		0.2-2.0	≥ 0.2	0.2-2.0	0.2-1.5
Diameter min arc	6-12	2-6	1-10		1-6	3-12		6-12
Cone of Emission	wide		narrow			wide		
Emission Mechanism	synchrotron	?	?		plasma waves		?	plasma waves
Exciting Agency	electrons trapped in plasmoid		magnetic field structures		electron stream? MHD pulse?	electron stream vel $c/3$	?	ascending shock vel 10^3 km sec^{-1}
Associated Phenomena	flares and proton streams		sometimes flares			often Type I or flares	Type III	Type IV

Table 6.1: Principal characteristics of solar radio bursts on metre and decametre wavelengths. (Wild *et al.*, 1963; Kundu, 1965; Krüger, 1972; Wild and Smerd, 1972; and references in the text.)

constitute a theoretical basis for the understanding of type I bursts.

It is relevant to study the physical conditions in the solar atmosphere above active regions, where noise sources are situated. Thus a brief review is given of the coronal conditions above active regions, which have been deduced from observations. This is followed by a summary of the most important observations. Succeeding sections are given to a critical appraisal of existing theories and construction of a new theoretical model.

2. SOLAR CORONA (Newkirk, 1967)

As eclipse photographs might suggest, the solar corona is largely inhomogeneous, consisting of structures of various shapes and sizes, with enhanced densities. Perhaps the most conspicuous of these are polar plumes, narrow rays and coronal streamers.

Polar plumes and narrow rays are essentially beams with enhanced brightness and density, protruding approximately radially outwards from the chromosphere. Polar plumes are generally associated with bright polar faculae on the solar surface and their densities are enhanced several times above that of the background corona (Newkirk, 1967).

Coronal streamers are larger structures which are also brighter and denser than the background corona and which extend beyond 0.5 - 1.0 solar radii. Two different types of coronal streamers can be distinguished (Mustel, 1962, 1963): one is the helmet streamer associated with quiescent prominences and the other is the active streamer associated with the active regions. The helmet streamers, whose main structural features at low heights are arches of quiescent prominences, are probably not important for the discussion of noise storm phenomena.

Active streamers are associated with more complex structures above the active regions. The diffuse condensation which surrounds an active region but which does not extend beyond $0.3 R_{\odot}$ above the limb, is often simply called "enhancement", and it seems to merge smoothly into the background corona. The "permanent" and "sporadic condensations" are smaller-sized density enhancements and they include loops, arches, isolated knots, and filaments.

Loops, with a typical diameter of 5×10^3 km, appear to follow magnetic field lines and seem to connect regions of opposite magnetic polarity within the same active region (Howard, 1971). Arches have larger cross-sections, they are more open and stable and they often connect different active regions. Isolated knots are more transitory, high-density enhancements, with a roughly spherical shape. They occur frequently together with loops and arches. Long, thin, and tortuous filaments are often seen above active regions. They are denser and cooler than the background corona. The part they play in solar flares and noise storm phenomena is not well understood.

It must be said that in the classification of coronal inhomogeneities, due to inherent complexities, some arbitrariness is often unavoidable and that the features described are changing more or less rapidly all the time.

3. THE ACTIVE REGION CORONA

In order to consider quantitative theories of type I bursts, it is necessary to adopt a simplified model of the solar corona above active regions. The basic physical quantities required are the electron density, the magnetic field strength, and the electron temperature. After what has already been said in the previous section, the following

models of the requisite parameters represent only an average, "smoothed out" picture of the corona in these regions.

(a) Electron Density

The conventional model of electron density distribution in the solar corona is given by the Baumbach-Allen formula:

$$n_e = 10^8 (1.55 \rho^{-6} + 2.99 \rho^{-16}) \text{ cm}^{-3}, \quad (6.1)$$

where $\rho = R/R_\odot$, R = radial distance from centre of the sun. On the basis of observations made with the K-coronameter, Newkirk (1961) proposed an electron density distribution in the quiet corona described by

$$n_q = n_0 10^{4.32/R} \text{ cm}^{-3}, \quad (6.2)$$

where $n_0 = 4.2 \times 10^4$. He suggested that electron densities along the axis of coronal streamers are about double those of the quiet corona at the same height. From radio observations of type II and type III bursts, Wild *et al.* (1963) proposed a streamer with twice the density given by the Newkirk coronal streamer. Such an electron density distribution, which is four times that of the Newkirk quiet corona and about ten times that of the Baumbach-Allen model, appears to be the best empirical model to adopt for the corona above active regions (see Fig. 6.2).

(b) Magnetic Field Strength

Even though the magnetic field probably determines the morphological structure of the corona, comparatively little is actually known about the coronal magnetic field (Howard, 1971). There is ample evidence that the coronal magnetic field is inhomogeneous on a scale-length of 10^3 km, possibly associated with fine magnetic field

Fig. 6.2: Average electron densities in the solar corona as a function of height. The plotted points give the measured height of origin of type II and type III bursts. $\odot$ — type III bursts. $\blacksquare$ — type II bursts. The bar represents the probable error of the 50 Mc/s point and is typical of the errors in all four points for type II bursts. (After Wild *et al.*, 1963.)

structures originating from the photosphere (see e.g., Sweet, 1971).

Potential field models for the coronal magnetic field above active regions have been proposed by Newkirk (1967) to match estimates derived from observations of radio bursts. Such models have been justified by Rust and Roy (1971), who show that coronal magnetic fields above slowly varying active regions can be accurately represented by vacuum potential fields derived from underlying photospheric field sources (principally the larger sunspots in a region). However, the observational data on coronal magnetic fields deduced from radio bursts usually suffer from the difficulty that the validity of the assumed radiation mechanism is frequently not known.

Nevertheless, a potential field of the type (A) (see Fig. 6.3) proposed by Newkirk (1967) appears to be a reasonable model for the coronal magnetic field above an active region. Let us take as our model the potential field of a dipole of magnetic moment 10^{32} e.m.u., with length 4×10^9 cm, embedded about the same distance below the photosphere. This corresponds to a magnetic field of strength 2.5×10^3 gauss on the solar surface, and at distances greater than about $0.1 R_{\odot}$ above the photosphere, the magnetic field strength can be approximated by

$$B = \frac{0.292}{\rho^3}, \quad (6.3)$$

where ρ is the height above the photosphere in units of solar radii. This model, together with twice the Newkirk model for electron density in a streamer can be used to obtain electron plasma frequency, electron gyrofrequency, their ratio, and the cutoff frequency of X-mode waves (see equation (5.1)) as functions of height above the photosphere (see Fig. 6.4).

Fig. 6.3: The strength of coronal magnetic field as a function of height above the photosphere as estimated by various techniques. Values referring to active regions appear as heavy lines. Simple models for the potential field above an active region (A), an extended dipole region (B), and the general solar field at the equator (C) appear for comparison. The curve H represents the field required for pressure equilibrium with the solid wind. (After Newkirk, 1967.)

Fig. 6.4: Electron plasma frequency f_p , electron gyrofrequency $\Omega_e/2\pi$, their ratio ω_p/Ω_e , and the cutoff frequency of X-mode waves as functions of height above the photosphere for the adopted models of electron density and magnetic field strength above active regions.

(c) Electron Temperature

On account of the high thermal conductivity of the solar corona, the kinetic temperature of electrons varies only slightly with height and it can be taken to be 2×10^6 °K above active regions. This corresponds to a thermal velocity of electrons $V_e = 7.8 \times 10^8$ cm sec⁻¹. Existence of ion-acoustic waves above active regions (Gordon, 1968) can be regarded as evidence for $T_e > T_i$ in these regions.

The adopted electron temperature and the adopted electron density distribution are sufficient to calculate the electron Debye length $\lambda_{De} = V_e / \omega_p$ and the collision frequency ν_c (see equation (3.45)) as functions of height above the photosphere (see Fig. 6.5).

Another important characteristic parameter is the plasma beta, which is the ratio of particle-kinetic pressure to magnetic pressure:

$$\beta \equiv \frac{\sum n_\alpha k T_\alpha}{B^2 / 8\pi} . \quad (6.4)$$

At the assumed temperature and on the supposition that electron pressure is dominant,

$$\beta \approx 6.75 \times 10^{-4} (\omega_p / \Omega_e)^2 , \quad (6.5)$$

whose variation with height above active regions is illustrated in Fig. 6.6. The Alfvén velocity, which is given, for a hydrogen plasma, by

$$v_A = 6.93 \times 10^8 \Omega_e / \omega_p \text{ cm sec}^{-1} , \quad (6.6)$$

is displayed as a function of height above the photosphere in Fig. 6.7.

4. OBSERVATIONS

Several reviews on the general nature of type I bursts and associated noise storm radiation are available; see for example Wild

Fig. 6.5: Electron Debye length and effective collision frequency as functions of height above the photosphere for adopted model of electron density distribution and $T_e = 2 \times 10^6$ °K.

Fig. 6.6: Plasma β as a function of height above the photosphere for adopted models of electron density distribution and magnetic field strength.

Fig. 6.7: Alfvén velocity as a function of height above the photosphere. The scale on the right is its value normalized with respect to V_e , taking $T_e = 2 \times 10^6 \text{ }^\circ\text{K}$.

et al. (1963), Kundu (1965) and Zheleznyakov (1970). However, more recently, heliographic observations, high-resolution observations with improved instruments and observations of chains of type I bursts have contributed significantly to our knowledge of the phenomenon, so that an updated review is called for. Although no attempt is made at being exhaustive in our review, all the important observations are included and are presented in an order which reflects their importance for our understanding and for the construction of theoretical models.

(a) Intensity

(1) The intensity of noise storm radiation varies from barely detectable values up to thousands of times the quiet sun intensity. Typical flux density observed is about a few times $10^{-21} \text{ W m}^{-2} \text{ Hz}^{-1}$, corresponding to brightness temperatures of about $10^9 \text{ }^\circ\text{K}$, for apparent sizes of the sources. The flux density from bursts is typically an order of magnitude greater than that of the storm radiation. Brightness temperatures of bursts $10^{11} \text{ }^\circ\text{K}$ or higher have occasionally been observed.

(b) Polarization

(2) Type I bursts and the background continuum are usually strongly circularly polarized to the same degree and in the same sense. Exceptions to this general rule can occur during the initial stages of a noise storm, when the polarization is highly variable, with varying degrees of polarization even within individual bursts (Chernov *et al.*, 1972). Otherwise, exceptions can be attributed to independent sources situated in opposite hemispheres or to the inhomogeneity of the background magnetic field in a large source.

(3) The degree of polarization of individual bursts usually increases gradually at the initial stage of storm development and after it attains 100%, it remains on this level till the end of the storm (Chernov *et al.*, 1972).

(4) The polarized radiation corresponds to ordinary mode waves of the magnetoionic theory (Payne-Scott and Little, 1951).

(5) There is no significant difference between the polarization of bursts occurring in chains and those occurring separately (Hanasz, 1966).

(c) Duration

(6) A noise storm as a whole can last up to hours or days. However, the spectrum of the background continuum can change continuously with time and during periods of high activity, it shows marked changes in profile in periods of about 20 minutes.

(7) Individual bursts have duration between about 0.1 to 1.0 sec, and it decreases with increase in frequency (Elgarøy, 1967). Their intensity-time profiles are usually symmetric (Elgarøy, 1961; Chernov *et al.*, 1972).

(8) The average lifetime of a chain of type I bursts is about 5 sec (Elgarøy and Ugland, 1970). The distribution of the number of chains w.r.t. lifetime follows an exponential law: $N \propto \exp(-t/30)$, where t is in seconds (Hanasz, 1966).

(d) Bandwidth

(9) The bandwidth of the background continuum radiation is usually about 100 MHz, while that of bursts is between 2 to 10 MHz,

typically about 4 MHz. The bandwidth of bursts increases with an increase in frequency (Elgaroy, 1967).

(e) Frequency Drift

(10) About half of type I bursts exhibit frequency drifts w.r.t. time, either toward higher or lower frequencies. The drift rate ranges between 1 MHz sec^{-1} to tens of MHz sec^{-1} , being between those of type II and type III bursts.

(11) About 50% of chains drift toward lower frequencies, 30% toward higher frequencies and 20% show no detectable drift (Elgaroy and Ugland, 1970). The absolute value of the frequency drift varies between $0.16 \text{ MHz sec}^{-1}$ and 0.25 sec^{-1} , corresponding to a "radial" velocity of the exciting agency between $\pm 0.27 V_e$, with an average absolute value of about $0.05 V_e$ (Elgaroy and Ugland, 1970; Hanasz, 1966; Wild and Tlamicha, 1964).

(12) Drift directions and drift rates can be different for bursts occurring in the same chain. In some cases, all bursts in a chain have no detectable frequency drift (Elgaroy and Ugland, 1970).

(f) The Source

(13) The shape of the source appears to be that of an elliptical disk (Malinge, 1963), the major axis of the ellipse being more or less in the N-S direction and the plane of the disk being roughly parallel to the photospheric surface.

(14) The angular size of storm centres varies from less than 1 min of arc to about 10 mins of arc and it decreases with increase in frequency of observation. At 169 MHz, a typical diameter of a storm centre is 3 or 4 mins of arc (Malinge, 1963). The angular size of

bursts is somewhat smaller, less than $1'.6$ has been observed (Goldstein, 1959). (One min of arc corresponds to a dimension of $0.062 R_{\odot}$ on the solar surface.)

(15) The mean height of sources of type I bursts, when investigated on 200 MHz, is between 0.2 to $0.3 R_{\odot}$ above the photosphere (Morimoto and Kai, 1961).

(16) In general, storm centres are located near the corresponding plasma level of Newkirk's model of the solar atmosphere above active regions.

(17) Chains of type I bursts appear most frequently at 100 MHz. They are not found in spectrographic records at frequencies greater than about 300 MHz (Elgaroy and Ugland, 1970). Assuming type I bursts originate at close to the local plasma frequency, and using the $2 \times$ Newkirk model of coronal streamer, one deduces that chains of type I bursts occur mostly between 0.3 and $0.7 R_{\odot}$ above the photosphere, with most frequent occurrence at about $0.5 R_{\odot}$.

(18) Although bursts and continuum radiation come from the same general region, they have been observed to come from distinct parts of the same source by radioheliographic observations (Sheridan, 1972).

(g) Centre-Limb Variations

(19) The number of occurrences of noise storms decreases from the central meridian toward the limb. This means the storm radiation is narrowly directed.

(20) The average height of sources of bursts and continuum radiation increases toward the limb (Morimoto and Kai, 1962; Malinge, 1963).

(21) The average degree of polarization of storm bursts and the

intensity of an average storm centre, both decrease from the central meridian (Kai, 1962; Malinge, 1963) (Figs. 6.8a and 6.8b).

(22) It appears that the average lifetime of the chains decreases toward the limb whilst the change in bandwidth between the start and end of chains and the average frequency drift, both increase toward the limb (Elgaroy and Ugland, 1970).

(h) Associated Phenomena

(23) Most noise storms are preceded within two hours by a flare and for important flares, the most probable time delay between noise storms and flares is about 30 minutes (Kundu, 1965).

(24) Noise storms are always associated with sunspot groups, with the area of a group greater than about 6×10^{-4} of the solar disk. When the area of a sunspot is greater than 7.5×10^{-4} of the solar disk, the probability of its association with an important storm centre is about 80% (Malinge, 1963).

(25) Some storm bursts appear to split into two components (Elgaroy, 1961; Yoh and James, 1967). Observed in the range 190-215 MHz, the mean frequency separation between the two elements is about 8 MHz (Elgaroy, 1961).

(26) Two or more chains running almost parallel at the same time are frequently present on spectrographic records (Elgaroy, 1961; Hanasz, 1966). Statistical analysis does not reveal any harmonic phenomenon, the two components of a pair often do not start simultaneously and their frequency separation varies between 10 to 50 MHz usually, with 20 MHz being a most common value (Hanasz, 1966).

(27) Weak type III bursts sometimes appear to "grow" out of type I chains (Wild and Tlamicha, 1964; Hanasz, 1966). However,

Fig. 6.8a: Centre-limb variation in the average degree of polarization of type I bursts. (After Kai, 1962.)

Fig. 6.8b: Centre-limb variation of the intensity of an average storm centre. (a) Intensity $< 20 \times 10^{-22} \text{ W m}^{-2} \text{ Hz}^{-1}$; (b) Intensity $> 20 \times 10^{-22} \text{ W m}^{-2} \text{ Hz}^{-1}$. (After Malinge, 1963.)

radioheliographic observation of a noise storm in 1968 showed that the type III source position was displaced about $0.5 R_{\odot}$ transversely from the type I (Stewart and Labrum, 1972).

A schematic drawing of a type I noise storm and type III bursts is given in Fig. 6.11.

5. REVIEW OF EXISTING THEORIES

Various interpretations to the above observations have been proposed. In this section, existing theories of type I bursts and associated noise storm radiation are reviewed and critically appraised.

Two theories, so far, appear to have survived in the literature: one involves coherent gyromagnetic radiation at low harmonics of the electron gyrofrequency (Fung and Yip, 1966), the other involves radiation from coherent plasma waves at about the local plasma frequency (Takakura, 1963; Trakhtengerts, 1966). These theories are antedated in their basic physical ideas, by earlier, less quantitative theories (see references cited in them). Because these theories are the most developed, qualitatively and quantitatively, our review and discussion will be confined to them. A summary of the aspects of type I phenomenon considered by these authors is given in Table 6.2.

It will be shown below that the theory based on coherent gyromagnetic radiation (Fung and Yip, 1966) is unsatisfactory and the theory based on plasma emission processes (Takakura, 1963; Trakhtengerts, 1966) is incorrect and incomplete in various ways. Our judgements are based on the criterion that observations, particularly those which distinguish type I phenomenon from phenomena of other spectral types, should be explained on established principles of plasma physics and on plausible assumptions.

Fig. 6.9: Dynamic spectra of type III bursts and type I noise storms (Jan, 1971 and May, 1972): (a) continuum radiation and type I burst chains with type III bursts 'growing' out of them; (b) same as (a), in addition, there are bursts and chains of bursts superimposed on the background continuum; (c) mainly isolated bursts superimposed on the background continuum. (By courtesy of K.V. Sheridan, C.S.I.R.O., Sydney.)

Fig. 6.10: Dynamic spectra (March, 1972) of type I bursts and chains of bursts, apparently without the background continuum (top two spectra). The bottom spectrum shows an arched chain, with type III bursts 'growing' out of it. (By courtesy of K.V. Sheridan, C.S.I.R.O., Sydney.)

Fig. 6.11: A schematic drawing of a type I noise storm and type III burst events in the solar corona.

	(a)	(b)	(c)	(d)	(e)*	(f)	(g)	(h)	(i)	(j)	(k)
Fung and Yip (1966)	✓*	✓	×	✓	✓	×	✓*	×	×	✓*	×
Takakura (1963)	✓	×	×	✓*	✓	✓*	✓	✓	✓	✓	✓*
Trakhtengerts* (1966)	✓	×	×	×	×	×	×	×	×	✓	✓

Table 6.2: Summary of aspects of type I phenomenon considered by the authors. Cases with asterisks are discussed in the text. Column (a) origin of type I bursts; (b) origin of the background continuum radiation; (c) origin of chains of type I bursts; (d) intensity of radiation; (e) polarization; (f) duration; (g) bandwidth; (h) origin of frequency drift; (i) centre-limb effects; (j) exciting agency; (k) origin of the exciting agency.

(a) Theory of Fung and Yip (1966)

Fung and Yip (1966) have developed a theory for type I bursts and continuum radiation, which can be summarized as follows. Electron streams gyrating in the solar corona above active regions emit gyromagnetic radiation at low harmonics of the electron gyrofrequency. Simultaneous solution of the dispersion and Doppler equations (see e.g. equations (2.81) and (3.18)) gives rise to so-called "single" and "double" frequency solutions when the conditions $s/\gamma \geq \omega_{0,X}/\Omega_e$ and $s/\gamma < \omega_{0,X}/\Omega_e$ respectively are satisfied. Here, as elsewhere, s is the harmonic of the radiation, γ is the Lorentz factor, Ω_e is the electron gyrofrequency and $\omega_{0,X}$ is the cutoff frequency of the 0, X-mode waves. "Single" frequency solutions are assumed to give rise to broad-band background continuum radiation, whilst "double" frequency solutions are assumed to give rise to narrow-band bursts of radiation. This theory is open to several criticisms, the main ones of which are related to polarization, burst radiation and model of exciting agency. These points are discussed in detail below.

(i) Polarization

The single-particle theory of gyromagnetic radiation (see e.g. equation (3.18)) predicts that the radiation, for given plasma parameters, is predominantly in the X-mode (see Fung and Yip, 1966, p.773). This already foreshadows a difficulty faced by gyromagnetic theories, since type I noise storms are polarized in the sense of the O-mode. Granted Fung and Yip's stream distribution (equation (3.8), *loc. cit.*) which will be questioned below, the the growth rates for both modes of radiation at the same harmonic are of the same order of magnitude, with X-mode growth rates significantly larger for large pitch angles of the stream. Large pitch angles are invoked by Fung and Yip (1966, p.780) to explain the narrow-bandedness of the bursts. Therefore, this radiation mechanism does not produce a predominance of O-mode radiation and in fact, when large pitch angles of the streams are assumed for type I bursts, the radiation produces an excess of X-mode waves relative to O-mode waves.

To explain the high degree of circular polarization in the sense of the O-mode, Fung and Yip (1966) propose that the radiation either occurs at frequencies below the cutoff frequency of X-mode waves or suffers strong gyroresonance absorption of the X-mode waves at the third harmonic (p.789, *loc. cit.*). These proposals might well be adequate in explaining the polarization of type I bursts at high frequencies, corresponding to low source altitudes. However, at low frequencies (say 80 MHz), corresponding to relative high altitudes, the ratio ω_p/Ω_e is probably large (greater than 4 say), emission purely in O-mode waves is confined to a very narrow frequency range (see Fig. 6.4), gyroresonance absorption at harmonics greater than 3 or 4 is weak (see section 4, chapter 3) and hence the above-mentioned proposals for

explaining the high degree of circular polarization in the sense of the O-mode seem implausible.

(ii) Burst Radiation

The "double" frequency solution condition, which is also the condition for occurrence of burst radiation in their theory (Fung and Yip, 1966) and the escape conditions for radiation give the following inequalities:

$$s\Omega_e < \gamma\omega_p < s\Omega_e / (1 - \eta_0 \beta_s \cos\theta) , \quad (6.7)$$

where $\beta_s \equiv u_s/c$, with u_s denoting the stream velocity parallel to the background magnetic field. From frequency drift rate measurements and other observations it seems that there is no reason for supposing the beams responsible for type I bursts are relativistic. If the electron beams are non-relativistic, then the inequalities (6.7) would place stringent conditions on the occurrence of burst radiation.

Fung and Yip (1966) predict maximum intensity of the burst radiation occurs near the level where $\omega_p \approx \Omega_e$ and significantly reduced intensities at higher levels. An examination of spectralgraphic records (see e.g. Fig. 6.10) appears to indicate that bursts of comparable intensities can occur at widely separated frequencies.

(iii) Model of Exciting Agency

As a model of the exciting agency, Fung and Yip (*ibid.*) adopt δ -function distributions for electron streams and they allow stream velocities up to 0.7 c. No justification, based on observations or otherwise, appears to have been given for neglecting velocity spreads in the electron streams or for considering relativistic velocities. In view of the discussions given in section 5 of chapter 3, the

hydrodynamic or reactive treatment of the amplification process of the radiation (associated with the δ -functions) also needs some justification.

An alternative model involving a bi-Maxwellian streaming distribution for electron streams has been used by Melrose (1973) to discuss coherent gyromagnetic radiation. He concludes that preferential gyroresonance absorption of X-mode waves cannot overcome preferential emission to give net polarization in the sense of the O-mode.

(b) Theory of Takakura (1963)

Takakura (1963) has developed a theory of type I bursts, based on the "plasma hypothesis", i.e. the radiation arises from nonlinear Rayleigh scattering of coherent electron plasma waves. This theory, which contains discussions on more features of the observed radiation, can be summarized as follows.

Two Alfvén wave packets, propagating in opposite directions, collide and produce a beam of electrons with a velocity about three times the Alfvén velocity $3v_A$. On the assumption that $v_A \approx V_e$, the beam generates coherent electron plasma waves, which are then scattered by thermal density-fluctuations to give rise to radiation at about the local plasma frequency. The amplitude of electric field of the electron plasma waves, which determines the intensity of the radiation, is taken to be that of the steady state, when energy gain of the electron plasma waves due to beam-plasma interaction balances energy loss due to collisional damping. The polarization characteristics are explained by the escape conditions of radiation and by gyroresonance absorption. For example, if $1 > X > 1 - Y$, where X and Y are given by equation (2.80), only O-mode radiation can escape to the earth, and if $1 - X > Y > \frac{1}{2}$, the

radiation is subjected to gyroresonance absorption at levels $Y = \frac{1}{2}, \frac{1}{3}$, etc. The duration of the burst is assumed to be determined by the deflection time of the beam, from which the beam velocity is determined to be about $1.2 - 1.6 \times 10^9$ cm sec⁻¹. The bandwidth was suggested to be due either to a variation in electron densities at the source or to a dispersion of velocities in the beam. The frequency drift was attributed to beam motion through the solar corona.

Although in this thesis, we will also develop (section 6) the "plasma hypothesis" for the generation of type I bursts, Takakura's theory is, nevertheless, criticized in several respects. These criticisms pertain to the intensity, polarization and duration of the bursts, and to the generation mechanism of the exciting agency. These are discussed separately and in detail below.

(i) Intensity

Takakura's calculation of the radiation intensity, based on the work of Ginzburg and Zheleznyakov (1958), neglects induced effects. As shown by Kaplan and Tsytovich (1969) induced emission and absorption processes become important as the energy densities of non-thermal electron plasma waves are raised. For example, from equations (3.44) and (4.87), it can be seen that at $W^{el} \sim W_1 (v_c/\omega_p)$, induced absorption of the electron plasma waves begins to dominate over collisional damping which is assumed by Takakura (p.466, *loc. cit.*) to be the sole absorption process determining the amplitude of electron plasma waves. In fact, at the energy densities of non-thermal electron plasma waves ($\sim 10^{-7}$ erg cm⁻³) considered by Takakura, induced effects cannot be neglected. In general, induced effects alter the naive picture of plasma turbulence and they affect plasma emission processes.

If induced effects on the radiation are only those of "positive" absorption, i.e., the radiation is damped by induced conversion to electron plasma waves, Kaplan and Tsytovich (*ibid.*) showed that the brightness temperature of the fundamental at the source cannot exceed about 10^9 °K. Hence the radiation must be amplified for higher brightness temperatures to have been observed.

(ii) Polarization

Takakura (1963) neglects the effects of a background magnetic field in his treatment of the "plasma emission processes". He predicts type I bursts at frequencies above the cutoff frequency of X-mode waves would be completely unpolarized at the source. Hence, to account for the polarization of type I bursts, he suggests that either the radiation occurs at frequencies below the cutoff frequency of the X-mode waves or the radiation is subjected to strong preferential gyroresonance absorption of the X-mode waves at levels $Y = \frac{1}{2}$ and $\frac{1}{3}$, as the radiation propagates to the earth.

According to this theory then, if the source is situated at a sufficiently high altitude where the condition $Y < \frac{1}{3}$ is likely to be satisfied, the type I bursts observed at earth should be unpolarized. The sources of type I bursts at low frequencies (24 - 38 MHz) are expected to satisfy the condition $Y < \frac{1}{3}$ (see Fig. 6.4) and the radiation is then expected to be unpolarized. The observation of strongly polarized type I bursts at these low frequencies by Yoh and James (1967), Warwick and Dulk (1969), and Sastry (1972), appears to suggest that causes of the strong polarization observed are not just those proposed by Takakura (1963).

(iii) Duration

Takakura (1963) suggests that the duration of type I bursts is determined by the deflection time of the electron beams. The deflection time is given by (*ibid.*)

$$\tau_D = m_e^2 u_s^3 / 8\pi e^4 n_e \ln \Lambda_e \approx 0.07 \tau_c (u_s/V_e)^3 \quad (\text{sec}), \quad (6.8)$$

where u_s ($> 2V_e$) is the beam velocity and τ_c is the collisional damping time given by (see equation (3.44))

$$\tau_c = 7.71 \times 10^{16} / \{f_p^2 (17.95 - \log f_p)\} \quad (\text{sec}), \quad (6.9)$$

with the local plasma frequency f_p in Hz.

Although the deflection time τ_D is capable of accounting for the duration of bursts at high frequencies (say 200 MHz), it appears to be too long compared with observed durations at low frequencies (Yoh and James, 1967; Warwick and Dulk, 1969; Sastry, 1972). For example, at 25 MHz, one finds $\tau_D \approx 8(u_s/V_e)^3$ secs, which is much larger than the typical observed value of 2 or 3 secs, for the duration of bursts at high source altitudes.

To circumvent this difficulty, Elgaroy and Eckhoff (1966) and Sastry (1972) suggest that type I bursts occur in coronal regions where the temperature is significantly less than 10^6 °K. Sastry (1972) suggests temperatures $T_e \approx 10^5$ °K. In view of the high thermal conductivity of the solar corona, it seems difficult to substantiate the hypothesis that coronal regions with temperatures as low as 10^5 °K are sufficiently prevalent to account for the common occurrence of type I bursts.

(iv) Generation Mechanism of Electron Beams

As pointed out by Trakhtengerts (1966), Takakura (1963) has not taken into account charge separation at the MHD wave front. When charge separation at the wave front is taken into account Trakhtengerts (*ibid.*) shows that the reflected particles are stable with respect to the excitation of electron plasma waves.

(c) Theory of Trakhtengerts (1966)

Trakhtengerts (1966) is principally concerned with the generation mechanism of electron beams which could give rise to type I bursts; little discussion is given over to the explanation of observed features of the bursts. He raises objections against Takakura's (1963) generation mechanism and proposes the following alternative.

A magnetohydrodynamic pulse (soliton) of finite amplitude propagates along the background magnetic field and reflects plasma particles from its wave front. On account of differences in mass between electrons and ions, more electrons than ions are reflected from the wave front and thus a charge separation is set up at the front. Contrary to Takakura's conclusions, the reflected electrons are shown to be stable with respect to the excitation of electron plasma waves. Under the conditions (1) $V_e/\Omega_e < d < V_e/v_c$ and (2) $1 \gg b/B_0 \gg (m_e/m_i)(V_e/v_A)^2$, where d is the width of the pulse, $b+B_0$ is the maximum amplitude of the magnetic field in the pulse, and other symbols have their usual meanings, the charge separation causes a counter current which is unstable with respect to the excitation of non-thermal electron plasma waves. Moreover, when two such MHD waves collide, the bursts of radiation that arise from the non-thermal electron plasma waves should appear as intense transient blips superposed on

a relatively weak background of radio emission, which consists of slowly drifting narrow-band bursts with large overlaps in frequency.

This theory can be questioned on a number of points. We discuss separately (i) the question of dissipation, (ii) the assumption $V_e \approx v_A$, (iii) the frequency drift of the bursts occurring in chains.

(1) Dissipation

In the absence of dissipation, it is possible to obtain from the MHD equations only solitary waves or solitons, which are represented by symmetrical potential barriers (Sagdeev, 1966) for disturbances propagating along the background magnetic field. However, in the presence of dissipation, in this case by the generation of microturbulence, a soliton solution changes into a shock solution (Tidman and Krall, 1971). It appears therefore the problem has not been solved self-consistently, since the effect of dissipation on the structure of the disturbance, viz. the soliton has to be taken into account.

(ii) Assumption $V_e \approx v_A$

Trakhtengerts (1966) makes the assumption $V_e \approx v_A$ and so the condition (2) above reduces to $1 \gg b/B_0 \gg m_e/m_i$, a condition which can probably be easily met. However, he also identifies v_A with the speed of propagation of the disturbance. Interpreting the frequency drift rates of chains of type I bursts as indicating the Alfvén speed v_A , Wild and Tlamicha (1964), Hanasz (1966) and Elgarøy and Ugland (1970) find the average value of v_A to be about $0.05 V_e$. It appears therefore that the assumption $V_e \approx v_A$ needs further justification.

(iii) Frequency Drift of Bursts Occurring in Chains

Following Trakhtengerts' theory, one would expect the directions of frequency drift of individual bursts occurring in a chain to be roughly aligned with the direction of frequency drift of the chain as a whole. However, drift directions and drift rates can be different for different bursts occurring in the same chain (Elgaroy and Ugland, 1970).

6. MODEL OF TYPE I BURSTS

In the previous section, some difficulties encountered by existing theories of type I bursts are indicated. In this section, an attempt is made to construct a model of type I bursts, which overcomes some of the objections raised in the previous section.

Our model is also based on the "plasma hypothesis", but it takes into account effects which have been ignored in a previous theory (Takakura, 1963). These effects are (a) the effect of a background magnetic field on the plasma emission processes and (b) the induced effects which alter radically the quantitative consideration of intensities of type I bursts given by Takakura (1963). The effect of a background magnetic field is generally believed to be manifested by the strong polarization of type I bursts (Kundu, 1965), whilst induced effects have been shown to play an important rôle both in the formation of spectra of plasma turbulence and also in the amplification and absorption processes of radiation (Kaplan and Tsytovich, 1969).

Indeed our understanding of induced effects in plasma emission processes has been enhanced by a quantitative theory of type III bursts (Smith, 1970). Although the radiation mechanism invoked here for type I bursts is the same as that invoked for type III bursts (Zheleznyakov and Zaitsev, 1970a,b; Smith, 1970), problems involved in

constructing theoretical models for the two phenomena are different in many respects. Firstly, interpreting the frequency drifts observed on spectrographs as indicating the radial motion of the source through the solar corona, one finds the speeds of electron streams associated with type I bursts are about several times the thermal velocity of electrons in the solar corona, whereas those associated with type III bursts are about $c/3$. This difference in electron stream speeds is expected to lead to different spectra of turbulence in electron plasma waves. Secondly, the background magnetic field is probably stronger in the source region of type I bursts than in that of type III bursts. The stronger magnetic field could lead to differences in the turbulence spectra and in the radiation processes.

To develop a fully quantitative theory of type I bursts, one needs to include a background magnetic field and obtain the turbulence spectra of electron plasma waves generated by electron streams, with average speeds several times V_e . Since this problem has not been solved, the model discussed below can only be semi-quantitative. However, it is possible to estimate certain quantities, such as the energy density of microturbulence in electron plasma waves and the number density of electrons in streams, which are required to account for certain features of type I bursts. Moreover, the inclusion of a background magnetic field will allow an estimate of the degree of polarization of the radiation in the source region.

(a) Intensity

The observed equivalent brightness temperature T_b of a source is related to the observed flux density of the radiation S_o by (Kundu, 1965; Bekefi, 1966)

$$T_b = \frac{2\pi^2 c^2 S_o}{\kappa \omega^2 \Omega_o} \quad (^\circ\text{K}) , \quad (6.10)$$

where Ω_o is the solid angle subtended by the apparent source area at the observer. However, due to scattering of the radiation by coronal inhomogeneities (Fokker, 1965; Steinberg *et al.*, 1970; Riddle, 1972) the apparent source size can be very much larger than the actual size of the source. The effect of coronal scattering can therefore lead to an underestimation of the true brightness temperature of the source by a large factor. For the fundamental radiation ($\omega \approx \omega_p$), Smith (1970) finds the factor to be between 10^4 to 10^5 . It would appear then the actual brightness temperature of a strong type I burst could be as high as 10^{16} $^\circ\text{K}$.

The brightness temperature of the source is related to the intensity of the radiation leaving the source region by the formula (Bekefi, 1966)

$$T_{bs} = \frac{4\pi^3 c^2}{\kappa \omega^2 \eta^2} I(\omega, \underline{\Omega}) \quad (^\circ\text{K}) , \quad (6.11)$$

where η is the refractive index and $I(\omega, \underline{\Omega})d\omega$ is the intensity of the radiation propagating in a direction $\underline{\Omega}$ in the frequency range between ω and $\omega + d\omega$.

To obtain the intensity of radiation leaving the source region, a radiative transfer equation needs to be solved for a particular source configuration. Assuming the magnetoionic wave modes are uncoupled in the source region, one deduces from equations (4.3), (4.21), (4.24) and (5.2), radiative transfer equations of the form

$$\frac{d}{d\ell} \left(\frac{I^\sigma}{\eta_\sigma^2} \right) = j^\sigma(\omega, \tau) - \frac{\mu^\sigma(\omega, \theta)}{v_g^\sigma(\omega, \theta)} \frac{I^\sigma(\omega, \theta)}{\eta_\sigma^2} , \quad (6.12)$$

where the derivative is taken along the ray path ℓ . The other quantities are given by

$$j^\sigma(\omega, \theta) \equiv \left(\frac{2}{\pi}\right)^{1/2} \frac{\omega_p^2}{(4\pi)^2 n_1 c^2} \frac{\partial(\omega \eta_\sigma)/\partial\omega}{(1 + Z_1 T_e/T_1)^2} R_1^{\sigma l}(\omega, \theta), \quad (6.13)$$

and, from equations (4.27) and (5.48),

$$\frac{\mu^\sigma(\omega, \theta)}{v_g^\sigma(\omega, \theta)} = \left(\frac{\pi}{2}\right)^{1/2} \frac{\omega_p^2}{c n_1 m_1 v_1^2} \frac{\partial(\omega \eta_\sigma)/\partial\omega}{(1 + Z_1 T_e/T_1)^2} R_2^{\sigma l}(\omega, \theta), \quad (6.14)$$

where the approximations $\omega = \omega' = \omega_p$ and $v_g^\sigma = c/\{\partial(\omega \eta_\sigma)/\partial\omega\}$ have been made. The functions $R_1^{\sigma l}(\omega, \theta)$ and $R_2^{\sigma l}(\omega, \theta)$ are related to the efficiency of the "nonlinear" radiation process and they are defined by

$$R_1^{\sigma l}(\omega, \theta) \equiv \int \frac{d^3 \underline{k}'}{(2\pi)^3} \left(\frac{W_E}{W_T}\right)^{el'} \frac{F^{\sigma l}(\omega, \theta, \theta')}{k' v_1 |\cos \theta'|} W^{el'}(\underline{k}') \exp \left\{ -\frac{(\omega - \omega')^2}{2k'^2 v_1^2 |\cos \theta'|^2} \right\}, \quad (6.15)$$

and

$$R_2^{\sigma l}(\omega, \theta) \equiv \int \frac{d^3 \underline{k}'}{(2\pi)^3} \left(\frac{W_E}{W_T}\right)^{el'} \frac{F^{\sigma l}(\omega, \theta, \theta')}{k' v_1 |\cos \theta'|} (\omega - \omega') W^{el'}(\underline{k}') \exp \left\{ -\frac{(\omega - \omega')^2}{2k'^2 v_1^2 |\cos \theta'|^2} \right\}, \quad (6.16)$$

when $\lambda_1 \equiv (\Delta k_\perp v_1 / \Omega_1)^2 \ll 1$; otherwise $k' |\cos \theta'|$ is to be replaced by k' , when $(\Delta k_\perp v_1 / \Omega_1)^2 \gg 1$. The function $F^{\sigma l}(\omega, \theta, \theta')$ is determined by equation (5.3). Given a particular distribution of non-thermal electron plasma waves $W^{el'}(\underline{k}')$, the integrals in equations (6.15) and (6.16) can be evaluated.

On defining an optical depth τ^σ by

$$d\tau^\sigma \equiv -\mu^\sigma dl / v_g^\sigma, \quad (6.17)$$

the equations of radiative transfer can be rewritten as

$$\frac{d}{d\tau^\sigma} \left(\frac{I^\sigma}{\eta_\sigma^2} \right) = S^\sigma - \left(\frac{I^\sigma}{\eta_\sigma^2} \right), \quad (6.18)$$

where the source functions S^σ are defined by

$$S^{\sigma}(\omega, \theta) \equiv \frac{j^{\sigma} v_g^{\sigma}}{\mu^{\sigma}} = \frac{3\omega_p^3 m_i v_i^2}{(2\pi)^3 c^2} \frac{R_1^{\sigma\ell}(\omega, \theta)}{R_2^{\sigma\ell}(\omega, \theta)}. \quad (6.19)$$

For a ray path of length ℓ , the optical depth is given by

$$\tau^{\sigma} = -\frac{\mu^{\sigma} \ell}{v_g^{\sigma}} = -\left(\frac{\pi}{2}\right)^{\frac{1}{2}} \frac{\omega_p^{\ell}}{c^2 n_i m_i v_i^2} \frac{\partial(\omega n_{\sigma})/\partial\omega}{(1+Z_i T_e/T_i)^2} R_2^{\sigma\ell}(\omega, \theta). \quad (6.20)$$

Consider a source region of total thickness less than 10^9 cm, with an amplification layer and an absorption layer.

Fig. 6.12: A source model.

The ray path (Fig. 6.12) has been taken to be rectilinear for the following reasons. (1) Bending of the ray by refraction is insignificant if the source has small dimensions $L \ll R_0$, so that the total change in refractive indices is negligible. (2) If $L < 10^9$ cm, scattering by coronal inhomogeneities is probably insignificant (Fokker, 1965). (3) If the source structure is highly transient and it changes in a time scale of the order of the time taken for the ray to traverse the layers, then the rays propagating initially in a backward direction and subsequently reflected forward will be unimportant.

For a ray emerging from a point at the base of the amplification region, it reaches the absorption region with intensity/unit frequency

range:

$$I^\sigma(\omega, \Omega) = \eta_\sigma^2 S_m^\sigma (1 - e^{-\tau_m^\sigma}) \approx -\eta_\sigma^2 S_m^\sigma e^{-\tau_m^\sigma}, \quad (6.21)$$

where the subscript 'm' designates quantities in the amplification region. Similarly, the intensity/unit frequency range for the radiation leaving the source region, after passing through the absorption layer is given approximately by

$$I^\sigma(\omega, \Omega) = -\eta_\sigma^2 S_m^\sigma e^{-\tau_m^\sigma - \tau_a^\sigma}, \quad (6.22)$$

where the assumptions $|\tau_m^\sigma|$, $|\tau_a^\sigma|$, $|\tau_m^\sigma - \tau_a^\sigma| \gg 1$ have been made and the subscript 'a' designates quantities in the absorption region.

The quantities S_m^σ , τ_m^σ , τ_a^σ and the thickness of the amplification and absorption regions all depend on the actual spectrum of microturbulence in electron plasma waves. Since this is not known, only estimates can be made. Let us assume that background magnetic field is weak ($\omega_p \gg \Omega_e$) and the differences due to the magnetoionic character of the waves can be neglected in a first approximation. Estimating the function given by (6.16), one finds

$$\frac{\mu^\sigma}{v_g^\sigma} \approx 10^{-1} \frac{\omega_p}{c} \frac{v_\phi}{V_e} \frac{W e \ell}{n_{im} V_i^2}, \quad (6.23)$$

where v_ϕ is the phase velocity of the electron plasma waves. On the assumption that equation (6.23) is numerically approximately equal for both the amplification region and the absorption region, except for a change in sign, net amplification of radiation would result if the thickness of the amplification layer ℓ_m is greater than the thickness of the absorption layer ℓ_a .

From equation (6.19), the source function can be estimated to be

$$S_m^\sigma \approx -\frac{4}{(2\pi)^3} \frac{\omega_p^2 m_e V_e^2}{c^2} \frac{v_\phi}{V_i}. \quad (6.24)$$

The equivalent brightness temperature of the radiation on emerging from the source can be found from equations (6.11), (6.20) - (6.24):

$$T_{bs} = -\frac{4\pi^3 c^3}{\eta^2 \omega^2 \kappa} \eta^2 S_m \exp\{-\tau_m - \tau_a\}$$

$$= \frac{2m_e v_e^2}{\kappa} \frac{v_\phi}{V_i} \exp\left\{10^{-1} \frac{\omega_p}{c} \frac{v_\phi}{V_e} \frac{W^{el}}{n_i m_i V_i^2} (\ell_m - \ell_a)\right\}, \quad (6.25)$$

On taking $v_\phi/V_e \approx 10$ and $T_i \approx T_e \approx 2 \times 10^6$ °K, equation (6.25) reads

$$T_{bs} \approx 3.4 \times 10^9 \exp\left\{0.06 W^{el} \frac{\omega_p}{n_e} (\ell_m - \ell_a)\right\}. \quad (6.26)$$

Hence to obtain a source brightness temperature of 10^{16} °K, one requires, from the previous equation,

$$W^{el}(\omega_p/n_e)(\ell_m - \ell_a) = 16.6 \ln(3 \times 10^6) \approx 3.63 \times 10^2 \text{ erg cm sec}^{-1}. \quad (6.27)$$

The assumption that $\ell_m - \ell_a \approx 10^8$ cm gives the condition

$W^{el} \omega_p/n_e \approx 3.63 \times 10^{-6} \text{ erg sec}^{-1}$. The energy densities in electron plasma waves required to give rise to radiation with a source brightness temperature of 10^{16} °K for assumed parameters at various plasma levels in the solar corona are given in Table 6.3.

MHz	200	169	100	80	50	27
$W^{el} (10^{-7} \text{ erg cm}^{-3})$	1.4	1.2	0.74	0.61	0.36	0.18

Table 6.3: Energy densities in electron plasma waves required to give rise to intense type I bursts, at various plasma levels.

The average intensity of the radiation at the second harmonic can be found from equations (4.7), (5.5) and (5.6) and it can be shown to be (Smith, 1970)

$$I(2\omega_p)\Delta\omega \approx 4\pi \times 10^{-2} \omega_p (W^{el})^2 (\ell_m + \ell_a)/n_e m_e c^2, \quad (6.28)$$

where $\Delta\omega$ is the bandwidth of the radiation. From equation (6.11), this intensity corresponds to a source brightness temperature,

$$T_{bs} = \frac{(2\pi)^4 \times 10^{-2}}{3} \frac{(W^{el})^2 (\ell_m + \ell_a)}{\kappa_m n_e \omega_p \Delta\omega} . \quad (6.29)$$

On assuming $\Delta\omega/\omega_p \approx 0.04$, $\ell_m + \ell_a \approx 2 \times 10^8$ cm, it is possible to estimate the brightness temperatures of the second harmonic radiation for the same energy densities in electron plasma waves which give rise to radiation in the fundamental at 10^{16} °K at the source. This is done for various *plasma levels* in the solar corona in Table 6.4.

MHz	200	169	100	80	50	27
T_{bs} (°K)	5.2×10^{12}	7.5×10^{12}	2.3×10^{13}	3.8×10^{13}	8.6×10^{13}	2.6×10^{14}

Table 6.4: Source brightness temperatures of the second harmonic radiation at various plasma levels.

Thus it appears that the source brightness temperatures of the second harmonic radiation are about 2-4 orders of magnitude smaller than that of the fundamental radiation, for the same energy densities in electron plasma waves.

The total energy in non-thermal electron plasma waves involved, if the source area is 10^{18} cm², can be estimated to be about 10^{19} ergs.

If the energy densities in non-thermal electron plasma waves are derived from interactions with an electron stream with speed $u_s \approx 10 V_e$, then we have

$$W^{el} \approx n'_m u_s \Delta u_s , \quad (6.30)$$

where Δu_s describes the amount of "flattening" of the stream distribution due to beam-plasma interactions. Taking $W^{el} \approx 10^{-7}$ erg cm⁻³ and

$\Delta u_s \approx 0.1 V_e$, one finds the number density of electrons in the stream n' to be about 200 cm^{-3} .

It is now justified *a posteriori* that collisional effects can be neglected in the above treatment of the nonlinear processes. The conversion of electron plasma waves to magnetoionic waves by nonlinear scattering off thermal plasma ions generally occurs in a time longer than, or at most comparable to, the scattering time of electron plasma waves into themselves (Tsytovich, 1970). From equation (5.44), it is seen that the nonlinear conversion time τ (for $e\ell \rightarrow \sigma$), with $W^{e\ell} \approx 10^{-7} \text{ erg cm}^{-3}$ and $T_e \approx T_i \approx 2 \times 10^6 \text{ }^\circ\text{K}$, can be estimated by

$$1/\tau \approx f_p \quad (\text{MHz}) \quad (6.31)$$

In the region of the solar corona of interest, we have $1/\tau_c \approx 0.27 \nu_c$ with the effective collision frequency ν_c decreasing from 50 sec^{-1} at $R/R_\odot = 0.1$ to 0.4 sec^{-1} at $R/R_\odot = 1.5$. It can be seen that the condition $\tau \ll \tau_c$ is generally satisfied and the neglect of collisions has been justified.

Finally, the weak nonlinear condition given by equation (1.2), can be written in the form

$$W^{e\ell} \ll (\omega/\omega_p)^2 W_p, \quad (6.32)$$

where ω is the (angular) frequency of the microturbulence and W_p is the kinetic energy density of the plasma particles. Since one has $W^{e\ell} \approx 10^{-7} \text{ erg cm}^{-3}$, $\omega \approx \omega_p$, and $W_p \approx n_e k T_e \approx 10^{-2} \text{ erg cm}^{-3}$, the weak nonlinearity assumption has also been justified.

(b) Polarization

As suggested by Fung and Yip (1966) and Takakura (1963), the strong circular polarization of type I bursts in the sense of the

O-mode might well be explained by

(i) Radiation occurring at frequencies below the cutoff frequency of the X-mode waves.

(ii) Strong preferential gyroresonance absorption of the X-mode waves at levels $Y = \frac{1}{2}$ and $\frac{1}{3}$.

However, at high photospheric altitudes and also during the initial stages of development of a noise storm at lower altitudes, one might reasonably expect the radiation to come from regions where $\omega_p > 3\Omega_e$. Moreover, when the frequencies of the radiation are above the cutoff frequency of the X-mode waves, i.e., $\omega > \omega_X$ (see the reverse of inequality (5.11)), any strong polarization of type I bursts in the sense of the O-mode that is observed must be attributed to causes other than those stated above. For such cases, it is postulated here that the strong polarization in the sense of the O-mode is due to *preferential amplification* of the O-mode waves in the source region.

On the assumption that the magnetoionic waves are uncoupled in the source for certain distributions of non-thermal electron plasma waves and for some angles of propagation of the radiation, amplification (and absorption) of O-mode waves is preferred (see e.g., chapter 5). This leads to small differences in optical thicknesses of the O-mode and X-mode waves. However, the intensity of radiation emerging from the source region is dependent exponentially on the optical thickness. The resultant intensity of the O-mode radiation can therefore be many times larger than that of the X-mode radiation, i.e., the net degree of polarization in the sense of the O-mode can be very high. For simplicity, let us illustrate this by the case $Y \ll 1 - X \ll 1$, which is applicable to the case of a very weak background magnetic field.

Taking care to include differences due to different magnetoionic modes, one obtains, from a slightly modified form of equation (6.14), the total optical thickness,

$$\tau^\sigma = - \left(\frac{\pi}{2} \right)^{\frac{1}{2}} \frac{\omega_p W^{el}(\lambda_m - \lambda_a)}{2c n_e m_e v_e^2 (1 + Z_1 T_e / T_i)^2} \frac{\partial(\omega \eta_\sigma) / \partial \omega}{(1 + \delta_\sigma^2)^{\frac{1}{2}}} F^{\sigma el}(\omega, \theta, \theta'), \quad (6.33)$$

where $F^{\sigma el}(\omega, \theta, \theta')$ is given by equation (5.3). On taking $\omega_p W^{el}(\lambda_m - \lambda_a) / n_e = 3.63 \times 10^2 \text{ erg cm sec}^{-1}$, from equation (6.27), $Z_1 = 1$, and $T_e \approx T_i \approx 2 \times 10^6 \text{ }^\circ\text{K}$, the total optical thickness reads

$$\tau^\sigma = -6.86 H^\sigma(\omega, \theta, \theta'), \quad (6.34)$$

where $H^\sigma(\omega, \theta, \theta')$ is defined by

$$H^\sigma(\omega, \theta, \theta') \equiv \frac{\{\sin^2 \theta' (\cos^2 \theta + T_\sigma^2) + 2(\cos^2 \theta - T_\sigma^2)^2 \cos^2 \theta' / \sin^2 \theta\}}{2\eta_\sigma \{(1 + T_\sigma^2)^2 + K_\sigma^2 T_\sigma^2\}^{\frac{1}{2}} (T_\sigma - Y \cos \theta)^2}. \quad (6.35)$$

Since $K_\sigma^2 = 0 \{Y^2 / (1-X)^2\} \ll 1$ and for $\frac{1}{2} Y \sin^2 \theta \ll (1-X) \cos \theta$,

$$T_\sigma \approx \mp \frac{\cos \theta}{|\cos \theta|} \left[1 \pm \frac{Y \sin^2 \theta}{2(1-X) |\cos \theta|} \right], \quad (6.36)$$

it can be found, after expansions in the parameter $Y/(1-X)$, that

$$H^\sigma(\omega, \theta, \theta') = \frac{\Theta(\theta, \theta')}{4(1-X)} \left\{ 1 \pm \frac{Y \Phi(\theta, \theta')}{(1-X) \Theta(\theta, \theta')} \right\}, \quad (6.37)$$

where

$$\Theta(\theta, \theta') \equiv \sin^2 \theta' (1 + \cos^2 \theta) + 2 \sin^2 \theta \cos^2 \theta' \quad (6.38)$$

and

$$\Phi(\theta, \theta') \equiv \{\sin^2 \theta (1 + 3 \cos^2 \theta') - (1 + \sin^2 \theta / 2) \Theta(\theta, \theta')\} / |\cos \theta|. \quad (6.39)$$

On neglecting small differences between the O-mode and X-mode waves in the source function, the degree of circular polarization in the sense of the O-mode, for radiation emerging from the source region can be estimated by

$$r_c = \{\exp(-\tau^O + \tau^X) - 1\} / \{\exp(-\tau^O + \tau^X) + 1\}, \quad (6.40)$$

where τ^0 and τ^X are given by equation (6.34) with $H^\sigma(\omega, \theta, \theta')$ determined by equation (6.37). Again from equations (6.34), (6.37), (6.38) and (6.39) the exponent reads

$$-\tau^0 + \tau^X = 3.43 Y \Phi(\theta, \theta') / (1-X)^2. \quad (6.41)$$

For illustration, take $Y=0.01$, $1-X=0.04$, $\theta'=0$ and $\theta=45^\circ$. In this case, $-\tau^0 + \tau^X = 22.8$, hence $r_c \approx 1$, i.e., the emergent radiation is fully polarized in the sense of the O-mode. Again, take $Y=0.01$, $1-X=0.04$, but $\theta'=30^\circ$ and $\theta=35^\circ$, one finds $-\tau^0 + \tau^X = 0.23$, hence $r_c \approx 12\%$, i.e., the emergent radiation is weakly polarized in the sense of the O-mode.

The above discussion of the polarization of radiation arising from a "plasma emission" process is semi-quantitative only, since the turbulence spectrum of electron plasma waves must be known to determine the optical thicknesses precisely. But it has been argued that preferential amplification of the O-mode waves, even in a weak background magnetic field, can give rise to strongly polarized type I bursts.

7. DISCUSSION

Including induced effects, a model of type I bursts based on the "plasma hypothesis" has been proposed. The physical parameters of the source, such as its dimensions and its temperature, have been chosen in agreement with the available observational data. The energy requirements deduced from the model can easily be met under the known conditions prevailing in the solar corona. An electron stream with speed $u_s \approx 10 V_e$ and number density 200 cm^{-3} has been shown to be capable of exciting a microturbulence in electron plasma waves with energy density of about $10^{-7} \text{ erg cm}^{-3}$, which is required to explain a strong type I burst.

The preferential amplification of O-mode waves in the source region has been postulated as the mechanism giving rise to strong circular polarization (in the sense of the O-mode) of type I bursts coming from regions where the background magnetic field is weak $Y \ll 1$. This mechanism rests on the assumption that coupling between O-mode and X-mode waves is weak in the source region. It can be shown (Cohen, 1960) that in the solar corona, mode coupling is weak for magnetoionic waves propagating in the QL-approximation under the conditions where (i) $X \ll 1$, (ii) $Y \ll 1$, and (iii) collisional effects are negligible. Mollwo (1970) has considered, in the solar corona, mode coupling of magnetoionic waves propagating close to the direction of the background magnetic field and under the assumption (i) $X \approx 1$ and (ii) $Y \approx 1$, and he shows that mode-coupling can be important. In the present context, it appears that the problem of mode-coupling under the conditions (i) $X \approx 1$ and (ii) $Y \ll 1$ needs to be considered.

Although Takakura's (1963) explanation for the duration of type I bursts has been questioned in section 5, no alternative is offered here, since no satisfactory explanation has been found. Scattering by coronal inhomogeneities can give rise to a duration of 0.2-0.6 sec at most (Steinberg *et al.*, 1970; Riddle, 1972) for an impulsive source and this appears too short in comparison with duration of type I bursts at high source altitudes.

The mechanism of accelerating electron streams responsible for type I bursts and the phenomenon of chains of bursts might well be connected with collisionless shock waves propagating parallel to the background magnetic field in the low- β coronal plasma. Parallel turbulent shocks have been considered by Parker (1961) and Kennel and Sagdeev (1967), but however, not in connection with particle acceleration and

non-thermal radiation. Such problems lie outside the scope of this thesis.

Finally, it is suggested here that the background continuum radiation of type I noise storm is probably not merely a superposition of type I bursts. Apart from the known observational evidence (see Kundu, 1965; Chernov *et al.*, 1972) in support of this view, a point which has not been stated before is this. The background continuum radiation usually has a bandwidth of about 100 MHz. If type I bursts were to give rise to this radiation, they must occupy a layer between two plasma levels whose frequency separation is about 100 MHz. Between the 200 MHz and the 100 MHz say, the thickness of the layer is about $0.25 R_{\odot}$ (see Fig. 6.4). Such a distance on the solar surface would subtend an angle of ~ 25 mins of arc at the earth. Since the source is probably disk-shape and since the angular size of a typical noise storm is usually not more than 10 mins of arc (even including the effects of coronal scattering), it appears that the required radial thickness of the source is much larger than reasonable expectations from observations. Hence a different radiation mechanism seems to be needed to account for the background continuum radiation of type I noise storms. Such a mechanism might be coherent gyromagnetic radiation (Fung and Yip, 1966) or the mechanism proposed by Friedman (1972).

CHAPTER 7

CONCLUSIONS

In this chapter, a synopsis of the theoretical and astrophysical results found in the previous chapters is presented. Other possible applications of the theory and suggestions for further research are briefly stated.

1. REVIEW OF RESULTS

(a) A general theory of weak nonlinear processes in a magneto-active plasma has been developed in the semiclassical formalism (chapters 2 and 4). For the specific examples considered, the following conclusions have been reached.

(i) On neglecting shielding effects, Thomson scattering of magnetoionic waves by non-relativistic electrons is determined by the zeroth harmonic scattering; first and higher harmonic scatterings are not important.

(ii) In the presence of a strong background magnetic field, under the conditions $\omega_p \ll \omega \ll \Omega_e$, Thomson scattering of magnetoionic waves by non-relativistic electrons in general has a reduced cross-section, being about ω^2/Ω_e^2 times the classical Thomson cross-section (save in a special case).

(iii) When shielding effects of a non-relativistic plasma are included, nonlinear scattering by thermal ions is the dominant scattering process. When the conditions $\omega_p \ll \omega \ll \Omega_e$ are satisfied, the

cross-section is reduced generally to about ω^2/Ω_e^2 times the classical Thomson cross-section (save in a special case).

(iv) The conditions under which the motion of relativistic electrons can be taken to be rectilinear in treating inverse Compton scattering have been derived. Such conditions are not satisfied in the case of inverse Compton scattering of magnetoionic waves by relativistic electrons moving in the direction of the background magnetic field. The results for this case are summarized by equations (4.46), (4.52) and Table 4.1. Under the simplifying assumptions $\omega' \gg \gamma\Omega_e \geq \omega_p$, the inverse Compton scattering of O-mode magnetoionic waves propagating perpendicular to the background magnetic field is enhanced by at least a factor $\gamma^2 \gg 1$ in comparison with all the other cases considered.

(b) A general theory of polarized radiation arising from micro-turbulences in electron plasma waves and electron cyclotron waves has been presented (chapter 5). Radiation arising from any one of the processes discussed is 100% polarized in the sense of the O-mode at the source if it is emitted in the frequency range $\omega_p < \omega < \omega_X$, where ω_X is the cutoff frequency of the X-mode waves. Bearing this clearly in mind, one reaches the following conclusions for the specific cases considered.

(i) For the case of electron plasma waves propagating nearly along the lines of the background magnetic field, radiation at the fundamental and radiation at the second harmonic have net degrees of polarization at the source in the sense of the O-mode. In a weak background magnetic field ($\omega_p \gg \Omega_e$), the second harmonic is always weakly polarized whilst the fundamental can be strongly or weakly polarized in the sense of the O-mode.

(ii) The radiation at $\omega \approx s\Omega_e$ ($s \geq 2$) arising from the non-linear conversion of electrostatic electron cyclotron waves has

generally a net degree of polarization in the sense of the X-mode.

(c) In chapter 6, an updated review of observations on type I solar radio bursts and their associated noise storm phenomena has been presented. Existing theories have been examined and their deficiencies have been indicated.

(i) Type I bursts have been explained on a model which is based on the "plasma hypothesis" and which includes previously neglected induced effects. The physical parameters assumed are consistent with the observational data.

(ii) The preferential amplification of O-mode waves has been shown to be capable of explaining the strong circular polarization (in the sense of the O-mode) of type I bursts from source regions where the background magnetic field is weak.

2. SUGGESTIONS FOR FURTHER RESEARCH

(i) To obtain a fully quantitative theory of type I bursts based on the "plasma hypothesis", one needs to solve the problem of formation of quasi-stationary turbulence spectra arising from the interaction of a suprathermal, non-relativistic electron stream with a magnetoactive plasma.

(ii) A justification of the mechanism of preferential amplification of one mode or the other, requires the solution of the problem of magnetoionic mode coupling in a region where $X \approx 1$ and $Y \ll 1$.

(iii) In view of the similarity of the radiation mechanism invoked for type I, type II, and type III solar radio bursts, one should attempt to explain the different polarization characteristics of these bursts.

(iv) The duration of type I solar radio bursts appears to be inadequately explained at present.

(v) The continuum radiation of type I noise storm is similar to stationary type IV B radiation in many respects. Its mechanism of radiation, together with that of stationary type IV B, needs further investigation.

(vi) The acceleration of electron streams responsible for type I bursts and the phenomenon of chains of type I bursts might well be connected with microturbulent, collisionless shock waves propagating parallel to the background magnetic field. Since parallel shocks may be important in type II bursts as well (Wild and Smerd, 1972), they deserve investigation.

(vii) The theory presented has been sufficiently general to be easily adaptable to other astrophysical and magnetospheric applications. It is straightforward for example to consider "whistlers" and evaluate the significance of their nonlinear interactions in the ionosphere.

(viii) The emission from electrostatic electron cyclotron waves in the earth's magnetosphere appears to be an important component of the radiation detected by airborne satellites (Kennel *et al.*, 1970).

(ix) The decametric radiation from Jupiter, identified as in the X-mode, has been attributed to coherent gyromagnetic radiation by electrons in the Jovian magnetosphere (Goldreich and Lynden-Bell, 1969). In view of the theoretical results found in this thesis, and in view of the large number of instabilities giving rise to electrostatic electron cyclotron waves, radiation by the nonlinear conversion of electrostatic electron cyclotron waves should be considered as a possible candidate radiation mechanism.

APPENDIX A

CALCULATION OF RESPONSE TENSORS

The infinite sums containing Bessel functions arise from power series expansions of the exponentials $e^{P(\phi')}$, $e^{-P(\phi)}$, etc. From (2.39), one has

$$P(\phi) = \frac{i\varepsilon_\alpha}{\Omega_\alpha} (\omega - k_{\parallel} v_{\parallel}) \phi - i\varepsilon_\alpha z \sin(\phi - \psi). \quad (\text{A.1})$$

From the definition of Bessel functions (Watson, 1944), one has

$$\exp\{-i\varepsilon_\alpha z \sin(\phi - \psi)\} = \sum_{s=-\infty}^{\infty} J_s(z) \exp\{-i\varepsilon_\alpha s(\phi - \psi)\}, \quad (\text{A.2})$$

and

$$v_{\perp} \exp\{-i\varepsilon_\alpha z \sin(\phi - \psi)\} = \sum_{s=-\infty}^{\infty} V_{\perp}^*(s, \underline{k}, \underline{v}) \exp\{-i\varepsilon_\alpha s(\phi - \psi)\}, \quad (\text{A.3})$$

where $V_{\perp}(s, \underline{k}, \underline{v})$ is given by (2.46). Substitution of (A.2) and (A.3) into (A.1) gives,

$$e^{-P(\phi)} = \sum_{s=-\infty}^{\infty} J_s(z) \exp\left\{-\frac{i\varepsilon_\alpha \omega_s}{\Omega_\alpha} \phi - i\varepsilon_\alpha \psi\right\}, \quad (\text{A.4})$$

and

$$v_{\perp} e^{-P(\phi)} = \sum_{s=-\infty}^{\infty} V_{\perp}(s, \underline{k}, \underline{v}) \exp\left\{-\frac{i\varepsilon_\alpha \omega_s}{\Omega_\alpha} \phi - i\varepsilon_\alpha \psi\right\}, \quad (\text{A.5})$$

where

$$\omega_s \equiv \omega - s\Omega_\alpha - k_{\parallel} v_{\parallel}. \quad (\text{A.6})$$

In cylindrical coordinates, one has Cartesian components,

$$\frac{\partial}{\partial \underline{p}} = \left(\cos\phi \frac{\partial}{\partial p_{\perp}} - \frac{\sin\phi}{p_{\perp}} \frac{\partial}{\partial \phi}, \sin\phi \frac{\partial}{\partial p_{\perp}} + \frac{\cos\phi}{p_{\perp}} \frac{\partial}{\partial \phi}, \frac{\partial}{\partial p_{\parallel}} \right). \quad (\text{A.7})$$

Application of (A.4) - (A.7), together with (2.41), gives

$$e^{P(\phi')} g_{sj}(\underline{v}', \underline{k}, \omega) \frac{\partial}{\partial p_{s'}} = \frac{q_\alpha}{\omega} \sum_{s'=-\infty}^{\infty} \exp \left\{ \frac{i\varepsilon_\alpha \omega s'}{\Omega_\alpha} \phi' + i\varepsilon_\alpha s' \psi \right\} \\ \times \left\{ \alpha_j^*(s', \underline{k}, \omega) \frac{\partial}{\partial p_\perp} + \beta_j^*(s', \underline{k}, \omega) \frac{1}{p_\perp} \frac{\partial}{\partial \phi'} + \gamma_j^*(s', \underline{k}, \omega) \frac{\partial}{\partial p_\parallel} \right\}, \quad (A.8)$$

$$e^{P_1(\phi')} g_{sj}(\underline{v}', \underline{k}_1, \omega_1) \frac{\partial}{\partial p_{s'}} = \frac{q_\alpha}{\omega_1} \sum_{\mu'=-\infty}^{\infty} \exp \left\{ \frac{i\varepsilon_\alpha \omega_1 \mu'}{\Omega_\alpha} \phi' + i\varepsilon_\alpha \mu' \psi_1 \right\} \\ \times \left\{ \alpha_j^*(\mu', \underline{k}_1, \omega_1) \frac{\partial}{\partial p_\perp} + \beta_j^*(\mu', \underline{k}_1, \omega_1) \frac{1}{p_\perp} \frac{\partial}{\partial \phi'} + \gamma_j^*(\mu', \underline{k}_1, \omega_1) \frac{\partial}{\partial p_\parallel} \right\}, \quad (A.9)$$

and

$$e^{P_2(\phi'')} g_{rl}(\underline{v}'', \underline{k}_2, \omega_2) \frac{\partial}{\partial p_{r''}} = \frac{q_\alpha}{\omega_2} \sum_{v'=-\infty}^{\infty} \exp \left\{ \frac{i\varepsilon_\alpha \omega_2 v'}{\Omega_\alpha} \phi'' + i\varepsilon_\alpha v' \psi_2 \right\} \\ \times \left\{ \alpha_l^*(v', \underline{k}_2, \omega_2) \frac{\partial}{\partial p_\perp} + \beta_l^*(v', \underline{k}_2, \omega_2) \frac{1}{p_\perp} \frac{\partial}{\partial \phi''} + \gamma_l^*(v', \underline{k}_2, \omega_2) \frac{\partial}{\partial p_\parallel} \right\}. \quad (A.10)$$

On substitution of (A.5), (2.37) reads,

$$\varepsilon_{ij}(\underline{k}, \omega) = \delta_{ij} + \frac{4\pi i}{\omega} \sum_\alpha \frac{|q_\alpha| n_\alpha}{\Omega_\alpha} \int_0^\infty p_\perp dp_\perp \int_{-\infty}^\infty dp_\parallel \sum_{s,s'=-\infty}^\infty \frac{q_\alpha}{\omega} \int_0^{2\pi} d\phi \\ \times \left\{ V_i(s, \underline{k}, \underline{v}) \exp \left[-\frac{i\varepsilon_\alpha \omega s}{\Omega} \phi - i\varepsilon_\alpha s \psi \right] \int^\phi d\phi' \exp \left[\frac{i\varepsilon_\alpha \omega s'}{\Omega_\alpha} \phi' + i\varepsilon_\alpha s' \psi \right] \right. \\ \left. \times \left[\alpha_j^*(s', \underline{k}, \omega) \frac{\partial f_\alpha^{(0)}}{\partial p_\perp} + \gamma_j^*(s', \underline{k}, \omega) \frac{\partial f_\alpha^{(0)}}{\partial p_\parallel} \right] \right\}. \quad (A.11)$$

Carrying out the trivial integrations over azimuthal angles ϕ' and ϕ and using well-known results on summation of Bessel function, e.g.

$$\sum_{s=-\infty}^\infty V_i(s, \underline{k}, \underline{v}) J_s(z) = v_\parallel \delta_{i3}, \quad (A.12)$$

one derives the dielectric tensor in the form (2.42).

Similarly, on application of (A.4), (A.5), (A.9) and (A.10), equation (2.38) reads,

$$\begin{aligned}
\varepsilon_{ij\ell}(\underline{k}_1, \omega_1; \underline{k}_2, \omega_2) &= \sum_{\alpha} \frac{4\pi i q_{\alpha}^3 n_{\alpha}}{\omega \omega_1 \omega_2 \Omega_{\alpha}^2} \int_{-\infty}^{\infty} dp_{\parallel} \int_0^{\infty} p_{\perp} dp_{\perp} \sum_{s, \mu, \mu', \nu, \nu'} \int_0^{2\pi} d\phi \\
&\times V_i(s, \underline{k}, \nu) \exp \left\{ -\frac{i\varepsilon_{\alpha} \omega s}{\Omega_{\alpha}} \phi - i\varepsilon_{\alpha} s \psi \right\} \int^{\phi} d\phi' J_{\mu}(z_2) \\
&\times \exp \left\{ \frac{i\varepsilon_{\alpha} \omega_{2\mu}}{\Omega_{\alpha}} \phi + i\varepsilon_{\alpha} \mu \psi_2 \right\} \exp \left\{ \frac{i\varepsilon_{\alpha} \omega_{1\mu'}}{\Omega_{\alpha}} \phi' + i\varepsilon_{\alpha} \mu' \psi_1 \right\} \left\{ \alpha_j^*(\mu', \underline{k}_1, \omega_1) \right. \\
&\times \left. \frac{\partial}{\partial p_{\perp}} + \beta_j^*(\mu', \underline{k}_1, \omega_1) \frac{1}{p_{\perp}} \frac{\partial}{\partial \phi'} + \gamma_j^*(\mu', \underline{k}_1, \omega_1) \frac{\partial}{\partial p_{\parallel}} \right\} J_{\nu}(z_2) \\
&\times \exp \left\{ -\frac{i\varepsilon_{\alpha} \omega_{2\nu}}{\Omega_{\alpha}} \phi' - i\varepsilon_{\alpha} \nu \psi_2 \right\} \int^{\phi'} d\phi'' \exp \left\{ \frac{i\varepsilon_{\alpha} \omega_{2\nu'}}{\Omega_{\alpha}} \phi'' + i\varepsilon_{\alpha} \nu' \psi_2 \right\} \\
&\times \left\{ \alpha_{\ell}^*(\nu', \underline{k}_2, \omega_2) \frac{\partial}{\partial p_{\perp}} + \gamma_{\ell}^*(\nu', \underline{k}_2, \omega_2) \frac{\partial}{\partial p_{\parallel}} \right\} f_{\alpha}^{(0)}(p_{\perp}, p_{\parallel}) .
\end{aligned}$$

Again, trivial integrations over azimuthal angles ϕ'' , ϕ' and ϕ give the result (2.43).

APPENDIX B

SUMMATION FORMULA (2.96)

On account of $\zeta_z(\lambda) \equiv e^{-\lambda} I_z(\lambda)$ remaining finite as $|z| \rightarrow \infty$, it follows from Cauchy's theorem that

$$\oint_c \cot \pi z \frac{\zeta_z(\lambda)}{z-q} dz = 0, \quad (B.1)$$

where c is a circular contour with radius $R \rightarrow \infty$. On evaluating the integral in (B.1) by the residue theorem and using the identity,

$$\pi \cot \pi z \equiv \sum_{s=-\infty}^{\infty} \frac{1}{z-s}, \quad (B.2)$$

it is easily shown that

$$\sum_{s=-\infty}^{\infty} \frac{\zeta_s(\lambda)}{q-s} = \pi \cot \pi q \zeta_q(\lambda). \quad (B.3)$$

Application of this result together with the simple algebraic identity,

$$\frac{s^2}{s^2-q^2} \equiv 1 - \frac{q}{2} \left(\frac{1}{q+s} + \frac{1}{q-s} \right), \quad (B.4)$$

gives the relation (2.96) trivially.

APPENDIX C

CALCULATION OF CURRENT DENSITIES DUE TO A CHARGE SPIRALLING IN $\underline{B}_0$

To derive the current density (3.13), it is noted that the solution of the relativistic equation for a charge spiralling in a background magnetic field $\underline{B}_0 = B_0 (0,0,1)$,

$$\frac{d\underline{p}}{dt} = \frac{q \underline{v}(t) \times \underline{B}_0}{c}, \quad (C.1)$$

can be written as

$$\underline{r}(t) = (R \sin\{\Omega^* t + \phi\}, \epsilon R \cos\{\Omega^* t + \phi\}, v_{\parallel} t), \quad (C.2)$$

where $R \equiv v_{\perp}/\Omega^*$, $\epsilon = q/|q|$, $\Omega^* = \Omega mc^2/E$ and the parameters ϕ , $v_{\perp}$, and $v_{\parallel}$ are determined by initial conditions.

The current density due to the motion of a charge reads

$$\underline{j}_i(\underline{r}, t) = q v_i(t) \delta\{\underline{r} - \underline{r}(t)\}, \quad (C.3)$$

the Fourier transform of which, with $\underline{r}(t)$ and $\underline{v}(t)$ determined by (C.2), reduces to

$$\underline{j}_i(\underline{k}, \omega) = \sum_{s=-\infty}^{\infty} q \int dt V_i(s, \underline{k}, \underline{v}) \exp\{i(\omega - s\Omega^* - k_{\parallel} v_{\parallel})t - is(\phi + \epsilon\psi)\}, \quad (C.4)$$

where $V_i(s, \underline{k}, \underline{v})$ is defined by (2.46) and equation (A.3) has been used. Final integration with respect to t gives the result (3.13).

On the other hand, when $\underline{k}$ is replaced by $\Delta\underline{k} = \underline{k} - \underline{k}'$ and ω is replaced by $\Delta\omega = \omega - \omega'$, instead of (3.13), it is possible to obtain an alternative form of the current density by making a different Fourier transform on (C.3) and using (A.2) and (A.3). After straightforward calculations, the current density reads

$$j_{\perp}(\Delta \underline{k}, \Delta \omega) = \sum_{s, s'=-\infty}^{\infty} 2\pi q V_{\perp}(s, \underline{k}, \underline{v}) J_{-s'}(z') e^{-i\varepsilon(s\psi + s'\psi)} \\ \times \delta\{\Delta\omega - (s+s')\Omega^* - \Delta k_{\parallel} v_{\parallel}\}, \quad (C.5)$$

where the initial conditions are chosen such that $\phi = 0$, and as usual

$$z' = k_{\perp}' v_{\perp} / \Omega^*, \text{ etc.}$$

APPENDIX D

CALCULATION OF CURRENT DENSITIES IN THOMSON AND NONLINEAR SCATTERINGS

In this appendix, the current density in Thomson scattering given by (4.9) with (4.10) to (4.13) and the current density in nonlinear scattering given by (4.14) with (4.15) are derived.

The equation of motion of a spiralling charge in the presence of an arbitrary perturbing force $\underline{F}(\underline{v}, \underline{r}, t)$ reads

$$\frac{d\underline{p}(t)}{dt} = \frac{q \underline{v}(t) \times \underline{B}_0}{c} + \underline{F}(\underline{v}, \underline{r}, t) . \quad (D.1)$$

Expanding in powers of $\underline{F}$, we write

$$\underline{r}(t) = \underline{r}^{(0)}(t) + \underline{r}^{(1)}(t) + \dots . \quad (D.2)$$

The current density which is proportional to $\underline{F}$ can be written as

$$\underline{j}^{(1)}(\underline{k}, \omega) = q \int dt \left\{ \frac{d\underline{r}^{(1)}(t)}{dt} - i \underline{k} \cdot \underline{r}^{(1)}(t) \frac{d\underline{r}^{(0)}(t)}{dt} \right\} e^{i(\omega t - \underline{k} \cdot \underline{r}^{(0)}(t))} , \quad (D.3)$$

where $\underline{r}^{(0)}(t)$ is determined by (C.1). Since the choice of initial conditions appears only in an unimportant phase factor in $\underline{j}^{(1)}(\underline{k}, \omega)$, the initial conditions are chosen such that

$$\underline{r}^{(0)}(t) = (R \sin \Omega^* t, R \cos \Omega^* t, v_{\parallel} t) , \quad (D.4)$$

where $R \equiv v_{\perp}/R^*$, $\epsilon = q/|q|$, $\Omega^* = \Omega/\gamma$, $\gamma^2 = (1 - v_{\perp}^2/c^2 - v_{\parallel}^2/c^2)$ and, $v_{\perp}$ and $v_{\parallel}$ are velocity components in the absence of the perturbing force $\underline{F}$.

From (D.1), the equation of motion to first order in $\underline{F}$ reads

$$\frac{d\underline{v}^{(1)}}{dt} = \frac{q \underline{v}^{(1)} \times \underline{B}_0}{m\gamma c} + \frac{\underline{F}(t)}{m\gamma} - \frac{1}{m\gamma} \frac{d}{dt} \left(\frac{\underline{v}^{(0)}(t)}{c^2} \int^t \underline{v}^{(0)}(t') \cdot \underline{F}(t') dt' \right) , \quad (D.5)$$

where $\tilde{F}(t) \equiv \tilde{F}(\underline{v}^{(0)}(t), \underline{r}^{(0)}(t), t)$. From (D.5), it is straightforward to show

$$r_i^{(1)}(t) = \frac{1}{m\gamma} \int^t dt' \int^t dt'' T_{ij}(t'-t'') \left\{ F_j(t'') - \frac{d}{dt''} \left[\frac{v_j^{(0)}(t'')}{c^2} \int^{t''} \underline{v}^{(0)}(t''') \cdot \tilde{F}(t''') dt''' \right] \right\}, \quad (D.6)$$

where

$$T_{ij}(t) \equiv \begin{pmatrix} \cos \Omega^* t & \sin \Omega^* t & 0 \\ -\sin \Omega^* t & \cos \Omega^* t & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (D.7)$$

Identifying $\tilde{F}$ with the force due to fluctuating electromagnetic fields, we have

$$\tilde{F}(\underline{v}, \underline{r}, t) = q \left\{ \underline{E}(\underline{r}, t) + \frac{\underline{v} \times \underline{B}(\underline{r}, t)}{c} \right\}, \quad (D.8)$$

and on using

$$F_i(\underline{v}, \underline{r}, t) = q \int \frac{d^3 \underline{k}}{(2\pi)^4} \frac{d\omega}{\omega} \{ (\omega - \underline{k} \cdot \underline{v}) \delta_{ij} + k_i v_j \} E_j(\underline{k}, \omega) e^{i(\underline{k} \cdot \underline{r} - \omega t)}, \quad (D.9)$$

$$\int^t T_{ij}(t-t') e^{-i\omega t'} dt' = i\tau_{ij}(\omega) e^{-i\omega t}/\omega, \quad (D.10)$$

$$\int^t dt' e^{-i\omega t'} \left\{ i \frac{d\underline{v}}{dt} + \omega \underline{v} \right\} dt' = i\underline{v}(t) e^{-i\omega t}, \quad (D.11)$$

and equations (A.2) and (A.3), it is reasonably easy to show

$$r_i^{(1)}(t) = -\frac{q}{m\gamma} \sum_{s=-\infty}^{\infty} \int \frac{d^3 \underline{k}}{(2\pi)^3} \frac{d\omega}{\omega} \frac{e^{-i\omega_s t} e^{i\underline{k} \cdot \underline{r}}}{\omega_s^2} E_j(\underline{k}, \omega) \\ \times \{ \tau_{il}(\omega_s) (\delta_{lj} \omega_s J_s(z) + k_l v_j^*(s, \underline{k}, \underline{v}^{(0)}) - v_i^{(0)}(t) v_j^*(s, \underline{k}, \underline{v}^{(0)})_{\omega/c^2} \}, \quad (D.12)$$

with $\omega_s = \omega - s\Omega^* - k_{\parallel} v_{\parallel}$. This expression, together with equations (D.3) and (D.4) yields (4.9).

By equation (3.7), the shielding field due to the current density (3.13) associated with the unperturbed motion of a charge reads

$$E_i^q(\underline{k}, \omega) = - \frac{8\pi^2 q i}{\omega} \frac{\lambda_{ij}(\underline{k}, \omega)}{\Lambda(\underline{k}, \omega)} \sum_{s=-\infty}^{\infty} e^{-i\epsilon s \psi} V_j(s, \underline{k}, \underline{y}) \delta(\omega - k_{\parallel} v_{\parallel} - s\Omega^*), \quad (D.13)$$

where $\underline{y} = d\underline{r}^{(0)}/dt$ is determined by (D.4). From (2.15) and (2.49), the nonlinear current density resulted from the interaction of the electric field $\underline{E}^{\sigma'}(\underline{k}, \omega)$ of a wave in mode σ' with the shielding field $\underline{E}^q(\underline{k}, \omega)$ given by (D.13), can be written as

$$j_i^{n.s.}(\underline{k}, \omega) = \frac{\omega}{2\pi i} \int \frac{d^3 \underline{k}' d\omega'}{(2\pi)^4} \epsilon_{ijl}(\underline{k}', \omega'; \underline{k} - \underline{k}', \omega - \omega') E_j^{\sigma'}(\underline{k}', \omega') E_l^q(\underline{k} - \underline{k}', \omega - \omega'),$$

from which, equation (4.14) emerges with (4.15).

Again by equation (3.7), the shielding field due to the current density (C.5) associated with the unperturbed motion of a charge reads

$$E_i^q(\Delta \underline{k}, \Delta \omega) = - \sum_{s=-\infty}^{\infty} \frac{8\pi^2 q i}{\Delta \omega} \frac{\lambda_{ij}(\Delta \underline{k}, \Delta \omega)}{\Lambda(\Delta \underline{k}, \Delta \omega)} e^{-i\epsilon s \psi} \delta(\Delta \omega - s\Omega^* - \Delta k_{\parallel} v_{\parallel}) \times \sum_{v=-\infty}^{\infty} V_j(s+v, \underline{k}, \underline{y}) J_v(z') e^{-i\epsilon v(\psi - \psi')}, \quad (D.14)$$

where $\Delta \omega \equiv \omega - \omega'$ and $\Delta \underline{k} = \underline{k} - \underline{k}'$.

The nonlinear current density, corresponding to (4.14), then reads

$$j_i^{n.s.}(\underline{k}, \omega) = \frac{2\pi i q^2}{m} \sum_{s=-\infty}^{\infty} \int \frac{d^3 \underline{k}' d\omega'}{(2\pi)^4 \omega'} A_{ij}^{n.s.}(s; \omega, \underline{k}; \omega', \underline{k}'; \underline{y}) E_j^{\sigma'}(\underline{k}', \omega') \delta\{\omega - \omega' - (k_{\parallel} - k'_{\parallel})v_{\parallel} - s\Omega^*\}, \quad (D.15)$$

where we have

$$A_{ij}^{n.s.}(s; \omega, \underline{k}; \omega', \underline{k}'; \underline{y}) = \sum_{v=-\infty}^{\infty} A_{ij}^{n.s.}(s+v, \underline{k}, \omega; v, \underline{k}', \omega'; \underline{y}), \quad (D.16)$$

with

$$\begin{aligned}
A_{ij}^{n.s.}(s, \underline{k}, \omega; s', \underline{k}', \omega'; \underline{v}) &= \frac{2im\omega'\omega}{q(\omega-\omega')} \varepsilon_{ijl}(\underline{k}', \omega'; \underline{k}-\underline{k}', \omega-\omega') \\
&\times \frac{\lambda_{lm}(\underline{k}-\underline{k}', \omega-\omega')}{\Lambda(\underline{k}-\underline{k}', \omega-\omega')} e^{-i\varepsilon(s\psi-s'\psi')} V_m(s, \underline{k}, \underline{v}) J_{s'}(z') \quad .(D.17)
\end{aligned}$$

APPENDIX E

CONDITIONS FOR SIMPLIFICATION OF THE INVERSE MAXWELLIAN TENSOR

In the regime (4.54), provided

$$|\Delta k_{\parallel}| \gg \frac{\Omega_e}{V_e} \gg \frac{\Omega_i}{V_i} \quad \text{and} \quad \lambda_{\alpha} = \left(\frac{\Delta k_{\perp} V_{\alpha}}{\Omega_{\alpha}} \right)^2 \ll 1, \quad (\text{E.1})$$

it is evident that

$$|y_{\alpha s}| = \left| \frac{\Delta \omega - s \Omega_{\alpha}}{\sqrt{2} \Delta k_{\parallel} V_{\alpha}} \right| \ll 1, \quad (\text{E.2})$$

for small harmonic numbers s and $\phi(y_{\alpha s}) \approx 2y_{\alpha s}$. Since $\lambda_{\alpha} \ll 1$, only small harmonic terms contribute significantly to the dielectric tensor (2.78) and it may be approximated by

$$\epsilon_{ij}(\Delta \mathbf{k}, \Delta \omega) = \delta_{ij} - \sum_{\alpha} \frac{\pi_{\alpha}^2}{|\Delta k_{\parallel}|^2 V_{\alpha}^2} \left\{ \sum_{s=-\infty}^{\infty} \left(1 - \frac{s \Omega_{\alpha}}{\Delta \omega} \right) e^{-\lambda_{\alpha}} \pi_{ij}^{\alpha}(\Delta \mathbf{k}, \Delta \omega, s) - \delta_{i3} \delta_{j3} \right\}, \quad (\text{E.3})$$

with the inequality,

$$\left| \frac{\Delta k_{\perp}}{\Delta k_{\parallel}} \right|^2 \equiv \tan^2 \theta \ll \min \left(1, \frac{\Omega_i^2}{|\Delta \omega|^2} \right), \quad (\text{E.4})$$

given by (4.54) and (E.1).

Summations over modified Bessel functions in (E.3), using formulae such as

$$\sum_{s=-\infty}^{\infty} e^{-\lambda} I_s(\lambda) = 1, \quad \sum_{s=-\infty}^{\infty} e^{-\lambda} I'_s(\lambda) = 1, \quad \sum_{s=-\infty}^{\infty} s^2 I_s(\lambda) e^{-\lambda} = \lambda \quad (\text{E.5})$$

and

$$\sum_{s=-\infty}^{\infty} s^{2n+1} I_s(\lambda) = \sum_{s=-\infty}^{\infty} s^{2n+1} I'_s(\lambda) = 0, \quad (n=0,1,2,\dots), \quad (\text{E.6})$$

give a dielectric tensor,

$$\varepsilon_{ij}(\Delta \mathbf{k}, \Delta \omega) = \begin{pmatrix} 1 - 1/\sum_{\alpha} |\Delta k_{\parallel}|^2 \lambda_{D\alpha}^2 & \sum_{\alpha} i\varepsilon_{\alpha} \Omega_{\alpha} / |\Delta \omega| |\Delta k_{\parallel}|^2 \lambda_{D\alpha}^2 & \sum_{\alpha} 2 \tan \theta / |\Delta k_{\parallel}|^2 \lambda_{D\alpha}^2 \\ -\sum_{\alpha} i\varepsilon_{\alpha} \Omega_{\alpha} / |\Delta \omega| |\Delta k_{\parallel}|^2 \lambda_{D\alpha}^2 & 1 - 1/\sum_{\alpha} |\Delta k_{\parallel}|^2 \lambda_{D\alpha}^2 & \sum_{\alpha} i\varepsilon_{\alpha} \Omega_{\alpha} \tan \theta / |\Delta \omega| |\Delta k_{\parallel}|^2 \lambda_{D\alpha}^2 \\ \sum_{\alpha} 2 \tan \theta / |\Delta k_{\parallel}|^2 \lambda_{D\alpha}^2 & -\sum_{\alpha} i\varepsilon_{\alpha} \Omega_{\alpha} \tan \theta / |\Delta \omega| |\Delta k_{\parallel}|^2 \lambda_{D\alpha}^2 & 1 + 1/\sum_{\alpha} |\Delta k_{\parallel}|^2 \lambda_{D\alpha}^2 \end{pmatrix}$$

(E.7)

where $\lambda_{D\alpha}^2 \equiv v_{\alpha}^2 / \pi_{\alpha}^2$.

In order that $\eta^2 = |\Delta \mathbf{k}|^2 c^2 / |\Delta \omega|^2 \gg |\varepsilon_{ij}|$, it is sufficient that

$$|\tan \theta| \leq |\Delta \omega| / \Omega_e \quad (E.8)$$

and

$$|\Delta \mathbf{k}| \geq v_1 / c \left(\sum_{\alpha} \lambda_{D\alpha}^2 \right)^{1/2}. \quad (E.9)$$

When (E.8) and (E.9) are satisfied, the inverse Maxwellian tensor is given by (4.61) with

$$\varepsilon^k(\Delta \mathbf{k}, \Delta \omega) = A = 1 + 1/\sum_{\alpha} |\Delta \mathbf{k}|^2 \lambda_{D\alpha}^2. \quad (E.10)$$

Again in the regime (4.54), provided

$$|\Delta k_{\parallel}| \ll \frac{\Omega_i}{v_i} \ll \frac{\Omega_e}{v_e} \quad \text{and} \quad \lambda_{\alpha} = \left(\frac{\Delta k_{\perp} v_{\alpha}}{\Omega_{\alpha}} \right)^2 \gg 1, \quad (E.11)$$

straightforward calculations of equation (2.78) give

$$\varepsilon_{ij}(\Delta \mathbf{k}, \Delta \omega) = \begin{pmatrix} 1 & \sum_{\alpha} i\varepsilon_{\alpha} \frac{\pi_{\alpha}^2}{|\Delta \omega| \Omega_{\alpha}} h(\lambda_{\alpha}) & 0 \\ -\sum_{\alpha} i\varepsilon_{\alpha} \frac{\pi_{\alpha}^2}{|\Delta \omega| \Omega_{\alpha}} h(\lambda_{\alpha}) & 1 - \sum_{\alpha} \frac{\pi_{\alpha}^2}{|\Delta k_{\parallel}|^2 v_{\alpha}^2} 2\lambda_{\alpha} h(\lambda_{\alpha}) & \sum_{\alpha} i\varepsilon_{\alpha} \frac{\pi_{\alpha}^2}{|\Delta \omega|^2} \tan \theta h(\lambda_{\alpha}) \\ 0 & -\sum_{\alpha} i\varepsilon_{\alpha} \frac{\pi_{\alpha}^2}{|\Delta \omega|^2} \tan \theta h(\lambda_{\alpha}) & 1 + \sum_{\alpha} \frac{\pi_{\alpha}^2}{|\Delta k_{\parallel}|^2 v_{\alpha}^2} \left(1 - \frac{e^{-\lambda_{\alpha}} I_0}{\lambda_{\alpha}} \tan^2 \theta \frac{|\Delta \omega|^2}{\Omega_{\alpha}^2} \right) \end{pmatrix}$$

(E.12)

where $h(\lambda_{\alpha}) \equiv \exp(-\lambda_{\alpha})(I_0 - I_1)$ and I_0 and I_1 are modified Bessel

functions with argument λ_α . The variation of $h(\lambda_\alpha)$ with respect to λ_α is given by fig. E.

Fig. E: Variation of $h(\lambda_\alpha)$ vs. λ_α .

On picking the largest term in (E.12), it is seen that $\eta^2 \gg |\epsilon_{ij}|$ provided

$$|\Delta \underline{k}|^2 \geq \omega_p^2 |\tan \theta| / c^2. \quad (\text{E.13})$$

Since effective nonlinear scattering stipulates $|\Delta \underline{k}| \leq \omega_p / v_e$,

$\lambda_e \leq (\omega_p / \Omega_e)^2$ and $\lambda_i \leq (m_i / m_e)(\omega_p / \Omega_e)^2$, under the condition

$$c^2 / v_e^2 \geq |\tan \theta| \gg 1, \quad (\text{E.14})$$

the inverse Maxwellian tensor again reduces to (4.61) and in addition if

$$|\Delta \omega| \leq \Omega_e v_e / c, \quad (\text{E.15})$$

$\epsilon^{\ell}(\Delta \underline{k}, \Delta \omega)$ is again determined by (E.10).

In the regime (4.55), where scattering by electrons is the dominant process, the conditions (2.104) give a dielectric tensor of the form (2.106), from which it is seen that $\eta^2 \gg |\epsilon_{ij}|$, when $v_s = v_i (Z_i T_e / T_i)^{1/2} \approx v_i$, provided

$$|\Delta \underline{k}| \geq (\Omega_i / v_s) (v_e / c) . \quad (E.16)$$

When (E.16) is satisfied, the inverse Maxwellian tensor is again given by (4.61) with

$$\epsilon^0(\Delta \underline{k}, \Delta \omega) = 1 / |\Delta \underline{k}|^2 \lambda_{De}^2 . \quad (E.17)$$

APPENDIX F

APPROXIMATE FORMS OF THE NONLINEAR RESPONSE TENSOR ϵ_{ijl}

In this appendix, useful approximate forms of the nonlinear response tensor $\epsilon_{ijl}(\underline{k}', \omega'; \underline{k}'', \omega'')$ given by (2.38) and (2.43) are derived.

An alternative form of equation (2.38) can be obtained on integration by parts:

$$\epsilon_{ijl}(\underline{k}', \omega'; \underline{k}'', \omega'') = \frac{4\pi i}{\omega} \sum_{\alpha} \frac{q_{\alpha} n_{\alpha}}{\Omega_{\alpha}^2} \int d^3 p \left\{ \int^{\phi} d\phi' e^{P_2(\phi'') - P_2(\phi')} \right. \\ \left. g_{rl}(\underline{v}', \underline{k}'', \omega'') \frac{\partial f_{\alpha}^{(0)}}{\partial p_r} \right\} g_{sj}(\underline{v}, \underline{k}', \omega') \frac{\partial}{\partial p_s} \left\{ \int^{\phi} d\phi' v_1' e^{-P(\phi') + P(\phi)} \right\}, \quad (F.1)$$

with quantities defined by (2.39) to (2.41).

Under the condition

$$\omega/|\underline{k}| \gg v_e, \quad (F.2)$$

with $\omega = \omega' + \omega''$ and $\underline{k} = \underline{k}' + \underline{k}''$, the exponential in (F.1) can be expanded in powers of $\underline{k} \cdot \underline{v}/\omega$ and the last integral in (F.1) can be evaluated approximately. To first order in $|\underline{k}| |\underline{v}|/\omega$, this leads to

$$\epsilon_{ijl}(\underline{k}', \omega'; \underline{k}'', \omega'') = - \sum_{\alpha} \frac{4\pi |q_{\alpha}| n_{\alpha}}{\omega^2 m_{\alpha} \Omega_{\alpha}} \int d^3 p \left\{ \int^{\sigma} d\phi' e^{P_2(\phi'') - P_2(\phi')} \right. \\ \left. g_{rl}(\underline{v}', \underline{k}'', \omega'') \frac{\partial f_{\alpha}^{(0)}}{\partial p_r} \right\} g_{sj}(\underline{v}, \underline{k}', \omega') \left\{ \tau_{is} + \tau_{is}^{\alpha} \frac{k_n}{\omega} \tau_{nm}^{\alpha} v_m \right. \\ \left. + \frac{k_s}{\omega} \tau_{in}^{\alpha} \tau_{nm}^{\alpha} v_m \right\}, \quad (F.3)$$

where $\tau_{is}^{\alpha} = \tau_{is}^{\alpha}(\omega)$ is given by (4.13) with $\epsilon \rightarrow \epsilon_{\alpha}$ and $\Omega \rightarrow \Omega_{\alpha}$. To the lowest order, equation (F.3) with (2.37) gives

$$\begin{aligned} \epsilon_{ijl}(\underline{k}', \omega'; \underline{k}'', \omega'') &= \frac{4\pi e}{\omega^2 m_\alpha} \left\{ \tau_{ij}^{(e)} \frac{k''_s}{\omega''} \sigma_{sl}^{(e)}(\underline{k}'', \omega'') \right. \\ &\quad \left. - \tau_{ij}^{(e)} \frac{k'_s}{\omega'} \sigma_{sl}^{(e)}(\underline{k}'', \omega'') + \frac{k'_s}{\omega'} \tau_{is}^{(e)} \sigma_{jl}^{(e)}(\underline{k}'', \omega'') \right\}, \quad (F.4) \end{aligned}$$

where only electronic contribution is retained.

If, in addition to condition (F.1),

$$\omega' / |\underline{k}'| \gg v_e \quad \text{and} \quad \omega'' / |\underline{k}''| \ll v_e, \quad (F.5)$$

imposition of symmetry (2.32) together with (F.4) gives

$$\begin{aligned} \epsilon_{ijl}(\underline{k}', \omega'; \underline{k}'', \omega'') &= -\frac{ie}{2m_e \omega^2} \left\{ \tau_{ij}^{(e)}(\omega) k''_s \{ \epsilon_{sl}^{(e)}(\underline{k}'', \omega'') - \delta_{sl} \} \right. \\ &\quad \left. + \frac{\omega_p^2}{\omega'} \frac{k''_s}{\omega''} \{ \tau_{il}^{(e)}(\omega) \tau_{sj}^{(e)}(\omega') - \tau_{is}^{(e)}(\omega) \tau_{lj}^{(e)}(\omega') \} \right\}, \quad (F.6) \end{aligned}$$

where $\epsilon_{sl}^{(e)}$ is the electronic contribution to the dielectric tensor.

On the other hand, condition (F.2), together with the conditions,

$$\omega' / |\underline{k}'| \ll v_e \quad \text{and} \quad \omega'' / |\underline{k}''| \gg v_e \quad (F.7)$$

gives, on calculating (F.3) and symmetrizing by (2.32),

$$\begin{aligned} \epsilon_{ijl}(\underline{k}', \omega'; \underline{k}'', \omega'') &= \frac{ie\omega_p^2}{2m_e \omega \omega' \omega''} \left\{ \frac{k_s}{\omega'} \tau_{sj}^{(e)}(\omega') \tau_{il}^{(e)}(\omega'') \right. \\ &\quad + \frac{k_s}{\omega''} \tau_{sl}^{(e)}(\omega'') \tau_{ij}^{(e)}(\omega') - \frac{k'_s}{\omega''} \tau_{sl}^{(e)}(\omega'') \tau_{ij}^{(e)}(\omega) \\ &\quad - \frac{k''_s}{\omega'} \tau_{sj}^{(e)}(\omega') \tau_{il}^{(e)}(\omega) + \frac{k'_s}{\omega} \tau_{is}^{(e)}(\omega) \tau_{jl}^{(e)}(\omega'') \\ &\quad \left. + \frac{k''_s}{\omega} \tau_{is}^{(e)}(\omega) \tau_{lj}^{(e)}(\omega') \right\}, \quad (F.8) \end{aligned}$$

which is the nonlinear response tensor for a cold electronic plasma.

LIST OF SYMBOLS

All symbols are defined in the text of the thesis at least when they first occur. Where no ambiguity can arise, a symbol is sometimes used with another meaning. Those symbols which are used more extensively are listed below in their usual meanings.

B_0	Background magnetic field (induction)
$\underline{B}$	Magnetic field (induction)
c	Speed of light in vacuo
$\underline{D}$	Electric displacement
e	Electron charge
$\underline{e}^\sigma$	Polarization vector of waves in mode σ
E	Energy of a particle
$\underline{E}$	Electric field
$\underline{E}^\sigma$	Electric field of wave in mode σ
f_p	Electron plasma frequency (Hz)
$f_\alpha(\underline{p})$	Particle distribution function
$\underline{F}^\sigma(\underline{k})$	Energy density flux of waves in mode σ in the range $d^3\underline{k}/(2\pi)^3$ at $\underline{k}$
$\hbar$	$\equiv h/2\pi$, h is Planck's constant
$I_n(\lambda_e)$	Modified Bessel function of the first kind
$I^\sigma(\omega, \underline{\Omega})$	Intensity per unit frequency range
$j^\sigma(\omega, \theta)$	Emissivity
$\underline{j}(\underline{k}, \omega)$	Current density
$\underline{j}^q(\underline{k}, \omega)$	Current density associated with Thomson scattering
$\underline{j}^{n.s.}(\underline{k}, \omega)$	Current density associated with nonlinear scattering
$J_s(z)$	Bessel function of the first kind
$\underline{k}, k_\perp, k_\parallel$	Wave vector, its component perpendicular ($\perp$), parallel ($\parallel$) to $\underline{B}_0$
ℓ	Length of ray path
L	Linear dimension of source
m_α	Mass of particle
n	Harmonic number
n_α	Number density of particles
$\underline{p}, p_\perp, p_\parallel$	Particle moment, its component perpendicular ($\perp$), parallel ($\parallel$) to $\underline{B}_0$

$\bar{P}^\sigma$	Time averaged power per unit volume radiated into waves of mode σ
q_α	Charge of particle
$\underline{r}$	Position vector
r_c	Degree of circular polarization
$R_\odot$	Solar radius
s	Harmonic number
S_o	Observed energy flux density of radiation
S_m^σ	Source function of amplifying layer
S_a^σ	Source function of absorbing layer
t	Time
T	Time interval
T_b	Observed brightness temperature
T_{bs}	Source brightness temperature
T_α	Temperature ($^\circ K$)
u_α	Streaming velocity
$u^{\sigma\sigma'\sigma''}(\underline{k}, \underline{k}', \underline{k}'')$	Probability of three-wave interaction
$\underline{v}, v_\perp, v_\parallel$	Particle velocity, its component perpendicular ($\perp$), parallel ($\parallel$) to $\underline{B}_0$
v_A	Alfvén speed
v_s	Sound speed
$v_g^\sigma(\underline{k})$	Group velocity
v_ϕ	Phase velocity
V	Spatial volume
v_e, v_i	Thermal velocity of electrons, ions
$w_\alpha^\sigma(s, \underline{p}, \underline{k})$	Probability of wave particle interaction
$w_\alpha^{\sigma\sigma'}(s, \underline{p}, \underline{k}, \underline{k}')$	Probability of scattering
w_{el}	Energy density in electron plasma waves
W_p	Kinetic energy density of plasma particles
W_{th}	Energy density in thermal fluctuations
W^σ	Energy density in waves of mode σ
$(W_E/W_T)^\sigma$	Ratio of electrical to total energy density of waves in mode σ
X	$:= \omega_p^2/\omega^2$
Y	$:= \Omega_e/\omega$
z	$:= k_\perp v_\perp/\Omega_\alpha$
Z_i	Valency of ions
α	Particle species
β	$:= \underline{v} /c$

γ	$\equiv (1-\beta^2)^{-1/2}$, Lorentz factor
$\gamma^\sigma(\underline{k})$	Growth or damping rate
δ_{ij}	Kronecker delta
ϵ_α	$\equiv q_\alpha/ q_\alpha $
$\epsilon_{ij}(\underline{k},\omega)$	Linear response (dielectric) tensor
$\epsilon_{ij\ell}(\underline{k}_1,\omega_1;\underline{k}_2,\omega_2)$	First nonlinear response tensor
$\epsilon^\ell(\underline{k},\omega)$	$\equiv \kappa_i \kappa_j \epsilon_{ij}(\underline{k},\omega)$
$\zeta_s(\lambda_e)$	$\equiv e^{-\lambda_e} I_s(\lambda_e)$
η_σ	Refractive index
θ	Wave normal angle w.r.t. B_0
θ_r	Ray normal angle w.r.t. B_0
κ	Boltzmann's constant
$\underline{\kappa}$	$\equiv \underline{k}/ \underline{k} $
λ_e, λ_i	$\equiv (k_\perp v_e/\Omega_e)^2, (k_\perp v_i/\Omega_i)^2$
$\lambda_{D\alpha}$	Debye length
$\ln \Lambda_e$	Coulomb logarithm
$\Lambda_{ij}(\underline{k},\omega)$	$\equiv \eta^2(\kappa_i \kappa_j - \delta_{ij}) + \epsilon_{ij}^h(\underline{k},\omega)$
$\Lambda(\underline{k},\omega)$	$\equiv \det\{\Lambda_{ij}(\underline{k},\omega)\}$
$\lambda_{ij}(\underline{k},\omega)$	Cofactor of $\Lambda_{ij}(\underline{k},\omega)$
λ_{ss}	Trace of $\lambda_{ij}(\underline{k},\omega)$
$\mu^\sigma(\underline{k})$	Absorption coefficient (in time)
ν_c	Effective collision frequency
π_α	Plasma frequency
σ	Wave mode
$\Sigma^{\sigma\sigma'}(\theta_r, \psi; \theta_r', \psi')$	Scattering cross-section
τ	Interaction time
τ_c	Mean time between collisions
τ_D	Deflection time of beam
τ_ϕ	Correlation (phase-shift) time
τ^σ	Optical depth
ϕ, ψ	Particle azimuthal angle, wave azimuthal angle
ω	Angular frequency
$\omega^\sigma(\underline{k})$	Angular frequency of waves in mode σ
ω_p	Electron plasma frequency
$\omega_{0,X}$	Cutoff frequency of O, X-mode waves
Ω_α	Gyrofrequency
Ω_o	Solid angle subtended by source at observer
$d\Omega$	Element of solid angle

REFERENCES

- Akhiezer, A.I., Akhiezer, I.A., Polovin, R.V., Sitenko, A.G., and Stepanov, K.N. (1967). - "Collective Oscillations in a Plasma", Pergamon Press, Oxford.
- Akhiezer, A.I., and Polovin, R.V. (1971). - Sov. Phys. Uspekhi 14, 278.
- Alfvén, H., and Falthammar, C.G. (1963). - "Cosmical Electrodynamics", Oxford University Press, London.
- Al'tshul', V.N., and Karpman, V.I. (1965). - Sov. Phys. JETP 20, 1043.
- Al'tshul', L.M., and Karpman, V.I. (1966). - Sov. Phys. JETP 2, 361.
- Bekefi, G. (1966). - "Radiation Processes in Plasmas", John Wiley & Sons, Inc., New York.
- Bell, T.F., and Buneman, O. (1964). - Phys. Rev. 133, 5A, 1300.
- Bernstein, I.B. (1958). - Phys. Rev. 109, 10.
- Blanken, R.A., Stix, T.H., and Kuckes, A.F. (1969). - Plasma Phys. 11, 945.
- Brambilla, M. (1970). - Plasma Phys. 12, 903.
- Briggs, R. (1964). - "Electron-Stream Interactions with Plasmas", M.I.T. Press, Cambridge.
- Buti, B. (1967). - Phys. Rev. 160, 188.
- Buti, B., and Jain, R.K. (1965). - Phys. of Fluids 8, 2080.
- Canobbio, E., and Croci, R. (1963). - Proc. 6th Conf. on Ion. Phen. in Gases, Paris, 3, 269.
- Canuto, V., Lodenquai, J., and Ruderman, M. (1971). - Phys. Rev. D3, 2303.
- Caroff, L.J., and Scargle, J.D. (1970). - Nature 225, 168.
- Chandrasekhar, S. (1943). - Revs. Mod. Phys. 15, 1.
- Chernov, G.P., Chertok, I.M., Fomichev, V.V., and Markeev, A.K. (1972). - Solar Phys. 24, 215.
- Cohen, M.H. (1960). - Astrophys. J. 131, 665.
- Crawford, F.W. (1965). - Nucl. Fusion 5, 73.
- Crawford, F.W. (1968). - "Cyclotron Harmonic Wave Phenomena in Plasmas", Paper in A Survey of Phenomena in Ionized Gases, p.109.

- Davidson, R.C. (1972). - "Methods in Nonlinear Plasma Theory", Academic Press, New York and London.
- Davidson, R.C., Hammer, D.A., Haber, I., and Wagner, C.E. (1972). - Phys. of Fluids 15, 317.
- Davis, W.D., De Silva, A.W., Dove, W.F., Griem, H.R., Krall, N.A., and Liewer, P.C. (1971). - Conf. Plasma Phys. Controlled Nucl. Fusion Res., I.A.E.A. III, 289.
- De Groot, S.R., and Mazur, P. (1969). - "Non-Equilibrium Thermodynamics", North-Holland Publishing Co., Amsterdam.
- Drummond, W.E., and Pines, D. (1962). - Nucl. Fusion Supp. 3, 1049.
- Eidman, V.Ya. (1958). - Sov. Phys. JETP 7, 91.
- Eidman, V.Ya. (1959). - Sov. Phys. JETP 9, 947.
- Eidman, V.Ya. (1962). - Sov. Phys. JETP 14, 1401.
- Einstein, A. (1917). - Phys. Zs. 18, 121.
- Elgaroy, O. (1961). - Astrophys. Norv. VII, 123.
- Elgaroy, O. (1967). - Astrophys. Lett. 1, 13.
- Elgaroy, O., and Eckhoff, H.Kr. (1966). - Astrophys. Norv. X, 127.
- Elgaroy, O., and Ugland, O. (1970). - Astron. & Astrophys. 5, 372.
- Ellis, G.R.A. (1962). - Aust. J. Phys. 15, 344.
- Fainberg, Ya.B. (1962). - J. Nucl. Energy, Part C, 4, 203.
- Fainberg, Ya.B., Shapiro, V.D., and Shevchenko, V.I. (1970). - Sov. Phys. JETP 30, 528.
- Fokker, A.D. (1965). - Bull. Astron. Inst. Neth. 18, 111.
- Fried, B.D., and Conte, S.D. (1961). - "The Plasma Dispersion Function", Academic Press, New York.
- Friedman, M. (1972). - Astrophys. Lett. 12, 129.
- Fukai, J., and Harris, E.G. (1972). - J. Plasma Phys. 7, 313.
- Fung, P.C.W., and Yip, W.K. (1966). - Aust. J. Phys. 19, 759.
- Furutani, Y., and Kalman, G. (1965). - Plasma Phys. 7, 381.
- Gaffey, J.D., Thompson, W.B., and Liu, C.S. (1972). - J. Plasma Phys. 7, 189.
- Gary, S.P., and Sanderson, J.J. (1970). - J. Plasma Phys. 4, 739.
- Gershman, B.N. (1960). - Sov. Phys. JETP 11, 657.
- Ginzburg, V.L. (1964). - "Electromagnetic Wave Propagation in a Plasma", Pergamon Press, New York.

- Ginzburg, V.L., and Eidman, V.Ya. (1959). - Sov. Phys. JETP 9, 1300.
- Ginzburg, V.L., and Zheleznyakov, V.V. (1958). - Sov. Astron. AJ, 2, 653.
- Ginzburg, V.L., and Zheleznyakov, V.V. (1965). - Phil. Mag. 11, 197.
- Ginzburg, V.L., and Zheleznyakov, V.V. (1970). - Comments Astrophys. Space Phys. 2, 67 and 197.
- Ginzburg, V.L., Zheleznyakov, V.V., and Eidman, V.Ya. (1962). - Phil. Mag. 7, 451.
- Goldreich, P., and Lynden-Bell, D. (1969). - Astrophys. J. 156, 59.
- Goldreich, P., Pacini, F., and Rees, M.J. (1972). - Comments Astrophys. Space Phys. IV, 23.
- Goldstein, M.L., and Lenchek, A.M. (1971). - Astrophys. J. 163, 515.
- Goldstein, S.J., Jr. (1959). - Astrophys. J. 130, 393.
- Gordon, I.M. (1968). - Astrophys. Lett. 2, 49.
- Gratzl, H. (1971). - Phys. of Fluids 14, 2006.
- Hall, L.S., and Heckrotte, W. (1964). - Phys. Rev. A134, 1474.
- Hanasz, J. (1966). - Aust. J. Phys. 19, 635.
- Harris, E.G. (1959). - Phys. Rev. Lett. 2, 34.
- Harris, E.G. (1961). - J. Nucl. Energy, Part C: Plasma Phys. 2, 138.
- Harris, E.G. (1967). - Phys. of Fluids 10, 238.
- Harris, E.G. (1969). - "Classical Plasma Phenomena from a Quantum Mechanical Viewpoint", Adv. in Plasma Phys. 3 (ed. A. Simon and W.B. Thompson), Interscience, New York.
- Howard, R. (1971). - "Solar Magnetic Fields", IAU Symposium No. 43 (ed. Howard).
- Kadomstev, B.B. (1965). - "Plasma Turbulence", Academic Press, New York.
- Kadomstev, B.B., and Petrviashvili, V.I. (1963). - Sov. Phys. JETP 16, 1578.
- Kai, K. (1962). - Publ. A.S. Japan 14, 1.
- Kalman, G., and Feix, M. (ed.) (1969). - "Nonlinear Effects in Plasmas", Proc. of the Second Orsay Summer Institute, Gordon & Breach, New York.
- Kaplan, S.A., and Tsytovich, V.N. (1968). - Sov. Phys.-Astron. 11, No. 6, 956.
- Kaplan, S.A., and Tsytovich, V.N. (1969). - Sov. Phys. Uspekhi 12, 42.

- Karpman, V.I. (1963). - Zh. Prikl. Mekhan. i Tekh. Fiz. 6, 34.
- Keilhacker, M., and Steuer, K.-H. (1971). - Phys. Rev. Lett. 26, 694.
- Kennel, C.F., Fredricks, R.W., and Scarf, F.L. (1970). - "High Frequency Electrostatic Waves in the Magnetosphere", 'Particles and Fields in Magnetosphere', ed. B.M. McCormac.
- Kennel, C.F., Sagdeev, R.Z. (1967). - J. Geophys. Res. 72, 3303 and 3327.
- Klozenberg, J.P., and Bernstein, I.B. (1970). - J. Plasma Phys. 4, 595.
- Komesaroff, M.M. (1970). - Nature 225, 612.
- Kovrizhnykh, L.M. (1968). - Proc. P.N. Lebedev Phys. Inst. 32, 155.
- Krall, N.A., and Rosenbluth, M.N. (1963). - Phys. of Fluids 6, 254.
- Krishan, S., and Selim, A.A. (1971). - Phys. of Fluids 14, 599.
- Kropotkin, A.P. (1968). - Sov. Phys. JETP 26, 1011.
- Kropotkin, A.P., and Pustovalov, V.V. (1966). - Sov. Phys. JETP 22, 928.
- Kropotkin, A.P., and Pustovalov, V.V. (1967). - Phys. of Fluids 10, 241.
- Krüger, A. (1972). - "Physics of Solar Continuum Radio Bursts", Akademie-Verlag, Berlin.
- Kundu, M. (1965). - "Solar Radio Astronomy", Wiley (Interscience), New York.
- Lacombe, C., and Mangeney, A. (1969). - Astron. & Astrophys. 1, 325.
- Lampe, M., Manheimer, W.M., McBride, J.B., Orens, J.H., Shanny, R., and Sudan, R.N. (1971). - Phys. Rev. Lett. 26, 1221.
- Lampe, M., Manheimer, W.M., McBride, J.B., Orens, J.H., Papadopoulos, K., Shanny, R., and Sudan, R.N. (1972). - Phys. of Fluids 15, 662.
- Landau, L. (1946). - J. Phys. U.S.S.R. 10, 25.
- Landau, L.D., and Lifshitz, E.M. (1960). - "Electrodynamics of Continuous Media" - Course of Th. Phys. Vol. 8 (Ch. XIII Electromagnetic Fluctuations), Pergamon Press, New York.
- Landau, L.D., and Lifshitz, E.M. (1969). - "Statistical Physics", Pergamon Press, New York.
- Landau, R.W., and Cuperman, S. (1970). - J. Plasma Phys. 4, 13.
- Liemohn, H.B. (1965). Radio Sci. 69D, 741.
- Livshitz, M.A., and Tsytovich, V.N. (1968). - Sov. Phys. JETP 26, 927.

- Malinge, A.-M. (1963). - Ann. d'Astrophys. 26, 97.
- Martres, M.J., Pick, M., Soru-Escaut, I., and Axisa, F. (1972). - Nature Phys. Sci. 236, 25.
- McBride, J.B. (1969a). - Phys. of Fluids 12, 844.
- McBride, J.B. (1969b). - Phys. of Fluids 12, 850.
- McKenzie, J.F. (1963). - Phil. Trans. Roy. Soc. London A255, 585.
- McKenzie, J.F. (1967). - Phys. of Fluids 10, 2680.
- Melrose, D.B. (1968a). - Astrophys. & Space Sci. 2, 171.
- Melrose, D.B. (1968b). - Astrophys. J. 154, 803.
- Melrose, D.B. (1971). - Astrophys. Space Sci. 10, 186.
- Melrose, D.B. (1972a). - Plasma Phys. 14, 1035.
- Melrose, D.B. (1972b). - Unpublished notes.
- Melrose, D.B. (1972c). - Plasma Phys. (in press).
- Melrose, D.B., and Sy, W.N. (1972a). - Astrophys. & Space Sci. 17, 343.
- Melrose, D.B., and Sy, W.N. (1972b). - Aust. J. Phys. 25, 387.
- Melrose, D.B. (1973). - Aust. J. Phys. (in press).
- Mizuno, K., Sugaya, R., and Tanaka, S., Phys. Lett. 36A, 441.
- Mollwo, L. (1970). - Solar Phys. 12, 125.
- Montes, C., Coste, T., and Diener, G. (1970). - J. Plasma Phys. 4, 21.
- Montes, C., and Peyraud, J. (1972). - J. Plasma Phys. 7, 69.
- Montgomery, D., and Bodner, S. (1971). - J. Plasma Phys. 5, 131.
- Montgomery, P.C., and Tidman, D.A. (1964). - "Plasma Kinetic Theory", McGraw-Hill Book Co., New York.
- Morimoto, M., and Kai, K. (1961). - Publ. Astron. Soc. Japan 13, 294.
- Morimoto, M., and Kai, K. (1962). - Publ. Astron. Soc. Japan 17, 220.
- Mustel, E.R. (1962). - Sov. Astron. AJ, 6, 333.
- Mustel, E.R. (1963). - Sov. Astron. AJ, 6, 488.
- Neufeld, J., and Wright, H. (1963). - Phys. Rev. 129, 1489.
- Newkirk, G. (1961). - Astrophys. J. 133, 983.
- Newkirk, G. (1967). - Ann. Rev. of Astron. Astrophys. 5, 213.

- Parker, E.N. (1961). - J. Nucl. Energy, Part C: Plasma Phys. 2, 146.
- Paul, J.W.M. (1971). - "Collisionless Shocks", Culham Lab. Preprint CLM-P290.
- Payne-Scott, R., and Little, R.G. (1951). - Aust. J. Sci. Res. A4, 508.
- Perkins, W.A., and Barr, W.L. (1966). - "Observation of a Velocity-Distribution Instability", Plasma Phys. & Controlled Nucl. Fusion Research II, Vienna, I.A.E.A., 115.
- Piddington, J.H. (1969). - "Cosmic Electrodynamics", Wiley-Interscience Publ., New York.
- Pines, D., and Schrieffer, J.R. (1962). - Phys. Rev. 125, 804.
- Polovin, R.V. (1962). - Sov. Phys.-Tech. Phys. 6, 889.
- Ramaty, R. (1969). - Astrophys. J. 158, 753.
- Riddle, A.C. (1972). - Proc. ASA 2, 98.
- Rogister, A., and Oberman, C. (1968). - J. Plasma Phys. 2, 33.
- Rosenbluth, M.N., MacDonald, W.M., and Judd, D.L. (1957). - Phys. Rev. 107, 1.
- Rust, D.M., and Roy, J.-R. (1971). - "Coronal Magnetic Fields Above Active Regions", 'Solar Magnetic Fields' - I.A.U. Symposium No. 43, (ed. R. Howard), p.569.
- Sagdeev, R.Z. (1960). - Rev. of Plasma Phys. 4, 23.
- Sagdeev, R.Z., and Galeev, A.A. (1969). - "Nonlinear Plasma Theory", W.A. Benjamin, New York.
- Sakurai, K. (1972). - Astrophys. J. 174, 135.
- Sastry, Ch.V. (1972). - Astrophys. Lett. 11, 47.
- Schwartz, M.J., and Lichtenberg, A.J. (1972). - Phys. of Fluids 15, 470.
- Shafranov, V.D. (1967). - Rev. of Plasma Phys. 3,
- Shapiro, V.D. (1963). - Sov. Phys. JETP 17, 416.
- Sheridan, K.V. (1972). - Private communication.
- Shima, Y., and Hall, L.S. (1965). - Phys. Rev. A139, 1115.
- Silin, V.P. (1960). - Sov. Phys. JETP 11, 1136.
- Sitenko, A.G. (1967). - "Electromagnetic Fluctuations in Plasma", Academic Press, New York.
- Sitenko, A.G., and Kirochkin, Yu.A. (1966). - Sov. Phys. JETP 9, 430.
- Sitenko, A.G., and Stepanov, K.N. (1957). - Sov. Phys. JETP 4, 512.

- Smith, D.F. (1970). - Adv. in Astron. & Astrophys. 7, 148.
- Smith, D.F. (1971). - Astrophys. J. 170, 559.
- Smith, D.F. (1972). - Astrophys. J. (in press).
- Smith, D.F., and Fung, P.C.W. (1971). - J. Plasma Phys. 5, pt. 1, 1.
- Steinberg, J.-L., Aubier-Giraud, M., Leblanc, Y., and Boischot, A. (1971). - Astron. Astrophys. 10, 362.
- Stenflo, L. (1972). - J. Plasma Phys. 7, 107.
- Stepanov, K.N. (1960). - Sov. Phys. JETP 11, 192.
- Stepanov, K.N., and Kitsenko, A.B. (1961). - Sov. Phys.-Tech. Phys. 6, 120.
- Stewart, R.T., and Labrum, N.R. (1972). - Solar Phys. (in press).
- Stix, T.H. (1962). - "The Theory of Plasma Waves", McGraw-Hill Book Co., Inc., New York.
- Stone, P.M., and Auer, P.L. (1965). - Phys. Rev. A138, 695.
- Sturrock, P.A. (1958). - Phys. Rev. 112, No. 5, 1488.
- Sugihara, R. (1963). - J. Phys. Soc. Japan 18, 1062.
- Sweet, P.A. (1971). - "Theories of Small-scale Magnetic Fields", 'Solar Magnetic Fields' - I.A.U. Symposium No. 43 (ed. Howard), p.451.
- Takakura, T. (1963). - Publ. Astron. Soc. Japan 15, 462.
- Tataronis, J.A., and Crawford, F.W. (1970a). - J. Plasma Phys. 4, 231.
- Tataronis, J.A., and Crawford, F.W. (1970b). - J. Plasma Phys. 4, 249.
- Tidman, D.A., and Dupree, T.H. (1965). - Phys. of Fluids 8, 1860.
- Traktengerts, V.Yu. (1966). - Sov. Astron. AJ, 10, 281.
- Trulsen, J., and Fejer, J.A. (1970). - J. Plasma Phys. 4, 825.
- Tsintsadze, N.L. (1964). - Sov. Phys.-Tech. Phys. 9, 1397.
- Tsyтович, V.N. (1962). - Radiofizika 5, 1078.
- Tsyтович, V.N. (1963). - Sov. Phys. JETP 17, 643.
- Tsyтович, V.N. (1967). - Sov. Phys. Uspekhi 9, No. 3, 805.
- Tsyтович, V.N. (1969). - Sov. Astron. AJ, 12, 806.
- Tsyтович, V.N. (1970). - "Nonlinear Effects in Plasma", Plenum Press, New York-London.

- Tsyтович, V.N. (1971). - Plasma Phys. 13, 741.
- Tsyтович, V.N., and Shapiro, V.D. (1965). - Nucl. Fusion 5, 228.
- Tsyтович, V.N., and Shvartsburg, A.B. (1966). - Sov. Phys. JETP 22, 554.
- Tsyтович, V.N., and Shvartsburg, A.B. (1967). - Sov. Phys.-Tech. Phys. 11, 1431.
- Twiss, R.Q. (1963). - Phil. Mag. 8, 1249.
- Vahala, G., and Montgomery, D. (1970). - J. Plasma Phys. 4, 677.
- Vedenov, A.A. (1968). - "Theory of Turbulent Plasma", Iliffe Books, London.
- Vedenov, A.A., Velikov, E.P., and Sagdeev, R.Z. (1961). - Nucl. Fusion 1, 82.
- Vedenov, A.A., Velikov, E.P., and Sagdeev, R.Z. (1962). - Nucl. Fusion Suppl. 465.
- Warwick, J.W., and Dulk, G.A. (1969). - Astrophys. J. 158, L123.
- Watson, G.N. (1966). - "A Treatise on the Theory of Bessel Functions", University Press, Cambridge.
- Wild, J.P., and McCready, L.L. (1950). - Aust. J. Sci. Res. A3, 387.
- Wild, J.P., Sheridan, K.V., and Neylan, A.A. (1959). - Aust. J. Phys. 12, 369.
- Wild, J.P., and Smerd, S.F. (1972). - Ann. Rev. Astron. Astrophys. 10, 159.
- Wild, J.P., Smerd, S.F., and Weiss, A.A. (1963). - Ann. Rev. Astron. Astrophys. 1, 291.
- Wild, J.P., and Tlamicha, A. (1964). - Nature 203, 1128.
- Yip, W.K. (1970). - Aust. J. Phys. 23, 161.
- Yoh, P., and James, J.C. (1967). - Astrophys. J. 149, 441.
- Zayed, K.E., and Kitsenko, A.B. (1968). - Plasma Phys. 10, 673.
- Zheleznyakov, V.V. (1970). - "Radio Emission of the Sun and Planets", Pergamon Press, New York.
- Zheleznyakov, V.V., and Zaitsev, V.V. (1970a). - Sov. Astron. AJ, 14, 47.
- Zheleznyakov, V.V., and Zaitsev, V.V. (1970b). - Sov. Astron. AJ, 14, 250.