THE DEPOSITION AND UTILIZATION OF INORGANIC
PYROPHOSPHATE IN TETRAHYMENA PYRIFORMIS

A THESIS

submitted for the degree of

DOCTOR OF PHILOSOPHY

in the

Australian National University

by

NISSIM MUNK

June 1968

STATEMENT

This Thesis embodies the results of research carried out in the Department of Biochemistry, of the John Curtin School of Medical Research, Australian National University, from March 1965 to June 1968, during the tenure of a Commonwealth Postgraduate Award.

CANDIDATE'S SIGNATURE

Barrie Monte

ACKNOWLEDGMENTS

It is a pleasure to express my thanks to my supervisor, Dr H. Rosenberg, for his helpful guidance and encouragement during this work. I also wish to thank him for his assistance during the preparation of the manuscript.

I am grateful to my Departmental colleagues who have given me the advantage of their knowledge, and in particular to Dr J.F. Morrison who was always available for critical discussion.

STATEMENT

In accordance with the regulations of the Australian National University, I wish to state that, the work described in this Thesis was carried out by myself under the supervision of Dr H. Rosenberg.

The photographic prints for this Thesis were prepared by Mr R. Whalen and his staff, whose help is greatly appreciated, and I also wish to thank Mrs A. Aldrich for her help in lettering for the illustrations.

CANDIDATE'S SIGNATURE

Nissim Muntz

ACKNOWLEDGEMENTS

It is a pleasure to express my thanks to my supervisor, Dr H. Rosenberg, for his helpful guidance and encouragement during this work. I also wish to thank him for his assistance during the preparation of the manuscript.

I am grateful to my Departmental colleagues who have given me the advantage of their specialised knowledge, and in particular to Dr J.F. Morrison who was always available for critical discussion.

I am indebted to Messrs. I.K. Reid and A.B. Hawkins in many ways, but particularly for their help in procurement of enzymes, chemicals and other materials required during this work.

The photographic prints for this Thesis were prepared by Mr R. Westen and his staff, whose help is greatly appreciated, and I also wish to thank Mrs A. Aldrich for preparing the lettering for the illustrations.

DMTA	dimethyltetraacetate
DNP	2,4-dinitrophenol
P ₂ O ₅	phosphorus pentoxide
UV	ultraviolet light
DEAE Cellulose	Diethylaminoethyl cellulose
DEAE Sephadex	Diethylaminoethyl derivative of Sephadex
CM Sephadex	Carboxymethyl derivative of Sephadex

PREFACE

A reference in the text to Methods applies to the relevant section of the chapter in which the reference is made.

Figures and Tables are presented on separate pages, a particular Figure or Table following in most instances the page on which the first reference to it has been made.

The abbreviations used in this Thesis for time, volume and mass and also for nucleotides are those recommended for the Biochemical Journal. In addition the following abbreviations are used:

Ca^{2+} , Mg^{2+}	calcium ion(s), magnesium ion(s), etc.
PPi	inorganic pyrophosphate
Pi	inorganic orthophosphate
TCA	trichloroacetic acid
Tris	2-amino-2-hydroxy methyl propane-1,3-diol
EDTA	ethylenediaminetetra-acetic acid
EGTA	ethyleneglycol-2-(2-aminoethyl) tetraacetic acid
DNP	2,4-dinitrophenol
P_2O_5	phosphorus pentoxide
UV	ultraviolet light
DEAE Cellulose	Diethylaminoethyl cellulose
DEAE Sephadex	Diethylaminoethyl derivative of Sephadex
CM Sephadex	Carboxymethyl derivative of Sephadex

For the sake of simplicity, enzymes have been referred to in the text by their trivial names. The numbers and official names of those enzymes which are mentioned in this Thesis and which appear in the Report of the Commission on Enzymes of the International Union of Biochemistry (1964) are listed below:

<u>Number</u>	<u>Systematic Name</u>	<u>Trivial Name</u>
1.1.1.49	D-glucose-6-phosphate: NADP oxidoreductase	Glucose-6-phosphate dehydrogenase
2.7.1.1	ATP:D-hexose 6-phospho- transferase	Hexokinase
2.7.1.2	ATP:D-glucose-6- phosphotransferase	Glucokinase
2.7.4.1	ATP:polyphosphate phosphotransferase	Polyphosphate kinase
2.7.4.3	ATP:AMP phosphotrans- ferase	Adenylate kinase
2.7.5.1	α -D-glucose-1-6 diphosphate: α -D-glucose- 1-phosphate phosphotransferase	Phosphoglucomutase
2.7.7.4	ATP:sulphate adenylyl transferase	Sulphate adenylyl transferase
3.1.3.1	Orthophosphoric monoester phosphohydrolase	Alkaline phosphatase
3.1.3.9	D-glucose-6-phosphate phosphohydrolase	Glucose-6-phosphatase
3.6.1.1	Pyrophosphate phosphohydrolase	Inorganic pyro- phosphatase
3.6.1.3	ATP phosphohydrolase	ATPase

<u>Number</u>	<u>Systematic Name</u>	<u>Trivial Name</u>
3.6.1.10	Polyphosphate polyphospho- hydrolase	Polyphosphatase
4.1.1.e	Pyrophosphate:oxaloacetate carboxy-lyase (phosphorylating)	Phosphoenol-pyruvate carboxytransphosphory- lase, or Phosphopy- ruvate carboxylase
4.1.1.32	GTP:oxaloacetate carboxylyase (transphosphorylating)	Phosphoenol- pyruvate carboxykinase, or Phosphopyruvate carboxylase

TABLE OF CONTENTS

	Page
Statement of Purpose	1
Chapter I General Introduction	4
Chapter II Conditions Governing the Deposition of Inorganic Pyrophosphate in <u>Tetrahymena</u> .	34
Chapter III Studies on the Inorganic Pyrophosphatase from <u>Tetrahymena</u> .	62
Chapter IV The Inorganic Pyrophosphatase Activity Associated With Pyrophosphate Granules Isolated from <u>Tetrahymena</u> .	82
Chapter V Is Pyrophosphate a Phosphoryl Group Donor in Biosynthetic Reactions in <u>Tetrahymena</u> ?	97
References	136

STATEMENT OF PURPOSE

The observation (Rosenberg, 1966) that the ciliate, Tetrahymena pyriformis, is capable of depositing large numbers of cytoplasmic granules composed of calcium and magnesium salts of inorganic pyrophosphate, is quite unique. Though inorganic pyrophosphate is the product of many vital metabolic reactions, its concentration in living matter is usually very low. This is due to the ubiquitous inorganic pyrophosphatases which hydrolyse inorganic pyrophosphate to orthophosphate. Though there are numerous examples that show that a great variety of microorganisms are capable of accumulating considerable quantities of condensed phosphates (Kuhl, 1960; Harold, 1966), we are aware of only two other instances in which pyrophosphate was shown to be accumulated in any significant amounts, namely, the cases of the butterfly Deilephila euphorbiae, and the moth Galleria melonella (Heller, Kapriack and Zubikowa, 1950; Niemierko, 1962).

The fact that Tetrahymena pyriformis is capable of depositing inorganic pyrophosphate raised the problem of the possible functions these deposits may have in this organism. The granules may represent a means of disposal of a reaction end-product, unlikely though

this may be in view of scarcity of phosphate in Nature. On the other hand, the function of these granules may be one of phosphate-storage, or even of energy-storage, similar to that attributed to polyphosphates in other microorganisms. It is easy to visualise how the 'energy-rich' anhydride bond of the pyrophosphate molecule could act as a source of chemical energy in several important metabolic reactions. If such reactions do exist in Tetrahymena, they would be not only of metabolic importance, but their demonstration would be of great interest from the point of view of biochemical evolution.

The observation of Miller and Parris (1964) that inorganic pyrophosphate is produced in good yields when hydroxyapatite is shaken in dilute buffer, at near neutral pH and room temperature, in the presence of potassium cyanate, suggests that pyrophosphate could have been available to primitive cells in primordial pools. Furthermore, in connection with the recent findings of Baltscheffsky and his associates (Baltscheffsky, von Stedingk, Heldt and Klingenberg, 1966; Baltscheffsky, 1967) that pyrophosphate is formed and utilised in subcellular energy transfer systems, it is tempting to speculate that pyrophosphate, with its relatively simple molecular structure, was a forerunner of the more elaborate molecules which participate in energy storage and transfer, such as adenosine triphosphate.

Therefore, the primary aim of the work described in this thesis was to attempt to demonstrate in Tetrahymena reactions in which pyrophosphate can act both as phosphate and as an energy donor.

Chapter I

GENERAL INTRODUCTION

CONTENTS

	Page
1. Introduction	1
2. The Chemistry of Condensed Phosphates	2
3. Isolation of the Naturally Occurring Polyphosphates	3
4. The Distribution and the Intracellular State of Polyphosphates in Ferns	9
Chapter I	
5. Enzymes Involved in the Metabolism of Polyphosphates	14
A. Biosynthesis	14
B. Breakdown of Polyphosphates	17
GENERAL INTRODUCTION	
6. Factors Controlling the Accumulation and the Degradation of Polyphosphates	21
7. Theories on the Functions of Polyphosphates	23
8. Participation of Inorganic Pyrophosphate in Enzymic Reactions	27
A. Production of Pyrophosphate in Biosynthetic Reactions	27
B. Pyrophosphate as a Phosphoryl-Group Donor in Biosynthetic Reactions	29

CONTENTS

	Page
1. Introduction	4
2. The Chemistry of Condensed Phosphates	4
3. Isolation of the Naturally Occurring Polyphosphates	8
4. The Distribution and the Intracellular State of Polyphosphates in Nature	9
5. Enzymes Involved in the Metabolism of Polyphosphates	14
A. Biosynthesis of Polyphosphates	14
B. Breakdown of Polyphosphates	17
6. Factors Controlling the Accumulation and the Degradation of Polyphosphates	21
7. Theories on the Functions of Polyphosphates	25
8. Participation of Inorganic Pyrophosphate in Enzymic Reactions	27
A. Production of Pyrophosphate in Biosynthetic Reactions	27
B. Pyrophosphate as a Phosphoryl-Group Donor in Biosynthetic Reactions	29

The term condensed inorganic phosphates has been applied to all pentavalent phosphorus compounds in which various numbers of tetrahedral phosphate groups are linked together by oxygen bridges (Thilo, 1962). These compounds can be classified into three broad groups:

- (a) linear condensed phosphates;
- (b) cyclic condensed phosphates;
- (c) cross-linked condensed phosphates.

1. Introduction

Inorganic pyrophosphate is the lowest homologue of the large family of condensed phosphates, in which phosphate molecules are joined together to each other through anhydride linkages. Since the available information about the deposition and the utilisation of condensed phosphates of high molecular weights may have some bearing on the present problem, it is proposed to review the literature on this class of compounds first. The chemistry, metabolism and function of condensed phosphates has been reviewed on several occasions and the material presented here has been collated from a number of sources (Schmidt, 1951; van Wazer, 1958; Huennekens and Whiteley, 1960; Kuhl, 1960; Thilo, 1962) and particularly from a recent review by Harold (1966). The second part of the literature survey will deal with the known biological energy-conserving reactions which involve pyrophosphate.

2. The Chemistry of Condensed Phosphates

The term condensed inorganic phosphates has been applied to all 'pentavalent phosphorus compounds in which various numbers of tetrahedral phosphate groups are linked together by oxygen bridges' (Thilo, 1962). These compounds can be classified into three broad groups:

- (a) linear condensed phosphates;
- (b) cyclic condensed phosphates;
- (c) cross-linked condensed phosphates.

(a) Linear Condensed Phosphates

This group of compounds, known as polyphosphates, have linear structures with the elementary composition of $M_{n+2}P_nO_{3n+1}$, where M signifies a univalent cation. The series ranges from the water soluble pyrophosphate (PPi), with the chain length of 2, to the insoluble crystalline Kurrol's and Manddrell's salts of chain lengths approaching 10^4 phosphate groups. The formula of the tri-polyphosphoric acid is shown in Fig. I.1. Only the lowest homologues are known as individual compounds. All other synthetic polyphosphates must be regarded as mixtures, containing various proportions of species of different molecular size. The polyphosphates contain, in addition to one strong acidic group per phosphorus atom, also two weakly dissociating hydrogen atoms per chain. The pK values of the four hydrogen atoms of pyrophosphoric acid in dilute aqueous solution at 25° are: $pK_1 \approx 1$, $pK_2 \approx 2$, $pK_3 = 6.6$, and $pK_4 = 9.6$ (van Wazer, 1958). The difference in the pK values of the two weakly dissociating hydrogen atoms tends to decrease as the chain length increases. The average chain length is commonly estimated by titration of the secondary acid functions associated with the end groups. A large number of metal ions bind tightly to polyphosphates and precipitate them from solutions.

Polyphosphates are stable in neutral aqueous solutions and in alkali, but are labile at acid pH values. Typically, complete hydrolysis to orthophosphate (Pi) requires about 15 min. at 100°

(a)

(b) Cyclic Condensed Phosphates

The cyclic condensed phosphates, designated metaphosphates, conform to the general formula $(\text{P}_n\text{O}_6)_m$, where n signifies a number of phosphorus atoms in the ring and m is the number of rings.

(b)

(c)

(c) Cross-linked Condensed Phosphates

Cross-linked condensed phosphates are defined as polyphosphates in which branching occurs (Fig. 1.1). Unlike polyphosphates and metaphosphates, these compounds are relatively rare and are therefore

Fig. 1.1. (a) Tripolyphosphoric acid; (b) Trimetaphosphoric acid; (c) a branched polyphosphate.

cross-linking occurs. Due to this property, they are unlikely to be encountered in biological material.

in 1 N HCl. The degradation of polyphosphates to Pi at intermediate acid pH values is thought to occur by the sequential hydrolytic removal of the terminal links, but at very high hydrogen ion concentrations it is thought that scission occurs along the chain, resulting in the formation of polyphosphates of smaller chain lengths.

(b) Cyclic Condensed Phosphates

The cyclic condensed phosphates, designated metaphosphates, conform to the general formula $M_n P_n O_{3n}$, where M signifies a univalent metal ion. The only compound of biological interest that belongs to this group is the trimetaphosphoric acid (Fig. I.1). The free trimetaphosphoric acid titrates like a strong monobasic acid, so that it must be assumed that there is only one dissociable hydrogen atom per phosphorus atom. Metaphosphates are stable in neutral aqueous solutions. Strong alkali convert metaphosphates quantitatively to the corresponding linear polyphosphates. In hot acids metaphosphates are hydrolysed to Pi via the corresponding linear polyphosphates.

(c) Cross-linked Condensed Phosphates

Cross-linked condensed phosphates are defined as polyphosphates in which branching occurs (Fig. I.1). Unlike polyphosphates and metaphosphates, these compounds are triesters and are, therefore, unstable in water. They are readily hydrolysed at positions where cross-linking occurs. Due to this property, they are unlikely to be encountered in biological material.

3. Isolation of Naturally Occurring Polyphosphates

The large number of methods devised for the separation of polyphosphates from biological materials are, in one way or another, modifications of the classical fractionation schemes for nucleic acids (Schneider, 1945; Schmidt and Thannhauser, 1945). In principle, these methods involve the extraction of polyphosphates into two fractions: 'acid-soluble' and 'acid-insoluble' polyphosphates. The 'soluble' polyphosphate is extracted with cold trichloroacetic or perchloric acids, and then precipitated from the solutions with heavy metal ions. After the removal of the 'acid-soluble' polyphosphates, ribonucleic acids and 'insoluble' polyphosphates are solubilised with hot alkali. The solution is neutralised with strong acid and deoxyribonucleic acids are removed by trichloroacetic acid precipitation. The ribonucleic acids are removed from the remaining supernatant by absorption on charcoal and the 'insoluble' polyphosphates are precipitated with barium.

Both paper- and column-chromatographic procedures have been developed for the separation of polyphosphates of different chain lengths (Ebel, 1952; Ebel, 1953; Ebel and Muller, 1958; Lindenbaum, Peters and Rieman, 1954; Peters and Rieman, 1956). Both of these methods can be used successfully for the separation of polyphosphates up to the chain length of about 15 phosphate groups, but are ineffective for the separation of polyphosphates of higher chain lengths.

Polyphosphates have been identified in cells in situ by the use of certain dyes that exhibit metachromacy, i.e. the ability to undergo changes in their absorption spectra under certain conditions. Only polyphosphates with chain length greater than 6 exhibit metachromacy (Wiame, 1947(a); Ebel and Muller, 1958). The most widely used procedure for cytological staining with a metachromatic dye is that of Laybourn (1924). A modified Gomori staining procedure for polyphosphates has been described, in which polyphosphate is precipitated in situ with lead, which is subsequently converted to black lead sulphide (Ebel, Colas and Muller, 1958).

A large number of early workers in the field have employed cytochemical methods as the sole criterion for the identification of polyphosphate deposits in microorganisms. Their results have to be assessed with great caution as other polymeric substances of high molecular weights also exhibit metachromacy.

4. The Distribution and Intracellular State of Polyphosphates in Nature

Polyphosphates are found in many cells, and were first isolated from yeast by Liebermann (1888). MacFarlane (1936) isolated from yeast a nucleic acid fraction with a high content of acid labile phosphates which could be separated from the nucleic acids by extraction with HCl. The compound was identified as trimetaphosphate by precipitation with heavy metal ions. Jeener and Brachet (1944) made the interesting observation that Pi-starved yeast

cells became strongly basophilic after a short incubation in Pi -containing media. The basophilic material was identified by Wiame (1946, 1947(a)), and independently by Schmidt, Hecht and Thannhauser (1946) as an inorganic polyphosphate of high molecular weight.

There followed numerous reports on the presence of these compounds in bacteria, fungi, yeast, protozoa, and more recently, in the higher plants and animals. A recent comprehensive review by Kuhl (1960) lists the names of organisms which have been shown to contain polyphosphates. This list was extended by many subsequent reports, but of relevance to the present work are polyphosphate deposits in the flagellate Crithidia fasciculata (Janakidevi, Dewey and Kidder, 1965). These workers isolated and characterised polyphosphates from this organism according to their chain lengths. The isolated polyphosphates ranged in chain length from 370 to 850 phosphate residues.

Although it was earlier generally believed that polyphosphates occurred only in lower organisms, recent reports have shown without any doubt that they are present also in tissues of multicellular organisms. The earliest reports of the occurrence of polyphosphates in higher organisms described large quantities of this compound in the butterfly Galleria malonella (Heller et al, 1950; Niemierko, 1962). Grossmann and Lang (1962) reported traces of short-chain polyphosphates in nuclei isolated from swine kidney. Penniall and his associates isolated and characterised polyphosphates

from rat liver nuclei (Penniall and Griffin, 1964; Griffin, Davidian and Penniall, 1965; Griffin and Penniall, 1966). These workers also showed that these nuclei synthesised polyphosphates in vitro, and that this latter synthesis was greatly stimulated by divalent metal ions, such as Ca^{2+} and Mg^{2+} .

The occurrence of polyphosphates in higher plants was reported by Miyachi (1961) in spinach leaves, and by Tewari and Singh (1964) in the dodder, Cuscuta reflexa. Vagabov and Kulaev (1964) reported the presence of polyphosphates in corn roots.

'Volutankugeln' was one of the early trivial names coined for the intracellular granules which were detected with ease by light microscopy in Spirillum volutans. The name, coined by Grimme (1902) was then shortened to 'volutin', and redefined in terms of staining reactions by Meyer (1904). Since 'volutin' granules stained metachromatically with basic dyes, they were also referred to as 'metachromatic' granules. From the work of Wiame (1946, 1947(b)), who developed chemical criteria for the recognition of 'volutin' granules, it became generally accepted that metachromatically staining granules contain polyphosphates. However, Martinez (1963) has recently challenged this view. He showed that the metachromatically-staining material in Spirillum was not polyphosphate, but rather a polymer composed of poly- β -hydroxy butyrate. Since it is known that polymeric substances other than polyphosphates, such as glycoproteins and phospholipids, may also exhibit metachromatic

properties, Martinez, as well as other authors (Drews, 1962), suggest that all cytological identifications of polyphosphates must be confirmed by more rigid chemical isolations and identifications.

Although it is generally accepted that in many cases at least part of the metachromatically staining material of cytoplasmic inclusions is made of polyphosphates, it is not at all clear what other components are present. The following are some of the suggestions found in the literature.

Kulaev and Belozerskii (1958) are strong proponents of the view that polyphosphates are associated in vivo with RNA. They have isolated this complex from Aspergillus niger. The nature of both natural and artificial polyphosphate-RNA complexes has also been investigated in some detail by Ebel and his associates (Ebel, Dirheimer, Stahl and Yacoub, 1963; Dirheimer, Muller-Felter, Weil, Yacoub and Ebel, 1963; Stahl and Ebel, 1963). They confirmed the findings of the Russian authors that natural complexes of polyphosphate and RNA could not be separated by electrophoresis or by precipitation with bases, but showed that these complexes were completely dissociated by charcoal. Covalent bonds are, therefore, not involved. The authors favour the view that RNA-polyphosphate complexes are formed during extraction and do not occur in vivo, but they do not regard the latter possibility as being rigorously disproved.

From electron microscopic evidence Drews (1960) suggested that in Mycobacterium phlei polyphosphate granules are deposited on a protein matrix. Recently, Voelz, Voelz and Ortigoza (1966), showed that the mode of polyphosphate deposition in Myxococcus xanthus depended on the physiological state of the cells. Depending on the nutritional conditions under which the cells were grown, polyphosphate granules were associated with either glycogen inclusions, were dispersed in the cytoplasm or were embedded within the nucleoid.

On the basis of differential staining of cell smears of Aerobacter aerogenes and Corynebacterium xerosis before and after treatment with various degradative enzymes, Widra (1959) proposed a composite representation of a polyphosphate granule. This representation consists of a polyphosphate core which is surrounded by a lipoprotein-RNA envelope. This structure is possibly stabilised by Mg^{2+} ions, which bridge the negatively charged polyphosphates and lipoproteins.

It is apparent from the examples given above that the actual state of polyphosphate granules in the intact cells is still unknown and can only be resolved after successful isolation of polyphosphate granules from cells. So far attempts at this, involving differential centrifugation in aqueous media of mechanically disrupted cells (Drews, 1962; Harold and Miller, 1961; Harold, 1962(a)), have not been successful. In these experiments it was

shown that 'insoluble' polyphosphate fractionated together with cell-wall debris. Harold and Miller (1961) showed conclusively that the binding of the 'insoluble' polyphosphate to the cell-wall debris was an artefact of isolation.

Recently an attempt has been made to separate polyphosphate granules from Chlorella cells by differential centrifugation of cell homogenates in non-aqueous solvents (Hase, Miyachi and Mihara, 1963). Granules isolated in such a manner were shown to contain 'insoluble' polyphosphate, traces of RNA and also small traces of proteins.

5. Enzymes Involved in the Metabolism of Polyphosphates

A. Biosynthesis of Polyphosphates

(a) Polyphosphate kinase. Polyphosphate kinase is the only known enzyme involved in the biosynthesis of polyphosphates. This enzyme has been shown to catalyse the reaction:

where the terminal phosphoryl group of ATP is transferred to a polyphosphate 'primer'. Such a mechanism was originally predicted by Hoffmann-Ostenhof and Weigert (1952), and later shown to be partly correct when Yoshida and Yamataka (1953) reported the synthesis of polyphosphate-like material in crude yeast extracts supplemented with ATP. These authors also observed that the synthesis of metachromatically staining material proceeded in the

absence of oxidisable substrates, such as glucose, when ATP was present, but that oxidisable substrates were essential for polyphosphate formation when P_i only was used in the incubation mixture.

Hoffman-Ostenhof, Kennedy, Keck, Gabriel and Schonfeller (1954) showed, independently from the Japanese workers, that yeast extracts catalysed the formation of ATP from ADP and polyphosphate. They assumed that the reaction was reversible, and showed that AMP could not replace ADP in their system.

Kornberg, Kornberg and Simms (1956) partially purified from Escherichia coli an enzyme which catalysed the synthesis of 'acid-insoluble' polyphosphate from ATP. By using differentially labelled ATP, these workers showed that only the terminal phosphate of ATP was incorporated into polyphosphate. Mg^{2+} was essential for the activity, and fluoride strongly inhibited the reaction. No measurable quantities of the postulated 'primer' could be detected. The authors also showed that the enzyme catalysed the incorporation of PP_i into polyphosphate, but that PP_i was not an obligatory intermediate in polyphosphate synthesis. They could not detect the formation of short-chain intermediates during polyphosphate synthesis. The synthesis of polyphosphate was strongly inhibited by ADP. In a subsequent publication it was shown that this strong inhibition by ADP was due to the reversibility of the reaction, as shown in Eq. I.1 (Kornberg, 1957(b)). The apparent K_m values for

ADP, ATP, and polyphosphate were shown to be $4.7 \times 10^{-5} \text{ M}$, $1.4 \times 10^{-3} \text{ M}$ and $2.6 \times 10^{-5} \text{ M}$, respectively. The author speculated about the possible physiological significance of the low affinity of ATP for the enzyme, compared to that of ADP. She suggested that the ATP/ADP ratio in the living cell can be responsible for the regulation of polyphosphate biosynthesis or degradation. As will be shown later, this suggestion could find some justification, since it is now known that polyphosphates are deposited in vivo when the conditions for growth are unfavourable, while during active growth, when ATP concentration is presumably low, there is little or no polyphosphate found in the cell.

Muhammed (1961) partially purified a polyphosphate kinase from Corynebacterium xerosis. This enzyme was similar in many respects to that of E. coli. ADP and AMP strongly inhibited the activity of the enzyme, which showed an absolute requirement for Mg^{2+} . No 'primer' could be detected, nor was the polyphosphate synthesised through short-chain intermediates, as was shown by the addition to the incubation mixture of short-chain [^{31}P]-polyphosphate, which made no difference to the specific activity of [^{32}P]-labelled polyphosphate formed. These observations could suggest that the nascent polyphosphate molecule is attached to the enzyme, and that the terminal phosphates of ATP are continuously being added to the growing end of the molecule. The polyphosphate molecule is released from the enzyme only after a high state of polymerisation

has been achieved. Polyphosphate kinase from C. xerosis differed from that of E. coli, with respect to the reversibility of the reaction. The synthesis of ATP from ADP and polyphosphate could not be demonstrated in this organism.

Polyphosphate kinase is also present in the green sulphur bacterium, Chlorobium thiosulphatophilum (Cole and Hughes, 1965), in Aerobacter aerogenes (Harold and Harold, 1963; Harold, 1964) and possibly also in Aspergillus niger (Nishi, 1960), Mycobacterium smegmatis (Winder and Denneny, 1955) and Azetobacter vinelandii (Zaitseva, Belozerskii and Novozhilova, 1960(a)).

B. Breakdown of Polyphosphates

Published reports indicate that a number of different enzymes may be involved in the breakdown of polyphosphates. Some of these enzymes catalyse the transfer of the terminal phosphate group of polyphosphate to a variety of acceptor molecules. Other enzymes (polyphosphatases) catalyse the hydrolytic cleavage of polyphosphate to Pi.

(a) Polyphosphate kinase. As was shown in the preceding section, the polyphosphate kinase-catalysed reaction was shown to be reversible in a number of instances. Therefore, this enzyme can be regarded as a degradative enzyme as well as a synthetic one. Its degradative or synthetic action may be controlled by the levels of free ATP and ADP in the cell.

(b) Polyphosphate-AMP-phosphotransferase. Winder and Denneny (1957) first reported the transfer of phosphate from polyphosphate to AMP in crude extracts of Mycobacterium smegmatis, and this finding was confirmed by Szymona (1962) for M. phlei. This enzyme was partially purified by Dirheimer and Ebel (1965) from Corynebacterium xerosis. The latter workers showed that the purified enzyme preparation was distinct from polyphosphate kinase, polyphosphate glucokinase and adenylate kinase. They studied the forward reaction only, thus:

The enzyme was specific for AMP and for a polyphosphate of high molecular weight. Dirheimer and Ebel (1964) suggested that in vivo this enzyme could provide continuous supply of ATP through the mediation of adenylate kinase. This would be of advantage, since the polyphosphate kinase-catalysed reaction in this organism is virtually irreversible.

(c) Polyphosphate glucokinase. Szymona and Ostrowsky (1964) reported the presence in Mycobacterium phlei of an enzyme which catalysed the phosphorylation of glucose by polyphosphate, according to the reaction:

The enzyme was purified by gel-filtration, but only partial separation from contaminating ATP-specific glucokinases was achieved

(Szymona, Uryson and Kulaev, 1967). However, the authors proved conclusively that the phosphorylation of glucose by polyphosphate occurs without the mediation of ATP and polyphosphate kinase, and that the two glucokinases differed by a number of criteria. The same workers screened 19 species of microorganisms for polyphosphate glucokinase. The enzyme was found in 5 species of Actinomyces. Other microorganisms, including some groups of fungi, Azetobacter vinelandii, Rhodospirillum rubrum, and Chlorella vulgaris did not reveal polyphosphate glucokinase activity. However, all the strains studied were found to possess ATP-specific glucokinases.

A similar enzyme was purified 10-fold from extracts of Corynebacterium xerosis (Dirheimer and Ebel, 1962). The enzyme phosphorylated both glucose and glucosamine, but was inactive towards other hexoses, pentoses and nucleosides. Metaphosphates and short-chain polyphosphates were inactive as phosphoryl donors. The enzyme had an apparent K_m for polyphosphate of $0.7 \times 10^{-4} M$. Clear evidence was presented to show that the phosphorylation occurred directly and did not involve ATP as an intermediate.

(e) Polyphosphate fructokinase. Extracts of Mycobacterium phlei grown on fructose can utilise both ATP and polyphosphate for the phosphorylation of fructose to fructose-6-phosphate (Szymona and Szumilo, 1966). Both fructokinase activities were absent from extracts of glucose-grown organisms. Fructokinase activity

was clearly separated from the glucokinases by gel-filtration but attempts to separate the ATP-dependent fructokinase from polyphosphate-dependent fructokinase resulted in the loss of the latter activity. The evidence presented is consistent with the conclusion that M. phlei has two distinct inducible fructokinases, dependent upon ATP and polyphosphate, respectively, but falls short of a rigorous proof of the existence of a specific polyphosphate fructokinase.

(f) Polyphosphatases. Muhammed, Rogers and Hughes (1959) studied a partially purified polyphosphatase from Corynebacterium xerosis. The enzyme did not attack either P_Pi, trimeta- or tetrametaphosphates, but hydrolysed substrates of chain length higher than hexapolyphosphate. The product of the reaction was P_i, and formation of short chain intermediates could not be detected. The enzyme did not exhibit polyphosphatekinase activity in the presence of either ADP or ATP, nor did it exhibit polyphosphate-AMP-phosphotransferase activity. The enzymic activity was inhibited by a variety of metal ions, including Mg²⁺. In this respect this enzyme differed from a polyphosphatase isolated from Aerobacter aerogenes (Harold and Harold, 1965), which required both Mg²⁺ and high concentrations of a neutral salt for its activity (0.3M KCl). In this case, again, no short-chain intermediates could be detected, suggesting that the polyphosphate was hydrolysed by the progressive removal of terminal phosphate groups.

Enzymes from yeast, which specifically hydrolyse short-chain polyphosphates and metaphosphates were investigated in some detail by Mattenheimer (1956), who presented evidence that distinct enzymes catalyse the hydrolysis of tripoly-, tetrapoly-, trimetapoly-, tetrametaphosphate and Graham's salt. A variety of methods yielded only limited separation of the various enzymes.

Grossmann and Lang (1962) reported the presence of similar polyphosphatases in cell nuclei of the livers of rats, calf and swine. Again, the authors claimed to have achieved some separation of these enzymes using methods similar to those of Mattenheimer. These polyphosphatases split off one phosphate residue at a time, while the metaphosphatases only opened the ring structure.

6. Factors Controlling the Accumulation and the Degradation of Polyphosphates

Results obtained from the early experiments on the accumulation of polyphosphates indicated that this phenomenon was closely related to the exhaustion of an essential nutrient from the growth medium, and the availability of an oxidisable substrate, certain metal ions, and (of course) phosphate. It was also observed that large amounts of polyphosphates could be synthesised by organisms that were transferred into phosphate-rich media after a period of phosphate starvation. Wilkinson and Duguid (1960) have reviewed some of the early observations on the deposition of polyphosphates as a function of cultural conditions.

Polyphosphate accumulation under conditions of nutrient imbalance fell into two distinct patterns. One pattern involved the cessation of nucleic acid synthesis owing to the exhaustion of an essential metabolite. When growth and nucleic acid synthesis were blocked, assimilation of P_i from the medium continued and resulted in a steady accumulation of polyphosphates. Resumption of growth and of nucleic acid synthesis induced a rapid degradation of the polyphosphate with concomitant transfer of polyphosphate phosphorus to the nucleic acids. By the use of chloramphenicol to dissociate cell growth from nucleic acid synthesis, evidence was obtained for a competitive relationship between the nucleic acid synthesis and the accumulation of polyphosphate. The resumption of nucleic acid synthesis reduced the rate of polyphosphate synthesis and at the same time stimulated its degradation (Smith, Wilkinson and Duguid, 1954; Mudds, Yoshida and Koike, 1958; Harold, 1960; Harold, 1963).

The second set of experimental conditions under which polyphosphate deposition was shown to occur consists of transferring cells which were previously starved of P_i to a P_i -enriched medium. Under such conditions a rapid and extensive accumulation of polyphosphates occurs in the cell. As growth and nucleic acid synthesis resume in such cells, the intracellular concentration of polyphosphate returns to normal levels. This phenomenon was designated 'Polyphosphat Übercompensation' by Liss and Langen (1962), and this was translated as 'polyphosphate overplus' by Harold (1964).

The former authors were the first to report that there was a marked increase in the rate of polyphosphate synthesis in cells that were transferred to phosphate-enriched medium after a period of phosphate-deprivation.

Deposition of polyphosphates under conditions of nutritional imbalance and the phosphate-overplus phenomenon in A. aerogenes and in Neurospora crassa have been described in detail by Harold and his associates (Harold, 1960; Harold, 1962(a), Harold, 1962(b); Harold and Sylvan, 1963; Harold and Harold, 1963; Harold, 1964). These authors have demonstrated, by using a number of mutants of A. aerogenes, that in this organism the deposition and the degradation of the polyphosphates are under the control of at least two different enzymes, the polyphosphate kinase and the polyphosphatase, respectively. A mutant, lacking the polyphosphate kinase, could not accumulate polyphosphate under any nutritional conditions. On the other hand, another mutant that lacked the enzyme polyphosphatase was normal with respect to the accumulation of polyphosphates, but was not capable of degrading them. These authors also showed that the levels of polyphosphate kinase, polyphosphatase and the alkaline phosphatase in a wild strain of A. aerogenes were low during the exponential-phase of growth, but that the levels of all three enzymes were elevated during phosphate-deprivation.

Similarly, it was shown that in Corynebacterium xerosis the accumulation of polyphosphates occurred during the lag-phase and

the stationary-phase of growth, while during the exponential-phase this compound was consumed (Muhammed et al., 1959; Hughes and Muhammed, 1962). As with A. aerogenes, polyphosphate deposition and degradation in this organism is under the control of polyphosphate kinase and polyphosphatase, respectively.

Evidence that the accumulation of polyphosphate is the function of the state of cell growth has also been gained from work with yeast (Jeener and Brachet, 1944; Schmidt et al., 1946; Wiame, 1947(b); Wiame, 1948; Katchman and Fetty, 1955; Langen and Liss, 1958; Langen, Liss and Lohmann, 1962), Hydrogenomonas (Kaltwasser and Schlegel, 1959; Kaltwasser, 1962), Azetobacter (Zaitseva and Frolova, 1951; Zaitseva et al., 1960(a); Zaitseva et al., 1960(b)). Mycobacteria (Mudd et al., 1958), and in Chlorella (Miyachi, Kanai, Mihara, Miyachi and Aoki, 1964).

Thus, it appears that the phosphate-overplus phenomenon and the accumulation of polyphosphates during nutritional imbalance, and their utilisation after resumption of growth, are widespread among microorganisms.

It has been demonstrated (Mudd et al., 1958; Zaitseva et al., 1960(a); Langen et al., 1962; Miyachi et al., 1964) that the so-called 'acid-soluble' and 'acid-insoluble' polyphosphate fractions exhibit different degrees of metabolic activities, but so far the exact function and interrelationship of these fractions is not yet clear.

7. Theories on the Functions of Polyphosphates

A number of functions have been attributed to polyphosphates. Hoffmann-Ostenhof and Weigert (1952) proposed, on theoretical grounds, the 'phosphagen' function for polyphosphates. More recently, Hoffmann-Ostenhof (1962) summarised the evidence in support of this theory. His arguments are as follows.

(a) Though polyphosphates, in the strict sense of the word, do not belong to the class of compounds generally accepted as phosphagens, i.e. 'those naturally occurring N-phosphorylated guanidine derivatives which participate with adenine nucleotides in phosphoryl group-transfer reactions' (Morrison and Ennor, 1960), the polyphosphates are, nevertheless, thermodynamically 'energy-rich' compounds (Meyerhof, Skatas and Kaplan, 1953; Yoshida, 1955).

(b) Various organisms have been shown to possess enzymes which catalyse the transfer of terminal phosphate groups of polyphosphates to acceptor molecules such as ADP, AMP, glucose, fructose and glucosamine (cf. subsection I.5).

(c) Polyphosphates are known to accumulate when ATP supply exceeds its demand, and conversely, polyphosphates are degraded in cells where the ATP breakdown exceeds that of ATP generation (cf. subsection I.6).

(d) Polyphosphates occur only in organisms in which phosphagens are lacking. This argument may have to be modified as more is learned about the physiological functions of polyphosphates recently

found also in multicellular organisms that are known to possess the 'true' phosphagens.

Harold (1966), on the basis of results obtained in his laboratory, challenged some of the arguments proposed by Hoffmann-Ostenhof in favour of the phosphagen theory of polyphosphates in microorganisms. Thus, Harold and Harold (1965) showed that a mutant of A. aerogenes, which lacked the polyphosphatase but still contained the polyphosphate kinase, was unable to degrade accumulated polyphosphates, and on this basis postulated that, at least in this organism, polyphosphates cannot function as phosphagen. Also, in N. crassa, the inhibition of oxidative and fermentative ATP generation by 2, 4, -dinitrophenol and iodoacetate, respectively, drastically lowered the ATP level, without stimulating polyphosphate degradation. Again, these observations argue against the existence of a rapid equilibrium between polyphosphates and adenine nucleotides in this organism (Harold, 1962(b)).

Other proposed functions for polyphosphates are those of phosphate reserves and phosphate economy (cf. Harold, 1966). As the P_i content of most natural environments tends to be low due to the insolubility of calcium phosphate, there is an obvious advantage in the ability to store phosphorus. However, storage of large quantities of free phosphate would not be possible, since it is known that P_i acts as an allosteric effector of a number of important enzymically catalysed reactions, particularly those involved in

glycolysis and gluconeogenesis (cf. review by Stadtman, 1966). Furthermore, large quantities of free P_i would invariably disturb the intracellular pH and osmotic pressure. Polyphosphates, on the other hand, owing to their strong affinity for metal ions, would not remain in free solution, and for this reason would be more suitable as phosphorus-storage compounds. If the storage hypothesis is correct, then the energy content of polyphosphates will be largely irrelevant to their physiological function (cf. Harold, 1966).

Some authors attribute to polyphosphates the function of regulators of metal ions concentrations in the cell (Katchman and Fetty, 1955; Katchman, Fetty and Bush, 1959; van Wazer and Campanella, 1950). However, the relatively low selectivity in the binding of ions by polyphosphates makes it somewhat difficult to believe in a regulatory function of this type (cf. Harold, 1966).

8. Participation of Inorganic Pyrophosphate in Enzymic Reactions

A. Production of Inorganic Pyrophosphate in Biosynthetic Reactions

Inorganic pyrophosphate is the product of many important and ubiquitous processes. A number of reviews have appeared on this subject (Kornberg, 1957(a); Hoffmann-Ostenhof and Slechta, 1958; Imsande and Handler, 1961; Hoffmann-Ostenhof, 1962; Kornberg, 1962). The mechanism, by which most of these reactions proceed,

has been shown to be a nucleophilic attack upon the α -phosphorus of ATP, or other triphosphate, by an electronegative group, splitting out PPi, thus (Kornberg, 1957(a)):

In most biological systems the biosynthesis of the important intermediates (cf. Table 1, Kornberg, 1962) is ensured by the constant removal of PPi. This is accomplished by very active inorganic pyrophosphatases, which catalyse the hydrolytic cleavage of PPi to Pi. The one known exception to this generalisation may be the case of cell nuclei, where pyrophosphatase activity is minimal. It is also noteworthy that the four enzymes which catalyse the pyrophosphorolysis of UDP-glucose, UDP-galactose, UDP-galactosamine, NAD and deamido NAD, occur almost solely within the nuclei (Imsande and Handler, 1961).

Boyer (1958) raised the objection that the hydrolysis of PPi wasted 'high energy' bonds, and suggested that, within the cell, PPi may not always be accessible to the pyrophosphatases. On the other hand, Kornberg (1962) pointed out that, while the removal of PPi may be energetically wasteful, it is justified in that it ensures the synthesis of important metabolites.

Although these reactions which involve the release of PPi are of great biological importance, they are not relevant to the work presented in this thesis and, therefore, will not be discussed further.

B. Inorganic Pyrophosphate as Phosphoryl-Group Donor in Biosynthetic Reactions

(a) Inorganic pyrophosphate-glucose phosphotransferase. In 1960 Rafter demonstrated that mouse-liver mitochondria catalysed the formation of glucose-6-phosphate from glucose and PPi (Rafter, 1960). Under the optimal conditions used by Rafter, glucose-6-phosphate formation represented only a very small fraction of the total PPi hydrolysed, but the hydrolytic and the phosphorylating activities were parallel. As shown below, this low phosphorylating activity was due to the low concentration of glucose present.

Rafter's observation was later studied in great detail by two independent groups in the U.S.A., those of Stetten and of Nordlie. Both of these groups have shown that microsomal preparations from rat liver and kidney contain a particulate enzyme that catalyses the following three reactions:

The three reactions were shown to be the function of the same enzyme on the basis of thermal inactivation, activation with

detergents, molybdate inhibition and kinetic data (Nordlie and Arion, 1964; Arion and Nordlie, 1964; Stetten, 1964; Stetten and Taft, 1964). The enzyme does not require Mg^{2+} for its activity, which is in contrast to the absolute requirement for Mg^{2+} exhibited by many other inorganic pyrophosphatases. The microsomal enzyme preparation lacks hexokinase activity. The enzyme exhibits low specificity towards the sugars and can transfer phosphoryl group from PPI to a number of heptoses, hexoses, pentoses, the triose glyceraldehyde and a number of alcohols (Stetten, 1965). While there is a competitive inhibition between glucose-6-phosphate and PPI (Eq. I.5), the effect of glucose on the pyrophosphatase activity is non-competitive (Eq. I.4). At high glucose concentrations (0.4M) the synthesis of glucose-6-phosphate from PPI predominates over the hydrolytic reaction in a ratio of 8.4:1 and at 0.2M glucose this ratio is 7.7:1 (Stetten, 1964; Stetten and Taft, 1964; Nordlie and Arion, 1964).

Both groups of workers have shown that the three enzymic activities of the microsomal preparations are elevated to the same degree when rats from which these preparations are obtained were fasted, made diabetic with alloxan or treated with glucocorticoids. The levels of all three enzymic activities returned to normal after insulin treatment or adrenalectomy (Nordlie, Arion and Glende, 1965; Fisher and Stetten, 1966). This evidence lends further support to the claim that the three activities are catalysed by the

same enzyme. Arion and Nordlie (1967) showed that the increase in the three activities brought about by glucocorticoid administration to rats is not abolished by actinomycin D treatment (10 μ g. per 100 g. body weight, daily for 3 days), thus precluding new protein synthesis.

The exact role of this enzyme in vivo is not yet understood. One suggestion is that its role is to phosphorylate hexoses when their concentrations rise (Fisher and Stetten, 1966). This suggestion is attractive, when it is recalled that in vitro high glucose concentrations favour glucose-6-phosphate formation from P_{Pi}, in preference to the hydrolytic cleavage of the latter. It is also known that the hexokinase activity is lowered in diabetic animals (Weber, Singhal, Stamm, Lea and Fisher, 1966).

(b) Phosphoenolpyruvic carboxytransphosphorylase. This enzyme was isolated and partially purified from Propionibacterium shermanii by Siu and Wood (1962) and was found to catalyse the following reversible reaction:

It was shown that equivalent amounts of phosphoenolpyruvate and Pi were formed from oxaloacetate and P_{Pi}. GDP and ADP could not replace Pi as phosphate acceptors, which was proof that the enzyme preparation was not contaminated with phosphoenolpyruvic carboxykinase.

(c) Phosphopyruvate synthetase. Hatch and Slack (1968) recently reported the isolation and partial purification, from leaves of certain tropical grasses, of an enzyme which catalyses the reversible conversion of pyruvate, ATP and Pi into phosphoenolpyruvate, AMP and PPi, thus:

The enzyme preparation was partially purified from other contaminating enzymes, and had very low activities of inorganic pyrophosphatase. Studies with [^{32}P]-labelled orthophosphate provided evidence that one of the phosphate groups of PPi was derived from Pi. Both PPi and AMP were required for the formation of pyruvate from phosphoenolpyruvate, and Pi could not replace PPi. The authors obtained a value for the equilibrium constant of 2×10^{-4} for the reaction proceeding towards phosphoenolpyruvate synthesis. However, in vivo, the removal of any of the products could ensure that the reaction would proceed in the direction of phosphoenolpyruvate formation. This indicates that in vitro the reaction would greatly favour pyruvate formation. This may be the case, since the authors reported that in crude extracts of the grasses studied the activities of phosphoenolpyruvate carboxylase, inorganic pyrophosphatase and adenylate kinase were in excess of the maximum photosynthesis rates.

(d) Pyrophosphate as Energy- and Phosphoryl-Group Donor in Energy-Coupled Reactions of Oxidative and Photosynthetic Phosphorylation. The first implication that PPi is involved in these reactions came from the work of Baltscheffsky and his associates,

who showed that P_{Pi} is formed during the illumination of chromatophores from the photosynthetic bacterium Rhodospirillum rubrum (Baltscheffsky, von Stedingk, Heldt and Klingenberg, 1966). They postulated at that time that there must exist in the electron transport chain of this organism an energy conservation site at which P_{Pi} is involved. They later demonstrated the presence of such a site, where P_i was phosphorylated to the P_{Pi} level. With the aid of the inhibitor oligomycin, it was shown that this phosphorylation occurs at the pre-adenosine diphosphate level, at which ADP is phosphorylated to ATP (Baltscheffsky and von Stedingk, 1966). More recently, Baltscheffsky (1967) reported that P_{Pi} can function in the chromatophores as a donor of energy for the reversal of the electron flow through the cytochrome chain. Thus, addition of P_{Pi} to chromatophores changed the steady state oxidation levels of the different electron carriers, as does ATP in respiring mitochondria (Klingenberg, 1961). The author also showed that P_{Pi} could act as an energy donor not only in such primitive systems as bacterial chromatophores, but also in respiring mitochondria from both lower and higher organisms.

From the evidence presented above it seems clear that, at least in some instances, the 'high energy' of the phosphoanhydride bond of P_{Pi} can be conserved in a direct transfer of one phosphoryl group to a suitable acceptor.

CONTENTS

	Page
Summary	34
1. Introduction	36
2. Materials and Methods	37
A. Materials	37
B. Methods	37
3. Results	41
A. Optimal Conditions for the Deposition of Inorganic Pyrophosphate Granules in <u>Tetrahymena</u>	41
B. The Hydrolysis of Inorganic Pyrophosphate Granules in <u>Tetrahymena</u>	44
C. Absolute Requirement for Metal Ions and Phosphate for the Deposition of Inorganic Pyrophosphate Granules in <u>Tetrahymena</u>	44
D. Uptake of Calcium and Magnesium Ions by <u>Tetrahymena</u>	48
4. Discussion	56

Chapter II

CONDITIONS GOVERNING THE DEPOSITION OF INORGANIC

PYROPHOSPHATE IN TETRAHYMENA

CONTENTS

	Page
Summary	34
1. Introduction	36
2. Materials and Methods	37
A. Materials	37
B. Methods	37
3. Results	41
A. Optimal Conditions for the Deposition of Inorganic Pyrophosphate Granules in <u>Tetrahymena</u>	41
B. The Hydrolysis of Inorganic Pyrophosphate Granules in <u>Tetrahymena</u>	44
C. Absolute Requirement for Metal Ions and Phosphate for the Deposition of Inorganic Pyrophosphate Granules in <u>Tetrahymena</u>	44
D. Uptake of Calcium and Magnesium Ions by <u>Tetrahymena</u>	48
4. Discussion	56

SUMMARY

1. Tetrahymena pyriformis were shown to deposit cytoplasmic granules, consisting of calcium, magnesium and pyrophosphate, when cultures reached late stationary phase of growth or when phosphate-deprived cells were allowed to resume growth in phosphate-enriched growth media.
2. The deposition of pyrophosphate granules has been shown to require the presence in the growth medium of two different divalent metal ions, preferably calcium and magnesium, as well as phosphate.
3. Isolated pyrophosphate granules were shown to contain the divalent metal ions in nearly constant proportions, regardless of the concentrations of these in the medium.
4. Calcium, magnesium and phosphate are taken up actively by Tetrahymena. Large quantities of these metal ions are taken up only under conditions that favour the deposition of pyrophosphate granules.
5. It was shown that pyrophosphate granules disappeared in cells that were subjected to phosphate deprivation. The hydrolysis of granules was accompanied by the release of calcium and magnesium

from cells. However, the kinetics of the release of these two metal ions differed.

6. It is proposed that the deposition of these granules in Tetrahymena is an enzymically catalysed process.

7. A number of possible physiological functions for pyrophosphate granules are suggested.

These organelles accumulated large quantities of hydroxyapatite ($\text{Ca}_3(\text{PO}_4)_2 \cdot \text{Ca}(\text{OH})_2$), but the process requires the presence of both phosphate and an oxidisable substrate. Although magnesium is not required for this process, mitochondria can accumulate also magnesium phosphate.

Pautard (1959) reported the presence of hydroxyapatite-like deposits in the protozoan Spizostoma spizum. Jones (1966, 1967) investigated the uptake of calcium and phosphate by this organism. Calcium is taken up only slowly and, once in the cell, is largely bound in a non-exchangeable form. Since calcium-uptake measurements were carried out in media containing phosphate and magnesium, no information is available whether phosphate and/or magnesium are essential for calcium uptake.

As mentioned in Chapter 1, inorganic pyrophosphate was shown to be accumulated in organs of certain insects (Niemelä, 1962).

Rosenberg (1966) showed that Tetrahymena can accumulate, under certain environmental conditions, large number of cytoplasmic granules. These granules were isolated from cell lysates and were

1. Introduction

Little is known about the accumulation in Nature of phosphate deposits other than polyphosphates. Even less is known about the role that cations play during the accumulation of these deposits.

Calcium phosphate deposits have been reported in mitochondria from a variety of tissues (Greenawalt, Rossi and Lehninger, 1964). These organelles accumulated large quantities of hydroxyapatite ($\text{Ca}_3 (\text{PO}_4)_2$, $\text{Ca} (\text{OH})_2$), but the process requires the presence of both phosphate and an oxidisable substrate. Although magnesium is not required for this process, mitochondria can accumulate also magnesium phosphate.

Pautard (1959) reported the presence of hydroxyapatite-like deposits in the protozoan Spirostomum ambiguum. Jones (1966, 1967) investigated the uptake of calcium and phosphate by this organism. Calcium is taken up only slowly and, once in the cell, is largely bound in a non-exchangeable form. Since calcium-uptake measurements were carried out in media containing phosphate and magnesium, no information is available whether phosphate and/or magnesium are essential for calcium uptake.

As mentioned in Chapter I, inorganic pyrophosphate was shown to be accumulated in organs of certain insects (Niemierko, 1962).

Rosenberg (1966) showed that Tetrahymena can accumulate, under certain environmental conditions, large number of cytoplasmic granules. These granules were isolated from cell lysates and were

shown to consist of calcium-magnesium pyrophosphate. Calcium and magnesium were found to be present in equimolar amounts.

The work reported in this chapter deals with the conditions that affect the deposition and utilisation of these cytoplasmic granules in Tetrahymena.

2. Materials and Methods

A. Materials

Chemicals used were of analytical grade or of the highest purity available. L-amino acids and certain vitamins used in the media were products of the Nutritional Biochemical Co., Cleveland, Ohio. Other media components were purchased from various sources.

[³²P]-Orthophosphate was purchased from the Australian Atomic Energy Commission. [⁴⁵Ca] and [⁸⁹Sr] were purchased as chlorides from the Radiochemical Centre, Amersham, England. Yeast inorganic pyrophosphatase was purchased from Sigma Chemical Co., St Louis, Missouri.

Tetrahymena pyriformis Wh 14 (mating type 2, variety 1) was originally obtained from Dr O.H. Scherbaum.

B. Methods

Growth media. Two types of growth media were used. One, referred to as the 'complex medium', contained 2.0 per cent (w/v) proteose peptone (Difco), 0.5 per cent (w/v) yeast extract (Difco) and 9×10^{-5} M Fe-chelate (Conner and Cline, 1964). These components

were dissolved in water at about 70° and the solution filtered through a layer of 'Hyflow' super cell. Conical flasks were filled with the medium to 0.3 of their nominal capacity, plugged, and autoclaved at 15 lb. for 15 min. Glucose was autoclaved separately as a 10 per cent (w/v) solution, and added under sterile conditions to give the final concentration of 1.0 per cent (w/v).

An alternative medium, which will be referred to as the 'defined medium', was used whenever a controlled environment was required. The composition of the defined medium was essentially that of the '2c' medium described by Kidder and Dewey (1951), with the following modifications: Tween 80, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, CaCl_2 and phosphate salts were omitted, and all nucleotides were substituted with corresponding nucleosides.

Maintenance of stock cultures. Tetrahymena stocks were maintained in upright screw-top tubes, which contained 5 ml. of the complex medium (less glucose). Transfers were made fortnightly.

The organism was grown axenically at 35° in the appropriate medium. The flasks were shaken on a 'Gyratory' shaker (New Brunswick Scientific Co.) at 170 oscillations per min. Unless otherwise stated, a flask containing 150 ml. of the complex medium was inoculated with 0.05 ml. of a stationary-phase cell suspension from stock tubes. Log-phase cells were then used for further inoculations. Cells were harvested by low speed centrifugation (5 min. at 500xg). All transfers of cells from the complex to the

defined growth medium were carried out under sterile conditions. Cell pellets were washed with 2-3 volumes of 0.9 per cent (w/v) NaCl solution. Growth curves of Tetrahymena in the two media are shown in Fig. II.1.

Histological demonstration of inorganic pyrophosphate. Cell suspensions were centrifuged in tapered centrifuge tubes in a MSE bench centrifuge (15 seconds at setting 5). The pellet was re-suspended in 2 ml. of 0.9 per cent NaCl solution. Smears were made on albumin-coated slides, which were then placed in an atmosphere of osmium tetroxide for 1 min. and dried in air. The cells were defatted by immersing the slides for several minutes in a mixture of chloroform-methanol (2:1 by volume). The cytoplasmic granules (which will be referred to subsequently as 'granules') were stained for inorganic pyrophosphate using the lead reagent (Ebel, Colas and Muller, 1958). The slides were placed for 10 min. in a solution of lead nitrate (5 per cent (w/v) lead nitrate in 0.1M acetate buffer, pH 3.5), washed with water and then placed into ammonium sulphide solution for 30 seconds. The slides were again washed with water and counterstained with eosin (1.0 per cent (w/v) eosin in 0.2 per cent (w/v) CaCl_2 solution) for 5 min. After a further wash with water the slides were dehydrated in aqueous alcohol solutions and then cleared in xylol prior to mounting.

Measurement of $^{45}\text{Ca}^{2+}$ and ^{32}P -orthophosphate-uptakes by Tetrahymena. Log-phase cells grown in the complex medium were

Fig. II.1. Growth curves of *Tetrahymena pyriformis* in the complex and the defined growth media.

Curve A : Complex growth medium inoculated at zero time with cells in the log-phase of growth.

Curve B : Defined growth medium supplemented with phosphate buffer, pH 7.0, 3 mM; CaCl_2 , 0.1 mM; MgCl_2 , 0.1 mM.

Washed cells in the log-phase of growth from the complex medium were used as inoculum at zero time.

harvested, washed and re-suspended in the defined medium. The cells were then shaken 3-6 hours in the Gyrotory shaker at 35° , to starve them of P_i , Ca^{2+} and Mg^{2+} . At the end of this period the cells were diluted with defined medium to a density of 2×10^5 cells per ml. 35 ml. of this suspension was transferred into 100-ml. conical flasks, and equilibrated at 35° for 5 min. The labelled salts were added and 1.0 ml. samples of cell suspension were removed at intervals. The cells were collected by filtration on oxoid filter membranes, which were then washed twice with 3 ml. of 0.9 per cent NaCl, dried, mounted on planchettes and counted in a Nuclear Chicago gas flow counter, fitted with a thin Mylar window.

Estimation of inorganic pyrophosphate. The cell-pellet from a 10 ml. sample of cell-suspension was extracted twice for 5 min. with 3 ml. of cold, 10 per cent (w/v) trichloroacetic acid (TCA). The TCA was extracted from pooled supernatants by shaking with cold ether (3x3 ml.). The pH of the aqueous extract was adjusted to 7.0 with 1N NaOH. Samples of this extract were incubated with yeast pyrophosphatase at 35° for 60 min. The incubation mixture in a total volume of 1.0 ml. contained: Tris-HCl buffer, pH 7.0, 50mM; $MgCl_2$, 3mM; yeast pyrophosphatase 1.0 unit (1 unit = amount of enzyme that will hydrolyse 1μ mole of PP_i per 1 min. at pH 7.2 and at 25°). Inorganic orthophosphate was estimated on samples before and after incubation with pyrophosphatase by a modified method of Fiske and Subbarow (Sumner, 1944).

Isolation of granules. Granules were isolated from cell-lysates by the method of Rosenberg (1966). In this method the cell pellet was suspended in cold 8M urea, and the suspension was then sonicated for 2-3 min. until the viscosity of the suspension was markedly decreased. The sonicated suspension was then centrifuged at 20,000xg for 15 min. and the supernatant discarded. The precipitate was washed once by resuspension in 8M urea, twice with 0.01M Tris-HCl buffer, pH 9.0, once with ethanol and finally with ether. The pellet was dried in a desiccator over P_2O_5 under reduced pressure.

Calcium and magnesium were estimated using a Perkin-Elmer Atomic Absorption Spectrophotometer with appropriate standards.

Cell counts. Cells were appropriately diluted with 0.9 per cent (w/v) NaCl solution and counted in a Celloscope Particle Counter (Lars Ljungberg & Co., Stockholm, Sweden), equipped with an aperture of 125 microns in diameter (cf. Plesner, Rasmussen and Zeuthen, 1964).

3. Results

A. Optimal Conditions for the Deposition of Inorganic Pyrophosphate Granules in Tetrahymena

Rosenberg (1966) showed that Tetrahymena grown in the complex medium and harvested in the stationary-phase of growth contained few granules. However, the cell content of granules was increased when the cells were grown for 40 hours in the defined medium of

Stockstad, Seaman, Davis and Hutner (1956), supplemented with 2mM phosphate.

Similar results were obtained when cells grown in the complex medium to the log-phase stage (3×10^5 cells/ml.) were transferred into the defined medium of Kidder and Dewey (see Methods). In Fig. II.2 are shown cells from a stationary-phase culture (24 hr., 5×10^5 cells/ml.) grown in the defined medium, supplemented with 3mM phosphate, 0.1mM CaCl_2 and 0.1mM MgCl_2 . These cells contained a large number of granules which were situated predominantly in the posterior part of the cell. In Fig. II.3 are shown cells from a stationary-phase culture (24 hr., 1×10^6 cells/ml.) of Tetrahymena grown in the complex medium supplemented with 2mM calcium and magnesium chloride. Only a few granules can be seen in these cells as compared to those shown in Fig. II.2.

It was also observed that Tetrahymena could form large number of granules in relatively short periods of time. This rapid accumulation was observed only in cells which were previously subjected to a period of phosphate- or metal ions-deprivation. The appearance of pyrophosphate in the cells was measured (Table II.1), and it was found that the level of free orthophosphate rose from 0.16 $\mu\text{mole phosphorus}/10^6$ cells at zero time to 1.94 $\mu\text{moles phosphorus}/10^6$ cells at 3 hr., while the level of the total pyrophosphate-phosphorus rose from undetectable amounts at zero time to 3.0 $\mu\text{moles pyrophosphate-phosphorus}$ at 3 hr. After reaching

Fig. II.3. Tetrahymena cells from a stationary-phase

Fig. II.2. Tetrahymena cells from a stationary-phase culture (24 hrs, 5×10^5 cells/ml) grown in the defined medium supplemented with 3 mM phosphate, 0.1 mM CaCl_2 and 0.1 mM MgCl_2 . Lead stain and eosin counterstain. x250.

Fig. II.3. Tetrahymena cells from a stationary-phase culture (24 hrs, 1×10^6 cells/ml) grown in the complex medium, supplemented with 2 mM CaCl_2 and 2 mM MgCl_2 . Lead stain with eosin counterstain. x250.

Table II.1. Inorganic Pyrophosphate and Orthophosphate in Tetrahymena After a Period of Calcium, Magnesium and Phosphate Deprivation

Cells from a log-phase culture (3×10^5 cells/ml.) grown in the complex medium, were harvested, washed and transferred into the defined medium (2×10^5 cells/ml.). 4 hours after transfer, CaCl_2 (0.1mM), MgCl_2 (0.1mM) and phosphate buffer, pH 7.0 (3mM) were added. 10-ml. samples of the cell suspension were removed just prior to the addition of salts (zero time) and at time intervals shown. Pyrophosphate and orthophosphate were determined as described under Methods.

* Time at which granules were first seen in cells.

** Cells contained large number of granules

Time of sampling (hour)	$\mu\text{moles Phosphorus}/10^6 \text{ cells}$	
	Inorganic phosphate = phosphorus	Inorganic pyrophosphate = phosphorus
0	0.16	<0.03
1	0.94	1.13
2*	1.62	2.03
3	1.94	3.00
4	1.53	2.92
5**	1.06	2.81

maximal levels, both pyrophosphate and phosphate levels decreased, but the rate of decrease of the former was slower than that of phosphate.

B. The Hydrolysis of Inorganic Pyrophosphate Granules in Tetrahymena

It was shown that hydrolysis of granules occurred in cells during phosphate deprivation. The hydrolysis of granules also occurred when cells were deprived of metal ions, but the rate of hydrolysis was much slower than that observed during phosphate deprivation. The disappearance of granules was reflected in the drop of the total pyrophosphate content of the cells (Table II.2).

C. Absolute Requirement for Metal Ions and Phosphate for the Deposition of Inorganic Pyrophosphate Granules in Tetrahymena

Cells grown in the defined medium which was supplemented with MgCl_2 and phosphate only, were devoid of granules (Fig. II.4), nor were granules detected in cells that were grown in the absence of either Mg^{2+} or phosphate (Table II.3). It was also shown that Tetrahymena cells formed granules when Sr^{2+} was substituted for either Ca^{2+} or Mg^{2+} . When $[\text{}^{89}\text{Sr}] \text{Cl}_2$ was used instead of either calcium or magnesium salts, isolated granules were found to be highly radioactive. However, both calcium and magnesium ions strongly inhibited the incorporation of $[\text{}^{89}\text{Sr}]^{2+}$ into granules (Table II.4).

Variations in the concentrations of calcium, magnesium and phosphate salts in the growth medium did not result in great

Table II.2. Pyrophosphate Levels in Cells Hydrolysing Granules

Cells were grown in the defined medium (600 ml.) supplemented with 3mM orthophosphate, and 0.1mM each MgCl_2 and CaCl_2 . Cells in the late stationary-phase of growth (7×10^5 cells/ml.) were harvested and resuspended in 4 ml. of the defined medium. Equal samples of this suspension were pipetted into each of four flasks containing 150 ml. of the defined medium and supplements, as shown. 10 ml. samples were removed at shown times and pyrophosphate was estimated as described under Methods.

Time of sampling (hour)	$\mu\text{moles pyrophosphate-phosphorus found per } 10^6 \text{ cells}$			
	A	B	C	D
	No further additions	PO_4^{3-} , 3mM; Ca^{2+} , 0.1mM	PO_4^{3-} , 3mM; Mg^{2+} , 0.1mM	Ca^{2+} , 0.1mM; Mg^{2+} , 0.1mM
0	3.20	3.02	3.17	3.16
3	1.80	2.94	2.86	1.75
6	0.82	2.56	2.78	0.91
9	0.41	2.03	2.58	0.36

Table II.3. Absolute Requirements for Metal Ions and Phosphate for the Deposition of Inorganic Pyrophosphate Granules in *Tetrahymena*

Condition	Supplement to medium (mM)			Deposition of granules
	Ca ²⁺	Mg ²⁺	PO ₄ ³⁻	
After Ca ²⁺ , Mg ²⁺ and PO ₄ ³⁻ depletion	0.1	0.1	0.0	+++

Fig. II.4. *Tetrahymena* cells from a stationary-phase culture (5×10^5 cells/ml) grown in the defined medium supplemented with 3 mM phosphate and 0.1 mM MgCl₂. CaCl₂ was omitted. Lead stain and eosin counterstain. x250.

Table II.3. Absolute Requirement for Metal Ions and Phosphate for the Deposition of Inorganic Pyrophosphate Granules in Tetrahymena

Cells were starved of Ca^{2+} , Mg^{2+} and orthophosphate in the defined medium for 6 hours. Then, CaCl_2 and MgCl_2 , 0.1mM, and phosphate buffer, pH 7.0, 3mM, were added. Cells were harvested 5 hours later and stained for pyrophosphate as described under Methods.

+++ Large numbers of granules present in cells.

- Granules absent from cells.

+ Few granules seen.

Condition	Supplement in medium (mM)			Deposition of granules
	Ca^{2+}	Mg^{2+}	PO_4^{3-}	
After Ca^{2+} , Mg^{2+} and PO_4^{3-} deprivation	0.0	0.0	0.0	-
A	0.1	0.1	3.0	+++
B	0.0	0.1	3.0	+
C	0.1	0.0	3.0	-
D	0.1	0.1	0.0	-

Table II.4. Inhibition of [^{89}Sr] $^{2+}$ Incorporation into Granules by Calcium and Magnesium Ions

Cells from a log-phase culture (600 ml., 3×10^5 cells/ml.) grown in the complex medium, were harvested and suspended in 10 ml. of the defined medium. 2.5 ml. of this cell-suspension was diluted into flasks A, B, C, and D, each containing 150 ml. of the defined medium. After 3 hr. cells were supplemented with phosphate (3mM), [^{89}Sr] $^{2+}$ (1 μCi , 10^6 cpm) and metal ions as shown. Six hours later the cells were harvested and granules were isolated as described under Methods. No granules could be isolated from cells in flask D. Granules were dissolved in 0.6 ml. of 1N HCl heated at 100° for 10 min., and 0.1 ml. samples of this solution were counted.

Condition	Concentration of metal ions in medium (mM)			Weight of granules (mg.)	[^{89}Sr] per mg. granules (cpm)
	Ca^{2+}	Mg^{2+}	Sr^{2+}		
A	0.1	0.1	0.1	5.0	1,360
B	0.1	0	0.1	4.7	12,800
C	0	0.1	0.1	4.6	13,640
D	0	0	0.1	-	-

changes in the composition of granules with respect to the divalent metal ions (Table II.5). It can be seen that even an one-hundred-fold difference in the ratio of calcium and magnesium ions in the medium resulted in comparatively small changes in the molar ratios of these ions in the isolated granules.

D. Uptake of Calcium and Magnesium Ions by Tetrahymena

The observation that phosphate, calcium and magnesium ions are necessary for the deposition of granules in Tetrahymena, prompted the further investigation of the pattern of Ca^{2+} and Mg^{2+} uptake by this organism. It was shown (Fig. II.5) that in the absence of Mg^{2+} or phosphate the cells take up Ca^{2+} from the medium at a comparatively slow rate, which levels off after 60 min. under these experimental conditions. It can be shown that this phase of Ca^{2+} uptake is not due to simple diffusion of Ca^{2+} into the cells, since at the point where the uptake ceases, the concentration of calcium in the cells ($0.06 \mu\text{g. atom Ca/mg. wet weight}$) exceeds that in the medium by a factor of 20, even assuming the entire volume of the cell to be water. When either Mg^{2+} or phosphate were added separately at the point where Ca^{2+} uptake ceased, there was a slight increase in the rate of Ca^{2+} uptake. However, when Mg^{2+} and phosphate were added simultaneously, there was a sharp increase in the rate of Ca^{2+} uptake, which continued at that high rate until all of the calcium was exhausted from the medium.

Table II.5. Metal Ions in the Growth Medium and the Composition of Inorganic Pyrophosphate Granules

Cells were grown in the defined medium to the late stationary-phase (7×10^5 cells/ml.). The medium was supplemented with 3mM phosphate and with metal ions as shown. Granules were isolated and calcium and magnesium estimated as described under Methods.

Ion in medium (mM)		Molar ratios in granules	
Ca ²⁺	Mg ²⁺	Ca ²⁺	Mg ²⁺
1.00	1.00	0.50	0.51
0.01, 0.10	1.00	0.39	0.55
1.00	0.01	0.65	0.32

Fig. II.5. $[^{45}\text{Ca}]$ -uptake by Tetrahymena. Cells from a log-phase culture (3×10^5 cells/ml) grown in the complex medium were harvested and transferred to the defined medium, shaken for 3 hr and then diluted with defined medium to a density of 2×10^5 cells/ml. $[^{45}\text{Ca}]\text{CaCl}_2$ was added at zero time (10^{-5} M, 1000 cpm/ $\mu\text{mole Ca}^{2+}$). At 120 min MgCl_2 and/or phosphate were added to a final concentration of 10^{-4} M and 10^{-3} M, respectively. 1.0 ml samples were removed at shown times and cells prepared for counting as described under Methods.

Cells from such an experiment (Fig. II.5) were examined for granules and pyrophosphate both by staining and by the enzymic assay using yeast pyrophosphatase. Cells were removed prior to the commencement of Ca^{2+} -uptake, just prior to Mg^{2+} and phosphate additions and at the termination of the experiment. No granules were detected by the staining method prior to the addition of Ca^{2+} or just prior to the additions of Mg^{2+} and phosphate. However, granules were detected in cells removed at the end of the experiment. It was shown (Table II.6) that the level of total pyrophosphate in the cells increased from undetectable levels at the commencement of the experiment to 1.62 $\mu\text{moles pyrophosphate-phosphorus}/10^6$ cells at the termination of the experiment. The increase in the pyrophosphate level between zero time and just prior to the addition of Mg^{2+} and phosphate most probably represents the increase in the pool of the soluble cellular pyrophosphate. As will be shown in Chapter III, inorganic pyrophosphatase from cell-extracts of this organism is inhibited by calcium.

Both phases of $[\text{}^{45}\text{Ca}]^{2+}$ -uptake are temperature-sensitive (Fig. II.6). Thus, Mg^{2+} and phosphate did not stimulate the uptake of $[\text{}^{45}\text{Ca}]^{2+}$ in precooled cells (curve B), nor was there any uptake observed in the precooled cells in the absence of the other two ions (curve C).

Glucose had no effect on the pattern of $[\text{}^{45}\text{Ca}]^{2+}$ -uptake by Tetrahymena, nor did potassium or sodium ions at concentrations

Table II.6. Calcium-uptake and the Deposition of Granules in Tetrahymena

For experimental details see Fig. II.5.

Condition	$\mu\text{moles phosphorus}/10^6 \text{ cells}$	
	Inorganic phosphate	Pyrophosphate-phosphorus
Prior to Ca^{2+} addition	0.20	< 0.03
Prior to Mg^{2+} and PO_4^{3-} addition	0.25	0.30
End of experiment	1.12	1.62

Fig. II.6. Effect of temperature on the uptake of $[^{45}\text{Ca}]^{2+}$ by Tetrahymena.

For the experimental details see Fig. II.5.

Curve A : Cells kept at 35° throughout the experiment.

Ca^{2+} was added at zero time. Mg^{2+} and PO_4^{3-} were added at 2 hr.

Curve B : Ca^{2+} was added at zero time. At 115 min cells were placed in a 5° bath. At 2 hr Mg^{2+} and PO_4^{3-} were added.

Curve C : Cells were precooled at 5° for 5 min prior to the addition of Ca^{2+} at zero time. At 115 min cells were transferred to a 35° bath.

from $5 \times 10^{-6} \text{ M}$ to $5.0 \times 10^{-3} \text{ M}$. 2,4-dinitrophenol ($4 \times 10^{-4} \text{ M}$) had no effect on $[^{45}\text{Ca}]^{2+}$ -uptake in the absence of Mg^{2+} and PO_4^{3-} , but completely inhibited $[^{45}\text{Ca}]^{2+}$ -uptake in the presence of these two latter ions. However, it was shown that 2,4-dinitrophenol also inhibited the uptake of phosphate by Tetrahymena in the presence or the absence of the two metal ions (Fig. II.7, curves D, E and F). As expected, the rate of phosphate-uptake and the total amount of phosphate taken up by Tetrahymena are greater when the two metal ions are present in the medium (Fig. II.7, curves A and B) than in their absence (curve C). This observation could be explained on the basis of the requirement of these metal ions for both the deposition of granules and for the maintenance of the electrochemical balance within the cell.

Therefore, there can be at least two explanations for the inhibition of calcium-uptake in the presence of the inhibitor. Firstly, 2,4-dinitrophenol could have inhibited the production of the hypothetical 'high energy' intermediates associated with oxidative phosphorylation reactions. These intermediates are thought to play an important part in the active transport of ions across biological membranes (Lehninger, Carafoli, and Rossi, 1967). The observation that calcium-uptake in the absence of Mg^{2+} and phosphate was not inhibited by dinitrophenol, indicates that the energy for this slower phase of calcium-uptake can be supplied through energy-yielding reactions not sensitive to dinitrophenol,

Fig. II.7. Effect of 2,4-dinitrophenol on the uptake of [^{32}P]-orthophosphate by Tetrahymena.

Where [^{32}P]-orthophosphate uptake was measured in the presence of DNP, cells were pre-incubated with the inhibitor for 10 min. The additions at zero time were : [^{32}P]-orthophosphate, 1 mM (1000 cpm/ μmole); CaCl_2 , 0.1 mM; MgCl_2 , 0.1 mM. For other experimental details see Fig. II.5.

A: Mg^{2+} , Ca^{2+} , PO_4^{3-} ; B: Ca^{2+} , PO_4^{3-} ; C: PO_4^{3-} ; D, E and F: 0.4 mM DNP and Ca^{2+} , Mg^{2+} and PO_4^{3-} as in A, B and C, respectively.

such as those involved in substrate level-linked oxidative phosphorylation.

Secondly, by uncoupling respiration from energy conserving reactions, dinitrophenol will indirectly interfere with biosynthetic reactions responsible for the assimilation of orthophosphate and eventually with reactions in which pyrophosphate is produced. It was shown in experiments dealing with the uptake of calcium and phosphate (also see results from magnesium-uptake experiments, Fig. II.9) that the high rate of uptake was maintained only when the conditions for the deposition of granules were favourable. Once these conditions are changed, e.g., when no pyrophosphate is available for the deposition of granules, the high rate of uptake of metals ceased.

Cells that were kept under conditions that favour the hydrolysis of granules were shown to release calcium into the medium. The net loss of calcium from cells was independent of the external calcium concentration (Fig. II.8). It was also shown that under the same conditions cells release only about the third of the original calcium taken up. These observations suggest that there may be two compartments of calcium in the cells. The calcium is lost at a relatively rapid rate from the smaller of these two compartments, while the calcium in the larger compartment apparently does not exchange with the calcium from the medium.

Fig. II.8. The release of calcium from cells hydrolysing granules. Cells which accumulated a large number of granules in the defined medium supplemented with $[\text{}^{45}\text{Ca}]\text{CaCl}_2$ (0.1 mM), MgCl_2 (0.1 mM) and phosphate (3 mM), were harvested, and one-half of the cells was transferred (at zero time) into unsupplemented defined medium (A). The rest of the cells were transferred into defined medium supplemented with unlabelled CaCl_2 , 1.0 mM (B). Final cell density in both media was 1×10^6 cells/ml. Samples of cell suspension were removed at times shown and were membrane-filtered. $[\text{}^{45}\text{Ca}]$ in cell was estimated by counting the membranes (cf. METHODS). Total calcium content in the cell-free filtrate was estimated by absorption spectrometry.

The pattern of magnesium-uptake by Tetrahymena cells was found to be very similar to that of calcium (Fig. II.9). In the absence of calcium and phosphate the cells took up magnesium but the rate of uptake levelled off after about 120 min. When either calcium or phosphate was added to the medium at the point where magnesium-uptake ceased, there was an increase in the rate of magnesium-uptake. This increase was more pronounced upon the addition of phosphate alone than upon the addition of calcium alone. However, when calcium and phosphate were added simultaneously at the point where magnesium-uptake levelled off, there was a rapid increase in the rate of magnesium-uptake by the cells.

When cells containing granules were kept under conditions that favoured the hydrolysis of granules and the release of Mg^{2+} from the cells was followed, it was found (Fig. II.10) that the kinetics of the release of this ion were different from those operating during the release of Ca^{2+} (Fig. II.8). Thus, it was shown (Fig. II.10) that though the total net release of magnesium (0.16 μmole of magnesium/ml. of cell suspension) was independent of the external magnesium concentration, the rate at which the magnesium was released from cells was dependent on the concentration of this metal ion in the medium. While during the first 6 hours of the experiment no magnesium was released from cells that were suspended in the medium supplemented with 1mM Mg^{2+} , during the same period 0.02 μmole of magnesium was released from cells

Fig. II.9. Mg^{2+} uptake by Tetrahymena. The experiment was identical with that described in Fig. II.5 except that Mg^{2+} uptake was measured. $MgCl_2$ (1.0×10^{-5} M) was added at zero time and 5.0 ml samples were removed at shown times. Cells were centrifuged and the pellet was washed twice with 5.0 ml of 0.9% w/v NaCl, re-suspended in 1.0 ml of 1 N HCl and boiled for 15 min at 100° . Magnesium was estimated as described under Methods.

no further additions;

 1.0×10^{-4} M CaCl_2 ;

 1.0×10^{-3} M PO_4^{3-} ;

 1.0×10^{-3} M PO_4^{3-}

and 1.0×10^{-4} M CaCl_2

Fig. II.10. The release of magnesium from cells hydrolysing granules. Cells were grown, harvested and transferred to media (A) and (B) as described previously (Fig. II.8). Medium (B) contained 1.0 mM Mg^{2+} . Samples of cell suspensions were removed at times shown, cells were centrifuged, and the pellet was washed once with 10 volumes of 0.9% (w/v) NaCl. solution. The washed pellet was suspended in 2.0 ml of 1 N HCl and 2.0 ml of 10% (w/v) TCA and heated at 100° for 15 min. The precipitated protein was centrifuged off. The magnesium content of the TCA treated extract and of the cell-free medium was estimated by absorption spectrometry (see METHODS).

that were kept in the Mg^{2+} -free medium. After this time period, the rate of the release of the metal ion from both lots of cells was almost identical. After the transfer into the phosphate-free medium, the cells were monitored for the disappearance of granules by the staining procedure. No granules were detected in cells 6-7 hours after transfer.

Data presented in Fig. II.8 and Fig. II.10 were replotted for comparison in Fig. II.11. A number of striking differences are apparent in the manner of release of calcium and magnesium from cells during the hydrolysis of granules. Firstly, the total amount of calcium released (by a given number of cells) was more than 2.5 times greater than that of magnesium released. Secondly, calcium was released from the cells as soon as they were transferred into the phosphate-free medium, the release being almost completed by the end of about 8 hours. By this time most of the granules disappeared from cells. The rate of calcium release from cells was only slightly affected by the presence of 1.0mM calcium in the medium. On the other hand, very little magnesium was released from cells during the first 6 hours after transfer into phosphate-free medium. Furthermore, this release was completely abolished if 1.0mM magnesium was present in the medium, indicating that this phase of magnesium release may be a passive process. When most of the granules disappeared from cells, magnesium was released at a much faster rate, and this phase of release was not affected by the presence of magnesium in the medium.

Fig. II.11. The release of metal ions from cells hydrolysing granules.

For experimental details see Fig. II.8 and Fig. 11.10.

- A - Medium without metal ions.
- B - Medium containing 1 mM Ca^{2+} or Mg^{2+} .
- — ● Release of Ca^{2+} .
- ▲ — ▲ Release of Mg^{2+} .

4. Discussion

Tetrahymena were found to deposit inorganic pyrophosphate granules when treated in either of two ways. Firstly, in the confirmation of the results obtained by Rosenberg (1966), cells grown to the late stationary-phase of growth were found to deposit granules (Fig. II.2). Alternatively, cells subjected to short periods of phosphate deprivation rapidly deposited large numbers of granules when transferred to media supplemented with phosphate (Table II.1).

It was shown that the process of the deposition of granules in vivo was reversible, i.e., under certain conditions granules were shown to be hydrolysed in the cells. The hydrolysis of granules was reflected in the drop of the total pyrophosphate content of cells (Table II.2). The hydrolysis of granules was also found to occur during metal ions deprivation, but these conditions were not as effective as phosphate deprivation. The latter was found to be the most effective single condition causing the loss of granules.

The deposition of granules in Tetrahymena appears to be an enzymically controlled process. This is reflected in the constant composition of granules with respect to the metal ions (Table II.5), and also indirectly, by the effect temperature has on the uptake of metal ions by this organism (Fig. II.6). Although the concentrations of calcium and magnesium ions in the medium were varied

through a large range of concentrations, the ratio of these two ions in the isolated granules remained almost constant. If the deposition of granules in Tetrahymena was a non-enzymic process this ratio would have been expected to change with the change in the concentrations of these two metal ions in the medium.

The argument that the deposition of granules in Tetrahymena is an enzymically controlled process is further supported by the observation that both ions, as well as orthophosphate, are essential for the formation of granules (Table II.3). In the absence of any one of these, granules were not deposited. Although calcium or magnesium ions could be substituted by strontium ions, the incorporation of strontium into the granules was found to be strongly inhibited by both calcium and magnesium (Table II.4). Thus, it appears that the deposition of granules in Tetrahymena can occur only when two divalent metal ions are present, and that the two metals naturally occurring in the granules, calcium and magnesium, are the preferred pair.

Evidence obtained from calcium- and magnesium-uptake experiments shows that large amounts of these two metal ions are taken up by Tetrahymena only under conditions that favour the formation of granules (Figs II.5 and II.9). In the absence of one of the ions and phosphate, only comparatively small amounts of the other metal ion are taken up. Only when both the metal ions and phosphate are present in the medium is there a rapid uptake of the metal ions, followed by the deposition of granules in the cell.

It appears that not all of the calcium and magnesium taken up by the cells are bound in the form of granules (Fig. II.8 and Fig. II.10). Although cells that contained granules were kept for 24 hours under conditions that favoured the hydrolysis of granules, and although the granules could not be detected in these cells at the end of this time period, the cells still retained about 70 per cent of the total calcium and magnesium taken up. This suggests that the calcium and magnesium taken up by Tetrahymena are compartmentalised into, at least, two compartments. One, which is made up of the metal ions bound in granules, is actively removed from the cell during the hydrolysis of granules, while the metal ions in the second compartment are more tightly bound and are only very slowly exchangeable with the metal ions in the medium. Jones (1967) had observed a similar compartmentalisation of calcium in Spirostomum ambiguum. In this organism, at least two compartments of calcium exist, one which is freely exchangeable and one which is non-exchangeable.

It was found that during the hydrolysis of granules the metal ions (presumably previously bound in granules) were released by the cells into the medium (Figs II.8, 10 and 11). The pattern of the release of magnesium and calcium was not the same. Calcium was released immediately upon the transfer of cells into the phosphate-free medium, while magnesium was released only when the hydrolysis of granules and the release of calcium had apparently

been completed. These differential rates of release of the two metal ions may be of great importance in understanding the mechanism by which the granules are solubilised within the cell by the possible action of the inorganic pyrophosphatase. Since Ca^{2+} is a potent inhibitor of the pyrophosphatase from Tetrahymena, and since this inhibition is partly reversed by high magnesium concentrations (cf. Chapter III), the rapid preferential removal of free Ca^{2+} from the cell and the maintenance of a high Mg^{2+} concentration within, would aid the hydrolysis of granular pyrophosphate by this enzyme.

It is difficult to speculate on the possible physiological roles these granules may play in vivo. There is the possibility that they function as a phosphate-store in Tetrahymena. This suggestion is supported by the observation that granules are hydrolysed during phosphate deprivation. Also, these granules may have energy storage functions, especially in the light of the recent findings of the Baltscheffsky's group that pyrophosphate is implicated in certain reactions of the oxidative and photosynthetic phosphorylation (cf. Chapter I). This, and the phosphate storage function, would have been of great importance to the ancestors of this organism in the early days of evolution.

Another possible function of these granules in Tetrahymena may be in a detoxication mechanism which enables this organism to cope with otherwise toxic concentrations of both calcium and

pyrophosphate. It was shown that when cells are transferred into phosphate-enriched medium after a period of phosphate-deprivation the pool of the total cellular pyrophosphate increased rapidly. This was most probably due to the resumption of various biosynthetic reactions in which pyrophosphate is a by-product. Under optimal conditions, this pyrophosphate would not accumulate in the cell, since it would be rapidly hydrolysed to orthophosphate by cellular pyrophosphatases. However, it can be envisaged that under certain conditions these enzymes will be partially inhibited. This inhibition could result from a sudden increase in the intracellular concentration of calcium ions, a potent inhibitor of the pyrophosphatase in Tetrahymena (cf. Chapter III). It was also shown in calcium-uptake experiments that cells are capable of taking up certain amounts of calcium ions even when phosphate and magnesium were not present in the medium. These amounts of calcium would be inhibitory to the pyrophosphatase, especially if the metal ion is localised within a small compartment in the cell, where its concentration will be much greater than that calculated with the assumption that it was distributed evenly throughout the whole cell (cf. Results). Partial inhibition of the pyrophosphatase by calcium ions would result in the accumulation of free pyrophosphate, i.e. $P_2O_7^{4-}$, which in itself is an inhibitor of the pyrophosphatase (cf. Chapter III). To ensure that inhibitory concentrations of either pyrophosphate or calcium are not accumulated under such

conditions, the cell has developed the mechanism for the deposition of granules. Once deposited, these granules are not an end-product of metabolism, but can be re-utilised by the cell, e.g., during phosphate deprivation.

CONTENTS

	Page
Summary	62
1. Introduction	63
2. Materials and methods	63
A. Materials	63
B. Methods	63
3. Results	67
A. Partial Purification of the Soluble Pyrophosphatase	67
B. Some Properties of the Enzyme	71
(i) The Effect of pH on the Activity	71
(ii) Activation by Metal Ions	73
(iii) The Effect of Substrate Concentration	73
C. The Effect of Culture Age on the Activity of the Pyrophosphatase	73
D. The Hydrolysis of Pyrophosphate Granules by the Soluble Pyrophosphatase	75
E. The Effect of Phosphate-Deprivation on the Specific Activity of the Pyrophosphatase	75
4. Discussion	75

Chapter III

STUDIES ON THE INORGANIC PYROPHOSPHATASE

FROM TETRAHYMENA

	Page
Summary	62
1. Introduction	63
2. Materials and Methods	65
A. Materials	65
B. Methods	65
3. Results	67
A. Partial Purification of Inorganic Pyrophosphatase from <u>Tetrahymena</u>	67
B. Some Properties of the Enzyme	71
(i) The Effect of pH on the Activity	71
(ii) Activation by Metal Ions	73
(iii) The Effect of Substrate Concentration	73
C. The Effect of Culture Age on the Activity of the Pyrophosphatase	73
D. The Hydrolysis of Pyrophosphate Granules by the Soluble Pyrophosphatase	75
E. The Effect of Phosphate-Deprivation on the Specific Activity of the Pyrophosphatase	75
4. Discussion	78

SUMMARY

1. Isolation and partial purification of a constitutive inorganic pyrophosphatase from Tetrahymena is reported. Some catalytic properties of this enzyme are discussed.

2. It was shown that the pyrophosphatase from Tetrahymena hydrolysed pyrophosphate associated with the isolated granules, but only in the presence of a large excess of Mg^{2+} .

3. Orthophosphate-deprivation did not affect the specific activity of this enzyme. It is suggested that this enzyme is not involved in the regulation of hydrolysis of granules in vivo.

1. Introduction

Inorganic pyrophosphatases are widely distributed in Nature. They have been reported in various bacteria, yeasts, higher plants and animal cells (cf. review by Kunitz and Robbins, 1961). These enzymes catalyse the hydrolytic cleavage of inorganic pyrophosphate to orthophosphate, thus:

These enzymes were generally found to be specific for inorganic pyrophosphate.

Although the equilibrium of the pyrophosphatase-catalysed reaction lies far towards orthophosphate formation, the reversal of the reaction had been demonstrated. Thus, Cohn (1958) had convincingly demonstrated exchange of Pi (both [^{18}O]- and [^{32}P]-labelled) into PPi using yeast pyrophosphatase. Robbins and Lipmann (1958) used sulphate adenylyltransferase system to trap PPi synthesised by pyrophosphatase:

Recently Josse (1966(a)) also demonstrated the reversal of the pyrophosphatase-catalysed reaction, by trapping the pyrophosphate formed to pyrophosphorylate thymidine-diphosphate-glucose to thymidine triphosphate and glucose-1-phosphate, thus:

These reactions were coupled to the strongly exergonic conversion of glucose-6-phosphate to 6-phosphogluconic acid in the presence of phosphoglucomutase and glucose-6-phosphate dehydrogenase.

Although the role of inorganic pyrophosphatases in the overall cell metabolism is not certain, they may be of considerable importance in the control of the equilibrium of several vital synthetic reaction sequences. Pyrophosphate is a by-product of many important enzymic syntheses, including the polymerisation of deoxyribo- and ribonucleic acids, and the syntheses of coenzymes, proteins and phospholipids. Although some of these reactions are themselves exergonic, the free energy change is often small. However, the coupling of these reactions with the hydrolysis of pyrophosphate (standard free energy change at neutrality = 5 kcalories) markedly shifts the over-all equilibrium in favour of synthesis (Kornberg, 1962).

The disappearance of inorganic pyrophosphate granules in Tetrahymena under certain conditions, as reported in the previous chapter, raised the possibility that the pyrophosphatase may be the means by which the mechanism of the deposition and the hydrolysis of granules in vivo is controlled.

The work reported in this chapter describes experiments designed to investigate this possibility.

2. Materials and Methods

A. Materials

Ammonium sulphate (Special Enzyme Grade) was obtained from Mann Research Laboratories Inc., New York. Streptomycin sulphate was a product of Glaxo-Allenburys Pty. Ltd., Sydney. Triton X-100 (t-octylphenoxy polyoxyethanol) was obtained from Rohm and Haas Company. N-ethylmorpholine (NEM) was twice vacuum-distilled and was kept in brown-glass bottles at -15° . Other materials used were as described in Chapter II.

B. Methods

Growth media and the maintenance of stock cultures of Tetrahymena were described in Chapter II.

Isolation of Granules. Granules were isolated from cells by two different procedures. One (A) was similar to that described in Chapter II, with the exception that the granules obtained after the treatment with 8M urea were washed a number of times in 0.05M Tris-HCl buffer, pH 9.0, and then suspended in the same buffer at pH 7.5. The steps using the organic solvents and the drying of granules over P_2O_5 were omitted. Granules were also isolated by another procedure (B), from cells grown to the late log-phase (7×10^5 cells/ml.) in the complex growth medium, supplemented with $CaCl_2$ and $MgCl_2$ (0.5mM). Cells were harvested, washed with 0.9 per cent (w/v) NaCl solution and then re-centrifuged. This step was repeated two or three times. The cell-pellet was then

re-suspended in 0.05M Tris-HCl buffer, pH 7.5, and the cells were disrupted in a Ribi Cell Fractionator (Ivan Sorvall Inc., Norwalk, Connecticut) at 5,000 lb/in². The viscous cell-lysate was diluted with buffer to a protein concentration of 20 mg/ml. The diluted suspension was then centrifuged at 30,000xg for 30 min. Under these conditions the only material to sediment were the granules. A fluffy layer on top of the white granules was carefully washed off with the buffer. The granules were scraped out, suspended in buffer and re-centrifuged at 10,000xg for 10 min. This step was repeated several times. Generally, the granules were used immediately after isolation or otherwise were kept at 0° overnight.

Standard assay of the inorganic pyrophosphatase of Tetrahymena.

The incubation mixture, in a total volume of 1.5 ml., contained the following reagents, except when otherwise stated: Tris-HCl buffer, pH 7.5, 50mM; sodium pyrophosphate, adjusted to pH 7.5 with NaOH, 2mM; MgCl₂, 4mM, and sufficient enzyme (usually 0.1mg. of protein) to hydrolyse not more than 25 per cent of the substrate in 10 min. The enzyme preparation used, except when otherwise stated, was the dialysed ammonium sulphate fraction (Fraction V, cf. Results). The enzyme and substrate were incubated at 35° for 10 min. and the reaction was stopped by the addition of 0.15 ml. of ice-cold, 50 per cent (w/v) TCA. The hydrolysis of PPi was measured by the appearance of Pi. Reproducibility between duplicate assays was ± 3 per cent.

Under these conditions the rate of pyrophosphate hydrolysis remained constant for at least 20 min., when about 40 per cent of the substrate was consumed. The rate of pyrophosphate hydrolysis was also strictly proportional to the amount of protein added within the range of 0.01mg.-0.25mg. protein.

A unit of enzymic activity was defined as that amount of inorganic pyrophosphatase which hydrolysed 1 μ mole of inorganic pyrophosphate in 10 min. under the conditions stated above.

Orthophosphate estimation. Orthophosphate was estimated by a modified method of Fiske and Subbarow (Sumner, 1944).

Protein estimation. Protein was estimated by either of two methods. The biuret method (Layne, 1957) was used with protein concentrations greater than 1.0mg/ml. The Lowry method (Lowry, Rosebrough, Farr and Randall, 1951) was used with more dilute solutions.

Pyrophosphate estimation. Pyrophosphate content of granules was estimated as orthophosphate after heating at 100° for 10 min. in 1N HCl. Pyrophosphate content of cell-extracts was estimated also as orthophosphate released after incubation with yeast inorganic pyrophosphatase, as described in Chapter II.

3. Results

A. Partial Purification of Inorganic Pyrophosphatase from Tetrahymena

All operations were carried out at 0-5° unless otherwise stated. TE buffer refers to 0.05 M Tris-HCl buffer, pH 7.5,

containing 1mM EDTA (ethylenediamine tetraacetic acid). Cells were grown in twelve-2-liter flasks, each containing 600 ml. of the complex growth medium. Each flask was inoculated with 1.5 ml. of a late log-phase cell suspension (7×10^5 cells/ml.). Cells were harvested after 16 hours (5×10^5 cells/ml.) by low-speed centrifugation. The cell pellet (55 g. wet weight) was washed once with 0.9 per cent (w/v) NaCl solution and re-centrifuged. It was then resuspended in 150 ml. of TE buffer and the cells were disrupted in a Ribi Cell Fractionator at 5,000 lbs/in.². The cell lysate was centrifuged at 3,000xg for 15 min. and the pellet was discarded. The supernatant (Fraction I) was diluted with TE buffer to give a protein concentration of 10.0 mg/ml. The diluted extract was centrifuged at 30,000xg for 30 min., the supernatant decanted and the pellet was re-suspended in 50 ml. of TE buffer, containing 0.1 per cent (w/v) Triton X-100 detergent. The thick suspension was gently homogenised in a glass homogeniser and then re-centrifuged at 30,000xg for 30 min. The pellet was discarded and the two 30,000xg supernatants were pooled (Fraction II).

Streptomycin sulphate precipitation of the inactive material.

Streptomycin sulphate, 5 per cent (w/v) in TE buffer, was added to Fraction II dropwise and with stirring over a period of 30 min. 0.5 g. of streptomycin sulphate was added for each g. of protein. The amount of streptomycin added (0.5 g/g. protein) was found to be sufficient for the maximal removal of nucleic acids and other inactive materials as measured by the absorption of 260m μ and

280 μ . The use of higher concentrations of streptomycin resulted in the loss of some pyrophosphatase activity. The suspension was centrifuged at 10,000 $\times$ g for 10 min. The residue was discarded and the supernatant liquid was collected (Fraction III).

Heat treatment. To fraction III, 1M MgCl_2 solution was added to a final concentration of 10mM. Batches of 300 ml. of this solution were placed in a 2-liter flask and were heated in a steam bath with constant stirring for 5-6 min., by which time the temperature of the solution reached 70 $^{\circ}$. The contents of the flask were immediately transferred into another pre-cooled flask. After the suspension was cooled on ice to 5 $^{\circ}$ it was centrifuged at 10,000 $\times$ g for 10 min. The supernatant fluid was retained (Fraction IV).

Ammonium sulphate fractionation. Ammonium sulphate was added to Fraction IV as a fine powder, with stirring, over a 30-min. period. The amount of ammonium sulphate added was calculated using a nomogram of Dixon (Dixon and Webb, 1966). Precipitates were collected by centrifugation at 10,000 $\times$ g for 10 min. The fraction which precipitated between 0-65 per cent saturation was discarded. The precipitate obtained between 65-85 per cent saturation was suspended in the minimum volume of TE buffer and dialysed overnight against two changes of 200 volumes each of the same buffer (Fraction V). This procedure, summarised in Table III.1, yielded a 10-fold purification of the enzyme. Fraction V

Table III.1. Partial Purification of Inorganic Pyrophosphatase from Tetrahymena

For further details see text.

Procedure	Fraction	Protein (mg/ml.)	Specific activity (units/mg. protein)	Yield (%)
Crude extract	I	15.4	0.83	100
30,000xg supernatant	II	8.8	0.82	98
Streptomycin precipitation	III	4.8	1.39	90
Heat treatment	IV	1.4	6.30	80
Ammonium sulphate precipitation	V	10.6	9.00	75

was stable for at least three months when kept at -15° . Although usually stored at -15° (to prevent microbial growth), the enzyme was equally stable at 0° .

It was found that TE buffer used did not extract the enzyme effectively, since a large proportion of the activity was still associated with the material precipitating at 30,000xg (Table III.2). This activity was, however, extracted nearly quantitatively by TE buffer containing the non-ionic detergent Triton X-100. It is possible that the pyrophosphatase associated with this pellet is a particulate microsomal pyrophosphatase which is released from these cytoplasmic particles only in the presence of the detergent.

B. Some Properties of the Enzyme

(i) The Effect of pH on the Activity

The enzyme showed a pH optimum of 7.5 (Fig. III.1). It will be noted that the activity of the enzyme was considerably lower in the presence of N-ethylmorpholine than in the presence of Tris-HCl buffer. The pH optimum of the pyrophosphatase from Tetrahymena resembles more those reported for the pyrophosphatases obtained from rat brain (Gordon, 1950), erythrocytes (Block-Frankenthal, 1954) and yeast (Kunitz and Robbins, 1961), all of which are most active around the neutral pH values, rather than pyrophosphatases obtained from E. coli (Schito and Pesce, 1965; Josse, 1966), firefly (McElroy, Coulombre and Hays, 1951) and

Table III.2. The Solubilisation by Triton X-100 of the Inorganic Pyrophosphatase

For details see text.

Fraction	Treatment	
	TE buffer	TE buffer and Triton X-100
	Units of activity in total fraction	
30,000xg supernatant	1,580	2,180
30,000xg pellet	610	23

Fig. III.1. The effect of pH on inorganic pyrophosphatase activity.

The assay mixture contained in a volume of 1.5 ml : buffers adjusted to pH values as shown, 50 mM; MgCl_2 , 4 mM; sodium pyrophosphate, adjusted to pH 7.5, 2 mM; and sufficient enzyme (Fraction V) to hydrolyse not more than 25% of substrate in 10 min. Ammediol: 2-amino-2-methyl-1,3-propanediol; NEM:N-ethylmorpholine-HCl. For other details see METHODS.

insect muscle (Gilmour and Calaby, 1953) which have a pH optimum on the alkaline side of neutrality.

(ii) Activation by Metal Ions

There was an absolute requirement for a divalent cation in amounts suggesting stoichiometric combination with the substrate (Fig. III.2) Mg^{2+} was found to be the preferred cation under the conditions of assay used, though both Mn^{2+} and Zn^{2+} were effective as well. Monovalent cations (Na^+ and K^+) were not effective. High Mg^{2+} concentrations had a slight inhibitory effect on the enzymic activity. Ca^{2+} was found to be a strong inhibitor of the pyrophosphatase activity even in the presence of relatively high concentrations of Mg^{2+} (Table III.3).

(iii) The Effect of Substrate Concentration

The effect of varying substrate concentrations was studied by varying the concentrations of PPi while holding the metal ion concentration constant (Fig. III.3). These results show that high concentrations of free pyrophosphate ($\text{P}_2\text{O}_7^{4-}$) almost completely abolished the activity of the pyrophosphatase. Under optimal conditions for the hydrolysis of pyrophosphate, orthophosphate (20mM) had no effect on the activity of the pyrophosphatase.

C. The Effect of Culture Age on the Activity of the Inorganic Pyrophosphatase

The specific activity of the pyrophosphatase in cell-lysates remained constant and was not affected by the stage of growth at

Table III.3. The Inhibition of Pyrophosphatase Activity by Calcium Ions

Fig. III.2. Activation of the pyrophosphatase by metal ions.

The concentration of pyrophosphate was kept constant at 2 mM, while the metal ion concentration was varied as shown. For other details see "Standard Assay" conditions, under METHODS.

Table III.3. The Inhibition of Pyrophosphatase Activity by Calcium Ions

Standard assay conditions (see Methods) were used with the concentration of pyrophosphate kept constant at 2mM and those of Mg^{2+} and Ca^{2+} varied as shown.

Metal ion concentration (mM)		Activity %
Mg^{2+}	Ca^{2+}	
4	0	100
4	0.25	50
4	0.50	15
4	0.75	10
8	0	100
8	0.25	65
8	0.50	25
8	0.75	10

Fig. III.3. The effect of substrate concentration on the activity. Standard assay conditions were used (see METHODS), with the concentration of Mg^{2+} kept constant at 2 mM.

which the cells were harvested (Fig. III.4). Therefore, the pyrophosphatase studied in these experiments appears to be a constitutive enzyme, i.e., an enzyme whose synthesis has remained constant under these growth conditions (Cohn, Monod, Pollock, Spiegelman and Stanier, 1953).

D. The Hydrolysis of Pyrophosphate Granules by the Inorganic Pyrophosphatase

Inorganic pyrophosphatase from Tetrahymena readily hydrolysed the inorganic pyrophosphate associated with the granules isolated by either of the two isolation procedures used (Table III.4). Although the results shown refer to granules obtained by the procedure (A) only, similar results were also obtained with granules isolated by the procedure (B). In either case the hydrolysis occurred only when Mg^{2+} was present in high concentrations.

E. The Effect of Phosphate-Deprivation on the Specific Activity of Inorganic Pyrophosphatase

It was previously shown (Chapter II) that under conditions of phosphate-deprivation granules disappeared from cells. Though it was shown that the specific activity of the inorganic pyrophosphatase in cells grown in the complex medium was not affected by the age of the culture, the possibility remained that the activity of the pyrophosphatase generally, and towards the granules in particular, increased under the conditions of phosphate deprivation. To test this possibility, the cells were grown under conditions optimal for the deposition of granules (Table III.5).

Fig. III.4. The effect of age of culture on the activity. Cells were grown in 2-liter flasks, each filled with 600 ml of the complex growth medium. Cells were harvested at times shown, the pellet was washed with 0.9% (w/v) NaCl solution, and then resuspended in 50 ml of TE buffer containing 0.1% v/v Triton X-100. Cells were disrupted by repeated freezing and thawing. The particulate material was removed by centrifugation at 3,000 x g for 15 min and the pyrophosphatase activity was estimated on the samples of the supernatant as described under standard assay conditions (see METHODS).

Table III.4. The Hydrolysis of Pyrophosphate Granules by the Inorganic Pyrophosphatase from Tetrahymena and Yeast

The incubation mixture contained: Tris-HCl buffer, pH 7.5, 50mM; 0.1 ml. suspension of granules in the same buffer (2.8 μ moles PPi). Other additions were as shown. Tetrahymena pyrophosphatase: 0.2 ml. Fraction V (2.0 mg. protein); yeast pyrophosphatase: 1.0 international enzyme unit. Distilled water was added to the final volume of 2.0 ml. Samples were incubated at 35° for 60 min. The reaction was stopped by the addition of 0.2 ml. of cold, 50 per cent (w/v) TCA.

Additions	<u>Tetrahymena</u> pyrophosphatase		Yeast pyrophosphatase	
	PPi hydro- lysed (μ moles)	Per cent of total PPi	PPi hydro- lysed (μ moles)	Per cent of total PPi
None	0.20	7.2		
20 μ moles Mg^{2+}	0.28	10.0		
PPase alone	0.20	7.2	0.18	6.4
PPase + 1.4 μ moles Mg^{2+}	0.33	11.9		
" + 2.8 " "	0.67	23.5		
" + 5.6 " "	1.06	38.0	1.21	43.0
" + 20.0 " "	2.10	75.0	2.30	82.0

Table III.5. The Effect of Orthophosphate-Deprivation on the Specific Activity of the Inorganic Pyrophosphatase from Tetrahymena

The cells that were grown to the late log-phase of growth (5×10^5 cells/ml.) in 600 ml. of the defined growth medium supplemented with orthophosphate (3mM), CaCl_2 and MgCl_2 (0.1mM), were washed and transferred at zero time into 600 ml. of the defined growth medium without any supplements. At times shown, 100 ml. samples of the cell suspension were centrifuged, the cells were washed and suspended in 25 ml. of 0.05M TE buffer, pH 7.5, containing 0.1 per cent (v/v) Triton X-100. The cells were disrupted by repeated freezing and thawing. The debris were removed by centrifugation at 3,000xg for 10 min. Pyrophosphatase activity was measured by the standard assay (see Methods).

Time of sampling (hour)	Specific activity (units/mg. protein)
0.00	0.86
0.10	0.84
0.25	0.89
0.45	0.81
3.00	0.79
6.00	0.85

The cells, containing large number of granules, were transferred to a medium which favoured the hydrolysis of granules. The specific activity of the inorganic pyrophosphatase was not affected by these manipulations (Table III.5).

Since phosphate-deprivation had no stimulatory effect on the pyrophosphatase, the possibility remained that this enzyme, or another pyrophosphatase in the cell, could be activated if the cells were grown on media containing pyrophosphate rather than orthophosphate. However, it was not possible to demonstrate any increase in the pyrophosphatase activity in cell-lysates prepared from cells grown in the defined medium supplemented with 1mM sodium pyrophosphate. Again, no change in the specific activity of the pyrophosphatase could be demonstrated when pyrophosphate granules were used instead of synthetic pyrophosphate in experiments similar to the one described in Table III.5.

4. Discussion

The properties of the isolated and partially purified inorganic pyrophosphatase from Tetrahymena in many ways resemble those of yeast pyrophosphatase (Kunitz and Robbins, 1961) and of E. coli pyrophosphatase (Schito and Pesce, 1965; Josse, 1966). The Tetrahymena pyrophosphatase is a constitutive enzyme (Fig. III.4 and Table III.5) with optimum activity around pH 7.5 (Fig. III.1). The enzyme was shown to require a divalent metal ion for its activity (Fig. III.2). Mg^{2+} was found to give optimal activities

under the assay conditions employed, though Zn^{2+} and Mn^{2+} could replace Mg^{2+} -requirement. The results suggest that the actual substrate is not free pyrophosphate ($\text{P}_2\text{O}_7^{4-}$), but rather a complex consisting of equimolar amounts of Mg^{2+} and pyrophosphate. Free pyrophosphate strongly inhibited the pyrophosphatase-catalysed reaction (Fig. III.3). Similarly, it was found that Ca^{2+} is an inhibitor of this reaction (Table III.3).

The pyrophosphatase from Tetrahymena was shown to hydrolyse the inorganic pyrophosphate associated with the isolated granules (Table III.4). However, this hydrolysis occurred only in the presence of a large excess of Mg^{2+} (20 μmoles of Mg^{2+} as compared to 2.8 μmoles of pyrophosphate). Similar results were observed with yeast pyrophosphatase. The failure of the pyrophosphatase to hydrolyse granules (except in the presence of a large excess of Mg^{2+}) can be understood when it is recalled that granules contain equimolar amounts of calcium and magnesium. As the hydrolysis of pyrophosphate proceeds, both Ca^{2+} and Mg^{2+} are presumably released, and the increasing concentrations of Ca^{2+} ion in the assay mixture promptly inhibit the pyrophosphatase. However, this situation is not likely to arise in vivo. Thus, it was shown (Chapter II, Figs II.8, 10 and 11) that during the hydrolysis of granules calcium was rapidly and preferentially released from the cells, while magnesium was retained and eventually released only when the hydrolysis of granules was completed. This preferential

release of calcium and the maintenance of magnesium within the cell may be necessary to ensure the complete dissolution of granules by the soluble pyrophosphatase.

The pyrophosphatase activity was not affected by such environmental changes as phosphate-deprivation (Table III.5) or addition of pyrophosphate to the medium. This would suggest that the utilisation of granular pyrophosphate may involve factors other than the pyrophosphatase.

With the results in hand it will be hazardous to speculate on what these other factors are that may control the hydrolysis of granules in vivo. One of the possibilities is that the dissolution of granules is affected by an inducible enzyme, possibly similar to the alkaline phosphatase isolated from E. coli. This enzyme had been shown to be induced during phosphate-deprivation (Horiuchi, Horiuchi and Mizuno, 1959; Torriani, 1960) and Heppel, Harkness and Hilmo (1962) had shown that it hydrolyses pyrophosphate, as well as a variety of other phosphomonoesters. Since the optimal activity of this enzyme is in the alkaline pH range (8.4-9.3), the possibility remains that if it is present in Tetrahymena, its activity was not detected under the assay conditions used for the assay of the pyrophosphatase.

Another suggestion is that, during phosphate deprivation certain enzymes, possibly pyrophosphatases, are released from cytoplasmic structures similar to, or identical with, lysosomes

(de Duve, 1963). The methods used to disrupt cells in experiments reported in this chapter, e.g., freezing and thawing, use of high pressures, and the treatment of cell-lysates with detergent, would have also disrupted these fragile organelles and so caused the release of latent enzymic activities and thus made it impossible to distinguish between the latent and soluble pyrophosphatase activities. The present results do not allow to choose with certainty between these alternative suggestions.

THE INHIBITORY PYROPHOSPHATASE ACTIVITY ASSOCIATED WITH

PYROPHOSPHATE GRANULES ISOLATED FROM *TRITRINUM*

CONTENTS

	Page
Chapter IV	
Summary	82
1. Introduction	83
2. Materials and Methods	83
THE INORGANIC PYROPHOSPHATASE ACTIVITY ASSOCIATED WITH PYROPHOSPHATE GRANULES ISOLATED FROM <u>TETRAHYMENA</u>	83
3. Results	84
4. Discussion	93

CONTENTS

	Page
Summary	82
1. Introduction	83
2. Materials and Methods	83
A. Materials	83
B. Methods	83
3. Results	88
4. Discussion	93

1. Introduction

SUMMARY

1. It was shown that an inorganic pyrophosphatase is associated with pyrophosphate granules isolated from Tetrahymena pyriformis.

2. The activity of this enzyme was confined towards the pyrophosphate enclosed within the granule, and externally added pyrophosphate was not hydrolysed.

3. The pyrophosphatase activity was unaffected by phosphate deprivation of the cells.

4. Attempts to dissociate this activity from isolated granules were unsuccessful.

2. Materials and Methods

A. Materials

The anion exchange resin Dowex-1x8 (Cl⁻ 100-200 mesh) was a product of Fluka A.G. Whatman Chromatography Paper (304) was a product of V. & R. Balston Ltd. Other materials used were as described in previous chapters.

B. Methods

Growth media and the maintenance of stock cultures of Tetrahymena were described in the previous chapters. Cells were deprived

1. Introduction

The results presented in the previous chapter suggested that the soluble pyrophosphatase from Tetrahymena was a constitutive enzyme. Thus, it is most unlikely that the dissolution of granules in vivo is under the control of this enzyme alone, although the cell may hydrolyse granules through the agency of this enzyme. On the other hand, there are indications that the cell is capable of regulating both the hydrolysis and the deposition of these granules. Therefore, there must be other mechanisms operative in the cell that do control these processes.

A chance observation indicated that, under certain conditions, orthophosphate was released from isolated granules incubated in the absence of any added enzymes. The work in this chapter describes experiments designed to investigate this activity and the possibility that it may be involved in the hydrolysis of granules in vivo.

2. Materials and Methods

A. Materials

The anion exchange resin Dowex-1x8 Cl⁻ (100-200 mesh) was a product of Fluka A.G. Whatman Chromatography Paper (3MM) was a product of W. & R. Balston Ltd. Other materials used were as described in previous chapters.

B. Methods

Growth media and the maintenance of stock cultures of Tetrahymena were described in the previous chapters. Cells were deprived

of phosphate for 3-6 hours, and were then transferred to the 'defined' medium supplemented with [^{32}P]-orthophosphate, 3mM (0.15 mCi/mmol Pi); CaCl_2 , 0.1mM, and MgCl_2 , 0.1mM. Cells were harvested in the late stationary-phase of growth and the granules were isolated by procedure (A) (cf. Chapter III).

Protein content of granules was estimated by the Lowry method (Lowry et al., 1951), and the pyrophosphate content of the granules was estimated as orthophosphate after heating granules in 2N HCl at 100° for 15 min.

Standard assay for pyrophosphatase activity associated with granules. The incubation medium contained: 0.15M acetate buffer, pH 5.8, 0.25 ml.; the suspension of [^{32}P]-labelled granules in distilled water, 0.10-0.20 ml. (4-5 μmoles of inorganic pyrophosphate containing 0.10-0.20 mg. protein); distilled water to 1.0 ml. The samples were incubated at 35° for 30 min., and the reaction stopped by immersing tubes into boiling water for 3 min. The tubes were chilled on ice and then centrifuged.

The orthophosphate released was measured by the isobutanol extraction method (cf. below) when [^{32}P]-labelled granules were used, or, by the colourimetric method of Sumner (1944) when unlabelled pyrophosphate granules were used in the incubation medium.

[^{32}P]-orthophosphate estimation by the isobutanol extraction method. To 0.6 ml. of the centrifuged incubation medium were added: 6.7N H_2SO_4 , 0.15 ml.; 5 per cent (w/v) ammonium molybdate solution,

0.25 ml., and isobutanol, 1.0 ml. The contents of the tube were immediately stirred vigorously on a Vortex Mixer (Scientific Industries Inc., New York) for 30 sec. and then centrifuged for 1-2 min. in a bench centrifuge. The isobutanol layer was carefully transferred into another tube containing 1.0 ml. of ice-cold 1N H_2SO_4 previously saturated with isobutanol. The tube was stirred for a further 30 sec., centrifuged, and the isobutanol layer removed for counting. 0.10-0.20 ml. samples were pipetted into aluminium planchettes, lined with thin paper disks (Green's 105 Lens Tissue) and then dried under an infrared lamp. The planchettes were counted in a Nuclear Chicago gas-flow counter, fitted with a thin Mylar window. Each isobutanol sample was counted in duplicate and the average of two readings was recorded.

It was found (Table IV.1) that [^{32}P]-orthophosphate was quantitatively recovered by this procedure and that [^{32}P]-labelled pyrophosphate did not interfere with the extraction of orthophosphate.

Preparation and purification of [^{32}P]-labelled inorganic pyrophosphate. [^{32}P]-labelled pyrophosphate was prepared essentially as described by Kornberg and Pricer (1951). 1.0 mCi (0.25 ml) of [^{32}P]-orthophosphate of high specific activity (15 mCi/ μgP) in dilute HCl solution was evaporated to dryness in a porcelain crucible over a steam bath. After the addition of 0.5 ml. of 0.05M Tris-HCl buffer, pH 7.0, and 1.0 ml. KH_2PO_4 solution, 1.0mM,

the contents were again evaporated to dryness, and the crucible was then heated at 400° for 1 hr. The labelled pyrophosphate was separated from unreacted orthophosphate and from condensed phosphates of high molecular weights (which invariably form during heating) on Dowex 1x8 Cl^{-} (100-200 mesh) by the method of Peters and Rieman (1956). The dimensions of the column were: 1.5 cm. (internal diameter) x 15.0 cm. The column was previously equilibrated with 200 ml. of 0.25M KCl, pH 5.0. The cooled contents of the crucible were dissolved in 5.0 ml. of 0.25M KCl, pH 5.0, and were transferred onto the column. The crucible was washed with two-5 ml. portions of the same solution and the washings were added onto the column. 220 ml. of the same KCl solution were used to separate the unreacted orthophosphate from pyrophosphate. The flow rate was maintained at 2.3 ml./min., and the effluent was collected in 10 ml. fractions which were monitored for radioactivity. Fractions containing the separated pyrophosphate (routinely fractions 8-14) were pooled and concentrated to 4-5 ml. on a Büchi rotatory evaporator (W. Büchi, Flawil, Switzerland).

The yield of the labelled pyrophosphate was 85-90 per cent, as calculated on the total radioactivity of the initial [^{32}P]-orthophosphate. Of the total radioactivity associated with the isolated pyrophosphate, 1-2 per cent was due to [^{32}P]-orthophosphate contamination as was shown by the isobutanol extraction method (Table IV.1).

Table IV.1. The Estimation of [^{32}P]-orthophosphate in the Presence of [^{32}P]-labelled Pyrophosphate by the Isobutanol Extraction Method

Each tube contained: 0.15M acetate buffer, pH 5.8, 0.2 ml.; [^{32}P]-orthophosphate or [^{32}P]-pyrophosphate as shown; distilled water to 0.6 ml. The [^{32}P]-orthophosphate was extracted into isobutanol as described in the text. The two sets of results shown for pyrophosphate represent results obtained for two separate preparations of pyrophosphate

Additions	Total radioactivity (cpm)	Radioactivity extracted into isobutanol (cpm)	Radioactivity extracted into isobutanol as % of total radioactivity
[^{32}P]-pyrophosphate (7.5mM)	36,700	350	0.95
	38,200	750	1.96
[^{32}P]-orthophosphate (6.0mM)	24,250	24,070	99.3
[^{32}P]-pyrophosphate	18,350	12,010	97.0
[^{32}P]-orthophosphate	12,250		

The [^{32}P]-labelled pyrophosphate was free from condensed phosphates of molecular weights greater than pyrophosphate as was shown by paper chromatography with appropriate standards. Ascending chromatography on Whatman 3MM paper was employed with a trichloroacetic acid- ammonia- acetone solvent (Fuchs and Czech, 1963). The radioactive spots on paper chromatograms were located by exposing the latter for a suitable time interval to either Kodak 'Blue Brand' or Ilford Industrial-G films. Phosphates were detected on the chromatograms as described by Rosenberg (1959).

3. Results

It was demonstrated (Fig. IV.1, curve A) that isolated granules exhibited a pyrophosphatase activity. The hydrolysis of granular pyrophosphate was not detected when the granules were incubated at 3° or in the presence of 0.5mM HgCl_2 , a non-specific enzyme inhibitor (curves E and G, respectively). Nor was pyrophosphate hydrolysed in the boiled control (curve F), i.e., where granules were heated for 3 min. at 100° . These results suggest that the hydrolysis of granules is an enzymic process. Mg^{2+} was not required for this hydrolysis (curve B), nor did orthophosphate (1mM - 20mM) affect the dissolution of granules (results not shown in Fig. IV.1). The optimal pH for the granular pyrophosphatase was found to be about pH 6.0 (Fig. IV.2). This is to be contrasted with pH optimum of 7.5 for the activity of the soluble pyrophosphatase from Tetrahymena (Fig. III.1, Chapter III).

Fig. IV.1. The hydrolysis of granular pyrophosphate in vitro. Incubation medium (1.0 ml.) contained: [³²P]-labelled granules, 0.2 ml (5.75 μmoles PPi, 9560 cpm/μmole PPi, containing 0.21 mg. protein); 0.15M acetate buffer, pH 5.8, 0.25 ml.; other additions as shown; distilled water to 1.0 ml. For isolation of granules, estimation of [³²P]-ortho-phosphate and other assay details see Methods. Additions: A-none; B- 2mM Mg²⁺; C- 1mM Ca²⁺; D- 8mM Ca²⁺; E- none, incubation carried out at 3°; F- none, granules boiled for 3 min. at 100° before the addition of buffer, referred to in text as boiled blank; G- 1mM HgCl₂.

Fig. IV.2. The effect of pH on the activity of granular pyrophosphatase. The reaction medium contained in a volume of 1.0 ml: buffers adjusted to pH values as shown, 50 mM; [32 P]-labelled granules in distilled water, 0.20 ml (4.1 μ moles PPi containing 0.23 mg protein) and distilled water to 1.0 ml. For other details see Methods.

It will be noted that there was a sharp decrease in the activity of the granular pyrophosphatase after 30 min. of incubation (Fig. IV.1, curves A and B). This decrease was usually observed when 15-20 per cent of the total granular pyrophosphate was hydrolysed. An explanation for this was thought to be the increase in the free calcium concentration in the incubation mixture during the course of hydrolysis. It can be calculated that the free calcium at the end of a 30 min. incubation period (curve A) would be about $0.5 \mu\text{mole}/0.6 \text{ ml.}$, since for each mole of pyrophosphate hydrolysed 1.0 mole of each calcium and magnesium are released. However, this explanation does not seem to be the correct one. When granules were incubated in the presence of added calcium (curves C and D) no complete inhibition of the pyrophosphatase activity was observed, though the overall activity was inhibited. Thus, after 30 min. incubation under the experimental conditions described, the activity of the granular pyrophosphatase was inhibited by 54 per cent and 76 per cent in the presence of 1mM and 8mM calcium, respectively.

In vivo the granules are probably surrounded by a membrane (Fig. IV.3) across which the free calcium could be rapidly removed into the cytoplasm and from the cytoplasm into the medium surrounding the cell. The rapid removal of calcium across such a membrane and away from the site of granules hydrolysis may explain the complete dissolution of granules observed in whole cells kept in

Fig. IV.3. Electron-micrograph of pyrophosphate granules isolated from Tetrahymena. Granules were isolated by treating cells with 8M urea, then washed with Tris-HCl buffer, pH 9.0, pelleted and fixed in 5% (v/v) glutaraldehyde in 0.1M phosphate buffer, pH 7.3. The granules were then post-fixed in osmium tetroxide in veronal-acetate buffer, pH 7.3. The dehydrated sections were stained in a saturated solution of uranyl acetate in 50% (v/v) ethanol. x 60,000. (Electron-micrograph was kindly provided by Dr. H. Rosenberg).

phosphate-free media. An attempt was made to examine whether the sharp decrease in the hydrolysis of granules in vitro could be modified by the addition of Mg-EGTA [ethyleneglycol bis- (2-amino-ethylether) tetraacetic acid] to the incubation mixture. The stability constants of Mg-EGTA and Ca-EGTA complexes at pH 5.8 are 0.18 and 3.4×10^4 , respectively (Ringbom, 1963). The large difference in the stability constants of these two complexes would ensure that all of the free calcium released as the granules are hydrolysed will be complexed in the form of Ca-EGTA. Results of such an attempt are shown (Fig IV.4). There was a slight increase in the amount of pyrophosphate hydrolysed in the presence of EGTA (curves B and C), but the increase was not as marked as would be expected if calcium per se was the only inhibitor. Samples removed from tubes A and B at the termination of the experiment were assayed for pyrophosphate after the removal of granules by centrifugation. There was a large increase in the concentration of pyrophosphate in the tube containing EGTA (1.8 μ moles of pyrophosphate/0.6 ml. as compared to 0.2 μ mole/0.6 ml. in tube A). This increase of pyrophosphate in the supernatant was presumably due to the breaking up of granules after the chelation of granular calcium with EGTA.

Further indication for the enzymic nature of the hydrolysis of granules was obtained from experiments in which the hydrolysis of [32 P]-labelled pyrophosphate was measured in the presence of

Fig. IV.4. The effect of EGTA on the hydrolysis of granules in vitro. For experimental details see Fig. IV.1. The incubation medium (0.6 ml) contained 4.1 μ moles PPI (9325 cpm/ μ mole PPI) containing 0.27 mg protein. Additions: A- none; B- 0.02M Mg-EGTA, pH 6.0, 0.20 ml. added at zero time; C- 0.02M Mg-EGTA, pH 6.0, 0.20 ml added at 30 min. EGTA- [ethyleneglycol bis-(2-aminoethylether) tetraacetic acid].

unlabelled granules. No hydrolysis of externally added labelled pyrophosphate was detected (Table IV.2), though the granular pyrophosphate was hydrolysed to the usual extent. The evidence obtained from these experiments suggests that the granular pyrophosphatase is associated with the membrane surrounding the granule in a manner which allows the enzyme to act only on the pyrophosphate within the granule.

Attempts were made to dissociate the pyrophosphatase activity from the granules. Granules were dialysed against 0.01M EDTA in 0.05M acetate buffer, pH 5.8. This treatment removed all the pyrophosphate associated with the granules. The empty 'membranes' were collected by centrifugation, washed in cold distilled water and resuspended in acetate buffer. These 'membranes' were devoid of pyrophosphatase activity when incubated with 2mM pyrophosphate, pH 6.0, with or without 4mM Mg^{2+} . Nor could the pyrophosphatase activity be demonstrated in 'membranes' that were sonicated for 30 sec. (MSE Ultrasonic Disintegrator, current 1.4 amps at 240 volts). Also, no pyrophosphatase activity was detected in 'membranes' which were incubated for 1 hr. at 30° with Triton X-100 (0.1 per cent v/v), followed by gentle stirring in a glass homogenizer.

It was previously shown that during phosphate deprivation granules disappeared from cells. Attempts were, therefore, made to find out whether granules isolated from phosphate-starved cells

Table IV.2. The Behaviour of Granular Pyrophosphatase Towards Externally Added Pyrophosphate

The incubation medium (0.6 ml.) contained 3.1 μ moles of granular pyrophosphate (unlabelled, containing 0.18mg. protein). The hydrolysis of 'cold' granules was followed by the release of orthophosphate by the method of Sumner (1944). The hydrolysis of the added [32 P]-labelled pyrophosphate was followed by the estimation of the [32 P]-orthophosphate extracted into isobutanol. For other detail see Methods.

Additions	PPi hydrolysed (μ mole/0.6 ml/ 30 min.)	[32 P] orthophosphate extracted into iso- butanol (cpm/0.6 ml. reaction mixture)
'Cold' granules alone	0.21	-
" + [32 P]-PPi (2mM, 10,000 cpm)	0.23	128
" + [32 P]-PPi + 4mM Mg ²⁺	0.20	143
[32 P]-PPi alone	-	131

displayed high pyrophosphatase activity. However, no significant change in the pyrophosphatase activity was detected in granules isolated from cells which were transferred into phosphate-free medium after being allowed to synthesise granules in phosphate-rich growth medium (Table IV.3). Neither did the addition to the incubation medium of samples of cell-lysates obtained from cells grown under similar conditions increase this pyrophosphatase activity.

4. Discussion

Evidence was presented in this chapter that there was an inorganic pyrophosphatase activity associated with isolated granules. A possibility existed that this activity was due to the contamination of the granules with traces of soluble pyrophosphatase, especially since the soluble pyrophosphatase (Fraction V, cf. Chapter III) survived treatment with 8M urea under the experimental conditions employed for the isolation of granules. However, a number of observations reported in this chapter make this suggestion very unlikely. First, the granular pyrophosphatase did not hydrolyse externally added pyrophosphate, but was active only towards the pyrophosphate associated with the granules (Table IV.2). Second, the granular pyrophosphatase differed from the soluble enzyme in its pH optimum (Fig. IV.2) by almost 1.5 pH units - a further strong evidence that two distinct enzymes are operative in this system. Third, the granular pyrophosphatase was

Table IV.3. The Effect of Phosphate Deprivation on the Activity of the Granular Pyrophosphatase

The cells were grown to the late stationary-phase of growth in the defined growth medium supplemented with orthophosphate (3mM, 0.15mCi [32 P]-orthophosphate/mmmole Pi), CaCl_2 and MgCl_2 (0.1mM). They were then harvested and transferred at zero time into defined growth medium containing no supplements. Cells were removed at times shown. Granules were isolated and the activity of the pyrophosphatase estimated as described under Methods. Cell extracts were obtained from cells grown under the same conditions used for the isolation of granules, except that (32 P]-orthophosphate was not added to the medium. After transfer into unsupplemented medium, samples of cell-suspension were removed at times shown, harvested and suspended in 0.05M Tris-HCl buffer, pH 7.0. The cells were disrupted by freezing and thawing. 0.1 ml. samples of the cell-lysate were added to the incubation mixture.

Time after transfer into Pi-free medium	Activity of granular PPase, μ moles PPI hydrolysed/mg. protein/30 min.	
	Granules alone	Granules + cell extract
0.0	1.50	1.47
0.25	1.48	1.36
1.0	1.23	1.38
2.0	1.36	1.27
4.0	1.39	1.39

not as sensitive to calcium inhibition (Fig. IV.1) as was the soluble pyrophosphatase.

The addition of Mg-EGTA to the incubation medium significantly increased the amount of the granular pyrophosphate hydrolysed (Fig. IV.4). This would suggest that the sharp cut-off in the activity of the enzyme after 30 min. of incubation may be due to the inhibition of the pyrophosphatase by the increasing concentrations of calcium ions which are liberated during the hydrolysis of pyrophosphate. Although the complexing agent had other effects on the granules, as evidenced by the increase of pyrophosphate in the medium, which was presumably not associated with granules, nevertheless, EGTA mimicked the removal of calcium by the cells during hydrolysis of granules in vivo. We have shown (Chapter II) that calcium is rapidly released from cells and that this release paralleled the dissolution of granules.

The attempts to dissociate the pyrophosphatase activity from the granules were unsuccessful. The inability to demonstrate the pyrophosphatase activity upon incubation of 'membranes' with a detergent or upon the disruption of 'membranes' by sonication would suggest that a certain degree of structural organisation is obligatory for this enzymic activity.

It was shown (Table IV.3) that phosphate deprivation did not affect the activity of the granular pyrophosphatase. Neither was the activity affected upon the addition to the incubation medium

of cell-lysate obtained from cells which were hydrolysing granules. It is difficult to believe that in vivo the activity of this enzyme is not regulated in some manner. If the activity of this enzyme is 'turned on' permanently, one is presented with the problem how the granules are deposited at all.

From the results presented so far, it can be argued that the dissolution of granules in vitro may be accomplished through the agency of both the granular and the soluble pyrophosphatases. However, this may not be the case in whole cells. It has to be remembered that the procedures employed for the isolation of granules would, undoubtedly, damage the membranes surrounding them. It is, therefore, suggested that in the cell, where these membranes are intact, the breakdown of granules may be accomplished through the agency of the granular pyrophosphatase only.

CONTENTS

	Page
Summary	97
1. Introduction	98
2. Materials and Methods	101
A. Materials	101
B. Methods	102
3. Results and Discussion	112
Chapter V	
A. Attempts to Demonstrate the Formation of Phosphorylpyruvate from Oxalate	112
B. Attempts to Demonstrate the Pyrophosphate-Glucose Activity in <u>Tetrahymena</u>	113
C. Attempts to Demonstrate a Pyrophosphate-ADP-Phosphotransferase Activity in <u>Tetrahymena</u>	118
4. Conclusion	121

IS PYROPHOSPHATE A PHOSPHORYL GROUP

DONOR IN BIOSYNTHETIC REACTIONS IN

TETRAHYMENA?

CONTENTS

	Page
Summary	97
1. Introduction	98
2. Materials and Methods	101
A. Materials	101
B. Methods	102
3. Results and Discussion	112
A. Attempts to Demonstrate the Formation of Phosphoenolpyruvate from Oxaloacetate in the Presence of ATP or Pyrophosphate	112
B. Attempts to Demonstrate a Pyrophosphate-Glucose Phosphotransferase Activity in <u>Tetrahymena</u> .	115
C. Attempts to Demonstrate a Pyrophosphate-ADP-Phosphotransferase Activity in <u>Tetrahymena</u> .	118
4. Conclusion	131

SUMMARY

1. The experiments described in this chapter were designed to demonstrate a pyrophosphate-dependent glucokinase or phosphoenolpyruvate carboxykinase activities in cell-free extracts of Tetrahymena.

2. Attempts were also made to demonstrate phosphorylation of ADP by pyrophosphate in similar extracts. Although these attempts were unsuccessful, the results are not fully conclusive, and the possibility that such an activity is present in Tetrahymena could not be definitely excluded.

1. Introduction

In the previous chapters it was demonstrated that Tetrahymena pyriformis deposited pyrophosphate granules under certain environmental conditions. These granules were rapidly deposited in cells which were transferred into phosphate-enriched media after a period of phosphate deprivation. Furthermore, it was shown that these cytoplasmic inclusions were not inert end-products of metabolism in this organism, since they were re-utilised by cells during phosphate deprivation. If it is assumed for the moment that the deposition of pyrophosphate granules represents a means of phosphorus storage in this organism when this compound is in plentiful supply, then the next question to be asked is: what is the fate of the pyrophosphate-phosphorus during the utilization of granules in vivo? One possible route is its hydrolysis to orthophosphate through the agency of the soluble and the granular pyrophosphatases, which were demonstrated in this organism, with the consequence that the pyrophosphate-phosphorus is then channelled into the general phosphate pool of the cell and from there into the various metabolic reactions. In this case the energy locked in the anhydride bond of the pyrophosphate molecule would be dissipated as heat. An attractive alternative route would be some enzymically catalysed reaction in which one of the phosphoryl groups of the

pyrophosphate is used to phosphorylate directly another molecule. In this case, the energy of the anhydride bond will not be dissipated as heat energy, but will be transferred to the phosphoryl-group acceptor molecule.

Siu and Wood (1962) reported the isolation from Propionibacterium shermanii of a phosphoenolpyruvic carboxytransphosphorylase (PEP carboxytransphosphorylase) which catalysed the carboxylation of phosphoenolpyruvate (PEP) to oxaloacetate. The authors demonstrated that in vitro the reaction was reversible. This PEP carboxytransphosphorylase differed from phosphoenolpyruvate carboxykinase (PEP carboxykinase) isolated from avian liver (Utter and Kurahashi, 1954; Kurahashi, Pennington and Utter, 1957) in that it was specific for pyrophosphate and neither ATP, GTP or ITP could replace pyrophosphate as the phosphoryl-group donor. Warnock and van Eys (1962) and more recently Shrago and Shug (1966) and Shrago, Brech and Templeton (1967) had demonstrated the presence of a PEP carboxykinase in Tetrahymena. The latter workers had shown that this enzyme was rather specific for ATP and Mn^{2+} , and the activity was lowered by about 50 per cent with GTP, ITP and Mg^{2+} . The activity was evenly distributed between the mitochondrial fraction and the 105,000xg supernatant fraction. They also showed that the activity of the soluble PEP carboxykinase was almost completely repressed in cells grown in the presence of glucose. On the other

hand, the activity of the enzyme was higher in cells grown under low oxygen tension. Warnock and van Eys (1962) unsuccessfully attempted to demonstrate pyrophosphate requirement for PEP carboxykinase isolated from Tetrahymena. However, these authors grew this organism on a proteose peptone-yeast extract growth medium supplemented with glucose. It was, therefore, decided to investigate whether the soluble, glucose-repressible, PEP carboxykinase from Tetrahymena could phosphorylate oxaloacetate to PEP with pyrophosphate as a phosphoryl- group donor (rather than ATP).

Nordlie and Arion (1964) and Stetten (1964) demonstrated a pyrophosphate-glucose phosphotransferase activity in microsomal fractions of rat kidney and liver. These authors had shown that the phosphorylation of glucose by pyrophosphate did not depend on the presence of adenine nucleotides. The same enzyme also exhibited a strong pyrophosphatase activity. Since Tetrahymena possessed both a soluble pyrophosphatase and a pyrophosphatase associated with granules it became desirable to investigate whether a pyrophosphate-glucose phosphotransferase activity was associated with either of these two pyrophosphatases.

It was suggested by Hogg (see Levy and Scherbaum, 1965), that Tetrahymena may have to revert temporarily to anaerobic metabolism during the night when the oxygen tension in waters of its natural habitat may fall due to the cessation of oxygen production by

photosynthetic organisms. Recent reports indicate that this suggestion may be valid. Levy and Scherbaum (1965) demonstrated that the rate of glyconeogenesis in Tetrahymena was greatly increased when the organism was kept under low oxygen tension. Shrago, Brech and Templeton (1967) also showed that the lowering of oxygen tension in culture media of Tetrahymena greatly stimulated the activity of PEP carboxykinase, a key enzyme in glyconeogenesis.

Obviously, then, if this organism has to revert to anaerobic, or partly anaerobic, metabolism during dark periods, it will have to generate ATP by means other than the oxidative phosphorylation pathway. An obvious alternative would be the glycolytic pathway of glucose metabolism. Another possibility would be for Tetrahymena to utilize the granular pyrophosphate for direct phosphorylation of ADP to ATP. If such a reaction existed in Tetrahymena it would be of utmost significance from both the biochemical and the evolutionary points of view. Experiments will be described in this chapter that were designed to demonstrate the presence of an enzyme that would catalyse such a reaction.

2. Materials and Methods

A. Materials.

Glucose-6-phosphate dehydrogenase (yeast) and hexokinase were the products of Calbiochem, Los Angeles, California. The specific activity of glucose-6-phosphate dehydrogenase was 200 EU/mg. protein

(1 EU= 1mg of enzyme preparation which will cause the turnover of 1 μ mole substrate / min at 30⁰). The specific activity of hexokinase was 250 EU/mg. protein at 30⁰. Adenylate kinase (muscle) (specific activity not stated) was obtained from Nutritional Biochemicals Corporation, Cleveland, Ohio.

Triphosphopyridine nucleotide, adenosine 5'-diphosphate (sodium salt) and adenosine 5'-triphosphate (disodium salt) were the products of P-L Biochemicals Inc., Milwaukee, Wisconsin. Dithiothreitol (Cleland's reagent) was a product of Calbiochem.

Activated charcoal (Norit A) was washed in 2N HCl, followed by water and ethanol prior to use.

All other materials used were as described in previous chapters.

B. Methods.

Isolation of PEP carboxykinase from *Tetrahymena*. PEP carboxykinase was prepared as described by Shrago, Brech and Templeton (1967). *Tetrahymena* were grown at 34⁰ in 2-liter conical flasks filled with 600 ml. of a peptone medium of the following composition: 2 per cent (w/v) proteose peptone (Difco) and 0.1 per cent (w/v) yeast extract (Difco). The flasks were shaken at 170 oscillations/min. on a New Brunswick 'Gyratory' shaker. Cells were harvested in the logarithmic-phase of growth (5×10^5 cells/ml.) by slow speed centrifugation (500xg for 2-3 min.). When the effect of partial anaerobiosis on

the enzymic activity was studied, the cells were kept for 1-2 hours at 28° without shaking prior to harvesting. After harvesting, the cell pellet was washed once with 0.9 per cent (w/v) NaCl solution and the cells were resuspended in a small volume of 0.1M Tris-HCl buffer, pH 7.2, containing 1.0 mM EDTA. The cells were disrupted by repeated freezing and thawing and the cell debris were centrifuged off at 500xg for 5 min. The supernatant was re-centrifuged at 30,000xg for 15 min. Both the 30,000xg supernatant and pellet fractions were assayed for PEP carboxykinase activity on the day of preparation.

The assay for PEP carboxykinase activity. The incubation mixture was similar to that described by Shrago, Brech and Templeton (1967), and contained in a volume of 3.0 ml: glycine-NaOH buffer, pH 9.5, 50 mM; ATP, pH 7.5, or sodium pyrophosphate, pH 9.0, 6.7mM; oxaloacetate, pH 8.0, 13.3mM; $MgCl_2$ or $MnCl_2$, 6.7mM and enzyme sample containing 1.5-3.0mg. protein. The reaction was started by the addition of the enzyme and the incubation was carried out at 30° for 10 min. The reaction was terminated by the addition of 1.0 ml. of 20 per cent (w/v) ice-cold TCA. The denatured protein was removed by centrifugation and the supernatant was immediately adjusted to neutrality with NaOH.

Estimation of PEP. The PEP formed was estimated by the method of Meyerhof and Oesper (1949). To 1.0 ml. of the

de-proteinized incubation mixture was added a slight excess of 0.1M iodine solution. Sufficient 2N NaOH was then added to remove the iodine colour. After 2 min. the solution was neutralised with 2N HCl. The treatment with NaOH and HCl was then repeated and the iodine was finally reduced with 0.05M sodium bisulphite solution. The iodoform was centrifuged off. This procedure, involving only a short period of alkalinity, splits PEP-phosphorus quantitatively without decomposing ATP or pyrophosphate (Table V.1). The control was treated in exactly the same manner except that water was substituted for iodine solution, which was added only after the sodium bisulphite.

The orthophosphate released was measured as phosphomolybdic acid extracted into isobutanol by the method of Berenblum and Chain (1938) as described by Ennor and Stocken (1950). PEP formed was equated to Pi liberated in this assay after deduction of Pi values in control tubes (no iodine treatment). The values obtained for these controls represented the Pi released due to the adenosine triphosphatase, pyrophosphatase and possibly other phosphatases present in the crude PEP carbosykinase preparations.

Table V.1 The Estimation of PEP in the Presence of ATP or Pyrophosphate.

PEP (6.0 μ moles) was added to the incubation medium, where shown. PEP carboxykinase and Mg^{2+} were omitted from the incubation mixture. For other details see Text.

Additions	Pi (μ moles /3.0 ml incubation medium)
ATP alone	0.03
PPi alone	Not detected
PEP alone	5.94
PEP + ATP	5.94
PEP + PPi	5.91

The assay for pyrophosphate-glucose phosphotransferase activity.
For the purpose of these studies Tetrahymena were grown as described above, harvested in the logarithmic-phase of growth and washed once with 0.9 per cent (w/v) NaCl solution. They were then suspended in a small volume of 0.1M Tris-HCl buffer, pH 7.2, containing 1.0mM EDTA, and disrupted in a Ribi Cell Fractionator at 10,000lbs./in.² (In some experiments cells were disrupted by repeated freezing and thawing). The cell-lysate was diluted with more buffer to about 10.0 mg. protein/ml. The cell debris were removed by centrifuging at 500xg for 5 min. The supernatant was then re-centrifuged at 30,000xg for 30 min. Both the supernatant

and the particulate fractions were kept at 4° and were assayed on the day of preparation.

The assay system for pyrophosphate-glucose phosphotransferase contained in a total volume of 1.5 ml.: Tris-HCl buffer, pH 7.2, 50mM; glucose, 0.05-0.4M; sodium pyrophosphate, 10.0-40.0mM, and a sample of cell-lysate containing 0.5-1.0 mg. protein. When the activity of the ATP-dependent glucokinase was measured under these conditions, ATP, (20mM) was substituted for pyrophosphate. Other modifications of the incubation medium are described in the legends to Tables. The samples were incubated for 30 min. at 30° and the reaction was stopped by heating at 100° for 3 min. The denatured proteins were removed by centrifugation. When buffers at acidic pH values were used, the pH of the de-proteinised solution was adjusted to pH 7.0.

The glucose-6-phosphate was estimated by the reduction of triphosphopyridine nucleotide (NADP) in the presence of glucose-6-phosphate dehydrogenase and the change in the optical density at 340m μ was measured. The assay medium for glucose-6-phosphate contained: Tris-HCl buffer, pH 7.2, 50mM; $MgCl_2$, 2mM; NADP, 0.4mM; glucose-6-phosphate dehydrogenase, 5 μ g. of protein (1.0 EU), and 1.0 ml. of the de-proteinised incubation mixture. The volume was made to 3.0 ml. with distilled water. The reaction was started by the addition of glucose-6-phosphate dehydrogenase.

Assay for pyrophosphate-ADP-phosphotransferase activity.

a) Preliminary experiments were carried out with cell-free extracts of Tetrahymena, prepared as described above, (the supernatant obtained by centrifuging at 30,000xg for 30 min) being used only. This fraction was dialysed for 2-3 hours against 200 volumes of the suspending buffer. The reaction medium contained in a volume of 1.0 ml.: Tris-HCl buffer, pH 8.0, 50mM; MgCl_2 , 5.0mM; [^{32}P] - labelled pyrophosphate, 2mM (30,000 cpm.); ADP, 2mM, and a sample of the 30,000xg supernatant containing 0.2-0.6 mg protein. The samples were incubated at 30° for 30 min, and the reaction was stopped with 0.1 ml of ice-cold 100 per cent (w/v) TCA. To this was added 1.0 ml of an aqueous charcoal suspension (50 mg./ml.) and the contents of the tube were stirred occasionally over a period of 15 min. The tubes were then centrifuged, the supernatant decanted off, and the charcoal pellet washed three times with 5 ml of cold distilled water. Finally, the charcoal pellet was suspended in 1.0 ml of 1.0 per cent (w/v) serum albumin solution, and samples of the suspension were pipetted onto planchettes, dried and counted in a Nuclear Chicago gas-flow counter fitted with a thin Mylar window.

b) Another assay for pyrophosphate-ADP-phosphotransferase, which also allowed the estimation of ATP formed by the adenylate kinase-catalysed reaction, was used. The reaction mixture, in a

volume of 1.5 ml., contained: Tris-HCl buffer, pH 8.0, 50mM; ADP, 2mM; MgCl_2 , 2mM; sodium pyrophosphate, pH 8.0, 2mM and a sample of enzyme preparation containing 0.1-0.5 mg. protein. Pyrophosphate was added to the reaction mixture as the magnesium complex (MgPPi^{2-} , 2mM). The reaction was blanked by measurement of ATP formed from ADP by the adenylate kinase-catalysed reaction by omitting pyrophosphate from the reaction mixture. The concentration of Mg^{2+} in these blanks was 3mM. The incubation mixture was incubated at 25° for 10 min (except when otherwise stated) and the reaction was terminated by heating at 100° for 3 min. The denatured protein was removed by centrifugation. ATP formed was estimated by the hexokinase-glucose-6-phosphate dehydrogenase coupled system. The reaction mixture contained in a volume of 3.0 ml: appropriate volumes of the de-proteinised reaction mixtures (0.5-1.0 ml); Tris-HCl buffer, pH 7.5, 0.05M; MgCl_2 , 2mM; glucose, 50 mM; NADP, 0.4 mM; hexokinase, 1 μg of protein (0.25 EU), and glucose-6-phosphate dehydrogenase, 1 μg of protein (0.20 EU). The reaction was started by the addition of glucose-6-phosphate dehydrogenase, and the reduction of NADP was measured as the increase in absorption at 340 m μ .

Assay for inorganic pyrophosphatase activity.

Since the purpose of this assay was to estimate the pyrophosphatase activity of the different enzyme fractions under the conditions used for the estimation of the ATP formed from ADP and pyrophosphate

by the pyrophosphate-ADP-phosphotransferase reaction, the composition of the reaction mixture was identical to that described on page 107-108, except that the ADP was omitted. The orthophosphate released was measured by the method of Sumner (1944). When [^{32}P]-labelled pyrophosphate was used the [^{32}P]-orthophosphate released was precipitated as molybdate-triethylamine complex (Sugino and Miyoshi, 1964) and the precipitate was filtered onto oxoid filter membranes, which were washed, dried and counted (La Nauze and Rosenberg, 1967).

Assay for adenosine triphosphatase activity.

The incubation medium contained in a volume of 1.5 ml.: Tris-HCl buffer, pH 8.0, 50mM; MgCl_2 , 3mM, ATP, 2mM and enzyme fraction containing 0.1-0.5 mg. protein. The incubation was carried out at 25° for 10 min. The reaction was stopped by heating at 100° for 3 min. and the orthophosphate released was measured by the method of Sumner (1944).

Preparation of myosin adenosine triphosphatase.

Myosin adenosine triphosphatase from rabbit muscle was prepared as described by Perry (1955), and purified by repeated precipitations. The myosin ATPase was stored in 0.5M KCl at 0° or, after dilution with an equal volume of glycerol, at -10° .

The estimation of the distribution of label in the β and γ phosphorus atoms of ATP. The incubation mixture containing ADP, [^{32}P]-labelled pyrophosphate and Mg^{2+} (see page 107) was

de-proteinised and treated with charcoal as described previously. The washed charcoal pellet was then stirred for 10 min with 10 ml of a cold ammonia-ethanol-water solution (0.3:40:60 v/v). The supernatant was removed by filtration and the charcoal re-extracted as above. The extracts were pooled, evaporated to dryness on a rotary evaporator and then redissolved in 1.0 ml of 0.05M Tris-HCl buffer, pH 8.0. The radioactivity associated with the β and γ phosphorus atoms was then estimated by either of two methods.

a) 0.5 ml of the Tris solution (see above) was incubated for 30 min. at 30° with glucose, 20 mM; MgCl_2 , 2mM; NADP, 2mM, hexokinase, 20 μg . of protein (5.0 EU) and glucose-6-phosphate dehydrogenase, 20 μg . of protein (4.0 EU). The other 0.5 ml of the Tris solution was incubated in exactly the same manner except that hexokinase was omitted. The incubation mixtures were then treated with charcoal. The amount of the label re-adsorbed on charcoal was taken to represent the label associated with the β phosphorus atom of ATP. The label associated with the γ phosphorus atom was then obtained by difference.

b) The label associated with the γ phosphorus atom of ATP was also estimated as [^{32}P]-orthophosphate released (Sugino and Miyoshi, 1964; La Nauze and Rosenberg, 1967) after incubating a sample of the material eluted from charcoal (see above) with myosin ATPase. The label associated with the β phosphorus atom was then obtained by difference or by measuring the radioactivity

re-adsorbed on charcoal after treating the material with the ATPase. The ATPase incubation mixture contained in a volume of 2.0 ml.: glycine-NaOH buffer, pH 9.1, 0.1M; CaCl_2 , 10.0mM; myosin ATPase preparation, 0.8 ml. containing 6.0 mg. protein, and a sample of the charcoal-eluted material. The incubation mixture was incubated for 10 min. at 25° .

Separation of adenine nucleotides.

The three adenine nucleotides, AMP, ADP and ATP were separated by either of two methods.

a) Electrophoresis was carried out by the method of Sato, Thomson and Danforth (1963) on Whatman 3MM paper, in a cooled-plate apparatus, for 40 min under a voltage gradient of 100V/cm. The buffer used contained: citric acid, 0.035M; sodium citrate, 0.0148M and zinc acetate 3.5 mM, pH 3.8. The electrophoregram was dried in a stream of warm air and the nucleotides were detected under a UV lamp. Under these conditions the mobilities (relative to AMP) were: AMP-1.0; ADP-2.4; ATP-2.9; Pi-3.6 and PPi-4.6.

b) Separation by ion-exchange chromatography. The adenine nucleotides were separated on Dowex 1x2 (mesh 200-400), formate. The column dimensions were: 1.7 cm (internal diameter) x 4.0 cm. The nucleotides were eluted from the column with a 4N formic acid-0.6M ammonium formate system, employing a linear elution gradient (cf. Hurlbert, 1957). The flow rate was maintained at

2.5 ml. per min. and 5-ml fractions were collected. The following successive fractions were pooled: 0-35ml, 40-70 ml and 110-160 ml. These contained AMP, ADP and ATP respectively.

All other methods used in this chapter were described in previous chapters.

3. Results and Discussion

A. Phosphoenolpyruvate Formation from Oxaloacetate in the Presence of ATP or Pyrophosphate.

Attempts to demonstrate a pyrophosphate-dependent PEP-carboxykinase activity in Tetrahymena were not successful (Table V.2). When ATP was the phosphoryl group donor the activities of the PEP carboxykinase in the two fractions, the 30,000xg supernatant and the 30,000xg pellet, were similar to those reported by Shrago, Brech and Templeton (1967).

However, when inorganic pyrophosphate was substituted for ATP no PEP formation could be demonstrated. The change in the pH of the incubation medium, or the substitution of the reagent pyrophosphate with granules (Table V.3) did not increase the amount of Pi liberated over the blank pyrophosphatase values. The enzyme preparations used in experiments for which results are shown in Table V.3, were obtained from cells that were hydrolysing granules in the defined medium from which orthophosphate was omitted.

Table V.2 PEP Formation from Oxaloacetate in the Presence of ATP or Pyrophosphate.

For experimental details see Methods.

OAA - oxaloacetate.

Additions	OAA	μmoles PEP formed / mg. protein / hr.	
		30,000xg supernatant	30,000xg pellet
ATP + Mn ²⁺	-	0.02	0.03
ATP + Mn ²⁺	+	1.19	1.43
PPi + Mn ²⁺	-	0	0
PPi + Mg ²⁺	-	0	0
PPi + Mn ²⁺	+	0	0
PPi + Mg ²⁺	+	0	0

Table V.3 The Effect of pH and Granular Pyrophosphate on the Activity of PEP Carboxykinase from Tetrahymena.

Incubation medium contained: buffers (0.05M) as shown, pH 5.8 - acetate buffer, pH 8.0 - Tris-HCl, pH 9.5 -glycine-NaOH; either reagent pyrophosphate (6.7 mM) or granules (5.2 μ moles P_{Pi} containing 0.30 mg. protein). Other components of the incubation medium were as described under the general assay procedures. For the isolation of granules, P_i estimation and other details see Methods.

Additions	μ Moles PEP formed / mg. protein / hr.					
	30,000xg supernatant			30,000xg pellet		
	pH			pH		
	5.8	8.0	9.5	5.8	8.0	9.5
ATP + Mn ²⁺	0.30	0.85	1.20	0	0.93	1.58
PPi + Mg ²⁺	0	0	0	0	0	0
PPi + Mn ²⁺	0	0	0	0	0	0
granules + Mn ²⁺	0	0	0	0	0	0
granules + Mg ²⁺	0	0	0	0	0	0

Since the activity of PEP carboxykinase from Tetrahymena is known to increase when the cells are grown under reduced oxygen tension (Shrago, Brech and Templeton, 1967) attempts were also made to demonstrate a pyrophosphate-dependent activity of this enzyme. As shown (Table V.4), no such activity could be demonstrated, while the ATP-dependent PEP carboxykinase activity was almost twice as high as that obtained from lysates of cells grown under vigorous aeration. (Tables V.2 and V.3).

B. Attempts to Demonstrate a Pyrophosphate-Glucose Phosphotransferase Activity in Tetrahymena.

No pyrophosphate-glucose phosphotransferase activity could be demonstrated in either the 30,000xg supernatant or the particulate fractions (Table V.5). However, some glucose-6-phosphate was formed when ATP was substituted for pyrophosphate, which reflects the presence of an ATP-dependent glucokinase in Tetrahymena. Stoichiometric amounts of glucose-6-phosphate were also formed when ATP and hexokinase were added to the glucose-6-phosphate assay system, which in turn indicated that the assay system was adequate. No pyrophosphate-glucose phosphotransferase activity could be demonstrated in cell-lysates of Tetrahymena when the following were tried (results not shown): change in the pH of the incubation medium (pH 5.5, 7.2, 8.0 and 9.5 were tried); pyrophosphate, Mg^{2+} and glucose concentrations in the incubation medium were varied from 10-40mM, 0-10mM and 0.05-0.4M, respectively;

Table V.4 The Effect of Subjecting Cells to Reduced Oxygen Tension on the Activity of PEP-carboxykinase.

The PEP carboxykinase preparation was obtained from cells that were kept without shaking at 28° for 90 min. prior to harvesting. For other experimental details see Table V.3.

Additions	μmoles PEP formed / mg. protein / hr.					
	30,000xg supernatant			30,000xg pellet		
	pH			pH		
	5.8	8.0	9.5	5.8	8.0	9.5
ATP + Mn ²⁺	-	-	2.10	-	-	1.60
PPi + Mn ²⁺	0	0	0	0	0	0
PPi + Mg ²⁺	0	0	0	0	0	0
granules + Mg ²⁺	0	0	0	0	0	0
granules + Mn ²⁺	0	0	0	0	0	0

Table V.5 Attempts to Demonstrate a Pyrophosphate-Glucose Phosphotransferase Activity in Cell-Extracts of Tetrahymena.

For experimental details see Methods.

Additions	G-6-P formed (μ mole / mg. protein / hr.)	
	30,000xg supernatant	30,000xg pellet
None	Not detected	Not detected
5mM Mg ²⁺	Not detected	Not detected
5mM Mn ²⁺	Not detected	Not detected
ATP substituted for P _P i in incubation medium, 5mM Mg ²⁺	0.39	0.17
0.3 μ mole ATP, 10 μ g. hexokinase into G-6-P assay system	0.26	0.26

synthetic pyrophosphate was substituted with granules; assays were done on cell extracts obtained by freezing and thawing. Although it was reported (Stetten, 1964) that optimal activity for microsomal pyrophosphate-glucose phosphotransferase was obtained in the presence of a detergent, such Triton X-100, no such activity could be demonstrated in different preparations of cell extracts from

Tetrahymena when these were incubated in the presence of this detergent (0.01-0.05 per cent v/v). Addition to reaction mixture of KCl in high concentrations (0.3M), which was shown (Szymona, 1962) to be required for the polyphosphate-dependent glucokinase isolated from M. phlei, was also without effect in cell extracts of Tetrahymena.

C. Attempts to Demonstrate a Pyrophosphate-ADP-Phosphotransferase Activity in Tetrahymena.

When cell-free extracts of Tetrahymena were incubated with ADP and [32 P]-labelled pyrophosphate, and the protein-free reaction product treated with charcoal, 10-25 per cent of the original radioactivity was recovered in the charcoal pellet (Table V.6). Mg^{2+} was essential for this reaction to occur. AMP or GDP could not replace ADP, and the amount of radioactivity in charcoal-adsorbable material was the same whether the reaction was carried out in the presence or in the absence of orthophosphate. When [32 P]-orthophosphate was substituted in the incubation mixture for labelled pyrophosphate, no radioactivity was associated with charcoal. However, when ATP was substituted for ADP in presence of pyrophosphate the amount of radioactivity adsorbed on charcoal was almost twice as great.

Table V.6 The Incorporation of Radioactivity Into Charcoal-adsorbable Material Upon the Incubation of Cell-Free Extracts of Tetrahymena with ADP and [32 P]-labelled Pyrophosphate.

[32 P]-labelled pyrophosphate: 38,000 cpm / 1.0ml incubation medium; [32 P]-labelled orthophosphate: 28,000 cpm / 1.0ml incubation medium. Each tube contained dialysed 30,000xg supernatant, 0.27 mg. protein. For other details see Methods.

Conditions	Radioactivity in Charcoal Adsorbable Material (cpm)
<u>[^{32}P]-pyrophosphate</u>	
Complete system	6540
Complete system (boiled cell-extract)	174
Complete system, + 50mM PO_4^{3-}	6320
ADP omitted	167
Mg^{2+} omitted	274
ADP omitted, + 2mM AMP	171
ADP omitted, + 2mM GDP	213
ADP omitted, + 2mM ATP	11650
ADP omitted, + 2mM ATP 50mM PO_4^{3-}	10575
<u>[^{32}P]-orthophosphate</u>	
Pyrophosphate omitted, + 2mM [32 P]-orthophosphate	212
Pyrophosphate omitted, + 2mM [32 P]-orthophosphate (boiled cell extract)	209

Electrophoregrams of deproteinised reaction mixtures in which either ADP or ATP were incubated with [32 P]-labelled pyrophosphate, showed that radioactivity was present in both the ADP and the ATP bands.

When the material was eluted from charcoal and incubated in the presence of glucose and hexokinase, only about 30-35 per cent of the original charcoal-adsorbable material could now be re-adsorbed on charcoal (Table V.7). Preliminary experiments showed that over 70 per cent of the activity responsible for the synthesis of charcoal-adsorbable material was recovered in the 105,000xg supernatant, and that most of this latter activity (over 75 per cent) could be precipitated with ammonium sulphate (0.50-0.80 saturation). The ammonium sulphate fraction also exhibited a pyrophosphatase activity (0.3 μ mole PPi hydrolysed/10min/mg protein) and an adenylate kinase activity (0.5 μ mole ATP formed/5 min/mg protein. No adenosine triphosphatase activity was detected in this fraction, though it was shown to be present in the 30,000xg supernatant fraction.

When such an enzyme fraction, which was prepared by precipitation with ammonium sulphate, was incubated with either ADP or ATP and [32 P]-pyrophosphate, it was again shown that labelled charcoal-adsorbable material was synthesised (Table V.8). In this Table are also shown the percentages of radioactivity associated with the β and γ phosphorus atoms of ATP. This radioactivity was distributed evenly between these two phosphorus

Table V.7 The Effect of Hexokinase on the Charcoal-Adsorbable Material.

The reaction mixture was of the same composition as described under Methods, except that the specific radioactivity of pyrophosphate was 82,500 cpm/ μ mole (163,000 cpm/1.0 ml reaction mixture). Enzyme preparation: 30,000xg supernatant. The charcoal-adsorbable material was eluted off and incubated with hexokinase and glucose-6-phosphate dehydrogenase as described under Methods.

Conditions	cpm
Radioactivity eluted off charcoal	24,300
Radioactivity re-adsorbed on charcoal after incubation:	
With hexokinase	4,170
Less hexokinase	12,050

atoms when ATP was incubated with [32 P]-pyrophosphate. The duration of the incubation had no effect on this distribution. On the other hand, the duration of incubation had an effect on the distribution of the label between the two phosphorus atoms of the ATP when ADP was incubated with pyrophosphate. The percentage of the total radioactivity was greater in the γ

phosphorus atom when the incubation period was only 5 min, but became more evenly distributed with longer periods of incubation.

To explain the results presented so far the following had to be assumed.

a) An enzyme was present in cell-free extracts of Tetrahymena that was responsible for the synthesis of ATP from ADP and pyrophosphate, thus:

where A-P-P, A-P-P-P, and P-P denote ADP, ATP and PPi respectively. Since 'cold' orthophosphate did not dilute the radioactivity adsorbed on charcoal, and also, since no radioactivity was detected on charcoal when [^{32}P]-orthophosphate was incubated with ADP, it had to be assumed that the phosphoryl group of pyrophosphate was directly transferred to ADP, i.e., orthophosphate was not an intermediate in this reaction. Above reaction could not account for the appearance of label in charcoal-adsorbable material when ATP and [^{32}P]-pyrophosphate were incubated, but this objection would be taken care of by the action of the ATPase present in the system, thus:

An adenylate kinase would tend to randomize the label into both the β and the γ phosphorus atoms of ATP and this distribution of label would be time-dependent. This would also account for the

Table V.8 The Distribution of Label Between the β and γ Phosphorus Atoms of ATP Upon Incubation of ADP and ATP with [^{32}P]-Pyrophosphate.

An ammonium sulphate fraction (0.5-0.8 saturation) of a 30,000xg supernatant was used, containing 0.2mg. protein. The amount of radioactivity associated with the γ phosphorus of ATP was estimated as [^{32}P]-orthophosphate released on incubation with myosin ATPase, (see Methods). The results shown are averages of three separate experiments.

Conditions	Incubation Time (Min)	Charcoal-Adsorbable Material (cpm)		Radioactivity per cent	
		Prior to ATPase Treatment	After ATPase Treatment	β	γ
ADP + PPi	15	1273	209	17	83
"	60	2542	1662	65	35
ATP + PPi	15	1851	867	47	53
"	60	5053	2203	44	56

observation that radioactivity was associated with both the ADP and ATP as shown on electrophoregrams.

This scheme, however, is not consistent with the following observations. Firstly, it does not explain why almost twice as much radioactivity was associated with charcoal when ATP, instead of ADP, was incubated with pyrophosphate. Secondly, it will not

explain the observation that the label was associated with charcoal when ATP and pyrophosphate were incubated with an enzyme preparation that did not exhibit an ATPase activity. Thirdly, it could not explain results obtained from an experiment in which 'sparker' amounts of ADP (0.1mM) were incubated with [³²P]-labelled pyrophosphate, 2mM; MgCl₂, 5mM; AMP, 1.0mM and a 30,000xg supernatant fraction. In the presence of the 'phosphotransferase' enzyme, sufficient ATP should have been formed from the 'sparker' amounts of ADP, and this in turn would have enabled the formation of more ADP through the adenylate kinase-catalysed reaction. It would have been expected, then, that through the agency of these two enzymes sufficient radioactive ADP and ATP would have been formed for the radioactivity to be detected on charcoal.

The results obtained so far, may also be explained by postulating the presence of an ATP-P_i exchange enzyme (cf. Stulberg and Novelli, 1962) in extracts of Tetrahymena, which would catalyse the following exchange reaction:

where E represents the enzyme. In this scheme it is proposed that the [³²P]-pyrophosphate from the medium would exchange with an enzyme-bound intermediate. Such a mechanism might explain why more radioactivity was associated with charcoal when ATP, rather than ADP, was incubated with pyrophosphate; it might also explain the formation of charcoal-adsorbable radioactive material

when an ammonium sulphate fraction, devoid of ATPase activity, was incubated with ATP and pyrophosphate; it would be consistent with the observation that radioactivity was associated with charcoal when ADP was incubated with pyrophosphate, and with the observed even distribution of label between the β and γ phosphorus atoms of ATP. On the other hand, if the exchange mechanism alone was operative, no net synthesis of ATP should take place when ADP is incubated with pyrophosphate, after the allowance is made for the ATP synthesised by the adenylate kinase-catalysed reaction.

The presence in extracts of Tetrahymena of a pyrophosphate-ADP-phosphotransferase and an ATP-pyrophosphate exchange activities together will also explain all these results.

It was shown (Figure V.1) that when ADP was incubated with pyrophosphate there was a net synthesis of ATP. With a 30,000xg supernatant fraction of a cell-free extract 0.70 μ mole of ATP was synthesised in 15 min. in the presence of pyrophosphate (curve C) and only 0.40 μ mole of ATP was synthesised in the absence of PPI, presumably due to an adenylate kinase activity (curve A). This represents an increase of 64 per cent in the amount of ATP formed. It was also shown that some purification was achieved for both the adenylate kinase and the 'phosphotransferase' activities when the 30,000xg supernatant was centrifuged at 100,000xg for 30 min (curves D and E). Only very low activities were associated with

Fig. V.1. The effect of pyrophosphate on the ATP formation from ADP by cell-free extracts of Tetrahymena. The incubation mixture contained, in a volume of 1.5 ml : Tris-HCl buffer, pH 8.0, 50 mM; ADP, 2 mM; $MgCl_2$, 3 mM. A and C : 30,000 x g supernatant, containing 0.33 mg protein; B : as A and C, but the enzyme preparation aged for 2 days at 4°. D and E : 100,000 x g supernatant, containing 0.20 mg protein. Additions : A and D - none; B, C and E - 2 mM $MgPPi^{2-}$.

both the 30,000xg and the 100,000xg pellets, an indication that both activities are due to soluble enzymes.

On ageing (2 days at 4°) there was no change in the activity of adenylate kinase but less ATP was synthesised in the presence of pyrophosphate (curve B). On longer standing though there was a slight decrease in the activity of the adenylate kinase, there was always 15-20 per cent more ATP synthesised in the presence of PPi. Addition of 0.1mM dithiothreitol, or 5mM mercaptoethanol, to the buffer in which the cells were disrupted had a slight protective effect on both the adenylate kinase and the phosphotransferase activities.

It was conclusively demonstrated (Table V.9) that the de-proteinised incubation mixtures had no effect on the assay system used for the estimation of ATP.

It will be recalled that the concentrations of Mg^{2+} added to the reaction mixtures in the presence or the absence of PPi differed. They were 5mM and 3mM, respectively. Since the apparent stability constants for $MgADP^{1-}$ and $MgPPi^{2-}$ complexes are $4 \times 10^3 M^{-1}$ and $5 \times 10^5 M^{-1}$, respectively (O'Sullivan and Perrin, 1964; Josse, 1966), it can be safely assumed that the concentration of Mg^{2+} 'available' to ADP both in the presence and the absence of $MgPPi^{2-}$ should have been the same. It can be calculated, using these stability constants, that the free Mg^{2+} ion concentration,

Table V.9 The Effect of the De-proteinised Reaction Mixtures on the Hexokinase-Glucose-6-Phosphate Dehydrogenase Coupled Assay System.

The ATP was estimated in the de-proteinised incubation mixtures by the hexokinase-glucose-6-phosphate dehydrogenase assay as described under Methods.

Blank= Tris-HCl buffer, pH 8.0, 50mM; ADP, 2mM; MgCl_2 , 3mM.

Test=similar to 'blank' but containing 2mM MgPPi^{2-} .

Additions	$E^{0}_{340\text{m}\mu}$
0.2 μmole ATP	0.425
0.5 ml of 'Blank'	0.230
0.5 ml of 'Blank' + 0.2 μmole ATP	0.660
0.5 ml of 'Test'	0.310
0.5 ml of 'Test' + 0.2 μmole ATP	0.715

$[\text{Mg}^{2+}]_{\text{free}}$ in the absence of MgPPi^{2-} was 1.32mM (Table V.10). If the magnesium ion concentration used in the adenylate kinase assay was not optimal for the activity of this enzyme, a possibility existed that the addition of MgPPi^{2-} may have affected the concentration of $[\text{Mg}^{2+}]_{\text{free}}$ and that the larger amounts of ATP synthesised following this addition were only a reflection of altered $[\text{Mg}^{2+}]_{\text{free}}$ concentration. As shown in Table V.10 this explanation is not correct, since variation of added Mg^{2+} between 2.0 and 4.0mM did not affect the amount of ATP synthesised.

Table V.10 The Effect of Mg^{2+} on the Adenylate Kinase Activity of Tetrahymena.

The incubation mixture contained, in a volume of 1.5 ml; Tris-HCl buffer, pH 8.0, 50mM; ADP, 2mM, and a 100,000xg supernatant fraction containing 0.15 mg protein. ATP was estimated on the de-proteinised samples as described under Methods.

Additions	$[Mg^{2+}]$ TOTAL (mM)	$[Mg^{2+}]$ FREE (mM)	μ mole ATP formed /1.0 ml incubation mixture	
			<u>5 min</u>	<u>10 min</u>
None	0	-	Not detected	Not detected
None	0.25	0.03	0.095	0.185
None	0.50	0.07	0.135	0.240
None	1.00	0.18	0.140	0.250
None	2.00	0.59	0.160	0.285
None	2.50	0.92	0.165	0.290
None	3.00	1.32	0.165	0.290
None	3.50	1.75	0.186	0.274
None	4.00	2.20	0.175	0.281
2mM Mg PPi^{2-}	3.00		0.201	0.331

$$K_{MgADP^{1-}} = \frac{[MgADP^{1-}]}{[Mg^{2+}]_{free} [ADP^{3-}]_{free}} = 4000M^{-1}$$

where:

$K_{MgADP^{1-}}$ = the apparent stability constant of the $MgADP^{1-}$ complex.

$$[Mg^{2+}]_{free} = [Mg^{2+}]_{total} - [MgADP^{1-}]$$

$$[ADP^{3-}]_{free} = [ADP^{3-}]_{total} - [MgADP^{1-}]$$

Numerous attempts were made to demonstrate an increase in the activity of the pyrophosphate-ADP-phosphotransferase in cell-extracts of Tetrahymena which were subjected to reduced oxygen tension. However, no such increase could be demonstrated. Oxygen tension in culture media was regulated by the rate at which the flasks were shaken, or by varying the cell density at a constant rate of shaking.

The results presented so far did not distinguish clearly between the two alternatives proposed earlier. Although most of the evidence presented could be interpreted by the operation of a ATP-PPi exchange, this alone would not result in a net increase in the synthesis of ATP when ADP was incubated with pyrophosphate. Yet, apparent net amount of ATP was synthesised over and above that produced by adenylate kinase, and therefore, the possibility cannot be excluded that a pyrophosphate-ADP-phosphotransferase activity is present in Tetrahymena. The obvious way to demonstrate conclusively the presence of this enzyme must involve its separation from adenylate kinase.

All attempts to separate these two activities by conventional column-chromatographic methods proved unsuccessful. The separation of the two activities was attempted with the following chromatographic media: DEAE-Cellulose, DEAE-Sephadex, CM-Sephadex, Sephadex G-100 and G-200. Crude cell-lysates, 30,000xg and 100,000xg supernatant fractions, as well as enzyme fractions

prepared by precipitation with ammonium sulphate were used in these fractionations. In all cases the adenylate kinase activity was eluted from the columns together with the 'phosphotransferase' activity.

The failure to separate these two activities prompted us to re-investigate whether the equilibrium of the adenylate kinase-catalysed reaction was in some way disturbed in the presence of $\text{Mg} \cdot \text{PPi}^{2-}$ under the assay conditions used. First, the behaviour of a commercial preparation of muscle adenylate kinase was tested under the same assay conditions as those used for the assay of adenylate kinase activity in extracts of Tetrahymena. It was found (Figure V.2) that in the presence of MgPPi^{2-} muscle adenylate kinase behaved in a similar manner as did adenylate kinase from Tetrahymena, i.e., more ATP was synthesised from ADP in the presence of MgPPi^{2-} than in its absence (curves A and B). Addition of sodium pyrophosphate (PPi^{4-}) decreased the amount of net ATP synthesised only slightly (curve C). Furthermore, under the assay conditions used in these experiments, the muscle adenylate kinase activity was not appreciably lowered when higher than optimal concentrations of Mg^{2+} were added to the incubation mixture (Figure V.3).

In order to gain a better understanding of this puzzling result, one question had to be resolved, namely, whether or not the increase in ATP formed was due solely to an effect on the

Fig. V.2. The effect of MgPPI^{2-} on the ATP synthesis from ADP by the muscle adenylate kinase. The incubation mixture contained, in a volume of 1.5 ml : Tris-HCl buffer, pH 8.0, 50 mM; ADP, 2 mM; MgCl_2 , 3 mM, and muscle adenylate kinase, 2 μg protein (specific activity not known). Additions : A - none; B - MgPPI^{2-} , 2 mM; C - MgPPI^{2-} , 2 mM, PPI^{4-} , 0.25 mM. Incubation temperature, 25°. Reaction was stopped by heating at 100° for 3 min., and the ATP formed was measured by the hexokinase-glucose-6-phosphate dehydrogenase coupled enzyme assay as described under Methods.

Fig. V.3. The effect of Mg^{2+} concentration on the production of ATP from ADP by muscle adenylate kinase. The incubation mixture contained, in a volume of 1.5 ml : Tris-HCl buffer, pH 8.0, 50 mM; ADP, 2 mM; muscle adenylate kinase, 2 μ g protein, and $MgCl_2$ as shown. Time of incubation : 10 min. For other details see Fig. V.2.

equilibrium of the myokinase-catalysed reaction. To answer this, the respective concentrations of all components were estimated in the reaction mixture incubated for 15 min in the presence and the absence of added MgPPi^{2-} . This was carried out after careful and complete separation of nucleotides (see 'separation of adenine nucleotides by column chromatography' under Methods) under conditions of nearly complete recovery. These experiments (Tables V.11 and V.12) showed that the difference in the ATP levels found in the presence of MgPPi^{2-} , with muscle adenylate kinase or with enzyme preparations obtained from Tetrahymena, reflected only a shift in the equilibrium of the reaction.

4. Conclusion

The results presented in this chapter described attempts to demonstrate the presence, in cell-free extracts of Tetrahymena, of enzymic activities that would catalyse the formation of glucose-6-phosphate, phosphoenolpyruvate and ATP from pyrophosphate and glucose, oxaloacetate and ADP, respectively.

The failure to demonstrate such enzymic activities does not definitely prove that they are absent from Tetrahymena. This applies in particular to the work with pyrophosphate-ADP-phosphotransferase activity. Although the results presented, strongly suggest that the net amount of ATP synthesised from ADP in the presence of pyrophosphate was due to the shift in the

Table V.11 The Effect of MgPPi^{2-} on the Production of ATP by Muscle Adenylate Kinase.

The incubation mixture contained in a volume of 1.5 ml: Tris-HCl buffer, pH 8.0, 50mM; MgCl_2 , 3mM; ADP, 2mM and muscle adenylate kinase, 4 μg of protein. The incubation was carried out at 25° for 15 min. The reaction was stopped by heating at 100° for 3 min. 1.0 ml samples were assayed for AMP, ADP and ATP by the column-chromatographic separation on Dowex 1x2 (formate) as described under Methods.

Results shown are averages of two separate experiments.

Additions	Adenine nucleotides per ml reaction mixture (μmoles)			Increase (+) or Decrease (-) (μmoles)	
	AMP	ATP	ADP	(ATP+AMP)	ADP
None	0.250	0.276	1.395	+0.526	-0.605
MgPPi^{2-} (2mM)	0.456	0.482	1.076	+0.938	-0.924

Table V.12 The Effect of MgPPi^{2-} on the Production of ATP by Adenylate Kinase from Tetrahymena.

The incubation mixture was identical with that described in Table V.11 except that a 30,000xg supernatant, containing 0.3 mg. protein was used. Incubation time was 30 min. For other details see Table V.11.

Results shown are the averages of two separate experiments.

Additions	Adenine nucleotides per ml reaction mixture (μmoles)			Increase (+) or Decrease (-) (μmoles)	
	AMP	ATP	ADP	(AMP+ATP)	ADP
None	0.583	0.568	0.767	+1.151	-1.170
MgPPi^{2-} (2mM)	0.671	0.655	0.599	+1.326	-1.338

equilibrium of a reaction presumably catalysed by adenylate kinase present in these extracts (Table V.12), the possibility that such a 'phosphotransferase' activity is, indeed, present in Tetrahymena, cannot be absolutely ruled out. The results obtained for the distribution of the label in the β and γ phosphorus atoms of ATP when ADP was incubated with pyrophosphate for short periods of time (Table V.8), cannot be explained by the assumption mentioned above. Nor are these results consistent with any ATP-PPi exchange, as already mentioned.

However, insufficient evidence is available at present to decide with any degree of certainty for or against the presence of such a 'phosphotransferase' activity in Tetrahymena. The ultimate resolution of this problem will depend on the successful separation of this activity from that of the adenylate kinase.

It was stated previously, that the deposition and the dissolution of pyrophosphate granules in Tetrahymena are likely to be enzymically controlled processes. The evidence for this were the observations that, firstly, both Ca^{2+} and Mg^{2+} were required for the formation of granules in vivo, and that the ratio of these two metal ions in isolated granules remained constant even when their concentrations in the medium were varied within a wide range. Secondly, it was observed that granules were rapidly dissolved when the cells were deprived of phosphate. It can be safely argued, then, that these granules are concerned with phosphate-storage. A similar function is attributed also to the polyphosphate deposits that are found in a large number of various microorganisms (cf. Chapter I). Furthermore, polyphosphates, in addition to providing phosphate-storage, act also in various species as direct phosphate donors for the phosphorylation of a number of important biological compounds, such as AMP, ADP and glucose. Therefore, in analogy with polyphosphates the possibility still exists that there is a similar role for pyrophosphate in

Tetrahymena, although the present experiments were unsuccessful in demonstrating this.

It was recently reported (Hatch and Slack, 1968) that an enzyme (phosphopyruvate synthetase), isolated from certain tropical grasses, catalysed the formation of PEP, AMP and PPi from pyruvate, ATP and Pi. A similar enzyme has recently been demonstrated in Propionibacterium shermanii (Evans and Wood, 1968).^{*} Reports about this interesting reaction appeared after the completion of the experimental work described in this Thesis, and its possible existence in Tetrahymena has not yet been investigated. A demonstration of such a reaction in this organism would also be of interest from the evolutionary point of view, since there are some doubts whether Tetrahymena belongs to the animal or the plant kingdom. (Holz, 1967)

* Evans, H.J. and Wood H.G. (1968). Fedn. Proc. Fedn. Am. Socs. exp. Biol., 27, 588. This reference came to hand too late to be included in the Reference List.

REFERENCES

- ARION, W.J. and NORDLIE, R.C. (1964). J. Biol. Chem., 239, 2752.
- ARION, W.J. and NORDLIE, R.C. (1967). J. Biol. Chem., 242, 2207.
- BALTSCHOFFSKY, H. and von STEDINGH, L.V. (1966). Biochem. Biophys. Res. Commun., 22, 722.
- BALTSCHOFFSKY, H., von STEDINGH, L.V., HELDT, H.W. and KLINGENBERG, M. (1966). Science, 153, 1120.
- BALTSCHOFFSKY, M. (1967). Biochem. Biophys. Res. Commun., 28, 270.
- BERENBLUM, I. and CHAIN, E. (1938). Biochem. J., 32, 295.
- BLOCH-FRANKENTHAL, L. (1954). Biochem. J., 57, 87.
- BOYER, P.D. (1958). Proc. Intern. Symp. Enzyme Chem., Tokyo and Kyoto 1957, (Tokyo), p.189.
- COHN, M. (1958). J. Biol. Chem., 230, 369.
- COHN, M., MONOD, J., POLLOCK, M.R. SPIEGELMAN, S. and STANIER, R.Y. (1953). Nature, London, 172, 1096.
- COLE, J.A. and HUGHES, D.E. (1965). J. gen. Microbiol., 38, 65.
- CONNER, R.L. and CLINE, S.G. (1964). J. Protozool., 11, 486.
- de DUVE, C. (1963). Ciba Found. Symp., Lysosomes, p.1, London: Churchill.
- DIRHEIMER, G. and EBEL, J.P. (1962). Compt. Rend. Acad. Sci. (Paris), 254, 2850.
- DIRHEIMER, G. and EBEL, J.P. (1964). Compt. Rend. Soc. Biol. (Strasbourg), 158, 1948.
- DIRHEIMER, G. and EBEL, J.P. (1965). Compt. Rend. Acad. Sci. (Paris), 260, 3787.

- DIRHEIMER, G., MULLER-FELTER, S., WEIL, J.H., YACOUN, M. and EBEL, J.P. (1963). Bull. Soc. Chim. Biol. Fr., 45, 875.
- DREWS, G. (1960). Arch. Mikrobiol., 35, 53.
- DREWS, G. (1962). Colloq. Intern. Centre Natl. Rech. Sci. (Paris). 106, 533.
- DIXON, M. and WEBB, C. (1966). Enzymes, 2nd ed., p.40. London: Longmans, Green and Co. Ltd.
- EBEL, J.P. (1952). Bull. Soc. Chim. Biol. Fr., 34, 321 and 498.
- EBEL, J.P. (1953). Bull. Soc. Chim. Fr., 20, 991, 998.
- EBEL, J.P., COLAS, J. and MULLER, S. (1958). Exptl. Cell. Res., 15, 28.
- EBEL, J.P., DIRHEIMER, G., STAHL, A.J.C. and YACOUN, M. (1963). Bull. Soc. Chim. Biol. Fr., 45, 865.
- EBEL, J.P. and MULLER, S. (1958). Exptl. Cell. Res., 15, 21.
- ENNOR, A.H. and STOCKEN, L.A. (1950). Aust. J. exp. Biol. med. Sci., 28, 647.
- FISHER, C.J. and STETTEN, M.R. (1966). Biochim. Biophys. Acta, 121, 102.
- FUCHS, R.J. and CZECH, F.W. (1963). Anal. Chem., 35, 769.
- GILMOUR, D. and CALABY, J.H. (1953). Enzymologia, 16, 34.
- GORDON, J.J. (1950). Biochem. J., 46, 96.
- GREENAWALT, J.W., ROSSI, C.S. and LEHNINGER, A.L. (1964). J. Cell. Biol., 23, 21.
- GRIFFIN, J.B., DAVIDIAN, N.M. and PENNIAL, R. (1965). J. Biol. Chem., 240, 4427.
- GRIFFIN, J.B. and PENNIAL, R. (1966). Arch. Biochem. Biophys., 114, 67.
- GRIMME, A. (1902). Zentbl. Bakt. Parasitkde. I. Abt. Orig., 32, 172.

- GROSSMANN, D. and LANG, K. (1962). Biochem. Z., 336, 351.
- HAROLD, F.M. (1960). Biochim. Biophys. Acta, 45, 172.
- HAROLD, F.M. (1962, a). Biochim. Biophys. Acta, 57, 59.
- HAROLD, F.M. (1962, b). J. Bacteriol., 83, 1047.
- HAROLD, F.M. (1963). J. Bacteriol., 86, 216.
- HAROLD, F.M. (1964). J. gen. Microbiol., 35, 81.
- HAROLD, F.M. (1966). Bacteriol. Rev., 30, 772.
- HAROLD, F.M. and HAROLD R.L. (1965). J. Bacteriol., 89, 1263.
- HAROLD, F.M. and MILLER, A. (1961). Biochim. Biophys. Acta, 50, 261.
- HAROLD, F.M. and SYLVAN, S. (1963). J. Bacteriol., 86, 222.
- HAROLD, R.L. and HAROLD, F.M. (1963). J. gen. Microbiol., 31, 241.
- HASE, E., MIYACHI, S. and MIHARA, S. (1963). Studies on Microalgae & Photosynthetic Bacteria. Special Issue of Plant and Cell Physiol., (Tokyo), 4, 619.
- HATCH, M.D. and SLACK, C.R. (1968). Biochem. J., 106, 141.
- HELLER, J., KAPRIAK, St. and ZUBIKOWA, I. (1950). Nature, London, 166, 187.
- HEPPELL, L.A., HARKNESS, D.R. and HILMOE, R.J. (1962). J. Biol. Chem., 237, 841.
- HOFFMANN-OSTENHOF, O. (1962). Colloq. Intern. Centre Natl. Rech. Sci. (Paris), 106, 641.
- HOFFMANN-OSTENHOF, O., KENNEDY, J., KECK, K., GABRIEL, O. and SCHONFELLINGER, H.W. (1954). Biochim. Biophys. Acta, 14, 285.
- HOFFMANN-OSTENHOF, O. and SLECHTA, L. (1958). Proc. Intern. Symp. Enzyme Chem., Tokyo and Kyoto 1957, (Tokyo), p.180.
- HOFFMANN-OSTENHOF, O. and WEIGERT, W. (1952). Naturwissenschaften, 39, 303.

- HOLZ, G.G. (1966). J. Protozool., 13, 2.
- HORIUCHI, T., HORIUCHI, S. and MIZUNO, D. (1959). Nature, London, 183, 1529.
- HUENNEKENS, F.M. and WHITELEY, H.R. (1960). In Comparative Biochemistry, vol. 1, 107. Ed. by Florkin, M. and Mason, J.S., New York: Academic Press Inc.
- HUGHES, D.E. and MUHAMMED, A. (1962). Colloq. Intern. Centre Natl. Rech. Sci. (Paris), 106, 591.
- HULBERT, R.B. (1957). In Methods in Enzymology, vol. 3, p.793. Ed. Colowick S.P. and Kaplan, N.O. New York: Academic Press Inc.
- IMSANDE, J. and HANDLER, P. (1961). In The Enzymes, vol. 5, part B, p.281. Ed. Boyer, P.D., Lardy, H. and Myrbäck, K. New York: Academic Press Inc.
- JANAKIDEVI, K., DEWEY, V.C. and KIDDER, G.W. (1965). J. Biol. Chem., 240, 1754.
- JEENER, R. and BRACHET, J. (1944). Enzymologia, 11, 222.
- JONES, A.R. (1966). J. Protozool., 13, 422.
- JONES, A.R. (1967). J. Protozool., 14, 220.
- JOSSE, J. (1966). J. Biol. Chem., 241, 1938, 1948.
- KALTWASSER, H. (1962). Arch. Mikrobiol., 41, 282.
- KALTWASSER, H. and SCHLEGEL, H.G. (1959). Arch. Mikrobiol., 34, 76.
- KATCHMAN, B.J. and FETTY, W.O. (1955). J. Bacteriol., 69, 607.
- KATCHMAN, B.J., FETTY, W.O. and BUSCH, K.A. (1959). J. Bacteriol., 77, 331.
- KIDDER, G.W. and DEWEY, V.C. (1951). In Biochemistry and Physiology of Protozoa, vol. 1, p.323. Ed. by Lwoff, A. New York: Academic Press Inc.

- KLINGENBERG, M. (1961). In Biological Structure and Function, vol. 2, p.227. Ed. by Goodwin T.W. and Lindberg. O. New York: Academic Press Inc.
- KORNBERG, A. (1957, a). Adv. Enzymol., 18, 191.
- KORNBERG, A. (1962). In Horizons in Biochemistry, p.251. Ed. by Kasha, M. and Pullman, B. New York: Academic Press Inc.
- KORNBERG, A., KORNBERG, S.R. and SIMMS, E.S. (1956). Biochim. Biophys. Acta, 20, 215.
- KORNBERG, A. and PRICER, W.E. (1951). J. Biol. Chem., 191, 535.
- KORNBERG, S.R. (1957, b). Biochim. Biophys. Acta, 26, 294.
- KUHL, A. (1960). Ergeb. Biol., 23, 144.
- KULAEV, I.S. and BELOZERSKII, A.N. (1958). Proc. Acad. Sci. USSR (English Transl.), 120, 128.
- KUNITZ, M. and ROBBINS, P.W. (1961). In The Enzymes, vol. 5, part B, p.169. Ed. by Boyer, P.D., Lardy, H. and Myrback, K. New York: Academic Press Inc.
- KURAHASHI, K., PENNINGTON, R.J. and UTTER, M.F. (1957). J. Biol. chem., 226, 1059.
- LANGEN, P. and LISS, E. (1958). Biochem. Z., 330, 455.
- LANGEN, P., LISS, E. and LOHMANN, K. (1962). Colloq. Intern. Centre Natl. Rech. Sci. (Paris), 106, 603.
- LA NAUZE, J.M. and ROSENBERG, H. (1967). Biochim. Biophys. Acta. 148, 811.
- LAYBOURN, R.L. (1924). J. Am. Med. Assoc., 82, 121.
- LAYNE, E. (1957). In Methods in Enzymology, vol. 3, p.450. Ed. by Colowick, S.P. and Kaplan, N.O. New York: Academic Press Inc.
- LEHNINGER, A.L., CARAFOLI, E. and ROSSI, C.S. (1967). Adv. Enzymol., 29, 259.
- LIEBERMANN, L. (1888). Ber. deutsch. chem. Ges., 21, 598.

- LINDENBAUM, S., PETERS, T.V. and RIEMAN, W. (1954). Anal. Chim. Acta, 11, 530.
- LISS, E. and LANGEN, P. (1962). Arch. Mikrobiol., 41, 383.
- LOWRY, O.H., ROSENBROUGH, N.J., FARR, A.L. and RANDALL, R.J. (1951). J. Biol. Chem., 193, 265.
- McELROY, W.D., COULOMBRE, J. and HAYS, R. (1951). Arch. Biochem. Biophys., 32, 207.
- MacFARLANE, M.G. (1936). Biochem. J., 30, 1369.
- MATTENHEIMER, H. (1956). Z. Physiol. Chem., 303, 107, 115, 125.
- MARTINEZ, R.J. (1963). Arch. Mikrobiol., 44, 334.
- MEYER, A. (1904). Bot. Z., 62, 113.
- MEYERHOF, O. and OESPER, P. (1949). J. Biol. Chem., 179, 1371.
- MEYERHOF, O., SKATAS, R. and KAPLAN, A. (1953). Biochim. Biophys. Acta, 12, 121.
- MILLER, S.L. and PARRIS, M. (1964). Nature, London, 204, 1248.
- MIYACHI, S. (1961). J. Biochem. (Tokyo), 50, 367.
- MIYACHI, S., KANAI, R., MIHARA, S., MIYACHI, S. and AOKI, S. (1964). Biochim. Biophys. Acta, 93, 625.
- MORRISON, J.F. and ENNOR, A.H. (1960). In The Enzymes, vol. 2, part A, p.89. Ed. by Boyer, P.D., Lardy, H. and Myrback, K. New York: Academic Press Inc.
- MUDD, S., YOSHIDA, A. and Koike, M. (1958). J. Bacteriol., 75, 224.
- MUHAMMED, A. (1961). Biochim. Biophys. Acta, 54, 121.
- MUHAMMED, A., RODGERS, A. and HUGHES, D.E. (1959). J. gen. Microbiol., 20, 482.
- NIEMIERKO, W. (1962). Colloq. Intern. Centre Natl. Rech. Sci. (Paris), 106, 615.
- NISHI, A. (1960). J. Biochem., (Tokyo), 48, 758.

- NORDLIE, R.C. and ARION, W.J. (1964). J. Biol. Chem., 239, 1680.
- NORDLIE, R.C., ARION, W.J. and GLENDE, E.A. (1965). J. Biol. Chem., 240, 3479.
- O'SULLIVAN, W.J. and PERRIN, D.D. (1964). Biochemistry, 3, 18.
- PAUTARD, F.G.E. (1959). Biochim. Biophys. Acta, 35, 33.
- PENNIAL, R. and GRIFFIN, J.B. (1964). Biochim. Biophys. Acta, 90, 429.
- PETERS, T.V. and RIEMAN, W. (1956). Anal. Chim. Acta, 14, 131.
- PLESNER, P., RASMUSSEN, L. and ZEUTHEN, E. (1964). In Synchrony in Cell Division and Growth, p.543. Ed. by Zeuthen, E., New York: Interscience Publishers.
- PERRY, S.V. (1955). In Methods in Enzymology, vol. 2, p.582. Ed. Colowick, S.P. and Kaplan, N.O. New York: Academic Press, Inc.
- RAFTER, G.W. (1960). J. Biol. Chem., 235, 2475.
- RINGBOM, A. (1963). In Complexation in Analytical Chemistry. New York: Interscience Publishers.
- ROBBINS, P.W. and LIPMANN, F. (1958). J. Biol. Chem., 233, 686.
- ROSENBERG, H. (1959). J. Chromatog., 2, 487.
- ROSENBERG, H. (1966). Exptl. Cell. Res., 41, 397.
- SATO, T.R., THOMSON, J.F. and DANFORTH, W.F. (1963). Anal. Biochem., 5, 542.
- SCHITO, G.C. and PESCE, A. (1965). Giorn. Microbiol., 13, 145.
- SCHMIDT, G. (1951). In Phosphorus Metabolism, vol. 1, p.442, Ed. by McElroy, W.D. and Glass, B. Baltimore: John Hopkins Press.
- SCHMIDT, G., HECHT, L. and THANNHAUSER, S.J. (1946). J. Biol. Chem., 166, 775.
- SCHMIDT, G. and THANNHAUSER, S.J. (1945). J. Biol. Chem., 161, 83.

- SCHNEIDER, W.C. (1945). J. Biol. Chem., 161, 293.
- SHRAGO, E., BRECH, W. and TEMPLETON, K. (1967). J. Biol. Chem., 242, 4060.
- SHRAGO, E. and SHUG, A.L. (1966). Biochim. Biophys. Acta, 122, 376.
- SIU, P.M.L. and WOOD, H.G. (1962). J. Biol. Chem., 237, 3044.
- SMITH, I.W., WILKINSON, J.F. and DUGUID, J.P. (1954). J. Bacteriol., 68, 450.
- STADTMAN, E.R. (1966). Adv. Enzymol., 28, 41.
- STAHL, A.J.C. and EBEL, J.P. (1963). Bull. Soc. Chim. Biol. Fr., 45, 887.
- STETTEN, M.R. (1964). J. Biol. Chem., 239, 3576.
- STETTEN, M.R. (1965). J. Biol. Chem., 240, 2248.
- STETTEN, M.R. and TAFT, H.L. (1964). J. Biol. Chem., 239, 4041.
- STOCKSTAD, E.L.R., SEAMAN, G.R., DAVIS, R.J. and HUTNER, S.H. (1956). Methods Biochem. Anal., 3, 23.
- STULBERG, M.P. and NOVELLI, D.G. (1962). In The Enzymes, vol. 6, p.401. Ed. Boyer, P.D., Lardy, H. and Myrbäck, K. New York: Academic Press, Inc.
- SUGINO, Y. and MIYOSHI, Y. (1964). J. Biol. Chem., 239, 2360.
- SUMNER, J.B. (1944). Science, 100, 413.
- SZYMONA, M. (1962). Acta Biochim. Polon., 9, 165.
- SZYMONA, M. and OSTROWSKY, W. (1964). Biochim. Biophys. Acta, 85, 283.
- SZYMONA, M., URYSON, S.O. and KULAEV, I.S. (1967). Biokhimiya (Russian), 32, 495.
- SZYMONA, O. and SZUMILO, T. (1966). Acta Biochim. Polon., 13, 129.
- TEWARI, K.K. and SINGH, M. (1964). Phytochemistry, 3, 341.

- THILO, E. (1962). Adv. Inorg. Chem. Radiochem., 4, 1.
- TORRIANI, A. (1960). Biochim. Biophys. Acta, 38, 460.
- UTTER, M.F. and KURAHASHI, K. (1954). J. Biol. Chem., 207, 787.
- VAGABOV, V.M. and KULAEV, I.S. (1964). Dokl. Akad. Nauk, SSSR, 158, 218.
- VOELZ, H., VOELZ, U. and ORTIGOZA, R.O. (1966). Arch. Mikrobiol. 53, 371.
- WARNOCK, L.G. and van EYS, J. (1962). J. Cell. Comp. Physiol., 60, 53.
- WAZER, van, J.R. (1958). In Phosphorus and Its Compounds, vol. 1. New York: Interscience Publishers.
- WAZER van, J.R. and CAMPANELLA, D.A., (1950). J. Am. Chem. Soc. 72, 655.
- WEBER, G., SINGHAL, R.L., STAMM, N.B., LEA, M.A. and FISHER, E.A. (1966). Adv. Enzyme Regulation, 4, 59.
- WIAME, J.M. (1946). Bull. Soc. Chim. Biol. Fr., 28, 552.
- WIAME, J.M. (1947 a). J. Am. Chem. Soc., 69, 3146.
- WIAME, J.M. (1947 b). Biochim. Biophys. Acta, 1, 234.
- WIAME, J.M. (1948). J. Biol. Chem., 178, 919.
- WIDRA, A. (1959). J. Bacteriol., 78, 664.
- WILKINSON, J.F. and DUGUID, J.P. (1960). Intern. Rev. Cytol., 9, 1.
- WINDER, F.G. and DENNENY, J.M. (1955). Nature, London, 175, 636.
- WINDER, F.G., DENNENY, J.M. (1957). J. gen. Microbiol., 17, 573.
- YOSHIDA, A. and YAMATAKA, A. (1953). J. Biochem. (Tokyo), 40, 85.

YOSHIDA, A. (1955). J. Biochem. (Tokyo)., 42, 163.

ZAITSOVA, G.N. BELOZERSKII, A.N. and NOVOZHILOVA, L.P. (1960 a).
Biochemistry (USSR) (English Transl.), 25, 147.

ZAITSOVA, G.N., BELOZERSKII, A.N. and NOVOZHILOVA, L.P. (1960 b).
Mikrobiologiya (English Transl.), 29, 254,

ZAITSOVA, G.N. and FROLOVA, L.Y. (1951). Biochemistry (USSR)
(English Transl.), 16, 174.