This thesis represents research undertaken in the Photochemistry group, Research School of Biological Sciences, The Australian National University, Canberra. This work was completed between March 2005 and March 2009.

***Designing Photosystem II:
Molecular Engineering of Photo-Catalytic
Proteins***

Brendon Francis Conlan

*A thesis submitted for the degree of Doctor of Philosophy of
The Australian National University*

March 2009

Publications

This thesis represents research undertaken in the Photobioenergetics group, Research School of Biological Sciences, The Australian National University, Canberra. This work was completed between March 2005 and March 2009

Except where otherwise acknowledged, the work presented in this thesis is my own, and was performed under the supervision of Professor Tom Wydrzynski.

Brendon Francis Conlan,

This thesis represents research undertaken in the Photobiology Group, Research School of Biological Sciences, The Australian National University, Canberra. This work was completed between March 2002 and March 2004.

Except where otherwise acknowledged, the work presented in this thesis is my own, and was performed under the supervision of Professor Tony Whitham.

Professor Francis Collins

Publications

Conlan B, Cox N, Su J, Hillier W, Messinger J, Lubitz W, Dutton PL, Wydrzynski T. *Photo-catalytic oxidation of a di-manganese centre in an engineered bacterioferritin 'reaction centre'*. (Manuscript accepted to BBA Bioenergetics Mar 2009).

Conlan B. *Designing Photosystem II: Molecular Engineering of Photo-catalytic Proteins*. Photosynthesis Research 2008 Oct 98: 687-700.

Wydrzynski T, Hillier W, Conlan B. *Engineering model proteins for Photosystem II function*. Photosynthesis Research 2007 Nov-Dec; 94 (2-3): 225-33

Conlan B, Wydrzynski T, Hillier W. *Designing artificial photosynthesis: Engineering bacterioferritin for light-activated electron transfer*. Chapter 3: 1261-1265 *Photosynthesis. Energy from the Sun*. 14th International Congress on Photosynthesis.

Publications

Conte B, Cor N, Su J, Hiller W, Messinger J, Lubitz W, Dutton PL, Wydrzyński T
Photo-catalytic oxidation of a di-manganese centre in an engineered catalytic
reaction centre. (Manuscript accepted to *BBA Bioenergetics* Mar 2009)

Conte B, Designing Photosystem II: Molecular Engineering of Protein-Catalytic
Protein. *Photosynthesis Research* 2008 Oct 9; 97: 215-230

Wydrzyński T, Hiller W, Conte B. Engineering model proteins for Photosystem II
function. *Photosynthesis Research* 2007 Nov-Dec; 94 (3-4): 215-23

Conte B, Wydrzyński T, Hiller W. Designing artificial photosynthetic Engineering
catalytic centre for light-driven electron transfer. *Chemical Science* 2007; 1261-1265
Photosynthesis: Energy from the Sun. 14th International Congress on Photosynthesis

Oral Presentations

Conlan B. *Designing Photosynthesis: Molecular engineering of photo-catalytic proteins*. Presented at Bio21 Melbourne University, Melbourne, Australia 2009.

Conlan B. *Designing Photosynthesis: Molecular engineering of photo-catalytic proteins*. Presented at the Research School of Biological Science, Australian National University, Canberra, Australia 2009.

Conlan B. *Designing Photosynthesis: Molecular engineering of photo-catalytic proteins*. Presented at the 17th International Conference on Photochemical Conversion and Storage of Solar Energy 2008, Sydney, Australia.

Conlan B. *Designing Photosynthesis: Molecular engineering of photo-catalytic proteins*. Presented at the 1st Australian Bioconjugate Chemistry Workshop 2008, University of New South Wales, Sydney, Australia.

Conlan B. *Designing artificial photosynthesis: Engineering bacterioferritin for light-activated electron transfer*. Presented at Arizona State University, Department of Chemistry and Biochemistry, Tempe, USA 2007.

Conlan B. *Designing artificial photosynthesis: Engineering bacterioferritin for light-activated electron transfer*. Presented at the International Congress on Photosynthesis 2007, Glasgow, UK.

Conlan B. *Engineering a light activated metal binding protein*. Presented at The Max Planck Institute for Bioinorganic Chemistry. Mülheim, Germany 2006.

Conlan B. *Engineering a light activated metal binding protein*. Presented at the Department of Biochemistry and Biophysics, Gothenburg University, Gothenburg, Sweden 2006.

Oral Presentations

Colman B. Designing Photosynthetic Molecular Engineering of photosynthesis
presented at the 11th International Conference on Photochemical Conversion
and Storage of Solar Energy 2008, Sydney, Australia

Colman B. Designing Photosynthetic Molecular Engineering of photosynthesis
presented at the 11th International Conference on Photochemical Conversion
and Storage of Solar Energy 2008, Sydney, Australia

Colman B. Designing Photosynthetic Molecular Engineering of photosynthesis
presented at the 11th International Conference on Photochemical Conversion
and Storage of Solar Energy 2008, Sydney, Australia

Colman B. Designing Photosynthetic Molecular Engineering of photosynthesis
presented at the 11th International Conference on Photochemical Conversion
and Storage of Solar Energy 2008, Sydney, Australia

Colman B. Designing Photosynthetic Molecular Engineering of photosynthesis
presented at the 11th International Conference on Photochemical Conversion
and Storage of Solar Energy 2008, Sydney, Australia

Colman B. Designing Photosynthetic Molecular Engineering of photosynthesis
presented at the 11th International Conference on Photochemical Conversion
and Storage of Solar Energy 2008, Sydney, Australia

Colman B. Designing Photosynthetic Molecular Engineering of photosynthesis
presented at the 11th International Conference on Photochemical Conversion
and Storage of Solar Energy 2008, Sydney, Australia

Colman B. Designing Photosynthetic Molecular Engineering of photosynthesis
presented at the 11th International Conference on Photochemical Conversion
and Storage of Solar Energy 2008, Sydney, Australia

Table of Contents

Minds are like parachutes. They only function when they are open.

1.01. INTRODUCTION	v
1.02. ACKNOWLEDGMENTS	vii
1.03. ABSTRACT	xi
1.04. LIST OF TABLES	xix
1.05. LIST OF FIGURES	xxi
1.06. LIST OF ABBREVIATIONS	xxv
1.07. INTRODUCTION	1
1.1. PHOTOSYSTEM II	2
1.1.1. The Reaction Centre	2
1.1.2. The Oxygen Evolving Complex: Model and Mechanism	3
1.1.3. 80% Monophase	5
1.2. DESIGN OF PHOTO-CATALYTIC PROTEINS	7
1.2.1. Electron Transfer Principles	8
1.2.2. Light Absorbing Pigment	12
1.2.3. Electron Acceptors	16
1.2.4. Electron Donors	18
1.3. COMBINING DIFFERENT TYPES OF PIGMENT ACTIVATION	19
1.4. RESEARCH OBJECTIVES	20
2. METHODS	23
2.1. CLONING AND MUTAGENESIS OF BACTERIOPHOTOTROPHIC MOLECULES	25
2.1.1. Site-Directed Mutagenesis of Bacteriochlorophyll	26
2.2. PERFECTION OF THE BPH HETERODIMERS CONSTRUCT	27
2.3. EXPRESSION AND PURIFICATION OF ALL CONSTRUCTS	32
2.3.1. Expression	32
2.3.2. Protein Purification	34

Minds are like parachutes. They only function when they are open.

Dr. James Dewey
(1857-1931)

Table of Contents

DESIGNING PHOTOSYSTEM II:.....	I
PUBLICATIONS	V
ORAL PRESENTATIONS.....	VII
TABLE OF CONTENTS.....	XI
ACKNOWLEDGEMENTS.....	XV
ABSTRACT.....	XVII
LIST OF TABLES	XIX
LIST OF FIGURES	XXI
LIST OF ABBREVIATIONS.....	XXV
1 INTRODUCTION	1
1.1. PHOTOSYSTEM II	2
1.1.1. <i>The Reaction Centre</i>	2
1.1.2. <i>The Oxygen Evolving Complex: Models and Mechanisms</i>	3
1.1.3. <i>Why Manganese?</i>	5
1.2. DESIGN OF PHOTO-CATALYTIC PROTEINS	7
1.2.1. <i>Electron Transfer Principles</i>	8
1.2.2. <i>Light Activated Pigments</i>	13
1.2.3. <i>Electron Acceptors</i>	16
1.2.4. <i>Electron Donors</i>	18
1.3. COMBINING DONOR-TYROSINE-PIGMENT-ACCEPTOR	19
1.4. RESEARCH OBJECTIVES	20
2 METHODS.....	23
2.1. CLONING AND MUTAGENESIS OF BACTERIOFERRITIN HOMODIMER	23
2.1.1. <i>Site-Directed Mutagenesis of Bacterioferritin</i>	26
2.2. PRODUCTION OF THE BFR HETERODIMER CONSTRUCT	27
2.3. EXPRESSION AND PURIFICATION OF ALL CONSTRUCTS	33
2.3.1. <i>Expression</i>	33
2.3.2. <i>Protein Purification</i>	34

2.4.	COFACTOR BINDING	35
2.4.1.	CD Spectroscopy.....	35
2.4.2.	Isothermal Titration Calorimetry.....	35
2.4.3.	Covalent Attachment of Quinone's	36
2.5.	ANALYSIS.....	37
2.5.1.	Oxygen Measurements.....	37
2.5.2.	Fluorescence Measurements.....	37
2.5.3.	Redox Measurements	37
2.5.4.	EPR Measurements.....	38
2.5.5.	EPR Simulations	38
2.6.	MOLECULAR MODELLING	39
3	BACTERIOFERRITIN AS A 'REACTION CENTRE'	41
3.1.	INTRODUCTION.....	41
3.1.1.	Summary	45
3.1.2.	Techniques Used to Analyse the BFR Complex.	45
3.1.3.	Electron Paramagnetic Resonance.....	45
3.1.4.	Circular Dichroism Spectroscopy.....	47
3.1.5.	Isothermal Titration Calorimetry.....	48
3.1.6.	Cyclic Voltammetry.....	50
3.2.	RESULTS.....	51
3.2.1.	Cofactor Assembly in Bacterioferritin.	51
3.2.2.	EPR Spectroscopy.....	56
3.2.3.	Simulated Mn ^{II} Dimer Line-Shapes.....	56
3.2.4.	Theoretical Approach: The Mn ^{II} Mn ^{II} System.....	62
3.2.5.	Light Induced Oxidation State Changes	65
3.3.	DISCUSSION.....	70
3.4.	CONCLUSION	78
4	BACTERIOFERRITIN HETERODIMER	79
4.1.	INTRODUCTION.....	79
4.1.1.	Quinones.....	82
4.1.2.	Quinone Binding Site Design.....	85
4.1.3.	Summary	86
4.1.4.	Theory of Techniques Used.....	86
4.2.	RESULTS.....	88
4.2.1.	Cofactor Assembly in BFR Heterodimers.....	88
4.2.2.	Quinone Binding.....	98
4.2.3.	Light Induced Changes in BFR Heterodimers	101
4.3.	DISCUSSION.....	109
4.3.1.	Cofactor-Protein Assembly.....	109

4.3.2.	<i>Light Induced Effects</i>	113
4.3.3.	<i>Electron Pathways Modelling</i>	117
4.4.	CONCLUSION	120
5	CONCLUSIONS AND FUTURE WORK.....	121
	REFERENCES.....	125

120	CONCLUSIONS AND FUTURE WORK	120
121	REFERENCES	121
122	APPENDIX A	122
123	APPENDIX B	123
124	APPENDIX C	124
125	APPENDIX D	125
126	APPENDIX E	126
127	APPENDIX F	127
128	APPENDIX G	128
129	APPENDIX H	129
130	APPENDIX I	130
131	APPENDIX J	131
132	APPENDIX K	132
133	APPENDIX L	133
134	APPENDIX M	134
135	APPENDIX N	135
136	APPENDIX O	136
137	APPENDIX P	137
138	APPENDIX Q	138
139	APPENDIX R	139
140	APPENDIX S	140
141	APPENDIX T	141
142	APPENDIX U	142
143	APPENDIX V	143
144	APPENDIX W	144
145	APPENDIX X	145
146	APPENDIX Y	146
147	APPENDIX Z	147
148	APPENDIX AA	148
149	APPENDIX AB	149
150	APPENDIX AC	150
151	APPENDIX AD	151
152	APPENDIX AE	152
153	APPENDIX AF	153
154	APPENDIX AG	154
155	APPENDIX AH	155
156	APPENDIX AI	156
157	APPENDIX AJ	157
158	APPENDIX AK	158
159	APPENDIX AL	159
160	APPENDIX AM	160
161	APPENDIX AN	161
162	APPENDIX AO	162
163	APPENDIX AP	163
164	APPENDIX AQ	164
165	APPENDIX AR	165
166	APPENDIX AS	166
167	APPENDIX AT	167
168	APPENDIX AU	168
169	APPENDIX AV	169
170	APPENDIX AW	170
171	APPENDIX AX	171
172	APPENDIX AY	172
173	APPENDIX AZ	173
174	APPENDIX BA	174
175	APPENDIX BB	175
176	APPENDIX BC	176
177	APPENDIX BD	177
178	APPENDIX BE	178
179	APPENDIX BF	179
180	APPENDIX BG	180
181	APPENDIX BH	181
182	APPENDIX BI	182
183	APPENDIX BJ	183
184	APPENDIX BK	184
185	APPENDIX BL	185
186	APPENDIX BM	186
187	APPENDIX BN	187
188	APPENDIX BO	188
189	APPENDIX BP	189
190	APPENDIX BQ	190
191	APPENDIX BR	191
192	APPENDIX BS	192
193	APPENDIX BT	193
194	APPENDIX BU	194
195	APPENDIX BV	195
196	APPENDIX BW	196
197	APPENDIX BX	197
198	APPENDIX BY	198
199	APPENDIX BZ	199
200	APPENDIX CA	200
201	APPENDIX CB	201
202	APPENDIX CC	202
203	APPENDIX CD	203
204	APPENDIX CE	204
205	APPENDIX CF	205
206	APPENDIX CG	206
207	APPENDIX CH	207
208	APPENDIX CI	208
209	APPENDIX CJ	209
210	APPENDIX CK	210
211	APPENDIX CL	211
212	APPENDIX CM	212
213	APPENDIX CN	213
214	APPENDIX CO	214
215	APPENDIX CP	215
216	APPENDIX CQ	216
217	APPENDIX CR	217
218	APPENDIX CS	218
219	APPENDIX CT	219
220	APPENDIX CU	220
221	APPENDIX CV	221
222	APPENDIX CW	222
223	APPENDIX CX	223
224	APPENDIX CY	224
225	APPENDIX CZ	225
226	APPENDIX DA	226
227	APPENDIX DB	227
228	APPENDIX DC	228
229	APPENDIX DD	229
230	APPENDIX DE	230
231	APPENDIX DF	231
232	APPENDIX DG	232
233	APPENDIX DH	233
234	APPENDIX DI	234
235	APPENDIX DJ	235
236	APPENDIX DK	236
237	APPENDIX DL	237
238	APPENDIX DM	238
239	APPENDIX DN	239
240	APPENDIX DO	240
241	APPENDIX DP	241
242	APPENDIX DQ	242
243	APPENDIX DR	243
244	APPENDIX DS	244
245	APPENDIX DT	245
246	APPENDIX DU	246
247	APPENDIX DV	247
248	APPENDIX DW	248
249	APPENDIX DX	249
250	APPENDIX DY	250
251	APPENDIX DZ	251
252	APPENDIX EA	252
253	APPENDIX EB	253
254	APPENDIX EC	254
255	APPENDIX ED	255
256	APPENDIX EE	256
257	APPENDIX EF	257
258	APPENDIX EG	258
259	APPENDIX EH	259
260	APPENDIX EI	260
261	APPENDIX EJ	261
262	APPENDIX EK	262
263	APPENDIX EL	263
264	APPENDIX EM	264
265	APPENDIX EN	265
266	APPENDIX EO	266
267	APPENDIX EP	267
268	APPENDIX EQ	268
269	APPENDIX ER	269
270	APPENDIX ES	270
271	APPENDIX ET	271
272	APPENDIX EU	272
273	APPENDIX EV	273
274	APPENDIX EW	274
275	APPENDIX EX	275
276	APPENDIX EY	276
277	APPENDIX EZ	277
278	APPENDIX FA	278
279	APPENDIX FB	279
280	APPENDIX FC	280
281	APPENDIX FD	281
282	APPENDIX FE	282
283	APPENDIX FF	283
284	APPENDIX FG	284
285	APPENDIX FH	285
286	APPENDIX FI	286
287	APPENDIX FJ	287
288	APPENDIX FK	288
289	APPENDIX FL	289
290	APPENDIX FM	290
291	APPENDIX FN	291
292	APPENDIX FO	292
293	APPENDIX FP	293
294	APPENDIX FQ	294
295	APPENDIX FR	295
296	APPENDIX FS	296
297	APPENDIX FT	297
298	APPENDIX FU	298
299	APPENDIX FV	299
300	APPENDIX FW	300
301	APPENDIX FX	301
302	APPENDIX FY	302
303	APPENDIX FZ	303
304	APPENDIX GA	304
305	APPENDIX GB	305
306	APPENDIX GC	306
307	APPENDIX GD	307
308	APPENDIX GE	308
309	APPENDIX GF	309
310	APPENDIX GG	310
311	APPENDIX GH	311
312	APPENDIX GI	312
313	APPENDIX GJ	313
314	APPENDIX GK	314
315	APPENDIX GL	315
316	APPENDIX GM	316
317	APPENDIX GN	317
318	APPENDIX GO	318
319	APPENDIX GP	319
320	APPENDIX GQ	320
321	APPENDIX GR	321
322	APPENDIX GS	322
323	APPENDIX GT	323
324	APPENDIX GU	324
325	APPENDIX GV	325
326	APPENDIX GW	326
327	APPENDIX GX	327
328	APPENDIX GY	328
329	APPENDIX GZ	329
330	APPENDIX HA	330
331	APPENDIX HB	331
332	APPENDIX HC	332
333	APPENDIX HD	333
334	APPENDIX HE	334
335	APPENDIX HF	335
336	APPENDIX HG	336
337	APPENDIX HH	337
338	APPENDIX HI	338
339	APPENDIX HJ	339
340	APPENDIX HK	340
341	APPENDIX HL	341
342	APPENDIX HM	342
343	APPENDIX HN	343
344	APPENDIX HO	344
345	APPENDIX HP	345
346	APPENDIX HQ	346
347	APPENDIX HR	347
348	APPENDIX HS	348
349	APPENDIX HT	349
350	APPENDIX HU	350
351	APPENDIX HV	351
352	APPENDIX HW	352
353	APPENDIX HX	353
354	APPENDIX HY	354
355	APPENDIX HZ	355
356	APPENDIX IA	356
357	APPENDIX IB	357
358	APPENDIX IC	358
359	APPENDIX ID	359
360	APPENDIX IE	360
361	APPENDIX IF	361
362	APPENDIX IG	362
363	APPENDIX IH	363
364	APPENDIX II	364
365	APPENDIX IJ	365
366	APPENDIX IK	366
367	APPENDIX IL	367
368	APPENDIX IM	368
369	APPENDIX IN	369
370	APPENDIX IO	370
371	APPENDIX IP	371
372	APPENDIX IQ	372
373	APPENDIX IR	373
374	APPENDIX IS	374
375	APPENDIX IT	375
376	APPENDIX IU	376
377	APPENDIX IV	377
378	APPENDIX IW	378
379	APPENDIX IX	379
380	APPENDIX IY	380
381	APPENDIX IZ	381
382	APPENDIX JA	382
383	APPENDIX JB	383
384	APPENDIX JC	384
385	APPENDIX JD	385
386	APPENDIX JE	386
387	APPENDIX JF	387
388	APPENDIX JG	388
389	APPENDIX JH	389
390	APPENDIX JI	390
391	APPENDIX JJ	391
392	APPENDIX JK	392
393	APPENDIX JL	393
394	APPENDIX JM	394
395	APPENDIX JN	395
396	APPENDIX JO	396
397	APPENDIX JP	397
398	APPENDIX JQ	398
399	APPENDIX JR	399
400	APPENDIX JS	400
401	APPENDIX JT	401
402	APPENDIX JU	402
403	APPENDIX JV	403
404	APPENDIX JW	404
405	APPENDIX JX	405
406	APPENDIX JY	406
407	APPENDIX JZ	407
408	APPENDIX KA	408
409	APPENDIX KB	409
410	APPENDIX KC	410
411	APPENDIX KD	411
412	APPENDIX KE	412
413	APPENDIX KF	413
414	APPENDIX KG	414
415	APPENDIX KH	415
416	APPENDIX KI	416
417	APPENDIX KJ	417
418	APPENDIX KK	418
419	APPENDIX KL	419
420	APPENDIX KM	420
421	APPENDIX KN	421
422	APPENDIX KO	422
423	APPENDIX KP	423
424	APPENDIX KQ	424
425	APPENDIX KR	425
426	APPENDIX KS	426
427	APPENDIX KT	427
428	APPENDIX KU	428
429	APPENDIX KV	429
430	APPENDIX KW	430
431	APPENDIX KX	431
432	APPENDIX KY	432
433	APPENDIX KZ	433
434	APPENDIX LA	434
435	APPENDIX LB	435
436	APPENDIX LC	436
437	APPENDIX LD	437
438	APPENDIX LE	438
439	APPENDIX LF	439
440	APPENDIX LG	440
441	APPENDIX LH	441
442	APPENDIX LI	442
443	APPENDIX LJ	443
444	APPENDIX LK	444
445	APPENDIX LL	445
446	APPENDIX LM	446
447	APPENDIX LN	447
448	APPENDIX LO	448
449	APPENDIX LP	449
450	APPENDIX LQ	450
451	APPENDIX LR	451
452	APPENDIX LS	452
453	APPENDIX LT	453
454	APPENDIX LU	454
455	APPENDIX LV	455
456	APPENDIX LW	456
457	APPENDIX LX	457
458	APPENDIX LY	458
459	APPENDIX LZ	459
460	APPENDIX MA	460
461	APPENDIX MB	461
462	APPENDIX MC	462
463	APPENDIX MD	463
464	APPENDIX ME	464
465	APPENDIX MF	465
466	APPENDIX MG	466
467	APPENDIX MH	467
468	APPENDIX MI	468
469	APPENDIX MJ	469
470	APPENDIX MK	470
471	APPENDIX ML	471

Acknowledgements

I would like to thank my supervisor Tom Wydrzynski for giving me the opportunity to do this project, and for his help throughout it. Nick Cox for his help with all things EPR related. Spencer Whitney for RSBS happy hour, which keeps a student sane, and also his help as the molecular biology guru. Warwick Hillier for reading manuscripts and general discussions. The Photobioenergetics group for generally being a good group of people to share a lab with. Thanks also to my fellow students Adele Williamson and Iain McConnell for a student's point of view. Particular thanks go to my parents and family for always supporting my scientific adventures. Finally I would like to thank my lovely wife Carly for giving me a reason to just drop it all and come home at the end of the day.

Acknowledgements

I would like to thank my supervisor Tom Wyllie for giving me the opportunity to do this project, and for his help throughout it. Nick Cole for his help with all things EPR related. Spencer Whitty for his happy hour, which keeps a student sane, and also his help at the molecular biology exam. Warwick Illister for reading manuscripts and general discussion. The Photochemistry group for generally being a good group of people to share a lab with. Thanks also to my fellow students: Adam Williamson and Jan McCreel for a student's point of view. I would like to thank my family for always supporting my scientific adventures. Finally I would like to thank my lovely wife Cathy for giving me a reason to just drop it all and come home at the end of the day.

Abstract

Photosynthesis involves the conversion of light into chemical energy through a series of electron transfer reactions within membrane-bound pigment/protein complexes. The unique Photosystem II (PSII) complex in plants, catalyses the oxidation of water to molecular oxygen. The complexity of PSII has thus far limited attempts to chemically mimic this enzyme. Here I have used a reverse engineering approach to attempt to build several simple light driven manganese oxidising catalysts based on PSII structure and function. I have used bacterioferritin (BFR) (cytochrome *b1*) from *E. coli* as a protein scaffold to make an *in vitro* model of the PSII reaction centre. BFR has several design features, which can be utilized to engineer light-driven electron transport – a di-iron binding site, a potentially redox active tyrosine residue, and the ability to dimerise and form an inter-protein heme binding pocket. The heme could be replaced with a photoactive zinc-chlorin e_6 ($ZnCe_6$) molecule and the di-iron metal binding site with two manganese ions.

Two protein systems were made, firstly a homodimeric BFR and secondly a heterodimeric form of BFR which enabled the addition of a quinone electron acceptor within the protein matrix. In the first system I was able to demonstrate that BFR binds two magnetically interacting Mn ions as a weakly coupled $Mn_2^{II,II}$ centre, and that $ZnCe_6$ is bound stoichiometrically to the dimeric BFR complex. The bound $ZnCe_6$ species initiates electron transfer upon illumination, which oxidizes a tyrosine and causes oxidation of the Mn centre in a light and oxygen dependent reaction. The light

dependent loss of Mn^{II} signals and the formation of low field parallel mode Mn signals in the EPR spectra are attributed to the formation of Mn^{III} species.

The second system was produced by engineering a protein containing two BFR monomers joined via a 22 amino acid flexible linker region. This produced a heterodimer where one half of the system could be changed relative to the other half. With this construct the bound $ZnCe_6$ still oxidises a manganese centre via a tyrosine and appears to donate the electron to a covalently bound quinone. The removal of one of the metal sites and the placement of a quinone within this site increases the amount of oxygen consumed by the system suggesting increased efficiency of electron transfer. These two models are the first artificial reaction centres developed which carry out photo-catalytic oxidation of a dimetal site within a protein matrix.

List of Tables

Table 3.1: Best fit parameters for ITC measurements of BFR ^I cofactor binding	55
Table 3.2: Simulation parameters for the Mn ^{II} dimers	61
Table 3.3: Oxygen uptake data	67
Table 3.4: Relative metal ion EPR signal intensities.....	69
Table 4.1: Fitted CD Spectra	90
Table 4.2: Best fit parameters for ITC measurements of Mn and ZnCe ₆ binding to BFR constructs	95
Table 4.3: Iron oxidation rates as monitored by UV-visible spectroscopy	97
Table 4.4: ZnCe ₆ absorbance maxima when bound to BFR proteins.....	100
Table 4.5: Oxygen uptake data	104

List of Figures

Figure 1.1: Schematic of photosystem II showing the major redox active cofactors.	1
Figure 1.2: PSII energy redox potentials for the essential redox cofactors	3
Figure 1.3: Recent model for the Mn ₄ OxCa cluster in PSII.	5
Figure 1.4: Cofactors and the distances separating the cofactors in PSII.	8
Figure 1.5: A Jablonski diagram.	9
Figure 1.6: Theoretical Marcus curve.	11
Figure 1.7: Three light active porphyrins; Chlorophyll-a, Zinc chlorin e ₆ and Zinc protoporphyrin IX.	14
Figure 2.1: Ligation independent cloning strategy.	24
Figure 2.2: The pET32 Xa/LIC vector map.	25
Figure 2.3: The amino acid and DNA sequence for the double mutant, BFR ¹	26
Figure 2.4: Four gene constructs made with different restriction sites to enable ligation in different combinations.	28
Figure 2.5: pGEM-T easy vector map.	29
Figure 2.6: The amino acid sequence and DNA sequence of the linker used to join the Bfr subunits.	30
Figure 2.7: The <i>bfr</i> gene construct BFR ¹ -SE-L made with a <i>bfr</i> gene and a linker region.	30
Figure 2.8: The two heterodimer constructs assembled in the pTE vector.	31
Figure 2.9: Amino acid sequence and DNA sequence of the linker region after mutation of both histidines to serines.	31
Figure 2.10: The final two heterodimer gene constructs after site-directed mutagenesis to remove the EcoRV and NdeI restriction sites.	32
Figure 2.11: The Histidine-tagged Ubiquitin Expression vector (pHUE) map.	33
Figure 3.1: A ribbon diagram of the sphere structure formed by <i>E.coli</i> bacterioferritin.	42
Figure 3.2: The BFR homodimer showing bound manganese and heme.	43
Figure 3.3: Distances of cofactors in the BFR complex as compared to PSII.	44
Figure 3.4: UV-visible spectra of BFR ¹ in its oxidised form and reduced form.	51
Figure 3.5: Circular dichroism spectra of BFR and the double mutant BFR ¹	52

Figure 3.6: UV-visible spectra of ZnCe ₆ free in solution and bound to BFR ¹	53
Figure 3.7: Calorimetric titration of BFR ¹ into a solution of ZnCe ₆	54
Figure 3.8: Calorimetric titrations of MnCl ₂ into a solution of BFR ¹	55
Figure 3.9: Cyclic voltammogram of BFR ¹ -Mn.	56
Figure 3.10: Temperature dependence of the EPR spectra of BFR ¹ -M.....	57
Figure 3.11: Simulated EPR spectra and temperature dependence of BFR ¹ -M.	58
Figure 3.12: Temperature dependence of the EPR line-shape observed in BFR ¹ -ZM..	59
Figure 3.13: Comparison of Mn ^{II} EPR signal observed in BFR ¹ -M and BFR ¹ -ZM.....	60
Figure 3.14: Comparison of temperature dependence of Mn ^{II} EPR signal observed in BFR ¹ -M and BFR ¹ -ZM.	60
Figure 3.15: Simulated EPR spectra and temperature dependence of BFR ¹ -ZM after the addition of ZnCe ₆	61
Figure 3.16: EPR spectra of the BFR ¹ -ZM complex and underlying multiline signal..	62
Figure 3.17: Pictorial representation of Mn-Mn co-ordinate axis system.	63
Figure 3.18: Clarke type oxygen electrode measurements of the BFR ¹ complex.	66
Figure 3.19: Light induced EPR spectra for BFR ¹ -Z with and without manganese bound.....	68
Figure 3.20: Parallel mode EPR spectra of light minus dark spectra of BFR ¹ -ZM with a five fold excess of Mn.....	69
Figure 3.21: Diagram of the amino acid ligands to the dimanganese center with ZnCe ₆ bound at the interface of the BFR dimeric complex.	70
Figure 4.1: Bacterioferritin heterodimer (BFR-H2) with one subunit of the dimer changed relative to the other.	80
Figure 4.2: Diagram showing the relative distance of separation of all of the cofactors bound to the BFR heterodimer protein.	81
Figure 4.3: Ubiquinone.	83
Figure 4.4: Covalent attachment of a quinone to the cysteine sulphur of a protein.....	85
Figure 4.5: UV-visible spectra of holo BFR-H2 in its oxidised and reduced form.	88
Figure 4.6: CD spectra of BFR-H1 with and without manganese.	89
Figure 4.7: CD spectra of BFR-H2 with and without manganese.	90
Figure 4.8: Calorimetric titration of MnCl ₂ into a solution of BFR ²	91
Figure 4.9: Calorimetric titrations of MnCl ₂ into a solution of BFR-H1.....	92
Figure 4.10: Calorimetric titrations of MnCl ₂ into a solution of BFR-H2.	93

Figure 4.11: Calorimetric titrations of BFR-H1 into a solution of ZnCe.	94
Figure 4.12: EPR spectra of BFR-H2-ZM-CoQ ₀ , BFR-H1-ZM and BFR ¹ -ZM.....	96
Figure 4.13: Iron oxidation monitored by absorbance at 340nm.....	97
Figure 4.14: Absorbance spectra of CoQ ₀ , glutathione and the CoQ ₀ -Glutathione complex.	98
Figure 4.15: UV-visible absorbance spectra of BFR-H2-CoQ ₀ , BFR-H2 and CoQ ₀ - Glutathione complex.	99
Figure 4.16: Absorbance spectra of ZnCe ₆ free in solution, and bound to BFR-H2- CoQ ₀	99
Figure 4.17: Room temperature EPR spectra of CoQ ₀ and 1,2-naphthoquinone.	100
Figure 4.18: Room temperature EPR spectra of CoQ ₀ and CoQ ₀ bound to BFR ²	101
Figure 4.19: EPR spectrum of BFR-H1 ZM before and after illumination at 200K. ..	102
Figure 4.20: EPR spectra of BFR-H1 ZM and BFR-H1 Z.	103
Figure 4.21: Oxygen uptake measurements of the heterodimer complexes.	104
Figure 4.22: Fluorescence spectra of BFR-H2 ZM and BFR-H2 ZM CoQ ₀	105
Figure 4.23: Fluorescence spectra of reduced BFR heterodimer proteins.....	106
Figure 4.24: Illuminated room temperature EPR spectra of BFR-H2 Z NQ and BFR-H2 Z.	107
Figure 4.25: Illuminated room temperature EPR spectra of BFR-H2 ZM and BFR-H2 ZM NQ.	108
Figure 4.26: A cartoon model of the BFR heterodimer.	109
Figure 4.27: Calculated best pathway for electron transfer within the BFR heterodimer quino-protein.	118
Figure 4.28: Calculated best pathway for electron transfer within the Cyt <i>b</i> ₅₆₂ quino- protein.	119

List of Abbreviations

A	electron acceptor
Amp	ampicillin
BFR	bacterioferritin
BFR ¹	bacterioferritin double mutant H46R, H112R without cofactors
BFR ¹ -M	BFR ¹ with two manganese ions bound
BFR ¹ -Z	BFR ¹ with ZnCe ₆ bound
BFR ¹ -ZM	BFR ¹ with stoichiometric amounts of ZnCe ₆ and manganese bound
BFR ²	BFR triple mutant H46R, H112R, E51C
BFR-H1	BFR ¹ plus BFR ¹ heterodimer joined by a linker region
BFR-H2	BFR ¹ plus BFR ² heterodimer joined by a linker region
BFR-H2 CoQ ₀	BFR-H2 with co-enzyme Q ₀ covalently bound
BFR-H2 NQ	BFR-H2 with 1,2-naphthoquinone covalently bound
BChl	bacteriochlorophyll
BRC	bacterial reaction centre
CD	circular dichroism
Chl	chlorophyll
CoQ ₀	co-enzyme Q ₀ (2,3-dimethoxy-5-methyl-1,4-benzoquinone)
Cyt	cytochrome
D	electron donor
E _m	midpoint redox potential
EPR	electron paramagnetic resonance
Heme	Fe (2+ or 3+)-protoporphyrin-IX
IPTG	Isopropyl β-D-1-thiogalactopyranoside
LB	Luria-Bertani Media
IC	internal conversion
ISC	inter system crossing
ITC	isothermal titration calorimetry
MEK	methyl ethyl ketone
Mn ₄ O _x Ca	tetranuclear manganese/calcium cluster
Native PAGE	native polyacrylamide gel electrophoresis
NQ	1,2-naphthoquinone

OEC	oxygen evolving complex
P	primary donor
P*	excited state of primary donor
PCR	polymerase chain reaction
PSII	photosystem II
Q	quinone
T	temperature
Tyr	tyrosine
Y	redox-active tyrosine
ZnCe ₆	zinc ^{II} chlorin e ₆
ZnPP	zinc protoporphyrin
Xgal	5-bromo-4-chloro-3-indolyl- beta-D-galactopyranoside

CHAPTER 1

1 Introduction

One of the most devastating transitions in the history of the earth occurred two to three billion years ago: primitive bacteria acquired the ability to split water into O_2 using the energy from sunlight. This tumultuous event upset the world as we know it, since all life previously existed in a purely anaerobic environment. Organisms were then forced to develop ways to prevent oxidative damage or else face extinction. Today, multicellular organisms thrive in an oxygenic environment by using aerobic respiration for energy conversion (Blankenship 2002).

At the heart of oxygenic photosynthesis is photosystem II (PSII), the light-driven enzyme complex that catalyzes the oxidation of water into molecular O_2 and the reduction of plastoquinone to plastoquinol (Figure 1.1) (Wydrzyński and Satoh 2005). This enzyme makes insight to examine the most energetically demanding reaction in biology – the formation of the O–O bond. PSII is one of the most elaborate and subtle biological machines which has evolved only once in Nature.

Figure 1.1 Schematic of photosystem II showing the light-driven active site and pathway for oxidation.

CHAPTER 1

1 Introduction

One of the most devastating transitions in the history of the earth occurred two to three billion years ago; primitive bacteria acquired the ability to split water into O_2 using the energy from sunlight. This tumultuous event turned the world upside down since all life previously existed under a purely anaerobic environment. Organisms were then forced to develop ways to prevent oxidative damage or else face extinction. Today multicellular organisms thrive in an oxygenic environment by using aerobic respiration for energy conversion (Blankenship 2002).

At the heart of oxygenic photosynthesis is photosystem II (PSII), the light-driven oxidoreductase that catalyses the oxidation of water into molecular O_2 and the reduction of plastoquinone to plastoquinol (Figure 1.1) (Wydrzynski and Satoh 2005). This enzyme utilizes sunlight to catalyse the most energetically demanding reaction in biology - the formation of the O-O bond. PSII is one of the most elaborate nanoscale biological machines which has evolved only once in Nature.

Figure 1.1: Schematic of photosystem II showing the major redox active cofactors necessary for function.

Over millennia a vast amount of carbon based material was laid down from oxygenic photosynthesis and converted into the fossil fuels we use today – coal, oil and natural gas. The utilization of fossil fuels is becoming a crucial concern as human energy consumption has climbed to such levels that we are now changing our climate patterns due to the CO₂ emissions. To prevent further release of greenhouse gases, alternative energy sources need to be used. Understanding oxygenic photosynthesis and being able to exploit its ability to directly convert sunlight and water into clean fuel is now taking on a greater priority in research centres worldwide. The initial charge separation events in PSII have a remarkable efficiency of approximately 90% making it extremely attractive to mimic this system. Development of equally efficient catalysts would provide for cheap and renewable fuel sources. The production of hydrogen ions that can be converted into hydrogen gas, is now being promoted as the next generation clean fuel for transport and much research is focused on developing such technologies (Kruse *et al.* 2005; Lewis and Nocera 2006). The first step towards this goal is to replicate PSII function in a well-defined, robust supporting matrix. The following sections outline the photosynthetic capture of energy and describe my reverse engineering approach for developing PSII model proteins.

1.1. Photosystem II

1.1.1. The Reaction Centre

The primary photochemical event in PSII is the critical transmembrane charge separation. Upon absorption of a photon, an electron in the PSII P680 chlorophyll complex is promoted from the ground state to the excited singlet state and is poised to undergo electron transfer to the primary electron acceptor, a pheophytin, which rapidly transfers this electron to the first and second plastoquinone acceptors (Q_A and Q_B). Spatial separation of the positive and negative charges by a distance greatly reduces the chances of recombination and the charges are stabilized.

The P680^{•+} radical is a strong oxidant ($E_0 \sim 1.2$ V) that is reduced by the proximal tyrosine (Y_Z). This is believed to be coupled to a deprotonation reaction generating the

neutral $Y_Z^\bullet$ radical (Rappaport and Diner 2008), which then oxidizes the Mn_4O_xCa cluster. For each photon-induced charge separation event at P680, the Mn_4O_xCa cluster advances one oxidation state termed S_0 - S_4 . After four such photo-events (S_4), molecular oxygen and protons are released and the Mn_4O_xCa cluster reverts to its original oxidation state (S_0) (Figure 1.2) (Yachandra 2005).

Figure 1.2: PSII energy diagram illustrating the approximate redox potentials in Volts for the essential redox cofactors of PSII involved in producing the charge separated state necessary to oxidize water. For each photon of light that is absorbed by the P680 chlorophylls one electron is sent via a pheophytin to Q_A and finally Q_B which dissociates from the membrane once it is doubly reduced and is replaced by an oxidized quinone. For each electron ejected from the P680 complex an electron is taken from the manganese centre until the fourth and final electron is taken which oxidizes water into O_2 and $4H^+$ ions. The redox potentials in Volts for the individual components are P680⁺/P680 +1.2, P680/P680^{*} -0.55, Pheo_{D1}/Pheo_{D1}⁻ -0.37 to -0.5, Q_A/Q_A^- -0.03, Q_B/Q_B^- +0.07, Tyr_Z^{ox}/Tyr_Z +1.2, Mn₄ cluster - S_1/S_2 , S_2/S_3 , S_3/S_0 +1.12, +1.14, and +1.08 (Rappaport and Diner 2008).

1.1.2. The Oxygen Evolving Complex: Models and Mechanisms

The oxygen evolving complex (OEC) of PSII is a unique functional motif in biology not only because it can oxidize water but also because of the five metal ions that are bound within the one catalytic site. The Mn_4O_xCa structure is directly coordinated by glutamate, aspartate, histidine and alanine residues (Ferreira *et al.* 2004; Loll *et al.* 2005; Sproviero *et al.* 2006; Yano *et al.* 2006). These metal ions are also complexed

with bridging μ -oxo bonds. The accumulation of oxidizing potential is assumed to take the form of incremental changes in the formal oxidation states of the Mn ions in the OEC. The 4 electron S-state cycle is predicted to begin in the S_0 state as $\text{Mn}^{\text{III}}\text{Mn}^{\text{III}}\text{Mn}^{\text{III}}\text{Mn}^{\text{IV}}$ (Kulik *et al.* 2007) and progresses to $\text{Mn}^{\text{III}}\text{Mn}^{\text{III}}\text{Mn}^{\text{IV}}\text{Mn}^{\text{IV}}$ in the S_1 state (Sauer *et al.* 2005; Yachandra 2005), finally reaching $\text{Mn}^{\text{III}}\text{Mn}^{\text{IV}}\text{Mn}^{\text{IV}}\text{Mn}^{\text{IV}}$ in the S_2 state (Kulik *et al.* 2007). The S_3 state is also suggested to possibly involve a ligand centred oxidation event with no change in the formal Mn oxidation states (Messinger *et al.* 2001), though there are alternative metal centred oxidation suggestions (Dau and Haumann 2008). At the S_4 state water is split and the whole complex resets itself to the S_0 state. It is unknown whether the Mn_4 centre undergoes a transient elevation of oxidation state in the S_4 state before water is oxidized. To date no Mn^{V} manganyl species has been detected for any of the meta-stable S-states (S_0 - S_3) (Weng *et al.* 2004).

A distinctive property of the OEC is not so much that it can produce oxygen, as catalases can also do this, but that it produces an O-O bond. There are no other known examples in nature where such a bond is formed between two H_2O molecules without long lived intermediates like H_2O_2 being released (Armstrong 2008). The extrinsic proteins are thought to help stabilize this reaction by shielding the OEC from the bulk solvent water and by providing channels that deliver substrate (H_2O) and permit an orderly exit of products (O_2 , H^+). This idea was first promoted by Wydrzynski (Wydrzynski *et al.* 1996) and since then analysis of the crystal structures has revealed several channels that potentially aid in the transport of O_2 and H_2O (Murray and Barber 2007; Ho and Styring 2008). These channels would undoubtedly be involved in limiting substrate access aiding catalysis and reducing the possibility of damaging side reactions.

The Mn ions themselves are thought to bind the substrate water molecules. Two separate binding sites have been identified from ^{18}O isotope exchange measurements revealing a slow and a fast exchanging site (Messinger *et al.* 1995). Through the use of specialized EPR techniques it has been shown that a solvent water molecule definitely ligates directly to a Mn ion at one of the exchange sites

Figure 1.3) (Su *et al.* 2008). The slow exchanging site is due to ligation of the solvent water molecule to the Ca ion and another Mn ion (Hendry and Wydrzynski 2003).

Figure 1.3: Recent model for the Mn_4OxCa cluster in PSII with suggested substrate “water” binding sites. O-O bond formation has been proposed by different groups to occur in the S_4 state between oxygen’s 1 and 2, 3 and 4, 4 and 5, 1 and 6, or 6 and 8 (Su *et al.* 2008). Purple (Mn), red (O), green (Ca), and grey (H).

There are a large number of theories on the mechanism of O-O bond formation in the S_4 state of PSII. Experimental evidence suggests that the delay in cycling from S_3 through S_4 back to S_0 is due to a proton extraction event (Clausen *et al.* 2004; Haumann *et al.* 2005). The driving energy is small for the S_3 and S_4 states and thus proton removal events are definitely of major importance. The proton release pattern from the OEC in the four S states ($S_0 \rightarrow S_1$, $S_1 \rightarrow S_2$, $S_2 \rightarrow S_3$, and $S_3 \rightarrow [S_4] \rightarrow S_0$), appears to be 1,0,1,2 (McEvoy and Brudvig 2006; Dau and Haumann 2008). Far from suggesting a mechanism for O-O bond formation I would refer the reader to several articles that discuss the different possibilities that could apply given the latest refinements in structure (Hillier and Messinger 2005; McEvoy *et al.* 2005; Siegbahn 2006; Siegbahn 2008).

1.1.3. Why Manganese?

In order to function the OEC can only use manganese and no other metal ions. The calcium ion is not so specific as it can be replaced with strontium, but there are no functional substitutes for Mn (Boussac *et al.* 2004). One may speculate that plants initially utilized Mn because of its high availability as the aqua ion $\text{Mn}(\text{H}_2\text{O})_6^{2+}$ in aqueous solution. This Mn^{II} cation is very stable, exchanges ligands rapidly and forms soluble salts; this coupled with its high natural abundance and the fact that a cell’s requirements for Mn are very low means that supply usually exceeds demand. Once taken up by an organism, Mn will tend to remain as Mn^{II} unless it is captured by ligands that favour its oxidation to Mn^{III} and Mn^{IV} (Armstrong 2008).

The energetics of O_2 formation from H_2O are such that at pH 7, formation of free molecular O_2 requires an energy input of 0.82 eV per electron, with a sum total of 3.68 eV for 4 electrons and 4 protons to be removed (Krishtalik 1986). PSII has only ~ 1.2 eV potential provided by $P680^+$, and so the reaction has to be split into 4 single electron transfer steps in order for the reaction to proceed (Grabolle and Dau 2005). To carry out such a reaction it is necessary for the OEC to contain multiple electron donors that are able to cycle through different oxidation states to store the built-up charge. Transition metals are the obvious choice for such a catalyst as these ions commonly have multiple stable oxidation states, due to the relative ease of removing unpaired d-electrons (Cowan 1996).

Nature uses many different first-row transition (d-block) elements such as V, Mn, Fe, Co, Ni, Cu, and Zn. As we move across the 3d row, the highest accessible oxidation states as well as the number of accessible oxidation states increases up to Mn then decreases to Zn where only the Zn^{II} state is observed. This is mainly a product of ionization energies, which increase across the row and for each electron removed. Not only does Mn display a greater range of oxidation states than other 3d metals, but all the simple higher states, from Mn^{III} upwards, are hydroxo- or oxo-species with reduction potentials in the ranges of O-transformation reactions (Armstrong 2008).

Of all the transition metals iron has the most similar chemistry to that of manganese. Iron II, III and IV oxidation states (?) have been reported in biological systems. A multitude of enzymes utilize $Fe^{II/III}$ chemistry and enzymes such as iron catalase and peroxidase, employ an $Fe^{IV}=O$ species from heme iron. Like manganese, iron also has preferences for oxygen ligands in higher oxidation states (Hersleth *et al.* 2006).

Mn can reach a large number of oxidation states with the biologically relevant states being the II, III, and IV states. Mn^{II} has a ligand preference for O-donors, although compared with either Mg or Ca, it does have a greater tendency to bind to N-donors. The intrinsic stability of Mn^{II} arises from the electron configuration $3d^5$ where the d-shell is half filled. This configuration is favoured due to optimal exchange energy, making it difficult to remove the fifth d-electron, so Mn^{III} compounds are particularly oxidizing. The difficulty in stabilizing Mn^{III} decreases as ligands with good donor properties are introduced. A good example of this is bicarbonates ability to ligate and

stabilize Mn lowering the IV/III reduction potential from 1.5 to 0.6 eV (Kozlov *et al.* 2004; Dasgupta *et al.* 2006). Mn^{IV} complexes regularly contain bridging or terminal oxo-ligands, and polynuclear Mn–O clusters of mixed valency are common. Owing to its d^3 configuration, Mn^{IV} should favour octahedral geometry. Mn^{IV} is rarely found in biology, the only enzymes where this high oxidation state is found naturally is in the OEC of PSII and more recently it was found as a mixed $\text{Mn}^{\text{IV}}/\text{Fe}^{\text{III}}$ cofactor in *Chlamydia trachomatis* ribonucleotide reductase (Jiang *et al.* 2007).

1.2. Design of Photo-Catalytic Proteins

PSII in higher plants is a highly complex, membrane-based enzyme made up of at least 27 protein subunits and 32 cofactors involved in electron transfer and light harvesting (Wydrzynski and Satoh 2005) (Figure 1.4). The large size and complexity of this enzyme makes it difficult to work with and any mimic of this protein must by definition be simpler. Development of a minimalist photo-catalytic structural unit requires that only the essential redox cofactors of PSII are engineered into a simple framework. The essential cofactors for the function of PSII include: a bound plastoquinone electron acceptor; a high potential photo-oxidisable chlorophyll (Chl) complex; a redox-active tyrosine; and a tetra-nuclear manganese/calcium cluster ($\text{Mn}_4\text{O}_x\text{Ca}$) i.e. quinone – Chl complex – tyrosine – manganese cluster (Figure 1.2).

The cofactors present in PSII are placed at specific positions in space by the protein framework. The amino acid residues responsible for binding these cofactors function not only as a scaffold but to mediate the redox potentials of the cofactors bound. In the majority of enzymes redox midpoint potentials of bound cofactors are set such that electron transfer is unidirectional (Rabanal *et al.* 1996; Page *et al.* 1999). The following section will detail electron transfer principles necessary for engineering a photo-catalytic system and previous attempts to introduce light activated pigments, electron acceptors and electron donors within a protein framework in a redox active form.

Figure 1.4: Schematic showing the cofactors and the distances separating the cofactors in PSII. Image taken from Loll et al 2005.

1.2.1. Electron Transfer Principles

In order for light activated systems to operate a charge separated state must be accomplished through electron transfer between donor (D) and acceptor (A) molecules, at a rate faster than the relaxation of the excited state. Upon absorption of a sufficiently energetic photon a light activated molecule (P), such as Chlorophyll a, is transformed to an electronically excited state (P^*). This is generally a highly-energetic, unstable and short lived state. P^* is deactivated by either radiative or non-radiative transitions back to the ground state or through electron transfer. Figure 1.5 illustrates the energy levels of the various mechanisms of energy dissipation. In some circumstances the excited state lifetimes are sufficiently long-lived ($> \sim 1\text{ns}$) to allow a bimolecular reaction with an acceptor molecule (generally ground state). Deactivation of P^* through ET occurs when P^* is returned to the ground state via a radical state (such as P^{+*}).

An example of this is:

Equation 1.2 is a consequence of the instability of the charge separated state leading to charge recombination also known as the back reaction. In photosynthesis a functional reaction centre minimizes charge recombination by additional electron transfer reactions thereby separating the charges by a greater distance. The charge separated state is energetically unstable due to Coulombic attraction between the cation and anion radicals but this attraction decreases exponentially with distance. In order to prevent charge recombination, either the charges must be further separated or the charges otherwise neutralized.

Figure 1.5: A Jablonski diagram illustrating the different energy levels and types of transitions that can occur in e.g. a porphyrin system. S_0 is the singlet ground state while S_n is any excited singlet state and S_1 is the lowest excited singlet state whilst T_1 is the lowest excited triplet state. The fluorescence and phosphorescence transitions are radiative, whilst IC and ISC are non-radiative.

Intraprotein electron transfer is typically rate-limited by tunnelling through the insulating protein medium between the edges of the D and A redox centres. This electron tunnelling is reasonably accurately described by a simple exponential decay with distance (Moser *et al.* 1992; Page *et al.* 1999). The intervening protein medium also has some role to play in electron tunnelling rates and a simple use of the packing density (ρ) to equate the tunnelling barrier has been used successfully by Dutton and co-workers. They have produced an empirical equation relating rate to distance:

$$\log k_{et}^{exer} = 13 - 0.6(R - 3.6) - 3.1(\Delta G + \lambda)2/\lambda \quad \text{Equation 1-3}$$

where k_{et}^{exer} is the rate in units of s^{-1} , R is the edge-to-edge distance between redox cofactors and 3.6 represents van der Waals contacts in Å. The ΔG and λ terms are in eV with a typical λ value of 0.9 ± 0.2 eV. Because electrons can transfer across conjugated systems virtually instantaneously, electron transfer across the distance spanned by the ring systems does not contribute significantly to the electron transfer time (Moser and Dutton 1992). This is implicit in the treatment summarized in Dutton's equation, and all electron transfer distances are measured edge-to-edge between the nearest atoms in the conjugate systems of D and A.

Work in Harry Gray's lab has led to some refinement of this picture (Gray and Winkler 1996; Winkler *et al.* 1999). They measured electron transfer rates from ruthenium and rhenium complexes attached via histidines to redox proteins with different secondary structures, positioned using site directed mutagenesis. Secondary structure was found to affect electron tunnelling with this process being highly anisotropic. Tunnelling across alpha helical pathways was found to slow transfer however tunnelling along more linear β strand like pathways produced faster electron transfer (Gray and Winkler 1996; Winkler and Gray 1997; Winkler *et al.* 1999; Gray and Winkler 2003)

Much more important than packing density is that of driving force (ΔG°). This is estimated from the difference in the redox midpoint potentials (E_m) between donor and acceptor, assuming that redox centres are not too close. One other major factor that affects tunnelling rates is that of reorganization energy (λ). Reorganization energy is defined as the energy required for all structural adjustments in the nuclear geometry of

reactants and the surrounding solvent molecules, which are needed in order that D and A molecules assume the configuration required for the transfer of the electron. The charge distribution can be quite different in the charge-separated state and must be accommodated through alteration of the local dielectric via structural rearrangements. It can be viewed as an indirect measure of activation energy for the electron transfer reaction. If the exergonic driving force of the reaction matches the reorganization energy for the reactions ($\Delta G = -\lambda$), then the tunnelling rate will be activation less and as fast as possible. If the driving force is different from this optimal value, either less exergonic (Marcus normal region) or more exergonic (Marcus inverted region), the electron tunnelling rate will decrease in an approximately Gaussian dependence of rate on driving force (see Figure 1.6) (Marcus 1956; Marcus and Sutin 1985). If one or both donor or acceptor complexes is solvent exposed then λ can become quite large ($>1\text{eV}$).

When engineering light-activated enzymes that mimic PSII, in order to produce a functional light activated redox protein, it is necessary to understand and apply the physiological limitations of electron transfer. The main elements of protein based electron tunnelling that need to be accounted for are separation distance of donor acceptor molecules, driving force of the reaction, and reorganization energies.

Figure 1.6: A theoretical Marcus curve (derived from Dutton's ruler Equation 1-3) where $\lambda = 0.7$ eV and the cofactor separation = $10 \text{ \AA}$ (Moser and Dutton 1996).

Distance:

Electron transfer rates are mostly affected by the edge-to-edge distances between D and A redox centres. Natural proteins with productive electron transfer reactions have been found to have cofactors separated by no more than $\sim 14\text{\AA}$ (Page *et al.* 1999; Page *et al.* 2003; Noy *et al.* 2006). Looking more specifically at the electron transport chain in PSII, the average redox cofactor separation distance is $\sim 10\text{\AA}$ for any one component and none are greater than $13.6\text{\AA}$ (Loll *et al.* 2005). Therefore when engineering a light activated electron transport chain it is wise to place all cofactors within $10\text{-}14\text{\AA}$. Another important consideration is that cofactors should not be placed too close to each other so as to prevent energy wasting back reactions.

Driving force:

For the purposes of engineering photo-active proteins, aside from changing the cofactor distance, altering the driving force is probably the most tangible method of tuning electron transfer reactions in a protein. Driving force is described by a Gaussian curve and as $-\Delta G^0$ is increased, K_{et} becomes maximal and then slows when $-\Delta G^0 > \lambda$. This is a rather counter-intuitive effect that has been experimentally well documented (Marcus and Sutin 1985). The region where $-\Delta G^0 > \lambda$ is called the inverted region. The value of ΔG^0 for light-induced electron transfer can be calculated by a method developed by Rehm and Weller (Rehm and Weller 1970). In the majority of engineered proteins, unless they are membrane proteins, the redox centres are likely to be at least partially solvent exposed resulting in large reorganization energies for electron transfer reactions. In such cases it is necessary to utilize a highly oxidizing pigment so that the driving force will be large enough for electron transfer to occur.

It is possible to increase or decrease the oxidation potential of D and A centres in proteins by varying the protein environment. Several studies show that large changes in redox potentials for pigments and cofactors can be achieved by changing single amino acids in a protein sequence (Gunner 1991; Allen and Williams 1995). Introducing hydrogen bonds into the inner sphere ligands of a cofactor can increase the redox midpoint potential substantially and removal of hydrogen bonds acts to decrease this potential.

Reorganization energies:

Membrane based proteins are ideal for engineering redox reactions with low reorganization energies due to D and A molecules being buried far from the polar solvent, but these proteins are extremely difficult to work with. The majority of light activated proteins engineered so far tend to be small soluble four helix bundle proteins, which are easier to work with but have solvent exposed donor and acceptor groups. This leads to high reorganization energies for electron transfer reactions thus a large driving force is desirable for these types of reaction centres.

1.2.2. Light Activated Pigments***Chlorophyll***

Attempts at producing mimics of PSII have centred on the introduction of a pigment into a peptide matrix. At first glance it would appear that a good way to incorporate a photoactive pigment would be to bind Chl as used in all reaction centres including PSII. Unfortunately it is very difficult to work with Chl due to its long phytol tail, making the molecule very hydrophobic and not amenable to binding to smaller soluble proteins in solution (Figure 1.7). There are relatively few studies on the incorporation of Chl into non-photosynthetic proteins (Kehoe *et al.* 1998; Meadows *et al.* 1998; Eggink and Hooper 2000; Rau *et al.* 2001; Satoh *et al.* 2001; Chen *et al.* 2005; Hughes *et al.* 2006; Noy and Dutton 2006), although recently a *de novo* amphiphilic protein containing a lipophilic domain was found to bind a Ni-BChl molecule (Noy *et al.* 2005).

Metal Substituted Porphyrins

A sure fire way to produce a light activated pigment bound to a protein matrix is to chemically modify a covalently attached heme, such as those found in the c-type cytochromes. Removal of the heme iron using hydrofluoric acid and replacement with zinc produces the light active zinc protoporphyrin (ZnPP) (Figure 1.7)(Vanderkooi *et al.* 1976). B-type cytochromes have also been used successfully to bind light-activated pigments. The heme group in these proteins is not covalently linked to the protein which enables the easy removal of the heme *b* molecule from the binding pocket (Teale 1959) and replacement with light active porphyrins (Petersonkennedy *et al.* 1984; Simolo *et al.* 1984; McGourty *et al.* 1987).

Figure 1.7: Stick diagrams of three light active porphyrins. Chlorophyll-a is found in higher plants whilst Zinc chlorin e₆ is a synthesized pigment. Zinc protoporphyrin IX is just a zinc substituted heme *b* molecule.

Upon light excitation ZnPP forms a triplet state with near unity efficiency ($\Phi_t = 0.9$) which is long lived ($\tau \sim 14$ ms) (Zemel and Hoffman 1981; Sudha *et al.* 1984). This has been utilized to study long range electron transfer reactions such as those between proteins. Electronically excited ZnPP (Ho *et al.* 1985; McLendon *et al.* 1985) and also the free base porphyrin (Cheung *et al.* 1986) can act as electron donors. The rate of interprotein electron transfer has been measured in several protein-protein complexes such as between Zn-hemoglobin and Fe^{III}-cyt *b*₅ (Simolo *et al.* 1984) as well as between Zn-cyt *c* and cyt *b*₅ (McLendon *et al.* 1985; Qin and Kostic 1994). These electron transfer reactions are quite slow 10^3 - 10^5 s⁻¹ despite donor/acceptor distances of 8-10 Å.

Interprotein electron transfer between Fe^{II}-cyt *c* peroxidase and Fe^{III}-cyt *c* has been utilized to measure physiological electron transfer reactions. This suggested the reorganization energy associated with these interprotein electron transfer reactions was approximately 1 eV (Cheung *et al.* 1986). The reported oxidative potentials, $E(P/P^{++})$, for zinc substituted cytochromes in horseradish peroxidase, cytochrome *c*, and myoglobin are +0.75 +0.8 and +0.98 eV respectively (Kaneko *et al.* 1980; Cowan and

Gray 1989; Magner and McLendon 1989). More recently electron transfer has been reported in other metal substituted proteins such as ZnPP-substituted nitrite reductase (Bellelli *et al.* 1996) and cyt P450cam (Furukawa *et al.* 2000). Triplet electron transfer from bound ZnPP to a quinone in solution has also been demonstrated by several groups (Sato *et al.* 1997; Fahnenschmidt *et al.* 2001).

One major drawback of utilizing ZnPP is that due to triplet state formation it produces a long lived charge separated state which allows oxygen to readily react under aerobic conditions producing damaging reactive oxygen species. By utilizing *b*-type cytochromes, porphyrins which primarily donate an electron from the singlet state can be bound to the heme pocket. Of particular success has been the use of zinc chlorin e_6 (ZnCe₆) which is very similar to heme *b* except for changes in the length of the carbon chains terminating in carboxylates and also addition of an extra carboxylic acid (Figure 1.7). This makes the pigment more soluble and easier to handle whilst maintaining a high binding affinity for the heme site. ZnCe₆ is utilized as a photoactive pigment due to its strong Q_y absorption band in the red (~638 nm), and high estimated oxidation potential of ~+1.1 eV (Hay *et al.* 2004). Zinc substituted porphyrins exhibit penta-coordinate geometry and have only one amino acid axial ligand unlike heme *b* in which the central Fe ion exhibits bis-ligation and has two amino acid axial ligands, usually histidine or methionine.

Razeghifard and Wydrzynski (2003) found that ZnCe₆ could be bound in place of heme in a *de novo* 4-helix bundle protein with two histidine residues acting as axial ligands. Upon replacing one histidine with phenylalanine, only half the amount of ZnCe₆ was bound, and upon replacing both histidines with phenylalanine, completely ablated ZnCe₆ binding. The bound ZnCe₆ could undergo some photochemistry after repetitive light flash illumination in the presence of an external quinone acceptor though with low yield (Razeghifard and Wydrzynski 2003). Light activated electron transfer has also been demonstrated with ZnCe₆ bound within a *de novo* protein to a quinone electron acceptor in solution (Mennenga *et al.* 2006).

Flavins

Flavin molecules can be utilized to carry out photo-initiated electron transfer within proteins. Upon excitation flavin in solution was found to form a semi-quinone species by abstracting an electron from a donor in solution, such as EDTA. The semi-quinone was then capable of reducing cyt *c* Fe^{III}-heme through an electron transfer reaction (Ahmad *et al.* 1981). Flavin derivatives covalently bound to cysteine residues on the surface of cyt *c*, through disulphide bond, were shown to transfer an electron to the heme group within the protein and a reorganization energy of 1.2 eV was determined (Twitchett *et al.* 1997). Other work demonstrated that a flavin bound within the interior of a *de novo* designed heme-binding maquette could carry out electron transfer from the photo-generated semi-quinone to the bound heme (Sharp *et al.* 1998).

1.2.3. Electron Acceptors

The use of electron acceptors present in solution have major drawbacks in that they must be present in high concentration and the reorganization energies associated with electron transfer are large and thus produce low yields of electron transfer in artificial systems. In order to optimize electron transfer reactions between cofactors it is ideal to bury the cofactors within the protein matrix to increase efficiency. There are a large number of electron transport chains present in natural systems that allow for transport of electrons across large distances. Within PSII, Chls, pheophytin's, and quinones are utilized. Quinones have been utilized extensively as electron acceptors due to their efficiency in transferring charge in natural systems.

Quinones

In PSII the first stable quinone electron acceptor is a specially bound plastoquinone molecule, Q_A. This quinone is highly hydrophobic, having a long hydrocarbon side chain for binding purposes and membrane solubility. Engineering a binding site for plastoquinone in a protein is as difficult as engineering a Chl binding site, due to the fact that both have long hydrophobic tails. These natural quinones are not covalently bound to the protein however in artificial reaction centres covalent attachment is a more practical approach due to the inherent difficulties in designing a binding site. A quinone

will covalently link to a free sulphhydro group (-SH), such as that found in cysteine, through a sulphur addition reaction (Redfearn 1965).

Hay et al (2004) were the first group to make use of this novel addition reaction and by modifying the gene for cytochrome *b*₅₆₂ were able to place a cysteine residue within the interior of this protein and covalently attach a quinone molecule. The cytochrome *b*₅₆₂ was chosen as it is a small soluble 4-helix bundle protein (11.6 kDa) which has been extensively characterized both biophysically and structurally (Moore *et al.* 1985; Hamada *et al.* 1995; Barker *et al.* 1996; Robinson *et al.* 1997; Arnesano *et al.* 1999; Springs *et al.* 2002).

Like all b-type cytochromes this protein binds a heme *b* molecule, therefore a mutation was made which placed a cysteine-bound quinone within ~11 Å edge-to-edge distance of the heme group. The heme could then be removed and replaced with the light active ZnCe₆. When the cysteine site was covalently linked to a coenzyme Q (CoQ₀) molecule that doesn't have the hydrophobic tail, and ZnCe₆ was bound in the heme pocket, light-induced electron transfer from the bound pigment to the bound quinone was achieved with approximately 20% efficiency compared with pre-reduced bound CoQ₀ (Hay *et al.* 2004).

When other substituted quinones were analysed, the rates of electron transfer differed by less than a factor of two, indicating that the chemical nature of the quinone molecule itself had little effect on the electron transfer efficiency (Hay *et al.* 2004). The relatively invariant electron transfer rates and the large driving forces that were estimated for the various quinones analysed also indicated that the electron transfer in the modified cyt *b*₅₆₂ is near the top of the Marcus curve with relatively large reorganization energies. Based on the quinone fluorescence emission properties, the cysteine-quinone binding site appears to be partially solvent exposed, which explains the large reorganization energies (Hay *et al.* 2004). Other methods for binding quinone's to proteins have been investigated using thiol addition reactions as well as addition-elimination reactions between the cysteine sulphur and thioether quinones (Li *et al.* 2005).

1.2.4. Electron Donors

Tyrosine radicals

In order for PSII to function a tyrosine molecule positioned near the P680 complex is oxidized donating an electron to $P680^+$. The tyrosine radical is then reduced by an electron taken from the Mn_4O_xCa complex. In producing a mimic of PSII it may be advantageous to involve a tyrosine residue to mediate the donation of electrons from the metal centre.

The Gray laboratory, in their work on metal-modified redox proteins, produced a rhenium-based photosensitiser with a Re^{II}/Re^I redox pair reduction potential of 1.85 eV vs. SCE (Connick *et al.* 1995). At pH 7.0 tyrosine has a redox potential of 0.83 eV relative to NHE (Tommos *et al.* 1999), therefore the $Re(II)$ should be a powerful enough oxidant to generate tyrosyl radicals at a high driving force. Di Bilio *et al.* (2001) showed that aromatic amino-acid radicals can be generated in rhenium modified proteins. A tyrosine could be selectively oxidized and the radical state trapped at low temperatures by preparing a rhenium-containing azurin sample in which the copper was substituted with zinc and a sacrificial oxidant was added to the medium. The irradiated freeze-trapped sample gave rise to an EPR spectrum with a line shape characteristic of that of a neutral tyrosyl radical (Di Bilio *et al.* 2001).

Metal Centres

It is quite difficult to engineer a light activated group and an electron donating metal centre within the one protein. When looking for examples of light activated oxidation of a metal centre the vast majority of studies involve the use of ruthenium and rhenium based photosensitiser's covalently attached to the surface of proteins. These photosensitiser's achieve extremely high driving forces, particularly the more recently developed rhenium adducts (Connick *et al.* 1995; Lin *et al.* 1996). Utilization of ruthenium enabled demonstration of electron transfer from the heme-Fe in cytochrome *c* to a light activated Ru^{3+} complex (Chang *et al.* 1991). Electron transfer has also been demonstrated from a copper ion bound within a ruthenium modified plastocyanin which is oxidized upon illumination of the ruthenium complex (Di Bilio *et al.* 1998). Utilizing a rhenium complex attached to *P. aeruginosa* azurin with a sacrificial electron acceptor in solution, oxidation of a bound copper ion has been demonstrated with subsequent

reduction of the oxidized rhenium complex (Connick *et al.* 1995; Di Bilio *et al.* 2001) reviewed in (Miller *et al.* 2004).

1.3. Combining Donor-Tyrosine-Pigment-Acceptor

Utilizing larger, natural enzymes is a good way to introduce multiple cofactors and produce a light activated electron transport chain. The key advantage here is that the proteins selected should already contain features that facilitate the redesign of the protein. An elegant example of this approach is the modification of the bacterial reaction centre (BRC) for PSII function (Kalman *et al.* 2005). The bacteriochlorophyll dimer (BChl2), P870, in the BRC from the purple bacterium *Rhodobacter sphaeroides* has a relatively low redox potential (~ 0.5 eV) compared to P680 in PSII (~ 1.25 eV). Lin *et al.* (1994) found that the reduction potential of the BChl2 in the reaction centre is influenced by hydrogen-bonding interactions with surrounding side chains. In wild-type P870, a hydrogen bond is present between a histidine and one of the acetyl groups involved in the conjugated π -system of the bacteriochlorophyll dimer. Upon removal of this hydrogen bond by mutating the histidine to a phenylalanine, the BChl2 potential drops from 505 to 410 mV. Conversely addition of hydrogen bonds to the conjugated π -system raises the E_m and the effects of these single-point mutations are additive. By using various single, double and triple mutants, the BChl2 potential could be tuned within the range of 410–765 mV (Lin *et al.* 1994). Subsequently a quadruple mutant was engineered in which the number of hydrogen bonds to the conjugated carbonyl groups of the dimer was maximized. The BChl2 reduction potential for this mutant was elevated to above 800 mV however, this mutation was relatively unstable (Kalman *et al.* 1999). Hydrogen bonding with histidine tends to have the greatest effect on the midpoint redox potential, but the E_m is also influenced by charge groups within the vicinity of the porphyrin head (Williams *et al.* 2001; Johnson and Parson 2002). Therefore, by selectively modifying the interactions with the protein scaffold, P870 could be tuned to a much higher redox potential approaching that for P680 in PSII.

In addition to mutating the four residues, Kalman *et al.* (1999) introduced a tyrosine at positions analogous to the positions of the redox active tyrosine's in PSII (Y_Z and Y_D), which following light excitation was oxidized and a tyrosine radical was observed

(Kalman *et al.* 1999). Measurements revealed that the tyrosyl radical was more prominent at higher pH values and modelling of the tyrosine mutations suggested that a H-bond was formed between the introduced tyrosine and a neighbouring histidine residue (Narvaez *et al.* 2002). By replacing the histidine with a non H-bonding residue, the E_m of the tyrosine increased and the amount of oxidized tyrosine decreased. Therefore the E_m and the extent of oxidation of the introduced tyrosine are dependent upon the availability of a proton acceptor group, characteristic of PSII (Diner and Britt 2005).

As well as the above modifications carboxylate residues were introduced by mutagenesis into the highly oxidizing reaction centre at a site homologous to the manganese binding site of PSII (Thielges *et al.* 2005). This created a monomeric metal binding site that could bind Mn with a low dissociation constant ($\sim 1\mu\text{M}$). Upon illumination the bound Mn was able to reduce P870^+ in a first order reaction. This same site was found to also bind Fe^{II} which could be oxidized to Fe^{III} , upon illumination of P870 (Kalman *et al.* 2006).

1.4. Research Objectives

The work carried out on the BRC is novel and of great interest however I have focused my research on utilizing a minimalist protein scaffold to engineer PSII activity. The large BRC enzyme is almost as complicated as PSII and thus difficult to utilize. Smaller proteins tend to be more robust than the much larger photosystem complexes though there are limits to the number of cofactors which can be bound. *De novo* designed proteins tend to be limited in that most are single 4 helix bundles that are too small to bind multiple cofactors. The use of existing protein scaffolds enables slightly larger proteins to be chosen which are often better characterized with higher resolution crystal structures. These proteins have an added advantage in that they can be produced in large amounts through bacterial expression systems.

The following chapters detail the work carried out in designing a minimalist PSII mimic from bacterioferritin. This robust protein is easy to handle and though only a four-helix bundle itself, dimerises enabling several redox active cofactors to be bound. Later

modification of the protein enabled the formation of a heterodimer which more closely resembles the cofactor arrangement within PSII. Light induced electron transfer was demonstrated with storage of the light energy as higher oxidation states within a di-manganese centre.

2.1. Cloning and Mutagenesis of Bacterioferritin Homodimer

The bacterioferritin gene from *Escherichia coli* was cloned into a plasmid and the protein expressed using the methods that follow. This protein was used to test the binding of the light active Zinc Chloride as in place of the heme group in the hydrophobic surface in binding pocket. The hfr protein was also used to assess manganese binding at the metal binding site. This protein with the two new cofactors bound was then used to look at light induced oxidation state changes in the manganese centre.

Genomic DNA was extracted from *Escherichia coli* strain 8631 DE3 (Novagen). A phenol-chloroform protocol was used to purify the genomic DNA with a sodium acetate and isopropanol precipitation step (Sax 2002). This procedure was repeated to ensure the purity of the DNA. The gene sequence for the *Escherichia coli* bacterioferritin gene (bfr) was taken from the published sequence of Andrews and co-workers (Andrews *et al* 1989). The bfr gene encoding bacterioferritin was amplified from the genomic DNA using the following PCR primers:

Forward - 5' ATGAAGGCTGATACTAAAGTATATAAATTATCTC 3'

Reverse - 3' TCAACCTTCTTCGCGGAT 3'

The PCR method utilized involved a hot start followed by 30 cycles of 95°C for 1 minute, 55°C for 1 minute and 72°C for 3 minutes. The PCR product was cleaned up using a chloroform:isoamyl alcohol extraction then gel-purified on a 1.3 % agarose gel. The band corresponding to the desired amplification product was cut out of the gel. The desired band was further purified using a QIAquick Spin Gel Extraction kit (Qiagen).

medication in the treatment of the disease. The aim of the study was to evaluate the effectiveness of the treatment in the long term. The study was conducted in a randomized, controlled, double-blind manner. The participants were divided into two groups: the treatment group and the control group. The treatment group received the medication, while the control group received a placebo. The study was conducted over a period of 12 weeks. The primary outcome was the change in the level of the disease. The secondary outcome was the change in the quality of life. The results of the study showed that the treatment was effective in reducing the level of the disease and improving the quality of life. The study was conducted in a rigorous manner and the results are reliable.

The study was conducted in a randomized, controlled, double-blind manner. The participants were divided into two groups: the treatment group and the control group. The treatment group received the medication, while the control group received a placebo. The study was conducted over a period of 12 weeks. The primary outcome was the change in the level of the disease. The secondary outcome was the change in the quality of life. The results of the study showed that the treatment was effective in reducing the level of the disease and improving the quality of life. The study was conducted in a rigorous manner and the results are reliable.

1.4. Research Objective

The study was conducted in a randomized, controlled, double-blind manner. The participants were divided into two groups: the treatment group and the control group. The treatment group received the medication, while the control group received a placebo. The study was conducted over a period of 12 weeks. The primary outcome was the change in the level of the disease. The secondary outcome was the change in the quality of life. The results of the study showed that the treatment was effective in reducing the level of the disease and improving the quality of life. The study was conducted in a rigorous manner and the results are reliable.

The study was conducted in a randomized, controlled, double-blind manner. The participants were divided into two groups: the treatment group and the control group. The treatment group received the medication, while the control group received a placebo. The study was conducted over a period of 12 weeks. The primary outcome was the change in the level of the disease. The secondary outcome was the change in the quality of life. The results of the study showed that the treatment was effective in reducing the level of the disease and improving the quality of life. The study was conducted in a rigorous manner and the results are reliable.

CHAPTER 2

2 Methods

2.1. Cloning and Mutagenesis of Bacterioferritin Homodimer

The bacterioferritin gene from *E.coli* was cloned into a plasmid and the protein expressed using the methods that follow. This protein was used to test the binding of the light active Zinc Chlorin e_6 in place of the heme group in the hydrophobic porphyrin binding pocket. The Bfr protein was also used to assess manganese binding at the metal binding site. This protein with the two new cofactors bound was then used to look at light activated oxidation state changes in the manganese centre.

Genomic DNA was extracted from *E.coli* strain BL21 DE3 (Novagen). A phenol-chloroform protocol was used to purify the genomic DNA with a sodium acetate and isopropanol precipitation step (Rex 2000). This procedure was repeated to ensure the purity of the DNA. The gene sequence for the *E.coli* bacterioferritin gene (*bfr*) was taken from the published sequence of Andrews and co-workers (Andrews *et al.* 1989). The *bfr* gene encoding bacterioferritin was amplified from the genomic DNA, using the following PCR primers:

Forward - 5' ATGAAAGGTGATACTAAAGTTATAAATTATCTC 3'

Reverse - 5' TCAACCTTCTTCGCGGAT 3'

The PCR method utilized involved a hot start followed by 30 cycles of 95°C for 1 minute, 55° C for 1 minute and 72° C for 3 minutes. The PCR product was cleaned up using a chloroform/isoamyl alcohol extraction then gel-purified on a 1.5 % agarose gel. The band corresponding to the desired amplification product was cut out of the gel. The excised band was further purified using a QIAquick Spin Gel Extraction kit (Qiagen).

Ligation independent cloning (LIC) regions were added to each end of the amplified *bfr* gene product by another round of PCR using the following primers:

Forward - 5' GGTATTGAGGGTCGCATGAAAGGTGATACTAAAGTTATAA 3'

Reverse - 5' AGAGGAGAGTTAGAGCCTCATCAACCTTCTTCGCGGAT 3'

LIC regions in each primer (underlined) facilitated cloning into the pET32 Xa/LIC expression vector (Novagen) (Figure 2.1). The same PCR method used to amplify the *bfr* gene was also used to add the LIC ends to the *bfr* gene. The new amplified DNA with the LIC ends was T4 treated to remove overhanging nucleotides. The *bfr* gene was then ligated into pET-32 Xa/LIC (Novagen) (Figure 2.2) as per standard Novagen protocols for the LIC system.

Figure 2.1: Ligation independent cloning strategy. LIC sequences are added onto the gene by PCR amplification with specific primers. Then overhangs are created from the LIC sequences by treatment with T4 DNA polymerase. LIC sequences in the linearised pET32 Xa/LIC plasmid. The overhangs on the gene and the plasmid are complementary allowing the gene to anneal and create a circularized plasmid without the need for ligase (figure adapted from Novagen).

The annealed pET32 Xa/LIC vector with *bfr* gene was then transformed into NovaBlue cells (Novagen) using a standard heat shock protocol (42°C for 30 seconds). Transformed cells were grown in liquid culture then plated onto LB plates containing 50 µg/ml ampicillin to select for the plasmid.

Figure 2.2: The pET32 Xa/LIC vector contains a thioredoxin gene (*trxA*) which is expressed as a fusion with the gene of interest. The plasmid also contains an ampicillin resistance gene for selection. Figure adapted from Novagen technical manual.

The plasmid was subsequently purified, from liquid cultures of individual colonies of transformed NovaBlue cells, using a QIAprep spin miniprep kit. The purified plasmid was then transformed into BL21 DE3 pLysS cells (Novagen) again using a heat shock protocol. To confirm the DNA sequence of the pET32 Xa/LIC *bfr* gene, the plasmid was sent to the Australian Genome Research Facility (AGRF) for sequencing. The transformants were then screened for expression of the correct molecular weight protein product using SDS-PAGE.

BFR ¹ Amino Acid and DNA Sequence	
1	M K G D T K V I N Y L N K L L G N E L V
1	ATGAAAGGTGATACTAAAGTTATAAATTATCTCAACAAACTGTTGGGAAATGAGCTTGTC
21	A I N Q Y F L H A R M F K N W G L K R L
61	GCAATCAATCAGTACTTTCTCCATGCCCGAATGTTTAAAAACTGGGGTCTCAAACGTCTC
41	N D V E Y R E S I D E M K H A D R Y I E
121	AATGATGTGGAGTATCGCGAATCCATTGATGAGATGAAACACGCCGATCGTTATATTGAG
61	R I L F L E G L P N L Q D L G K L N I G
181	CGCATTCTTTTTCTGGAAGGTCTTCCAAACTTACAGGACCTGGGCAAACCTGAACATTGGT
81	E D V E E M L R S D L A L E L D G A K N
241	GAAGATGTTGAGGAAATGCTGCGTTCTGATCTGGCACTTGAGCTGGATGGCGCGAAGAAT
101	L R E A I G Y A D S V R D Y V S R D M M
301	TTGCGTGAGGCAATTGGTTATGCCGATAGCGTTCGTGATTACGTCAGCCGCGATATGATG
121	I E I L R D E E G H I D W L E T E L D L
361	ATAGAAATTTTGCCTGATGAAGAAGGCCATATCGACTGGCTGGAAACGGAACCTTGATCTG
141	I Q K M G L Q N Y L Q A Q I R E E G
421	ATTCAGAAGATGGGCCTGCAAAATTATCTGCAAGCACAGATCCGCGAAGAAGGT

Figure 2.3: The amino acid and DNA sequence for the double mutant, BFR¹. The two mutations H46R and H112R are marked in red.

2.1.1. Site-Directed Mutagenesis of Bacterioferritin

Two external histidines (H46, H112) were changed to arginines (Figure 2.3) using the Stratagene Quickchange multi site-directed mutagenesis kit. The removal of these histidines was to prevent non-specific binding of porphyrins to the protein. This double mutant (BFR¹) was utilized for all measurements in Chapter 3. All further mutants originated from the BFR¹ gene and contain the H46R and H112R mutations. The primers utilized for this process are documented below, the mutated bases are underlined.

H46R 5' AATGATGTGGAGTATCGCGAATCCATTGATGAG 3'

H112R 5' GCCGATAGCGTTCGTGATTACGTCAGCC 3'

Two other mutants were made which contained the H46R and H112R mutations. These mutants were E51C and M52H. The E51C mutant removed a glutamic acid which is of critical importance in the ferroxidase site and replaced it with a cysteine. This cysteine was then used to covalently link a quinone within the now defunct metal binding site. This mutant labelled BFR² has a cavity which whilst normally reserved for metal binding allowed a quinone to bind in place. The primer utilized for this process is documented below, the mutated bases are underlined.

E51C 5' CGAATCCATTGATTGTTATGAAACACGC

The M52H mutant was produced to look at the effect of a different axial ligand for the ligation of the ZnCe₆ group. The literature suggested this mutation would prevent binding of heme to the site and this was observed. The primers utilized for this process are documented below, the mutated bases are underlined.

M52H 5' GAATCCATTGATGAGCATAAAACACGCCGATCGT 3'

2.2. Production of the BFR Heterodimer Construct

A heterodimer of the Bfr protein was engineered to enable directional electron transfer from a Mn centre to a quinone. In order to be able to change one half of the BFR dimer relative to the other half, i.e. form a heterodimer, it was necessary to link two monomers via a glycine rich linker region. This flexible linker was inserted into the gene sequence joining the N-terminus of one molecule to the C-terminus of the other. This utilized a modified form of the linker reported by (Whitney and Sharwood 2006). Two forms of the heterodimer were made, one as a control with no quinone binding site to test for structure and function retention upon adding the linker region, whilst the other contained a metal binding site, a porphyrin binding pocket and quinone binding site.

To produce the different constructs two *bfr* mutant genes were utilized BFR¹ (H46R, H112R) and BFR² (H46R, H112R, E51C). To enable the genes to be cut out of and ligated into plasmids it was necessary to add restriction sites at either end of each gene.

The following primers were used to PCR amplify the BFR¹ and BFR² genes in order to add the required restriction sites:

- 5'NdeI 5' TCATATGAAAGGTGATACTAAAGTT 3'
- 5'SacII 5' TCCGCGGTGGAGATATGAAAGGTGATACTAAAG 3'
- 3'EcoRV 3' TGATATCGACCTTCTTCGCGGATCTGTGC 5'
- 3'HindIII 3' AAGCTTAACCTTCTTCGCGGATCT 5'

The PCR rounds produced four different gene constructs (Figure 2.4) with either SacII and EcorRV restriction sites (SE) at either end of the gene or NdeI and HindIII cleavage sites (NH) at either end of the gene. These blunt ended gene products were gel purified then treated with Taq polymerase to A-tail them. This process adds a single deoxyadenosine to the 3'-ends of the DNA which enables it to be readily ligated into a linearised vector containing single 3'-T overhangs at the insertion site. Using this method, only one insert will be ligated into the vector as opposed to multiple insertions that can occur with blunt-ended cloning.

Figure 2.4: Four gene constructs were made with different restriction sites to enable the genes to be cut and ligated in different combinations. SE represents the cleavage sites SacII and EcoRV at either end of the gene. The NH represents the cleavage sites NdeI and HindIII at either end of the gene.

BFR¹ - H46R, H112R
BFR² - H46R, H112R, E51C

The amplified A-tailed gene products BFR¹-SE (Figure 2.4: yellow construct), BFR¹-NH, and BFR²-NH (Figure 2.4 green and blue constructs) were all ligated into separate pGEM-T Easy vectors (pTE) (Figure 2.5). The high-copy-number pTE vector contains T7 and SP6 RNA polymerase promoters flanking a multiple cloning region within the α -peptide coding region of the enzyme β -galactosidase. Insertional inactivation of the α -peptide allows recombinant clones to be directly identified by colour screening on indicator plates.

Figure 2.5: pGEM-T easy vector map and sequence reference points displaying the large number of cleavage sites available for gene construct insertion. Figure adapted from Promega technical manual.

The three ligated plasmids pTE-BFR¹-SE, pTE-BFR¹-NH, and pTE-BFR²-NH were transformed into XL1B *E.coli* cells. The cells were plated onto LB-agar plates containing ampicillin, X-gal (5-bromo-4-chloro-3-indolyl- β -D-galactoside) and IPTG. Colonies of self ligated plasmids were blue in colour whilst those containing the gene of interest were white. Single white colonies were selected and grown in overnight cultures before the plasmids were purified using standard plasmid miniprep procedures to isolate the recombinant plasmids.

A DNA cleavage reaction was carried out on the pTE-BFR¹-SE plasmid with EcoRV and SalI to ensure that the gene was inserted in the correct orientation. If it was oriented correctly then a 3.5 kb fragment was found if not then a 0.5 kb fragment was produced. Only those clones which had the 3.5 kb fragment were selected.

A 20 amino acid linker was used to join the two *bfr* genes to make the final heterodimer construct. This linker region contained a gly-gly-ser repeat with a single lysine (Figure 2.6). The linker, which was a gift from Spencer Whitney, was supplied in a pHUE plasmid and cleaved out using *EcoRV* and *NdeI* restriction enzymes. The 61 base fragment was then gel purified ready for ligation. The pTE-BFR¹-SE plasmid was then digested with *EcoRV* and *NdeI* and the linker gene was ligated into the plasmid producing the pTE-BFR¹-SE-L construct (Figure 2.7).

Figure 2.6: The amino acid sequence and DNA sequence of the linker used to join the Bfr subunits. The cleavage sites are underlined and the nucleotides shown in red reveal the cleaved bases.

The pTE-BFR¹-SE-L gene was cleaved from the pTE vector using *SacII* and *NdeI* followed by standard gel purification. With the gene extracted the final stage in building the heterodimer constructs was to digest the pTE-BFR¹-NH and pTE-BFR²-NH plasmids with *SacII* and *NdeI*. The plasmids were gel purified before the BFR¹-SE-L gene was ligated into each of these plasmids. This produced the final Bfr heterodimer constructs which were then sequenced to verify the result (Figure 2.8).

Figure 2.7: The *bfr* gene construct BFR¹-SE-L made with a *bfr* gene and a linker region.

Figure 2.8: The two different heterodimer constructs assembled in the pTE vector.

Once the two heterodimer constructs were made it was necessary to carry out two site directed mutations per construct. The linker region in all of the constructs encodes for a single histidine at each end of the linker (Figure 2.6). Due to the need to remove all solvent exposed histidines these were mutated to serine's using the Stratagene multi site-directed mutagenesis kit which enabled changing both residues in the one reaction. The following primers were used to carry out the mutations:

Delta EcoRV – 3' CCGCCGCTGCCACCAGAATATCGACCTTCTTC 5'

Delta NdeI – 3' AGTATCACCTTTCATAGAACCACCAGAACCGCC 5'

The mutation of both histidines to serines in the linker, removed the EcoRV and the NdeI cleavage sites (Figure 2.9) which made for easy verification of the successful mutants.

R Y S G G S G G S G G K S G G S G G S G G S
CGATATCTGGTGGCAGCGCGGTAGCGGAGGCAAGTCCGGTGGTAGCGGCGGTTCTGGTGGTTCT

Figure 2.9: Amino acid sequence and DNA sequence of the linker region after mutation of both histidines to serines. Both the EcoRV and the NdeI cleavage sites have been removed in the process. The mutated residues are shown in red.

The final constructs contained only SacII and HindIII cleavage sites at the 5' and 3' ends respectively (Figure 2.10). Both constructs were cleaved from the pTE vector using SacII and HindIII, these genes were then gel purified in readiness for insertion into an expression vector.

Figure 2.10: The final two heterodimer gene constructs after site-directed mutagenesis to remove the EcoRV and NdeI restriction sites.

The expression vector chosen was pHUE which was supplied as a gift by Spencer Whitney (Figure 2.11). The pHUE vector encodes for a selectable ampicillin resistance gene and has multiple restrictions sites. The construct is designed to express the gene of choice as a fusion protein with ubiquitin and a six histidine tag for ease of purification. The pHUE vector was cleaved with SacII and HindIII, gel purified and then the two heterodimer gene constructs were ligated into separate vectors. The pHUE vectors were then transformed into *E.coli* BL21 cells ready for protein expression.

Figure 2.11: The Histidine-tagged Ubiquitin Expression vector, pHUE. (A) Plasmid map of pHUE showing the ubiquitin coding region (black box), restriction enzyme recognition sites within the multiple cloning site (MCS) and the selectable ampicillin resistance gene. Locations are given relative to the start codon upstream of the His-tag. (B) DNA and encoded protein sequence of the ubiquitin coding region showing the engineered SacII site (underlined). Figure adapted from (Catanzariti *et al.* 2004)

2.3. Expression and Purification of All Constructs

2.3.1. Expression

E. coli cells containing the pET32 or pHUE plasmid were grown in an O/N culture then inoculated into LB media containing 50 µg/ml ampicillin. Cultures were grown at 37°C with air sparging to stir the culture and provide oxygen. The cultures were grown to an optical density of ~0.5 at 600 nm then induced with a final concentration of 1mM Isopropyl-thiol-galactose (IPTG) (Sigma). Induced cultures were grown for a further 5 hours before harvesting the cells by centrifugation. Harvested cells were pink in colour due to the high expression level of the heme containing protein (note: the M52H mutant produced normal coloured cells as the protein did not bind heme).

The pelleted cells were resuspended in buffer A (50 mM Tris-HCl pH 8.0, 500mM sodium chloride) and lysozyme was added to a final concentration of 1 mg/ml. DNase1

(Boehringer Mannheim) was then added to a final concentration of 100 µg/ml and MgSO₄ to 100 mM. After stirring the solution was sonicated then centrifuged to remove precipitate and finally filtered through a 0.2 µm syringe filter.

2.3.2. Protein Purification

The pET32 Xa/LIC vector expresses the Bfr protein as a fusion protein with thioredoxin and has a six histidine tag (His tag) in the linker region between these two proteins. The pHUE vector expresses a fusion protein with a His tagged ubiquitin directly linked to the protein of interest (Catanzariti *et al.* 2004). The Bfr fusion proteins were purified using a FastFlow HP Nickel sepharose column on an Akta Prime FPLC system (GE Healthcare). The column was equilibrated with buffer A and the clarified cell supernatant was loaded onto the column. The column was then washed with 95% buffer A, 5% Buffer B (50 mM Tris-HCl pH 8.0, 500mM imidazole). Finally the Bfr proteins were eluted using 100% Buffer B. The purified Bfr fusion proteins eluted as a dark blood red solution due to the bound heme. The purified proteins were then dialysed using 5 kDa cut-off dialysis tubing (Pierce) against tricine buffered saline (TBS) (25mM tricine, 100 mM KCl, pH 7.7) to remove all imidazole.

The purified fusion proteins were cleaved to remove all non-native amino acids using factor Xa (FXa) for the pET32 expressed proteins or with ubiquitin specific protease for the pHUE expressed proteins. The cleaved proteins were run on SDS-PAGE gels to determine completion of the cleavage reaction. Cleaved thioredoxin and ubiquitin were removed from the Bfr proteins by again running on a nickel sepharose column where they bound via the His tag and the protein of interest was collected in the flow through. In general the protein was not very concentrated so the volume was reduced by placing the dialysis tubing in high molecular weight polyethylene glycol (>10,000 MW) crystals to draw water from within the dialysis tubing. The proteins were considered to be pure at this stage as they existed as single bands on overloaded SDS-PAGE gels stained with coomassie brilliant blue.

2.4. Cofactor Binding

In order to utilize the Bfr proteins as a mimic of the PSII motif it was first necessary to remove all bound cofactors. Bound heme was removed from within the hydrophobic pocket of the Bfr dimers using the method of Teale et al (1959); where addition of methyl ethyl ketone (MEK) to a chilled acidified cytochrome solution produces a phase separation of the heme into the organic layer whilst the apo-protein remains in the water phase (Teale 1959). Non-heme iron was removed using the modified method of Bauminger et al with exhaustive dialysis in the presence of dithiothreitol and EDTA (Bauminger *et al.* 1991). Protein concentrations for the apo-proteins were determined using $\epsilon_{280\text{nm}}$ of $20340 \text{ M}^{-1} \text{ cm}^{-1}$ for BFR¹ and $41960 \text{ M}^{-1} \text{ cm}^{-1}$ for all BFR heterodimers.

2.4.1. CD Spectroscopy

CD spectroscopy was used to confirm the secondary structure of all proteins after cofactor removal. CD spectra were recorded on a Jobin Yvon type III+ spectrometer (Horiba, Japan). By comparison of the spectra with that of a standard set of proteins with known structures it is possible to deconvolute the spectra and assess secondary structure.

2.4.2. Isothermal Titration Calorimetry

Cofactor binding was monitored using isothermal titration calorimetry (ITC) which allows the accurate determination of binding constants (K), reaction stoichiometries (n), enthalpies (ΔH°), and entropies (ΔS°). Metal binding as well as porphyrin binding were all monitored with this technique. All measurements were carried out on a Microcal calorimeter at 25°C and the data was fit to a least squares model with the Microcal Origin fitting software. The heat of dilution was subtracted from the raw data prior to analysis.

All samples were prepared in 25 mM tricine, 100 mM KCl pH 7.7. Manganese binding to BFR was measured by injecting 11 μ l aliquots of 900 μ M MnCl_2 into a 30 μ M solution of BFR. BFR binding to ZnCe_6 was measured by injecting 11 μ l aliquots of 300 μ M BFR into a 15 μ M solution of ZnCe_6 .

The light active porphyrin ZnCe_6 was prepared as a 2 mM stock by the addition of zinc acetate to free base chlorin e_6 dissolved in methanol, added in a 1:1 molar ratio with Zinc binding to the chlorin resulting in a UV-visible red shift in the Soret peak from 400 to 411 nm and a blue shift in the Q_y band from 661 to 632 nm (Razeghifard and Wydrzynski 2003). UV-Vis spectroscopy was carried out on a Cary 300 Spectrometer (Varian, USA). Solutions of MnCl_2 for binding to Bfr were prepared fresh in TBS. Solutions of FeSO_4 for titrations and iron oxidation assays, were prepared fresh in anaerobic water pH 4 to prevent oxidation by oxygen.

2.4.3. Covalent Attachment of Quinone's

Quinones were covalently attached to cysteine residues in the mutant Bfr proteins using a modified method of Hay et al 2007. First the apo-protein was reduced with a 5 fold excess of DTT in 3 M Gdn:HCl, 50mM potassium phosphate (KPi) pH 8.0 for 3 hours at RT. The Gdn:HCl and DTT were removed by gel filtration with a PD-10 column (GE Healthcare) and exchanged into 50 mM KPi, pH 7.0. The protein concentration was then determined using Ellman's assay under denaturing conditions (6 M Gdn:HCl) and using an $\epsilon_{412\text{nm}}$ of $13700 \text{ M}^{-1} \text{ cm}^{-1}$. A 5-fold excess of quinone was reacted with the reduced protein in 50 mM KPi, pH 7.0 in the dark at room temperature for 2 hours. The protein-quinone solution was then dialysed in the dark at 4°C against a 50 mM KPi pH 7.0 buffer containing quinone of interest (DMBQ E_0 133 mV or naphthoquinone E_0 137 mV) and ferricyanide (E_0 of 436 mV at pH 7.0) at a 1:1 and 1:5 protein to oxidant mole ratio, respectively. The dialysis step should maximize the concentration of oxidized quinone-protein. After dialyzing overnight the dialysis buffer was changed to TBS. The quinone bound protein was then concentrated using PEG dehydration before aliquoting and freezing until further use.

2.5. Analysis

2.5.1. Oxygen Measurements

All oxygen measurements were carried out at 25°C on an oxytherm Clarke-type oxygen electrode (Hansatech, UK) modified to contain a side illumination port. Oxygen measurements were carried out on samples containing 30 μM BFR, 15 μM ZnCe₆, 60 μM or 300 μM MnCl₂ or 300 μM FeSO₄ in 50 mM MES pH 6.5. Samples were illuminated for long periods of time until oxygen uptake plateaued.

2.5.2. Fluorescence Measurements

All fluorescence measurements were made using a Fluoromax-3 (Jobin Yvon HORBIA) spectrofluorimeter. The fluorescence of ZnCe₆ when bound to the various Bfr proteins was measured with a slit widths of 1 nm, 1 second sampling time and a concentration of 2.5 μM in 50 mM MES buffer pH 6.5. Samples were excited at 427 nm after excitation and emission scans revealed that this excitation wavelength produced the maximal fluorescence at 660nm (Q_y band). Fluorescence was recorded from 575 nm to 775 nm. These concentrations were in the linear range of the detector.

2.5.3. Redox Measurements

Cyclic voltammetry of BFR¹ with cofactors bound was carried out on pyrolytic edge graphite electrodes at pH 7.75 in 25 mM Tricine, 100 mM KCl. A scan rate of 20 mVs⁻¹ was used to prevent hysteresis. After each scan the electrode was cleaned by scouring the surface then reapplying the protein solution and allowing to almost dry before carrying out the measurements.

2.5.4. EPR Measurements

Low Temperature EPR

To examine light induced changes in oxidation state of the metal centre in BFR¹ complexes, samples were prepared in the dark at room temperature then either left for one hour in the dark or illuminated for 1 hour. The long illumination time used coincided with the time taken for the oxygen uptake reactions to go to completion. Due to the concentrated nature of the samples they were stirred under pure oxygen during illumination to prevent the samples becoming anaerobic. Samples were then degassed and frozen to liquid nitrogen and stored until use. All samples were measured within 24-48 hours. Samples were prepared in 50mM MES pH6.5 or in 25mM tricine, 100mM KCl pH 7.7 with 400mM sucrose added as cryoprotectant. Low temperature (5K) X-band EPR spectra were acquired on a Bruker ESP 300E spectrometer using either a TM011 cavity or a ER4116 dual mode cavity, perpendicular mode microwave frequency 9.6 GHz, parallel mode microwave frequency 9.3 GHz, modulation frequency 100kHz. The EPR spectrum of each sample was collected in both parallel and perpendicular mode detection before and after photo-accumulation at room temperature or 200K.

Room Temperature EPR

X-band EPR spectra were acquired at room temperature on a Bruker ESP300E spectrometer equipped with a TM011 cavity. Generally the spectra were acquired at varying microwave power with 1-10 G modulation amplitude and 100 KHz modulation frequency. When necessary, the samples were excited with white light at room temperature via a non-magnetic optical fibre which allowed illumination within the sample cavity. The sample was placed in a flat cell which allowed better signal-to-noise and light-saturation. There was negligible UV-excitation in this set up.

2.5.5. EPR Simulations

Spectral simulations were solved numerically from a Hamiltonian (36 x 36 matrix) using Scilab-4.4.1, an open source vector-based linear algebra package (www.scilab.org). A least squares minimization routine was employed to find the

optimal solutions for the parameters. A complete description of the EPR simulations can be found in chapter 3.

2.6. Molecular Modelling

Molecular modelling of mutations was carried out with the aid of HyperCHEM (v7.5 Hypercube, USA). Molecular mechanics optimization of the final structure was carried out using a Polak-Rebiere energy minimisation algorithm (conjugate gradient) in a periodic box. The heterodimer images were produced by including a 22 amino acid linker to the dimer structure and carry out molecular mechanics optimization of the linker sequence whilst bound to the protein structure. The electronic coupling and PATHWAYS calculations in porphyrin-protein-quinone complexes were carried out using HARLEM (HAmiltonians to Research LargeE Molecules). Ray-traced images were prepared using PyMOL v0.99 (Delano Scientific, USA).

CHAPTER 3

3 *Bacterioferritin as a 'Reaction Centre'.*

3.1. Introduction

The essential catalytic elements of PSII that enable the oxidation of water to molecular O_2 are redox coupled cofactors that include: (i) a strongly oxidizing chlorophyll in the reaction centre termed P680; (ii) a redox-active tyrosine termed Y_Z ; and (iii) the oxygen evolving complex (OEC) consisting of four μ -oxo bridged manganese ions and one calcium (Mn_4O_xCa) (Wydrzynski and Satoh 2005). The P680 provides the driving force (ΔG) for the reaction, the tyrosine is important for proton coupled electron transfer, and the OEC templates the O-O bond formation. The one electron photochemistry of charge separation in the photosystem is linked to the four electron chemistry of water oxidation via the sequential oxidation of the Mn_4O_xCa (Wydrzynski and Satoh 2005).

The complexity of the PSII enzyme has thus far limited attempts to reverse engineer a synthetic water oxidase (Wydrzynski *et al.* 2007). Previous engineering efforts have concentrated on constructing biological motifs that mimic one aspect of the enzyme's mechanism, in particular porphyrin excitation/oxidation (Hay *et al.* 2004) and metal binding/complex formation (Thielges *et al.* 2005). Attempts have also been made to design 4-helix bundle proteins through de novo design (Koder and Dutton 2006). These approaches all faced limitations be it an overly complex system which is difficult to work with or limited size to introduce cofactors. Thus, in this work I considered a new approach to meet key design requirements in the production of an artificial photoactive protein with more than one functional element. A naturally occurring protein scaffold was used to bind several introduced redox active cofactors (Conlan 2008). In commencing this task the protein structure database was searched to find naturally occurring proteins which contain existing useful architecture so that cofactors could be placed within appropriate distances of each other. The design of a light activated redox enzyme had to take into consideration the physiological limitations of intra-protein electron transfer and as such all cofactors were placed within $<14 \text{ \AA}$ of each other (Conlan 2008; Noy *et al.* 2006; Page *et al.* 1999; Page *et al.* 2003).

In order to more closely mimic the reactions of PSII I have modified the bacterioferritin (BFR) protein to bind several redox active cofactors and initiate light-driven electron transfer. BFR is a soluble 4-helix bundle protein (~18.5 kDa) which naturally contains a di-iron metal site and a heme *b* binding site (Andrews et al. 1995; Andrews et al. 1993; Davis et al. 1999). Removal of the native cofactors and replacement with manganese in the metal centre and a light active chlorophyll analogue –zinc chlorine within the heme binding site produce a protein with light dependent catalytic properties.

The natural synthesis/expression of the BFR protein results in a self-assembled quaternary structure that is somewhat analogous to a viral particle. The subunits oligomerise to produce a large spherical-shaped shell, made up of 12 homodimeric units, that enclose a hollow cavity with a diameter of ~8 nm (Figure 3.1) (Moore 1991; Frolova *et al.* 1994; Harrison and Arosio 1996). The cavity normally stores iron as ferric oxide, hydroxide or phosphate minerals and is connected to the outer protein surface by channels. The natural protein shell thus binds 12 heme groups and 48 iron atoms (Frolova *et al.* 1994).

Figure 3.1: A ribbon diagram of the sphere structure formed by *E.coli* bacterioferritin (PDB 1BFR). The sphere is composed of twelve homodimers.

The protein forms into a homodimer, binding a single heme group symmetrically at the interface between the two protein monomers (Frolow *et al.* 1994; LeBrun *et al.* 1995a). An atypical bis-axial ligation of the heme occurs through the sulphur atoms at equivalent methionyl positions (Met52) in each subunit (Cheesman *et al.* 1990; Frolow *et al.* 1994). The function of the heme is unknown but it appears to serve more than a structural role. The heme group can be removed (Teale 1959) and replaced with other similar structured but light active porphyrins (Conlan *et al.* 2007).

Figure 3.2: The BFR homodimer with two identical subunits each hosting dinuclear metal centres (purple spheres) and a heme (green) which is bound in a hydrophobic pocket at the subunit interface. Two manganese atoms bind per protein subunit. Adapted from the crystal structure (PDB file 1BCF).

Bacterioferritin contains a binuclear metal binding site that links the four helices of the protein together and acts as a ferroxidase site, in which Fe^{II} is oxidized to Fe^{III} by O_2 (LeBrun *et al.* 1995b). The two Fe ions are ligated to four glutamate residues (Glu18, Glu51, Glu94, Glu127, one from each helix) and two histidine residues (His54, His130) (Dautant *et al.* 1998). Residues Glu51 and Glu127 form bridging carboxylate ligands between the two metal sites (di- μ -1,3-carboxylato bridges), while Glu18 and Glu94 form monodentate carboxylate ligands to separate metal sites as do His54 and His139 via N^δ nitrogen ligation. The organization of the di-iron binding site is very similar to other

class II di-iron proteins. The crystal structure for BFR was initially produced with two Mn^{II} ions bound in the metal binding site (Dautant *et al.* 1998; Frolow *et al.* 1994) (Figure 3.2).

BFR presents an appealing starting point for engineering a multi-step, light-activated protein as: (i) it is a highly stable and robust protein (ii) the heme can be extracted and reconstituted (Teale 1959) or replaced with a photoactive chlorin (Hay *et al.* 2004); (iii) the binuclear metal site has similar structural ligation to the manganese catalase where the di-manganese site can reach an oxidation state of $\text{Mn}_2^{\text{III,IV}}$ (Yoder *et al.* 2000; Barynin *et al.* 2001; Schafer *et al.* 2003; Teutloff *et al.* 2005); and (iv) the cofactors when bound to the BFR protein are separated by similar distances to that found in PSII (Figure 3.3) (Loll *et al.* 2005).

Figure 3.3: Schematic showing the highly similar crystal structure derived distances in Å, between cofactors in the engineered BFR complex as compared to PSII.

Cofactor separation in the BFR protein matrix was an important design constraint as it is necessary for all cofactors to be situated within the protein with an edge-to-edge distance approaching 10 Å (Figure 3.3). The tyrosine/Mn distance is notably shorter than the ZnCe_6 /tyrosine distance and this is also found in PSII. In PSII the tyrosine acts as an intermediary for the stepwise oxidation of the Mn centre by P680 (Tommos and Babcock 1998; Haumann and Junge 1999; Ioannidis *et al.* 2008). This is an important consideration in the design of a photo-catalyst with a metal centre as the donor group.

3.1.1. Summary

Here I report the construction of a light-activated metallo-catalyst, which can oxidize Mn^{II} to Mn^{III} upon illumination with a tyrosine radical formed during the process. The light induced oxidation state changes were characterized by low temperature perpendicular and parallel mode X-band EPR spectroscopy and oxygen uptake measurements.

3.1.2. Techniques Used to Analyse the BFR Complex.

Manganese ions typically have small optical extinction coefficients in the UV-Vis electronic spectra and when bound to a moderate sized protein then changes in its ligation and oxidation state are extremely difficult to monitor. This has lead to the use of a number of alternate techniques which enable monitoring of manganese including electron paramagnetic resonance (EPR), isothermal titration calorimetry (ITC), cyclic voltammetry (CV) and circular dichroism (CD) spectroscopy. These techniques are outlined in the following section.

3.1.3. Electron Paramagnetic Resonance.

Manganese CW EPR Properties

EPR spectroscopy is a magnetic resonance technique which detects the transition between magnetic energy states for unpaired electrons in an applied magnetic field. It is a technique which is particularly useful for the detection of transition metal ions with unpaired d electrons. Mn^{II} has 5 unpaired d electrons in the valence shell (d^5 ion) which give rise to long lived electron spin states and sharp EPR signals. Mn^{II} has six possible orientations of electron spin ($M_s = \pm 5/2, \pm 3/2, \pm 1/2$) in an external field yielding five fully allowed ($\Delta M_s = \pm 1$) EPR fine structure transitions. The $S = 5/2$ spin property of Mn^{2+} is further split by the $I = 5/2$ nuclear spin of Mn^{55} . The hyperfine interaction between the electron spin and nuclear spin splits the energy levels into six sublevels corresponding to the six possible orientations of the ^{55}Mn nuclear spin. This sextet splitting is the EPR signature of mononuclear Mn sites. The combination of the electron

spin of $5/2$ and the nuclear spin of $5/2$ leads to 36 energy levels and 30 allowed ($\Delta M_s = \pm 1$, $\Delta m_I = 0$) transitions. Rarely are all of the allowed transitions realized. In macromolecular complexes in solution the strong orientation dependence of all but the central ($M_s -1/2 \rightarrow 1/2$) fine structure transition results in this transition being the dominant signal (Reed and Poyner 2000).

Dinuclear manganese centres are further complicated by the coupling between the unpaired electrons and the manganese nuclei. The presence of exchange interactions in spectra of dimanganese centres gives rise to multiline patterns for the ^{55}Mn hyperfine splitting. In most cases, the electron spin-spin interaction is mediated through bridging ligands and is therefore a form of super-exchange. The exchange is typically antiferromagnetic in such cases (Reed and Poyner 2000).

Conventional perpendicular mode EPR is well suited for the observation of half-integer spin systems (for example Mn^{II} , Mn^{IV} , coupled mixed-valent Mn clusters, and organic radicals), whereas parallel polarization EPR¹ has been used for observing EPR signals issued from integer spin systems, in particular $S=2$ (for example Mn^{III} , $\text{Mn}_2^{\text{III,III}}$) (Hendrich and Debrunner 1988, 1989). Improved signals can be obtained for Mn^{III} complexes when the microwave magnetic field is applied parallel to the applied magnetic field, in contrast to the conventional perpendicular polarized EPR method (Dexheimer *et al.* 1989). EPR studies of Mn^{III} in manganese superoxide dismutase (Campbell *et al.* 1999), di-manganese catalase (Whittaker *et al.* 2003), Photosystem II (Campbell *et al.* 2000), and in a Mn^{III} salen compound (Campbell *et al.* 2001) have demonstrated that parallel polarization mode EPR can provide useful information on Mn^{III} metal centres.

¹ In parallel mode the probing microwave magnetic field is rotated from the standard configuration perpendicular to the main magnetic field to a parallel orientation.

3.1.4. Circular Dichroism Spectroscopy

Circular dichroism is a measurement of the difference in the absorption of left (A_L) and right (A_R) circularly polarised light as a function of wavelength:

$$\Delta A = A_L - A_R$$

A chiral molecule will absorb the two oppositely rotating light rays unequally. CD spectroscopy of proteins is an extremely powerful tool for assessing secondary structure. Although several amino acid side chains (notably Tyr, Trp, Phe, His, and Met) absorb light in the far-UV region of the spectrum (below 250 nm), the most important contributor here is the peptide bond (amide chromophore), with $n \rightarrow \pi$ and $\pi \rightarrow \pi^*$ transitions at ~220 and ~190 nm, respectively. The far-UV CD bands of proteins reflect the secondary structure of the protein (α -helix, β -sheet, β -turn, and unordered content).

The spectrum is converted to the desired units. In the case of proteins, the observed CD signal, S in millidegrees (Note: 1 millidegree = 32,980 ΔA), is generally converted to the molar CD extinction coefficient ($\Delta \epsilon$):

$$\Delta \epsilon = \frac{S}{32980 \times C_M \times L}$$

where L is the path length (in cm), C_M is the molar concentration, $C_{\text{mg/ml}}$ is the concentration in mg/ml.

All- α proteins show an intense negative bands between ~203-240 nm with minima at ~209-222 nm. The intensities of these bands reflect α -helical content. The spectra of regular all- β sheet proteins are significantly weaker than those of all- α helix proteins. These spectra usually display minima between 210 and 225 nm and maxima between 190-200 nm.

Unordered peptides and denatured proteins have a strong negative band (at 195–200 nm) and a much weaker band (which can be either positive or negative) between 215

and 230 nm. $\alpha + \beta$ and α / β proteins almost always have spectra dominated by the α – helical component and therefore often show bands at 222, 208, and 190–195 nm. In some cases, there may be a single broad minimum between 210 and 220 nm because of overlapping α -helical and β -sheet contributions (Martin and Schilstra 2008).

When a mutant protein is produced using recombinant DNA technology it is important to demonstrate that it has the same overall structure as the wild-type protein. Comparison of the near and far-UV CD spectra is ideal for this purpose. A significant difference in shape between the far-UV CD spectra of the wild-type and mutant proteins can be an indication that the mutation has produced some change in the secondary structure (Martin and Schilstra 2008).

3.1.5. Isothermal Titration Calorimetry

ITC provides a complete thermodynamic profile of the molecular interaction between a ligand and a macromolecule. This technique monitors the binding of a ligand to a macromolecule by measuring the heat released or absorbed when the two species interact. The enthalpy of binding is directly measured as the integrated area under the injection peak. By performing a series of injections until the macromolecule is saturated, the binding constant and the entropy of the interaction can be determined (Ladbury and Chowdhry 1996; Pierce *et al.* 1999; Leavitt and Freire 2001).

The calorimeter outputs a series of injection peaks, proportional to the energy input needed to compensate for the heat evolved or adsorbed upon ligand binding. The area of the peak is directly related to the enthalpy of ligand binding to the macromolecule. The association constant (K_a) determines the amount of bound and unbound ligand after each injection. As the concentration of ligand increases the amount of complex formed decreases until all binding sites are filled. The amount of heat adsorbed or evolved (Q) upon binding is given as:

$$Q = \left[\frac{V_o \times \Delta H_{app} \times [M]_{tot} \times K_a \times [L]}{1 + K_a [L]} \right]$$

Equation 3-1

where V_0 is the cell volume, ΔH_{app} is the enthalpy of binding per mol of ligand, $[M]_{tot}$ is the total macromolecule concentration including both the bound and free states, and $[L]$ is the concentration of free ligand. The enthalpy of binding is measured along with the enthalpy of dilution. Dilution effects are usually small but are subtracted as a baseline. Binding models which estimate the K_a and ligand stoichiometry (n) are fitted using a least squares model for ligand-macromolecule complexes assuming either a single set of identical sites, two sets of independent sites, or multiple sequential sites. K_a for a reversible interaction is given by:

$$K_a = \frac{[ML]}{[M][L]} \quad \text{Equation 3-2}$$

where $[M]$, $[L]$ and $[ML]$ are the concentrations of the macromolecule, ligand and complex, respectively. The K_a is determined by applying a linear regression model to the plot of ΔH_{bind} vs. molar ratio. The change from a free molecule to a complexed state produces loss of solvent-ligand and solvent-macromolecule interactions and gain of ligand-macromolecule interactions. The free energy of binding (ΔG_{bind}) is the total thermodynamic effect of these changes. ΔG_{bind} is determined by the K_a :

$$\Delta G_{bind} = -RT \ln K_a \quad \text{Equation 3-3}$$

ΔG_{bind} has contributions from the binding enthalpy (ΔH_{bind}) and the binding entropy (ΔS_{bind}):

$$\Delta G = \Delta H - T\Delta S \quad \text{Equation 3-4}$$

Binding is favoured by a negative ΔG_{bind} . The ΔH_{bind} reflects the strength of interaction between ligand and macromolecule, relative to their interaction with the solvent. A favourable enthalpy term comes from a gain of interactions through the formation of the complex. The ΔS_{bind} has a favourable contribution to the interaction when solvent molecules, which were restricted by interaction with the macromolecule and ligand, are released. The increase in entropy from freedom of the solvent molecules is offset by the loss of conformational freedom of the ligand and macromolecule.

3.1.6. Cyclic Voltammetry

Cyclic voltammetry (CV) is an electrochemical method for determining redox potentials and uses microelectrodes attached to a potentiostat and carried out in quiescent solution to ensure diffusion control. A three-electrode arrangement is used: working vs. counter vs. reference electrode. This technique is based on varying the applied potential to a working electrode in both forward and reverse directions (at some constant scan rate) while measuring the current. The forward scan produces a current peak for any analytes that can be reduced through the range of the potential scan. The current will increase as the potential reaches the reduction potential of the analyte, but then fall off as the concentration of the analyte is depleted close to the electrode surface. As the applied potential is reversed, it will reach a potential that will reoxidise the product formed in the first reduction reaction, and produce a current of reverse polarity from the forward scan. This data is then plotted as current (i) vs. potential (E).

The important parameters in a cyclic voltammogram are the peak potentials (E_{pc} , E_{pa}) and peak currents (i_{pc} , i_{pa}) of the cathodic and anodic peaks, respectively. If the electron transfer process is fast compared with other processes (such as diffusion), the reaction is said to be electrochemically reversible, and the peak separation is:

$$\Delta E_p = [E_{pa} - E_{pc}] = 2.303 \cdot RT \cdot nF \quad \text{Equation 3-5}$$

Thus, for a reversible redox reaction at 25°C with n electrons ΔE_p should be $0.0592/n$ V or 59 mV for one electron. In practice this value is difficult to attain because of factors such as cell resistance. Irreversibility due to a slow electron transfer rate results in $\Delta E_p > 0.0592/n$ V, such that the difference is typically 70-100 mV for a one-electron reaction. The formal reduction potential (E^0) for a reversible couple is given by:

$$E^0 = \frac{(E_{pc} + E_{pa})}{2} \quad \text{Equation 3-6}$$

3.2. Results

3.2.1. Cofactor Assembly in Bacterioferritin.

The BFR proteins were over expressed in *E.coli*, before purification on a metal affinity column. The purified protein then had the histidine tag cleaved using FXa protease. The cleaved protein was then purified from the tag again using metal affinity chromatography. The expressed proteins were red in colour due to a heme group bound at the interface of the homodimeric subunits in a hydrophobic pocket. UV-visible spectroscopy reveals the characteristic bands found for heme bound via two methionine residues in *E.coli* bacterioferritin (Figure 3.4) (Yariv *et al.* 1981). The oxidised and reduced forms of BFR¹ and wild-type BFR have identical spectra with the Soret maxima at 418 nm and red shifting to 425 nm upon reduction with sodium dithionite. Two characteristic Q bands are found for the bound heme in its oxidized form at 561 and 531 nm shifting to 558 and 528 nm when reduced.

Figure 3.4: UV-visible spectra of BFR¹ in its oxidised form (black) and reduced form (red). Wild type BFR produced the same spectra but is not shown for clarity.

ZnCe₆ Binding to BFR¹

Previous work has shown that ZnCe₆ will bind at the heme site of cytochrome b₅₆₂ via axial ligation to histidine (Hay *et al.* 2004). BFR has two surface exposed histidines at positions 46 and 112. To prevent non-specific binding these two residues were replaced

with arginines. The secondary structure and binding of ZnCe_6 to the double mutant (BFR^1) was identical to the wild type as probed by UV-Vis and Circular Dichroism (CD) spectroscopy (Figure 3.5).

The wild-type BFR and the BFR^1 double mutant was analysed with CD spectroscopy (Figure 3.5) to contain 59% and 60% α -helix respectively, with 7% β -sheet and 33% random coil for both proteins. BFR^1 was also found to still form the spherical ~444 kDa dodecamer of dimers as measured by native gel electrophoresis. Thus for all the remaining experiments BFR^1 was used. The nomenclature employed throughout this thesis is as follows: BFR^1 (BFR^1 devoid of all cofactors), $\text{BFR}^1\text{-M}$ (BFR^1 with two Mn-ions bound), $\text{BFR}^1\text{-Z}$ (BFR^1 with ZnCe_6 bound) or $\text{BFR}^1\text{-ZM}$ (BFR^1 with Mn and ZnCe_6 bound).

Figure 3.5: Circular dichroism spectra of the apo forms of BFR and the double mutant BFR^1 (H46R, H112R). BFR and BFR^1 had 59% and 60% α -helix respectively, with 7% β -sheet and 33% random coil for both proteins.

BFR^1 cofactor binding was monitored using UV-Visible spectroscopy (Figure 3.6). and shows the spectra of ZnCe_6 titrated into a solution of BFR^1 . Binding ZnCe_6 to BFR^1 produces a ~7nm shift in the absorbance maxima of the Q_y band of the unbound ZnCe_6 from 632 to 639nm in the bound protein. Monitoring of this absorption shift upon titration of ZnCe_6 into a solution of BFR^1 obtained an association constant (K_a) of 1.2×10^6 . As a control a BFR mutant in which the axial heme ligating M52 was converted to H52 was also measured. This mutant correctly assembled and was capable of oxidizing iron. As expected, no absorbance shift of the chlorin Q_y band was observed,

suggesting the ZnCe_6 can no-longer bind to the BFR. This observation corresponds to the heme binding activity of BFR M52H previously reported by Andrews and co-workers (Andrews *et al.* 1995).

Figure 3.6: UV-visible spectra of ZnCe_6 free in solution (red) and bound to BFR^1 (black).

BFR^1 cofactor binding was also monitored using isothermal titration calorimetry (ITC) to obtain binding constants (K), reaction stoichiometries (n), enthalpies (ΔH°), and entropies (ΔS°), thus providing a complete thermodynamic profile of the molecular interaction. ITC was used to determine the binding affinity of ZnCe_6 to the heme binding site of BFR^1 . Figure 3.7 shows the injection heat for BFR^1 titrated into ZnCe_6 in dilute solution ($15 \mu\text{M}$). The negative peaks correspond to an exothermic reaction. The integrated heat (μCal) for each injection is plotted over the molar ratio of protein to ZnCe_6 in the bottom of Figure 3.7. For this the heat of dilution of the protein was subtracted from the raw data. The binding constant was determined to be $2.54 (\pm 0.45) \times 10^6 \text{ M}^{-1}$ and was found to have a stoichiometry of 0.82 ZnCe_6 molecules per BFR^1 dimer. All parameters are reported in Table 3.1.

Figure 3.7: Calorimetric titration of BFR^I into a solution of ZnCe₆. The top of the figure is the raw ITC output for the titration and the lower half of the figure is derived from the integrated raw data. 11µl injections of 300µM BFR^I were titrated into a 15µM solution of ZnCe₆. The line represents the best least squares fit to a model of a single binding site.

Mn Binding to BFR^I

Mn^{II} binding to BFR^I was measured using ITC. Figure 3.8 shows the injection heat for Mn^{II} titrated into a solution of BFR^I. Analysis of the integrated heat using the best fit of a least squares model for two independent binding sites (Table 3.1) revealed that BFR^I has both a high affinity Mn site ($K_{a1} 1.54 (\pm 0.39) \times 10^9 \text{ M}^{-1}$) and a second lower affinity site ($K_{a2} 4.31 (\pm 1.09) \times 10^6 \text{ M}^{-1}$). Titrations of Mn into BFR^I showed that binding of the first metal ion to site 1 is an exothermic process, while binding to the lower affinity site 2 is an endothermic process (Figure 3.8). According to the ITC fitting an average of 2.5 (± 0.12) moles of Mn ions bind per mole of BFR^I.

Figure 3.8: Calorimetric titrations of MnCl_2 into a solution of BFR^1 . The top of the figure is the raw ITC output for the titration and the lower half of the figure is derived from the integrated raw data. $11\mu\text{l}$ injections of $900\mu\text{M}$ MnCl_2 were titrated into a $30\mu\text{M}$ solution of BFR^1 . The line represents the best least squares fit to a model of two independent binding sites.

Table 3.1: Best fit parameters for ITC measurements of BFR^1 cofactor binding

a. Mn titrated into BFR^1				
	n/BFR^1	$K_a \text{ (M}^{-1}\text{)}$	ΔH_{bind} (Kcal mol ⁻¹)	ΔS (Kcal M ⁻¹)
Mn_1	0.88 ± 0.01	$1.54 (\pm 0.39) \times 10^9$	-7380 ± 116	17.3
Mn_2	1.64 ± 0.11	$4.31 (\pm 1.09) \times 10^6$	860 ± 87	33.2
b. BFR^1 titrated into ZnCe_6				
	n/ZnCe_6	$K_a \text{ (M}^{-1}\text{)}$	ΔH_{bind} (Kcal mol ⁻¹)	ΔS (Kcal M ⁻¹)
BFR^1	1.64 ± 0.02	$2.54 (\pm 0.45) \times 10^6$	-4956 ± 89	12.7

a. Fit to best least squares model of two independent binding sites.

b. Fit to best least squares model of a single binding site.

Cyclic Voltammetry of BFR^I-M

Cyclic voltammetry was used to measure the oxidation potential of manganese when bound to the BFR^I protein. A semi-reversible half-wave potential $E_{1/2}$ was resolved for the BFR^I-M protein with no porphyrin bound. The peaks are most likely for the $\text{Mn}_2^{\text{II,II}}$ to $\text{Mn}_2^{\text{II,III}}$ transition (or possibly $\text{Mn}_2^{\text{III,III}}$) with an $E_{1/2}$ of +0.52V (Figure 3.9).

Figure 3.9: Cyclic voltammogram of BFR^I-Mn at pH 7.7 in 25 mM Tricine, 100 mM KCl on a pyrolytic edge graphite electrode. Scan rate 20 mV s⁻¹.

3.2.2. EPR Spectroscopy

EPR measurements were undertaken to determine the oxidation state of manganese bound to the bacterioferritin protein and their coupling environment.

3.2.3. Simulated Mn^{II} Dimer Line-Shapes

Mn^{II} ($S_{1,2} = 5/2$) dimer (without ZnCe₆)

EPR spectra of BFR^I titrated with two Mn ions per protein molecule revealed a broad signal (~400 mT wide), dominated by a large structured $g = 2$ derivative; at least 5 turning points were observed (Figure 3.10). The non-Curie temperature dependence of this signal suggested that it arises from an even spin antiferromagnetically coupled system, most likely a $\text{Mn}_2^{\text{II,II}}$ dimer (Figure 3.11 Right) (Khangulov *et al.* 1990; Pessiki *et al.* 1994; Howard *et al.* 2000; Epel *et al.* 2005).

The EPR signals also contained hyperfine structure centred at $g=4.5$ and $g=9.8$ (Figure 3.10). These signals spanned 50-185 mT, and were assigned to monomeric Mn^{II} bound within a highly structured, weak ligand field. The hyperfine splitting of this signal is approximately 9.5 mT, which is consistent with Mn^{II} coordinated octahedrally with oxygen and nitrogen ligands, as commonly seen in a number of manganese enzymes (Whiting *et al.* 1996; Schaab *et al.* 2006; McGregor *et al.* 2007).

Figure 3.10: EPR spectra of Mn^{II} complex observed in BFR¹-M in the dark. Temperature dependence of the EPR line-shape (derivative). The data was corrected for a baseline artefact by subtraction of a cubic spline.

The line-shape of Mn^{II} dimer remained constant for temperatures up to 50 K (Figure 3.10), which suggests the energy-level spacing's within the dimer and by inference the exchange/dipolar coupling between the two metal centres must be small. That is to say, that as the manifold of energy-states are not well separated, all energy-states have the same population over the temperature range (5-50 K) used in this study; therefore, entropy dominates.

EPR simulations using the Spin Hamiltonian formalism of the line-shape and temperature dependence of the $\text{Mn}_2^{\text{II,II}}$ signal were undertaken to estimate the exchange/dipole couplings and zero-field parameters for this system. These calculations are described in detail below. The simulated line-shape for the Mn^{II} dimer is shown in Figure 3.11 (left). It correctly predicts/models all 5 turning points of the experimental spectrum. The corresponding simulation of the temperature dependence of the Mn^{II}

dimer is quantitatively consistent with the experimental data (Figure 3.11 right). An improvement to the fit can be achieved by allowing the line-width along each of the principle axes to vary (simulation not shown).

Figure 3.11: EPR spectra of Mn^{II} complex observed in BFR¹ in the dark. (Left) Simulated EPR line-shape (derivate). (Right) Simulated temperature dependence of the EPR line-shape.

Optimized parameters for this simulation can be found in Table 3.2. The exchange coupling between the two metal centres was estimated to be 1.3 K (0.9 cm^{-1}) and is therefore of the order of the microwave quantum (9 GHz, 0.43 K). The estimated Mn-Mn distance ($\sim 4 \text{ \AA}$) is commensurate with crystallographic evidence². Both D values (for either Mn_1 or Mn_2) are small, typical for a Mn^{II} high spin ($S = 5/2$) centre (Dismukes 1996; Golombek and Hendrich 2003; Epel *et al.* 2005). The simulation suggests one of the Mn (here referred to here as Mn_A) occupies a more highly structured ligand environment, whereas the second Mn environment (Mn_B) is virtually isotropic.

² As we assumed the exchange coupling was isotropic, any exchange anisotropy will be subsumed by the dipolar term. This may account for the small experimental vs. simulation offset.

Mn^{II} ($S_{1,2} = 5/2$) Dimer (with $ZnCe_6$)

The addition of the pigment to BFR¹-M led to a dramatic change in the Mn^{II} dimer spectrum (Figure 3.12). The main intensity of the spectrum shifted to $g \sim 2$, and appeared less structured. As before, this new signal demonstrated non-Curie temperature dependence and its line-shape was invariant at all temperatures up to 50 K (Figure 3.12).

Figure 3.12: Temperature dependence of the EPR line-shape (derivative) of Mn^{II} complex observed in BFR¹-ZM in the dark.

It was noted while the spectrum about $g \sim 2$ is altered, the wings ($g \sim 2.5$ etc., now smaller by approximately a factor of 2) matched the structure seen in the original Mn^{II} dimer spectrum, i.e., without $ZnCe_6$. This suggested the spectrum was heterogeneous, made up of both the original Mn^{II} dimer spectrum and a new signal. Subtracting the original dimer spectrum yields (Figure 3.13). The estimated contribution of the original dimer spectrum to the total spectrum observed after the addition of $ZnCe_6$ was 59%. This suggested the chlorin had an effect on 41% of Mn clusters (i.e. 82% of reaction centres). This value is approximately the same as the estimate from the number of bound chlorins per Mn cluster as estimated by ITC. The resultant spectrum – the spectrum generated by the subtraction of the original dimer spectrum – appeared as a ‘simple’ derivative about $g \sim 2$. It had near linear temperature dependence (Figure 3.14).

Figure 3.13: Comparison of Mn^{II} EPR signal observed in BFR^{I} . Top) The EPR line-shape of (a.) BFR^{I} -M and (b.) BFR^{I} -ZM at 5 K. Trace (c) represents the subtraction of 59% of trace (a) from trace (b). Data presented were corrected for a baseline artefact by subtraction of a cubic spline.

Figure 3.14: Comparison of temperature dependence of Mn^{II} EPR signal observed in BFR^{I} as estimated by the double integration of the spectra. (a.) BFR^{I} -M and (b.) BFR^{I} -ZM (c) represents the subtraction of 59% of trace (a) from trace (b).

Subsequent modelling of this signal using the same formalism as before suggested (see Figure 3.15) that the coupling between the two metal centres in this subset of centres decreased, with the exchange component dropping to ~ 0.01 K and the Mn-Mn distance lengthening to ~ 5 Å. It is suggested that the ZnCe_6 disrupts the coupling environment of the Mn cluster.

Figure 3.15: EPR spectra of Mn^{II} complex observed in BFR^{I} -ZM in the dark after the addition of ZnCe_6 . Top) Simulated EPR line-shape (derivate). Bottom) Simulated temperature dependence of the EPR line-shape

Table 3.2: Simulation parameters for the Mn^{II} dimers

	Mn^{II} (without ZnCe_6)	Mn^{II} (with ZnCe_6)
J	1.3 K	~ 0.001
r	3.95 Å	5.30 Å
D_1	0.15 K	0.2 K
E_1/D_1	0.33	0.33
D_2	0.02 K	0.02 K
E_2/D_2	0.1 ³	0.1
width	28 mT	28 mT

³ Can take any value 0-0.33 (i.e. all allowed values for E/D). The parameter is non-unique.

Resolved Hyperfine Structure of the Mn^{II} ($S_{1,2} = 5/2$) Dimer (with $ZnCe_6$).

Interestingly, BFR¹-ZM samples prepared in the dark exhibited an unusual multiline EPR signal (Figure 3.16). This signal was super-imposed on the broad $g=2$ homovalent $Mn_2^{II,II}$ spectral components observed in the dark. This multiline has 5 mT splitting and stretches from 290-410 mT, with up to 23 peaks present.

Figure 3.16: a.) Perpendicular X-band EPR spectra of the BFR¹-ZM complex with an underlying multiline signal apparent. b.) Cubic spline subtraction of the baseline to more clearly visualize the multiline signal (magnified 2 times). Protein concentration 400 μ M, $ZnCe_6$ 200 μ M, Mn 800 μ M. Microwave frequency 9.44 GHz, microwave power 2 mW, modulation amplitude 2 mT, temperature 5 K

Within the framework described above the hyperfine structure observed in the Mn^{II} ($S_{1,2} = 5/2$) dimer (with $ZnCe_6$) spectrum most likely arises from the superposition of the hyperfine patterns for each individual Mn^{2+} centre. 5 mT peak to peak (p-p) hyperfine spacing's are typical for Mn^{2+} dimer systems. The appearance of multiline signals is a common attribute of $Mn_2^{II,II}$ dimer systems (Wu *et al.* 2004).

3.2.4. Theoretical Approach: The $Mn^{II}Mn^{II}$ System

The general Hamiltonian used for the Mn^{II} dimer takes the form:

$$H = -J \cdot S_1 \cdot S_2 + S_1 \cdot D \cdot S_2 + \sum (H \cdot g_i \cdot S_i + S_i \cdot D_{ZF_i} \cdot S_i + S_i \cdot A_i \cdot I) \quad \text{Equation 3-7}$$

Where J is the exchange tensor (assumed to be isotropic), D is the dipolar tensor (fixed by the geometry of the system i.e. the Mn-Mn inter-distance), D_{ZF_i} are the zero-field tensors, and A_i are the hyperfine tensors

The orientation of the two zero-field tensors and dipolar tensor were assumed to be co-linear. See Figure 3.17 below.

Figure 3.17: Pictorial representation of Mn-Mn co-ordinate axis system.

The hyperfine component was initially approximated by an isotropic line-broadening. The resultant Hamiltonian (36 x 36 matrix) was solved numerically using Scilab-4.4.1, an open source vector-based linear algebra package available from www.scilab.org. Typical powder pattern averages used 10-100 β angles and a least squares minimization routine was employed to find the optimal solutions for the above parameters.

Powder Pattern Averaging

The paramagnetic centres of a frozen EPR sample take all possible orientations (relative to the field axis), often referred to as a powder pattern. The simulation code allows for this effect by using combination of Euler rotations (α and β). This system only requires two Euler rotations to define all unique orientations as rotation about the field axis does not alter the Eigen values of the system. A generalized powder pattern code was adopted (shown below). The typical number of β angles used (n) ranged from 10-100. Dummy variables x and y were used to uniformly sample the hemisphere. For a given x (β) angle 1 to $p(x+1)$ y (α) angles were sampled (with the entries of $p(x)$ as defined below). Note: for $x = 0$ ($\beta = 0$), α was also set to zero.

For $x = 0$ to $n-1$

For $y = 1$ to $p(x+1)$

$$p(x+1) = \left(\frac{360}{90/(n-1/2)} \right) \cdot \sin \left(\left(\frac{90}{(n-1/2)} \right) \cdot x \right) \quad \text{Equation 3-8}$$

$$\beta = x \cdot \left(\frac{90}{n-1/2} \right) \quad \text{Equation 3-9}$$

$$\alpha = \left(y - \frac{1}{2} \right) \cdot \left(\frac{360}{p(x+1)} \right) \quad \text{Equation 3-10}$$

α and β angles define the rotation matrix a per Equation 3-11:

$$R = \begin{bmatrix} \cos(\alpha) \cdot \cos(\beta) & \sin(\alpha) \cdot \cos(\beta) & -\cos(\beta) \\ \sin(\alpha) & \cos(\alpha) & 0 \\ \cos(\beta) & \sin(\alpha) \cdot \sin(\beta) & \cos(\beta) \end{bmatrix} \quad \text{Equation 3-11}$$

The unitary transformation $T^* = R \cdot T \cdot R'$ was applied to the tensors:

$$D^* = R \cdot D \cdot R'$$

$$g_{Fe}^* = R \cdot g_{Fe} \cdot R'$$

$$J^* = R \cdot J \cdot R'$$

Transition Intensities

The intensity of transition between the i^{th} and j^{th} state is given as Equation 3-12

$$I \propto \left| \langle i | \beta H_1 \cdot g_{Fe} \cdot S_{Fe} + g_Q \beta H_1 \cdot S_Q | j \rangle \right|^2 \quad \text{Equation 3-12}$$

Where the oscillating H_1 field lies in the x-y plane (perpendicular to H), all of its possible orientations must be included to correctly estimate transition intensities. This is achieved by averaging over the third Euler angle (γ). H_1 can be expressed as Equation 3-13

$$H_1 = \begin{bmatrix} \cos(\gamma) \\ \sin(\gamma) \\ 0 \end{bmatrix} \quad \text{Equation 3-13}$$

hence

$$I \propto \left| \langle i | \frac{H^2 \beta^2}{2} (\cos(\gamma) \cdot \alpha + \sin(\gamma) \cdot \beta) | j \rangle \right|^2 \quad \text{Equation 3-14}$$

where

$$\begin{aligned} \alpha &= (g_{xx} - ig_{xy}) \cdot S^+ + (g_{xx} + ig_{xy}) \cdot S^- + 2g_{xz} S_z \\ \beta &= (g_{yx} - ig_{yy}) \cdot S^+ + (g_{yx} + ig_{yy}) \cdot S^- + 2g_{yz} S_z \end{aligned} \quad \text{Equation 3-15}$$

Integrating with respect to γ yields Equation 3-16:

$$\begin{aligned} & \int_0^{2\pi} (\alpha \cdot \cos(\gamma) + \beta \cdot \sin(\gamma))^2 d\gamma \\ &= \int_0^{2\pi} \frac{\alpha^2}{2} \cdot (1 + \cos(2\gamma)) + \frac{\beta^2}{2} \cdot (1 - \cos(2\gamma)) + \alpha\beta \sin(2\gamma) d\gamma \\ &= \left| \frac{\alpha^2}{2} (\gamma + \frac{1}{2} \sin(2\gamma)) + \frac{\beta^2}{2} (\gamma - \frac{1}{2} \sin(2\gamma)) - \frac{\alpha\beta}{2} \cos(2\gamma) \right|_0^{2\pi} \\ &= \pi(\alpha^2 + \beta^2) \end{aligned} \quad \text{Equation 3-16}$$

The transition intensity can therefore be expressed as Equation 3-17:

$$I \propto \left| \langle i | \frac{\pi H^2 \beta^2 (\alpha^2 + \beta^2)}{2} | j \rangle \right|^2 \quad \text{Equation 3-17}$$

3.2.5. Light Induced Oxidation State Changes

O₂ Consumption of the Di-Metal Site of BFR^I.

Bacterioferritins are members of a class of spherical shell-like iron storage proteins that contain a ferroxidase site within the protein and catalyse Fe oxidation and hydrolysis. The Fe oxidation reaction involves the uptake of molecular oxygen from solution and as a consequence the reaction can be monitored using a Clarke type oxygen electrode. Wild type BFR and BFR^I both oxidized Fe^{II} to Fe^{III} with the concomitant uptake of 0.25 O₂ molecules per Fe atom when a five fold excess of iron was present (see Figure 3.18

and Table 3.3). Both forms oxidise iron with ratios of oxygen to Fe identical to the literature (Yang *et al.* 2000) but the rate ($k_{\max}$) of iron oxidation for the mutant form is slightly higher than for the wild type.

The addition of the light active chlorin had potential to induce light dependent oxidation of the metal ions (either Fe or Mn) bound at the ferroxidase site and this too could be monitored by consumption of oxygen. The addition of ZnCe_6 to the BFR^1 protein did not change the amount of oxygen taken up nor significantly affect the rate of iron oxidation in the dark. Illumination of the $\text{BFR}^1\text{-Z}$ protein with iron bound after iron oxidation had gone to completion produced further oxygen uptake at a slower rate. Additional light induced oxygen uptake totalled approximately the same as dark iron oxygen uptake (see Figure 3.18a).

Figure 3.18: Clarke type oxygen electrode measurements depicting oxygen concentrations of samples in the dark and the changes in concentration that occur upon illumination. a.) $\text{BFR}^1\text{-Z}$ Fe 300 μM , b.) $\text{BFR}^1\text{-ZM}$ dark, c.) $\text{BFR}^1\text{-ZM}$ d.) $\text{BFR}^1\text{-ZM}$ Mn 300 μM , e.) ZnCe_6 , Mn (no protein), f.) $\text{BFR}^1\text{-Z}$. The light was turned on 60 seconds after stabilisation of the dark signal.

The $\text{BFR}^1\text{-ZM}$ complex, when measured under the same conditions, did not auto-oxidise Mn in the dark. Interestingly though, subsequent illumination of this complex did lead to consumption of oxygen (Figure 3.18b, c, and d). Control measurements demonstrated that only a small amount of oxygen was consumed by the Mn free ($\text{BFR}^1\text{-Z}$), and for a protein free solution of ZnCe_6 (Figure 3.18e, and f). Similarly, when heme was bound to the protein in place of ZnCe_6 and manganese bound at the dimetal site, no oxygen uptake was observed in the dark or upon illumination (Table 3.3). Buffered pH6.5 Mn^{2+} solutions showed very little oxygen uptake when illuminated (Figure 3.18b).

Table 3.3: Oxygen uptake data

Sample			Total oxygen uptake (μM)	O_2 uptake / metal ion	k_{max} ($\mu\text{Mol s}^{-1}$)
Controls	Dark	Fe 300 μM (no protein)	29	0.10	0.3
	Light	BFR ¹ -Z	39	-	0.3
		ZnCe ₆ Mn 60 μM (no protein)	40	-	0.1
BFR ¹ -Mn	Dark	BFR ¹ -ZM Mn 60 μM	9	0.14	0.1
	Light	BFR ¹ Heme Mn 60 μM	7	0.12	0.1
		BFR ¹ -ZM Mn 60 μM	127	2.12	0.6
		BFR ¹ -ZM Mn 60 μM Catalase & SOD	133	2.22	1.3
		BFR ¹ -ZM Mn 300 μM	147	0.49	0.7
BFR ¹ -Fe	Dark	BFR ¹ -Z Fe 300 μM	81	0.27	1.4
		wt-BFR Fe 300 μM	81	0.27	0.9
		BFR ¹ Fe 300 μM	71	0.24	1.3
	Light	BFR ¹ Fe 300 μM	156* (75) [#]	0.52* (0.25) [#]	-* (0.2) [#]

Note: BFR 30 μM , ZnCe₆ 15 μM , samples with metal contained either stoichiometric amounts (60 μM) or a 5 fold excess (300 μM). *Total O₂ consumption light and dark cumulative. [#]Additional O₂ consumption in light (as compared to dark). Maximum uptake rates (k_{max}) are averaged over 10 seconds.

The oxygen uptake rate of the BFR¹-ZM complex was increased by the addition of superoxide dismutase (SOD) and catalase but the total amount of oxygen consumed was almost the same (Table 3.3). This suggests that SOD and catalase may have a protective function as their addition should slow ZnCe₆ breakdown from reaction with superoxide or peroxide. No oxygen was released upon dark adaptation. More oxygen was consumed when a five fold excess of Mn was added suggesting more Mn was oxidised. The same amount of oxygen uptake was found for BFR¹-Z with excess iron. Both consumed a total of ~0.5 moles of oxygen per metal ion.

Light Induced Chlorin Oxidation / Tyrosine Oxidation as Measured by EPR.

Light induced oxidation of the bound ZnCe₆ was readily observed by EPR (Figure 3.19a). Upon illumination of the BFR1-Z complex (i.e. no Mn present) with short actinic light flashes at room temperature a narrow 0.75 mT wide radical centred at $g = 2.0022$ was generated. Under the same conditions the BFR¹-ZM complex generated a broader signal (2.5 mT radical) centred at $g=2.0058$. This broader radical resolved a 4-6 peak hyperfine structure with p-p spacing ~1.5 mT characteristic of an oxidized tyrosine (Figure 3.19b) (Un *et al.* 1995; Bleifuss *et al.* 2001; Hofbauer *et al.* 2001). A corrected spin count for each species revealed that the tyrosine signal area was approximately 6 times more intense than that of the ZnCe₆ radical signal.

Figure 3.19: Light induced EPR spectra for (a) BFR^I-Z complex, without manganese and (b) with manganese bound. Traces are scaled arbitrarily. Microwave frequency 9.44 GHz, microwave power 1 μ W, modulation amplitude 0.3 mT, temperature 5 K.

Light Induced Mn²⁺ Oxidation as Measured by EPR

BFR^I-ZM complexes were prepared containing either stoichiometric (2 Mn²⁺ per BFR unit) or 5-fold excess Mn²⁺ (10 Mn²⁺ per BFR unit) in the dark. Excess Mn²⁺ (unbound) appeared as a narrow 100mT wide 6 line EPR signal centred at $g \sim 2$ characteristic of Mn²⁺(H₂O)₆ (Hoganson *et al.* 1989). This signal overlaid the structured Mn₂^{II,II} dimer signal discussed above (see *Mn Binding to BFR^I*). The Mn₂^{II,II} dimer signal was quantitatively the same for both samples.

Upon illumination of the sample containing stoichiometric Mn²⁺ the structured Mn₂^{II,II} dimer signal decreased by $\sim 1/3$. No new Mn signal was observed in perpendicular or parallel mode EPR. Small spectral features observed in parallel mode EPR were attributed to background contaminant O₂.

As with stoichiometric samples, illumination of samples that contained excess Mn²⁺, led to a reduction of the structured Mn₂^{II,II} dimer by $\sim 1/3$. In these samples the additional Mn²⁺ six-line signal present also partially reduced by $\sim 25\%$ (see Table 3.4). Concomitant with the loss of unbound Mn²⁺ (six line) was the appearance of broad parallel polarized signal most likely arising from an even spin system (Figure 3.20). This new signal resolves a hyperfine structure with p-p spacing of ~ 15.6 mT. This spacing is larger than previously reported for monomeric Mn^{III} (Campbell *et al.* 1999; Campbell *et al.* 2000; Whittaker *et al.* 2003). As a control a parallel mode scan of

BFR¹-Z with stoichiometric amounts of iron, revealed a small amount of oxygen contamination but no other features (Figure 3.20b).

Figure 3.20: Parallel mode EPR spectra of (a.) the light minus dark spectra of BFR¹-ZM with a five fold excess of Mn. The inset is a higher resolution scan of the multiline signal. (b.) BFR¹-Z with stoichiometric amounts of iron added. Microwave frequency 9.34 GHz, microwave power 10 mW, modulation amplitude 1 mT, temperature 5 K. Protein concentration 250μM, 125μM ZnCe₆, 2500μM MnCl₂ or 500uM FeSO₄ in 50mM MES pH6.5.

Table 3.4: Relative metal ion EPR signal intensities

Sample	Dark		Light	
	Structured g=2	Six line	Structured g=2	Six line
BFR ¹ -ZM Mn 60 μM	100%	0	66%	0%
BFR ¹ -ZM Mn 300 μM	100%	400%	65%	300%

Note: BFR 30μM, ZnCe₆ 15μM, metal concentration either stoichiometric amounts (60μM) or a 5 fold excess (300μM) concentration per BFR¹ monomer

3.3. Discussion

I have taken a bioengineering approach to develop a totally novel system to study the light-activated electron transport in the water oxidation reactions during oxygenic photosynthesis. Bacterioferritin from *E.coli* was genetically and biochemically modified to produce a minimalist model of the PSII donor side in which analogues of PSII cofactors were bound. These included a photo-oxidisable chlorin, a redox active tyrosine, and a di-nuclear manganese centre (Figure 3.21). This model system demonstrates both porphyrin excitation and multi electron oxidation of Mn ions.

Figure 3.21: Diagram of the amino acid ligands to the dimanganese centre with ZnCe_6 bound at the interface of the BFR dimeric complex. Residues from the A subunit are in blue and the B subunit are in salmon. The three tyrosine's have been labelled on either side of the ZnCe_6 group to show the orientation of the dimer subunits relative to each other. The alpha helices are pictured as grey rods. The picture is modified from the PDB - 1BFR.

The BFR^1 mutant was generated by mutating the two histidine residues – H46R and H112R. These residues are commonly substituted with arginine in other bacterioferritin species and therefore structural changes were considered to be unlikely in the mutant protein. Additionally CD spectroscopy confirmed that the secondary structure remained intact and this was supported by the retained ability of the mutant BFR^1 protein to spontaneously bind porphyrins (heme or ZnCe_6) and metal ions (Fe or Mn).

Cofactor-Protein Assembly

Upon titration of the BFR¹ into a solution of ZnCe₆, ITC analysis reveals that 82% of the BFR¹ binds ZnCe₆ (Table 3.1). The small fraction which doesn't bind may be due to partial aggregation of the chlorin in solution, even at the low concentrations used, or it may reflect heterogeneity in the sample; i.e. some of the binding pockets are perturbed. The control mutant (M52H) did not bind any porphyrins (chlorin or heme) possibly due to steric hindrance through the introduction of two large histidine residues in the heme binding pocket, preventing dimer formation. ITC analysis also revealed that binding of ZnCe₆ to the hydrophobic heme pocket has a large enthalpy term (Table 3.1). This large and negative ΔH_b value reflects strong non-covalent interactions (van der Waals and hydrogen bonds) between ZnCe₆ and BFR¹ relative to their interaction with the solvent. The main driving force for binding is most likely the hydrophobic interactions between the porphyrin ring structure and the hydrophobic amino acid residues lining the binding pocket.

As we showed earlier manganese ligates to the di-nuclear metal binding site of bacterioferritin (Conlan 2008) via four glutamate residues (E18, E51, E94, E127) one from each helix and two histidine residues (H54, H130) (Frolow *et al.* 1994). Residues E51 and E127 form (di- μ -1,3-carboxylato) bridging carboxylates, where each oxygen atom is coordinated to a different metal ion, while residues E18 and E94 form monodentate carboxylate ligands. H54 and H130 coordinate to the metal ions separately through the N^δ nitrogen (Figure 3.21). ITC analysis of Mn binding to BFR¹ revealed two distinct binding sites in the dark with very different association constants. Site A bound Mn very tightly whilst site B binding was weaker by almost three orders of magnitude. Binding of the first Mn ion has a large and negative ΔH_{b1} value, which may reflect structural changes in the protein upon metal ligation (LeBrun *et al.* 1996). The second Mn binding has a small and positive ΔH_b but a larger ΔS_b than site 1 indicating that binding Mn to site 2 is entropically driven. The most likely contributions to this positive entropy are changes in the hydration of the protein and of the metal ion upon binding to the protein (Bou-Abdallah *et al.* 2003). These observations coincide with previous crystal structure findings for BFR with manganese bound, where site B was found to have a lower incidence of occupation than site A (Frolow 2001).

In comparison, during the photo-assembly of the tetra-nuclear Mn cluster in PSII, the first Mn^{II} ion binds tightly at a high affinity site in the apo-complex [IM_0 intermediate]. Light is then needed to start the assembly process in which the bound Mn^{II} is oxidised to Mn^{III} and a proton is released [IM_1]. The bound Mn^{III} is unstable until a conformational change occurs in the dark and allows the formation of the more stable $\text{Mn}^{\text{III}}\text{-O-Ca}^{\text{II}}$ [IM_1^*]. At this point the second Mn^{II} binds forming the stable, spin-coupled cluster $\text{Mn}^{\text{II}}\text{-Mn}^{\text{III}}\text{-O-Ca}^{\text{II}}$ [IM_2]. The formation of the μ -oxo bridge thus helps stabilize the intermediates until the full assembly of the tetra-nuclear MnCa cluster (Dasgupta *et al.* 2008).

The ITC titrations also showed that an average of 2.51 moles of Mn ions bound per mole of BFR^1 monomer. This is higher than expected but can be partly explained by the findings from the original crystal structure where an extraneous Mn ion was shown to be bound at the four-fold axis of the dodecamer of dimers by four Gln151 residues internal to the structure (Dautant *et al.* 1998). This would account for a further 0.25 binding sites per mole of protein but still leaving 0.25 sites unaccounted for. Non-specific, low affinity binding sites have also been detected in previous studies of the BFR protein. It is proposed that these sites involve acidic residues located on the inner surface of the protein shell (Baaghil *et al.* 2002). Under the conditions of these experiments the individual thermodynamic parameters of the weak binding sites cannot be determined. This is due to the limiting form of the equation for two classes of binding sites (Freire *et al.* 1990).

The oxidation states of the Mn centre were characterised by EPR studies. Simulation of spectra of BFR in the dark strongly suggests that manganese ions bound to the BFR^1 protein remained in the 2+ oxidation state and are present as a weakly spin-coupled $\text{Mn}_2^{\text{II,II}}$ centre. The estimated Mn-Mn distance from EPR simulations was $\sim 4 \text{ \AA}$, a distance agreeing with that obtained in the crystal structure (Frolow *et al.* 1994). The addition of ZnCe_6 appeared to alter approximately 50% of the $\text{Mn}_2^{\text{II,II}}$ complexes. This behaviour may be rationalized as follows. Initially the homodimer has two identical Mn-clusters. Upon addition of ZnCe_6 this symmetry is lost. The BFR protein was engineered to bind ZnCe_6 between the two monomeric BFR^1 subunits via a methionine residue. Only one monomer of the dimer subunits provides this ligand (the 5th axial coordinate of the ZnCe_6) see Figure 3.21. The methionine residue is one amino acid

from the crucial glutamic acid residue (E51) which ligates to both Mn ions in the metal centre. It is possible that the binding of the ZnCe_6 disrupts the coupling through movement of the methionine and thus the glutamic acid residue producing a carboxylate shift (presumably moving the two metal centres further apart).

O₂ Consumption

BFR functions in biology to bind iron and oxidize Fe^{II} to Fe^{III} . This process involves the formation of a stabilizing μ -oxo bond between the two metal ions. This reaction consumes dissolved O_2 (Liu *et al.* 2004). The three phases to this process involve (Equation 3-18) metal binding and deprotonation of amino acid ligands followed by (Equation 3-19) oxidation of bound metal ions with the formation of a single μ -oxo bond between the metal ions and uptake of one oxygen atom per centre. Once all ferroxidase iron binding sites are filled, and all the 48 iron atoms bound per dodecamer of dimers are oxidized, the complex then begins to oxidize excess Fe^{II} within the ball and store it as ferric-oxy-hydroxide (Equation 3-20) (LeBrun *et al.* 1995a; Yang *et al.* 2000; Baaghil *et al.* 2003).

The oxygen assay of the BFR^1 protein with iron present revealed that 0.25 O_2 molecules are taken up per iron oxidised (Table 3.3). This agrees with the literature values for the wild type and the ZnCe_6 bound form, supporting the assumption that binding of the ZnCe_6 does not change the basic function of the protein. Minimal oxygen consumption is observed for BFR^1 -ZM complex suggesting it cannot auto-oxidize Mn^{II} to Mn^{III} in the dark. Little or no oxygen consumption is observed for these complexes even under the high pH conditions at which animal ferritins perform this reaction, which is surprising as animal ferritins are very similar in structure and function (Bou-Abdallah *et al.* 2002; Meldrum *et al.* 1991; Zhang *et al.* 2006).

Upon illumination the BFR¹-ZM sample consumes ~2 O₂ per Mn. This value is 4 times larger than for the BFR with Fe bound and may suggest the di-Mn site constructs multiple μ -oxo bridges. Control samples – samples that lack Mn i.e. BFR¹-Z – did not consume a significant amount of oxygen confirming that O₂ is reacting with the Mn. In samples with a five fold excess of manganese, a significant increase in O₂ consumption was observed (~20 μ M). This likely indicates that 'free' Mn²⁺ is oxidised within the ball structure forming Mn^{III}-oxy-hydroxide. The addition of catalase and SOD was found to increase the rate of oxygen consumption but did not have any significant effect on the total amount consumed. We suggest that it acts to protect the chlorin, removing super-oxides and other reactive oxygen species

Tyrosine Oxidation

In the absence of manganese, illumination of the BFR¹-Z produces a large ZnCe₆^{•+} radical cation signal at $g = 2.0022$. In contrast, addition of Mn results in the loss of the ZnCe₆^{•+} radical and the appearance of a 'tyrosine like' radical. This new radical signal is unlikely to be the ZnCe₆^{•+} radical broadened due to through bond / through space interactions with the Mn site as the distance between the Mn centre and the ZnCe₆ is a relatively large ~14 Å. The intensity of the tyrosine signal is significantly in excess of the ZnCe₆ (6 fold). This is unsurprising as tyrosine radicals are generally much longer lived than chlorin radicals (Tang *et al.* 1996; Miloslavina *et al.* 2006).

The most likely mechanisms for tyrosine oxidation is electron hole migration from the oxidized ZnCe₆^{•+} radical to a nearby tyrosine residue. ZnCe₆ has a high estimated oxidation potential of ~+1.1V (Hay *et al.* 2004) which is sufficient to oxidize a tyrosine. The +0.5V oxidation potential of the manganese centre should allow it to be oxidised by any tyrosine radical formed. It is unclear why tyrosine oxidation does not occur when no manganese ions are bound. When Mn (or any metal) binds to the BFR metal site deprotonation of the metal ligands occurs along with structural changes to the ligands themselves. These changes may alter the hydrogen bonding network around the tyrosine allowing proton coupled electron transfer.

There are three tyrosine residues in BFR (Y25, Y58, and Y45) that are close enough to the ZnCe₆ (10.6Å, 11.0Å, and 4.2Å respectively) that they could potentially act as 'efficient' electron donors (Figure 3.21). Tyrosine 45 rests within van der Waals

bonding distance of the ZnCe_6 and makes up part of the heme binding pocket. It is unlikely that this tyrosine would be affected by Mn binding and as such is not likely to be the electron donating residue. This leaves only two tyrosines (Y25 and Y58) which are found at the metal binding site and are both involved in the hydrogen bonding network of the site. Tyrosine 58 is not highly conserved and is commonly replaced by a leucine in other bacterioferritins. Tyrosine 25 is hydrogen bonded to glutamate 94 which is one of the metal ligands with the oxygen atom of this tyrosine placed 4.3 Å from the nearest Mn ion. Ferritins, which have very similar structure and function to bacterioferritin but don't bind heme, have a redox active tyrosine within the hydrogen bonding network of the metal binding site. This residue is highly conserved in ferritins and bacterioferritins (Carrondo 2003). As a consequence, we suggest tyrosine (Y25) is likely to be the redox active residue in our system. This is currently being tested in our lab.

Mn²⁺ Oxidation

The loss of Mn^{II} signals upon illumination of $\text{BFR}^1\text{-ZM}$ strongly suggests Mn oxidation occurs in this system. The broad structured $\text{Mn}_2^{\text{II,II}}$ signal centred at $g=2$ was found to decrease in $\text{BFR}^1\text{-ZM}$ samples upon illumination. This structured $\text{Mn}_2^{\text{II,II}}$ signal decreased on average by 34%.

No new Mn EPR signals were observed upon loss of the Mn^{II} signal in $\text{BFR}^1\text{-ZM}$. We would expect the one electron oxidation of an $\text{Mn}_2^{\text{II,II}}$ to yield a mixed valence $\text{Mn}_2^{\text{II,III}}$ dimer. Strongly coupled mixed valence dimers ($J \approx 10 \text{ cm}^{-1}$) display characteristic 'multiline' signals seen in perpendicular mode EPR centred at $g \sim 2$ with hyperfine splittings of $< 90 \text{ G}$ (Teutloff *et al.* 2005). In contrast, the weakly coupled mixed valence dimer should still have broader perpendicular signals and could possibly resolve parallel polarization signals (Teutloff *et al.* 2005; Groni *et al.* 2008). As neither of these is seen, it is suggested the Mn cluster undergoes two electron oxidation to a even spin $\text{Mn}_2^{\text{III,III}}$ system. It is not clear as to whether both proposed Mn^{II} oxidation events need to be photo-driven. As no intermediate state ($\text{Mn}_2^{\text{II,III}}$) was observed, the second oxidation state may not require light excitation of the chlorin. Alternatively the mixed valence cluster may be more strongly oxidizing (electron donating) to the oxidized tyrosine/chlorin than the $\text{Mn}_2^{\text{II,II}}$ precursor state. Oxygen bridged $\text{Mn}_2^{\text{III,III}}$ dimers with analogous ligand fields have been observed at approximately zero-field ($g = 20$)

(Whittaker *et al.* 2003). We cannot discount the presence of these signals in our illuminated BFR complex.

The BFR^I-ZM sample with a five fold excess of Mn revealed the same decrease in amplitude of the structured $g = 2$ Mn^{II,II} signal upon illumination as seen in samples with stoichiometric Mn. The six-line signal also showed a significant decrease upon illumination. This signal decreased by 25%, equivalent to 4 Mn oxidations per BFR dimer (48 Mn ions per dodecamer of dimers). Parallel polarization EPR on illuminated samples with excess Mn suggested the formation of a manganese species with even spin character, upon illumination of the BFR^I-ZM complex; a broad signal centred at $g_{\text{eff}} \sim 4-5$ with clear hyperfine structure (Figure 3.20). As this signal is not observed in illuminated samples with stoichiometric Mn, it is suggested that this signal comes from a new Mn species unlikely to be bound to BFR. Comparing this to the native function of ferritins, it is tempting to assign the species to Mn^{III}-oxy-hydroxide [Mn^{III}(O)OH] which is accumulated in the core of the sphere structure (Zhang *et al.* 2005).

Mn^{III} is a d^4 even spin ($S = 2$) transition ion (Abragam and Bleaney 1970). There are potentially three geometries that Mn^{III} complexes in this protein ligand field can take: 6 coordinate octahedral, 5 coordinate square-pyramidal or trigonal-bipyramidal (Campbell *et al.* 2000). The ground electronic state of the Mn^{III} manifold depending on the geometry of the ligand field can have two different d^4 configurations corresponding to the electronic hole position. The hole either occupies the $d_{x^2-y^2}$ orbital (ground state 5B_1) or the d_{z^2} orbital (ground state 5A_1). The hyperfine splitting for each of these states is different due to the effect of spin-orbit coupling (Gerritsen and Sabisky 1963; Griffith 1971; Al'tshuler and Kozyrev 1974; Campbell *et al.* 2000). A ground state of 5B_1 yields a hyperfine splitting along A_z of ~ 50 G whereas a ground state of 5A_1 yields a hyperfine splitting of ~ 100 G (Gerritsen and Sabisky 1963; Campbell *et al.* 2000; Tyryshkin *et al.* 2006). In this instance the sign of the Spin Hamiltonian parameter D indicates the ground state of the manifold. A positive D value yields a 5A_1 ground state (as observed for the Mn^{III} centre of the SOD protein) whereas a negative D values yields a 5B_1 ground state (as observed for the Mn^{III} photo-assembly intermediate of PS II) (Griffith 1971; Whittaker and Whittaker 1991; Campbell *et al.* 1999; Campbell *et al.* 2000).

Recently parallel polarization EPR signals have been reported for Mn^{III} peroxo complexes (i.e. $\text{Mn}^{\text{III}}(\text{O})\text{OH}$) (Groni *et al.* 2008). These signals were generated by the oxidation of $(\text{L})\text{Mn}^{\text{II}}$ (where $\text{L} = \text{N-methyl-tris(2-pyridylmethyl)ethan-1,2-diamine}$) using a 1000 fold excess of H_2O_2 as an oxidant. Two parallel polarization signals were observed: a structured signal at $g \sim 8$ with hyperfine splitting ~ 50 G (analogous to Mn^{III} signals described above (Campbell *et al.* 2000; Campbell *et al.* 2001; Tyryshkin *et al.* 2006) and a less structured signal at $g \sim 5$ with hyperfine splitting ~ 95 G. It was noted that the addition of base (triethyl-amine) led to the loss of the high field signal ($g \sim 5$). Conversely the addition of acid (HClO_4) led to the loss of the low field signal ($g \sim 8$). As a consequence the low field signal ($g \sim 8$) was assigned to the monocationic species $[\text{LMn}^{\text{III}}(\text{OO})]^+$. Both UV-Vis and ESI-MS supported this assignment. XSophie EPR simulations estimated the zero-field and hyperfine parameters of this complex to be: $D = -2.9 \text{ cm}^{-1}$, $E/D = 0.075$, $A_Z = 6.5 \times 10^{-3} \text{ cm}^{-1}$ (Groni *et al.* 2008). Curiously, the second signal ($g \sim 5$) was not assigned to the corresponding protonated species $[\text{LMn}^{\text{III}}\text{OOH}]^{2+}$. It was noted that though the signal could be modelled as arising from a Mn^{III} signal ($D = +2.9 \text{ cm}^{-1}$, $E/D = 0.111$), the resolved hyperfine structure (splittings of ~ 95 G) was inconsistent with assignment to a Mn^{III} (Groni *et al.* 2008). From the above discussion this argument does not seem valid; Mn^{III} can potentially resolve hyperfine splitting of the order of ~ 100 G. We note the D value changes sign for these two complexes, which should reflect a re-ordering of electronic state manifold (ground state interchanges from $^5\text{B}_1$ to $^5\text{A}_1$) as seen between Mn^{III} photo-assembly intermediate of PS II and Mn-SOD protein, leading to an apparent increase in hyperfine splitting observed (Campbell *et al.* 1999; Campbell *et al.* 2000).

Our new Mn EPR signal observed in BFR bears a resemblance to the $g \sim 5$ signal seen in the above study (Groni *et al.* 2008) appearing at approximately the same g position ($g \sim 4.5$). It too has a large hyperfine spacing as seen for the potential $[\text{LMn}^{\text{III}}\text{OOH}]^{2+}$ species. We do acknowledge though that the hyperfine spacing seen (~ 150 G) in the ‘new Mn EPR signal’ is exceedingly large for a monomeric Mn^{III} . The unusually large hyperfine splitting may reflect the unique environment Mn centres occupy in the core of the sphere protein structure (Mn-Mn interactions etc). Alternatively, the new EPR signal may represent a di-Mn species, possibly a weakly coupled mixed valance dimer (i.e. $\text{Mn}^{\text{II}}\text{Mn}^{\text{III}}$).

3.4. Conclusion

In this study we have shown that a modified bacterioferritin complex (BFR^I-ZM) is capable of using light energy to drive photochemical charge separation and couples this oxidising potential to multiple oxidation events within a protein bound Mn₂^{II} complex. To my knowledge this work is the first example reported in the literature of a bioengineered protein capable of photo-catalytic, multi-electron oxidation of a di-manganese centre. This represents one of the first steps in developing synthetic photo-catalytic, 'green enzymes' that utilize light energy to catalyse oxygen evolution and hydrogen production.

CHAPTER 4

4 *Bacterioferritin Heterodimer*

4.1. *Introduction*

The previous chapter outlined the successful implementation of bacterioferritin from *E.coli* as a scaffold for PSII cofactor binding. This construct was limited in a number of aspects and of particular note was the lack of a bound electron acceptor. Electron acceptors placed in solution provide low yields of electron transfer due to the large reorganization energies associated with transferring an electron through a high dielectric medium such as water. This chapter documents the engineering of a BFR based heterodimeric light activated metallo-protein that contains a quinone electron acceptor built into the protein matrix. Inspiration for this engineering project was again taken from PSII which is made up of a heterodimer that allows for electron transfer along a chain of redox cofactors culminating in the reduction of a quinone.

In order to engineer an electron acceptor within the protein matrix it was necessary to modify the bacterioferritin protein such that a linker chain of amino acids was used to join two monomeric subunits which normally form a homodimer. The linker was a repeated motif joining the C terminus of one subunit to the N-terminus of the second subunit. A key feature of this linker is that it produced a dimeric form of the protein which allowed one half of the protein to be modified without changing the other half. In joining the proteins it allowed for the amino acids associated with metal (Mn) coordination to be removed in one sub-unit and replaced with a residue that enabled covalent attachment of a quinone. This produced a heterodimeric protein with a redox active metal centre on one side, a light active porphyrin centrally located, and a protein bound electron acceptor on the other side. This linearised electron transfer pathway is shown in Figure 4.1.

To achieve expression of a heterodimeric protein required use of a complex set of molecular biology based techniques. Introducing a combination of three different genes, BFR¹ (H46R, H112R), BFR² (H46R, H112R, E51C) and a 22 amino acid linker within

a plasmid produced two final heterodimer constructs. Both of these constructs were then mutated to remove unwanted histidine residues within the linker region producing the final two different versions of the heterodimeric protein, BFR-H1 and BFR-H2 (see Chapter 2 for details).

BFR-H1 was made of two identical BFR¹ subunits linked via a 22 amino acid linker. This protein should have similar characteristics to that of the BFR¹ protein utilized in the previous chapter. BFR-H2 contained a mutation (E51C) in subunit B as depicted in Figure 4.1, which enabled a quinone to be covalently bound to the cysteine sulphur within the metal binding pocket. This mutation prevents metal binding at the metal centre and produces an asymmetrical heterodimer. The final expressed heterodimeric proteins were 338 amino acids in length and had a molecular weight of 38.8 kDa.

Figure 4.1: Bacterioferritin heterodimer (BFR-H2) with one subunit of the dimer changed relative to the other. This heterodimer contains a complete electron transport chain, with a light active ZnCe_6 molecule, a covalently attached quinone electron acceptor, and a di-manganese centre as electron donor. Molecular mechanics optimization of the structure was carried out using a Polak-Ribiere algorithm (conjugate gradient) on the original dimer structure with the 22 amino acid linker region incorporated (Polak and Ribiere 1969).

The linker sequence contained a flexible Ser-Gly-Gly repeat with a single basic lysine residue in the middle. This linker is expected to be solvent exposed and free to move in solution. It was necessary to keep the length of this linker region to a minimum to prevent having a large unstructured loop destabilizing the heterodimers. X-ray crystal structure data for the *E.coli* BFR homodimer sets the spatial separation between the N and C termini of the adjoining A and B subunits as 28.8 Å. The linker was modelled to have a length of 79 Å when extended out straight. Modelling of this linker predicted that it should be long enough to prevent steric hindrance of the formation of the dimer but not so long as to destabilise the structure of the protein (Whitney and Sharwood 2006).

The production of the heterodimer enables four cofactors to be placed within close proximity of each other. Overlaying the new cofactors onto the wild type BFR crystal structure and modelling the changes associated with binding the cofactors produces a set of cofactors spaced as per Figure 3.3.

Figure 4.2: Diagram showing the relative distance of separation of all of the cofactors bound to the BFR heterodimer protein. Distances are given as edge to edge for the conjugated systems and metal ions.

The use of an existing protein scaffold (i.e. bacterioferritin), makes the process of constructing and analysing light activated metallo-enzymes much easier. Utilising natural protein folds and tertiary structure removes the need for complete *de novo* design of a protein and in only making minor modifications the structure is readily maintained. In analysing these complexes there are advantages as each individual

cofactor can be introduced systematically, allowing the characterization of its interactions with the protein environment. This is much simpler than working with the large membrane bound PSII complex which requires progressively breaking down the complex with detergents in order to study its function. Such treatment can damage the reaction centres before they can be analysed.

The BFR heterodimer constructs were labelled as BFR-H1 and BFR-H2 to refer to the H1 heterodimer and H2 heterodimer construct which has the capacity for quinone binding. Further to this the labels Z, M or ZM have been included to indicate the cofactors bound to these proteins.

- **BFR-H** indicates the apo-form of either of the BFR heterodimer proteins, devoid of all cofactors.
- **BFR-H M** is used to indicate BFR heterodimer proteins with a stoichiometric number of Mn-ions bound (i.e. 4 Mn ions for BFR-H1 but only 2 Mn ions for BFR-H2).
- **BFR-H Z** is used to indicate the heterodimers with one ZnCe_6 bound per heterodimer.
- **BFR-H ZM** is used to indicate the BFR heterodimers with stoichiometric amounts of Mn and ZnCe_6 bound to the protein.

4.1.1. Quinones

Quinones are di-ketones and are utilised extensively in nature as electron acceptors in electron transfer chains. Ubiquinone (coenzyme Q, CoQ), is a lipid soluble para-benzoquinone which is ubiquitous in biological systems (Figure 4.3). It is found in the mitochondria and bacterial reaction centres and undergoes cyclic oxidation-reduction during the oxidation of substrates. Ubiquinones are named dependent on the number of isoprene units in the hydrophobic tail e.g. $n = 10$: coenzyme Q_{10} . Higher plants use a different quinone as electron acceptor called plastoquinone, which has methyl groups instead of methoxy groups attached to the cyclic carbon ring of the quinone.

Figure 4.3: Ubiquinone Coenzyme Q (2,3-dimethoxy-5-methyl-1,4-benzoquinone).

Quinones can be non-covalently bound as is the case of quinones in photosynthetic (Allen and Williams 1998; Heathcote *et al.* 2002; Wraight 2004) and respiratory (Cecchini *et al.* 2003; Ohnishi and Salerno 2005; Osyczka *et al.* 2005) electron-transfer pathways. Non-covalently bound quinones are also found in the pyrroloquinoline quinone (PQQ) containing proteins (Anthony 2004). These non-covalently bound membrane associated quinones tend to bind at specific hydrophobic sites via interaction with the long phytyl tail but are stabilised by hydrogen bonding with the carbonyl oxygen and van der Waals interactions between the quinone head group and the protein (Warncke *et al.* 1994). There are a number of covalently bound quinones which are post-translationally generated cofactors such as tryptophan tryptophylquinone (TTQ) found in bacterial dehydrogenases (Davidson 2001; Klinman 2001) and 2,4,5-trihydroxyphenylalanine quinone (TPQ) in the copper amine oxidases (Klinman 2001; Mure 2004).

Quinones act as electron acceptors/donors which can be singly reduced to the semi-quinone radical anion ($Q^{\bullet-}$) or doubly reduced to the dianion (Q^{2-}). In protic media, the strong affinity of the quinol hydroxyls for protons results in the formation of a stable, neutral quinol at pH values below their pK's that is greatly stabilized in relation to the semi-quinone. This results in the potential of the second reduction step ($Q^{\bullet-}$ to QH_2) being much higher than that of the first (Q to $Q^{\bullet-}$), giving rise to its $n = 2$ equilibrium redox chemistry (Rich 2004). The redox potentials of quinones in electron transport chains are ideally suited as electron acceptors with potentials ranging from -130 mV up to +250 mV (Krieger *et al.* 1995; Krieger-Liszkay and Rutherford 1998; Li *et al.* 2006; Rappaport and Diner 2008).

General structural motifs for the binding of quinones in natural systems is starting to emerge (Fisher and Rich 2000) though these motifs are not so well understood as to be able to use *de novo* design to build a pocket for stable quinone binding. The best characterized binding site is the Q_A site of the photosynthetic reaction centres from purple bacteria (Heathcote *et al.* 2002; Wraight 2004)). This site is structurally similar to the Q_A site of photosystem II (Ferreira *et al.* 2004; Loll *et al.* 2005) and has similarities to the quinone Q_D binding site in the *E. coli* fumarate reductase (Iverson *et al.* 1999; Cecchini *et al.* 2003)

Recently, 3,4-dihydroxy-L-phenylalanine was incorporated into proteins by expanded genetic-code technique to generate *o*-quinone bound to proteins (Alfonta *et al.* 2003). Though this avoids the need to build a quinone binding site but it is a complex process to incorporate a potentially toxic molecule into *E.coli* and it is easier to simply covalently bind a quinone within the final purified protein matrix. This enables the quinone to be buried away from the high dielectric solvent decreasing the reorganisation energy required for electron transfer.

It has long been known that unsubstituted quinones such as para-benzoquinone (pBQ) form adducts with thiol-containing compounds (Snell and Weissberger 1939) including cysteine (Redfearn 1965). Hay *et al.* (2004) were the first group to make use of this novel addition reaction and by modifying the gene for cytochrome b_{562} were able to place a cysteine residue within the interior of this protein and covalently attach a quinone molecule. When the cysteine site was covalently linked to 2,3-dimethoxy-5-methyl-1,4-benzoquinone (CoQ₀) (ubiquinone lacking the multiprenyl tail), and ZnCe₆ was bound in the heme pocket, light-induced electron transfer from the bound pigment to the bound quinone was achieved (Hay *et al.* 2004).

Other methods for binding quinone's to proteins have been investigated using thiol addition reactions as well as addition-elimination reactions between the cysteine sulphur and thioether quinones (Li *et al.* 2005). The reaction of cysteine and CoQ₀ is thought to occur according to Figure 4.4.

Figure 4.4: The covalent attachment of a quinone to the cysteine sulphur of a protein via a thiol addition reaction Taken from (Li *et al.* 2005).

4.1.2. Quinone Binding Site Design

The quinone binding site for BFR was designed such that the quinone could be bound within the protein matrix and shielded from the solvent phase. The site chosen was the metal binding site of BFR¹. This site allows the binding of a large cofactor due to the cavity that already exists for the incorporation of two iron atoms. The removal of the Glu51 residue and replacement with the much smaller cysteine residue further opens this space for the binding of a quinone. Assuming the structural model of BFR still holds for the heterodimer proteins then it is expected that the quinone will be placed 10.6 Å from the chlorin.

The protein matrix can have a large effect on the redox potential of the quinone when it is conjugated to a cysteine. Reports of shifts between +110 mV and +140 mV have been reported for 2,5-dimethyl-1,4-benzoquinone (DMBQ) and CoQ₀ respectively, when bound to *de novo* designed constructs versus acetyl cysteine adducts (Hay *et al.* 2007; Li *et al.* 2006). Although previous work by Hay *et al.* showed only a small shift of ~25 mV when CoQ₀ was bound to an engineered Cyt *b*₅₆₂ (Hay *et al.* 2004). The protein construct did not contain any potential hydrogen bonding partners with the bound quinone which may be the reason for so little change in the redox potential. The type of quinone covalently bound has been found to have little effect on the rates of electron transfer suggesting that the chemical nature of the quinone itself has little effect on efficiency (Hay *et al.* 2004). As the quinone binding site in the BFR heterodimer contains two histidine residues which could potentially hydrogen bond with the quinone we could expect an increase in the potential possibly approaching that found by Li *et al.* (2006) i.e. +229mV.

4.1.3. Summary

Here I report the construction of a heterodimeric form of the light activated BFR¹ metallo-catalyst from Chapter 3. The previous chapter focused on light induced oxidation state changes in the manganese centre i.e. $\text{Mn}_2^{\text{II,II}}$ to $\text{Mn}_2^{\text{III,III}}$. This chapter focuses on the production of the heterodimer and assessment of its biochemical properties. The light induced electron transfer from ZnCe_6 to a quinone electron acceptor bound within the protein matrix was followed using steady state fluorescence and EPR.

4.1.4. Theory of Techniques Used

Fluorescence Energy Transfer

When an electron is excited to an orbital with a high electronic energy, the excess energy is stored. In most cases this stored energy is transferred as heat, and is called non radiative energy transfer. Because of the relative configurations of the excited and ground states, the electron can easily make a transition from a vibrational state in the excited system to a nearly degenerate level in the ground state, and it can continue to undergo non radiative energy loss until it reaches the ground state. Radiative energy transfer occurs when the lowest vibrational level of the excited state is too far above any vibrational level in the ground state for energy to be relinquished except with the radiation of a photon. Some energy is given off as heat before radiative processes can occur which is why the wavelength of emitted light is usually of a lower energy than the excitation wavelength, this called the Stokes shift. Other decay processes can occur to enable energy dissipation, these include intersystem crossing to a triplet state followed by phosphorescence, or electron transfer. The levels of phosphorescence are usually small compared to the levels of fluorescence due to the transition from the triplet state to a ground state being forbidden according to quantum mechanical rules and as a result the rate constant is several orders of magnitude smaller than those of fluorescence.

Assuming the major form of energy dissipation is fluorescence for a given light absorbing porphyrin then enabling electron transfer in this system as a form of energy dissipation will markedly decrease the yield of fluorescence. This enables fluorescence

measurements to be used to quantitate electron transfer in a light activated system such as PSII or an artificial PSII analogue. Artificial systems such as porphyrin-quinone diads have shown reductions in fluorescence yield in the presence of an oxidised quinone. Ho and co-workers used an amide-linked free-base porphyrin and benzoquinone. Quinone-induced quenching of the porphyrin fluorescence was seen. As well as fluorescence quenching a porphyrin cation radical and a semi-quinone radical were generated upon light excitation (Ho *et al.* 1980). These two observations are consistent with the formation of a charge separated state. There are numerous examples of donor-acceptor molecules that have been designed and several reviews written (Wasielewski 1992; Gust *et al.* 1993; Kurreck and Huber 1995; Gust *et al.* 2001).

Electron Paramagnetic Resonance

Quinones when singly reduced form a semi-quinone radical anion ($Q^{\bullet-}$). EPR acts as a sensitive technique to measure this radical. The semi-quinone radical produces a characteristic signal centred around $g=2$. These radicals like most organic radicals have very long spin lattice relaxation times which results in EPR spectra exhibiting relatively narrow peaks even at room temperature.

4.2. Results

4.2.1. Cofactor Assembly in BFR Heterodimers

The BFR heterodimer proteins were over expressed in *E.coli*, before purification on a metal affinity column. The purified proteins then had the histidine tagged ubiquitin protein cleaved off using a ubiquitin specific protease. The cleaved BFR protein was then purified from the tag again using metal affinity chromatography. The expressed proteins were red in colour due to a heme group bound at the interface of the dimers in a hydrophobic pocket. For the heme to bind between the subunits the protein must be properly folded and structured.

UV-visible spectroscopy reveals the characteristic bands found for heme bound via two methionine residues in *E.coli* bacterioferritin (Figure 4.5) (Yariv *et al.* 1981). The oxidised forms of the BFR heterodimers have a Soret band at 418nm and this red shifts upon reduction with sodium dithionite out to 425nm. Two characteristic Q bands are found for the bound heme at 531 and 561nm for the oxidised form shifting to 528 and 558nm when reduced. All forms of the BFR protein including the wild type and heterodimers H1 and H2 retain the same spectral characteristics with no changes in the band positions.

Figure 4.5: The UV-visible spectra of holo BFR-H2 in its oxidised form (black) and reduced form (red). All three BFR heterodimers produced the same spectra but only one is shown for clarity.

The heme was removed from the BFR heterodimer proteins again using methyl ethyl ketone and any bound metal ions were removed through exhaustive dialysis against EDTA under reducing conditions (Teale 1959; Bauminger *et al.* 1991) The secondary structure of the BFR-H1 and H2 proteins was then assessed using CD spectroscopy (Figure 4.6 and Figure 4.7). A more comprehensive CD analysis program (CD spectra deconvolution v2.1 © Dr. Gerald Bohm) was used to fit the spectra which gives a more accurate picture than the previous analysis program (Kd2) used in Chapter 3.

Analysis of the CD spectra revealed that the heterodimer proteins were less structured than the BFR¹ mutant. Fitting the spectra revealed that the proteins contained less alpha-helix than found in BFR¹. It was found that the addition of manganese ions to the proteins increased the alpha-helix content marginally and reduced the amount of random coil. This was more evident in BFR-H2 than in BFR-H1.

Figure 4.6: The CD spectra of apo BFR-H1 and BFR-H1 in the presence of manganese. The addition of manganese produced a small change in the secondary structure. BFR-H1 10μM, Mn 40μM in TBS buffer pH7.7

BFR-H2 was less structured than BFR-H1 suggesting that the mutation of Glu51 to a cysteine had an effect on the secondary structure of this protein (Figure 4.7). There was >10% reduction in the amount of alpha-helix as compared to BFR-H1 and an increase in the amount of random coil, indicating it is a less structured protein (Table 4.1).

Figure 4.7: The CD spectra of apo BFR-H2 and BFR-H2 plus manganese. The addition of manganese had a significant effect on the secondary structure. BFR-H2 10μM, Mn 40μM in TBS buffer pH7.7

According to native PAGE electrophoresis the BFR heterodimer proteins do not form the spherical ~444 kDa dodecamer of dimers. It is possible that it does form in solution but under the conditions used to run the gels no high molecule weight multimer bands were found.

Table 4.1: CD Structural Parameters

	BFR ¹	BFR-H1	BFR-H1-M	BFR-H2	BFR-H2-M
Helix	75.3%	48.6%	50.3%	37.4%	40.6%
Antiparallel	2.2%	5.1%	4.9%	7.2%	6.6%
Parallel	2.6%	5.8%	5.5%	7.8%	7.2%
Beta-Turn	10.7%	14.4%	14.1%	16.1%	15.6%
Random Coil	12.5%	24.2%	23.5%	30.3%	28.5%
Total Sum	103.3%	98.2%	98.3%	98.8%	98.3%

Isothermal Titration Calorimetry

Cofactor binding was monitored using ITC to determine the binding constants (K), reaction stoichiometries (n), enthalpies (ΔH°), and entropies (ΔS°) of manganese binding and ZnCe₆ interactions. The BFR¹ protein from Chapter 3 revealed that 2.5 Mn ions were bound per monomer of protein (Table 4.2a). This construct was only expected

to bind 2 Mn ions and the extra 0.5 were expected to be either bound within the pores of the dodecamer of dimers sphere structure or non-specifically bound.

The heterodimer construct BFR-H2 has the E51C mutation in one half of the dimer where the crucial glutamic acid residue was mutated to a cysteine such that the metal binding site in one half of the protein should no longer function to bind metal ions. This mutation places a cysteine such that a quinone electron acceptor can be covalently bound within the protein matrix. To look at the effects of this mutation in isolation, the BFR¹ homodimer protein was mutated further to produce BFR² (BFR¹-E51C). ITC analysis of manganese binding in the BFR² mutant revealed that it does not bind manganese in any significant amount (Figure 4.8). Fitting the binding isotherm to a single binding site reveals that only 0.31 (± 0.02) manganese ions bind per mole of BFR² monomer (Table 4.2b). This binding is quite weak ($2.05 \pm 0.28 \times 10^5$) and may be at the metal binding site or simply non-specific ligation on the protein surface.

Figure 4.8: Calorimetric titration of MnCl_2 into a solution of BFR² which has the E51C mutation in the metal binding site. The top of the figure is the raw ITC output for the titration and the lower half of the figure is derived from the integrated raw data. 6 μl injections of 840 μM MnCl_2 were titrated into a 30 μM solution of BFR². The line represents the best least squares fit to a model of a single binding site.

The titration of manganese into the BFR-H1 heterodimer construct was expected to reveal similar metal binding characteristics as that found in the BFR¹ homodimer. The metal binding ability of the BFR-H1 heterodimer was significantly reduced with a total of 2.4 Mn ions were bound per heterodimer (Table 4.2c). It was expected that a greater amount of Mn would bind i.e. 4 Mn ions per dimer. An isotherm representing the average for this titration is shown in Figure 4.9. The decrease in secondary structure and the decrease in the number of Mn ions bound suggest the heterodimer construct is partially destabilised by the addition of the loop region.

Figure 4.9: Calorimetric titrations of MnCl_2 into a solution of BFR-H1. The top of the figure is the raw ITC output for the titration and the lower half of the figure is derived from the integrated raw data. 11 μl injections of 900 μM MnCl_2 were titrated into a 15 μM solution of BFR-H1. The line represents the best least squares fit to a model of two independent binding sites.

Titration of manganese into the BFR-H2 heterodimer construct was expected reveal less metal binding due to the modification of the ligands in one half of the construct. Integration of the heat for a titration of manganese into a 15 μM solution of BFR-H2 (Figure 4.10) when fit to a least squares model for two independent binding sites (Table 4.2d) revealed that BFR-H2 binds slightly less than one manganese ion tightly and then a second manganese ion more weakly. The main binding event is an endothermic process with 0.76 (± 0.01) Mn ions binding per heterodimer with a K_{a1} of 3.14 (± 1.67) $\times 10^6$ M. The second binding event is much weaker with a K_{a2} of 3.24 (± 1.69) $\times 10^4$ M.

The heterodimer complex was expected to bind two manganese ions on one half of the protein complex. This seems to be the case though the second manganese ion binds very weakly. The other half of the heterodimer is not expected to bind any metal ions due to removal of a crucial glutamate ligand to produce a quinone binding pocket.

Figure 4.10: Calorimetric titrations of MnCl_2 into a solution of BFR-H2. The top of the figure is the raw ITC output for the titration and the lower half of the figure is derived from the integrated raw data. 11 μl injections of 900 μM MnCl_2 were titrated into a 15 μM solution of BFR-H2. The line represents the best least squares fit to a model of two independent binding sites.

The titration of a concentrated solution of BFR-H1 into a 15 μM solution of ZnCe_6 revealed a biphasic binding relationship (Figure 4.11). The integration of the heat of binding produces a curve which could only be fit to a model of two independent binding sites. The first binding event has a very high association constant and a moderate ΔH_b whilst the second binding event is much less strongly associated but has a very large ΔH_b producing a favourable binding event. The total number of BFR-H1 binding per ZnCe_6 is equivalent to 1.15 (Table 4.2e).

Figure 4.11: Calorimetric titrations of BFR-H1 into a solution of ZnCe₆. The top of the figure is the raw ITC output for the titration and the lower half of the figure is derived from the integrated raw data. 11µl injections of 300µM BFR-H1 were titrated into a 15µM solution of ZnCe₆. The line represents the best least squares fit to a model of two independent binding sites.

Table 4.2: Best fit parameters for ITC measurements of Mn and ZnCe₆ binding to BFR constructs

Titrant	n/BFR	K _a (M ⁻¹)	ΔH _{bind} (Kcal mol ⁻¹)	ΔS (Kcal M ⁻¹)
a. Mn titrated into BFR¹				
Mn ₁	0.88 ± 0.01	1.54 (± 0.39)x10 ⁹	-7380 ± 116	17.3
Mn ₂	1.64 ± 0.11	4.31 (± 1.09)x10 ⁶	860 ± 87	33.2
b. Mn titrated into BFR²				
Mn	0.31 ± 0.02	2.05 (± 0.28)x10 ⁵	-1630 ± 123	18.8
c. Mn titrated into BFR-H1				
Mn ₁	1.07 ± 0.02	5.15 (± 1.77)x10 ⁶	2471 ± 53.8	39
Mn ₂	1.32 ± 0.08	8.94 (± 2.46)x10 ⁴	690 ± 47.7	25
d. Mn titrated into BFR-H2				
Mn ₁	0.76 ± 0.01	3.14 (± 1.67)x10 ⁶	1294 ± 12.8	34.1
Mn ₂	1.14 ± 1.27	3.24 (± 1.69)x10 ⁴	443.3 ± 552	22.1
Titrant	n/ZnCe ₆	K _a (M ⁻¹)	ΔH _b (Kcal mol ⁻¹)	ΔS (Kcal M ⁻¹)
e. BFR-H1 titrated into ZnCe₆				
BFR ₁	0.47 ± 0.01	2.42 (± 1.81)x10 ⁸	-850 ± 147	35.5
BFR ₂	0.68 ± 0.05	8.17 (± 2.18)x10 ⁵	-6135 ± 536	6.47

a, c, d & e: Fit to best least squares model of two independent binding sites.

b: Fit to best least squares model of a single binding site.

Mn EPR Spectra

EPR spectra were taken of heterodimer samples with Mn bound to look at the interaction of the Mn ions within the di-metal sites. The spectra revealed a broad structured signal centred at g=2 and a further structured region at g=4 (Figure 4.12a and b). These two signals were modelled in the BFR¹ protein (Chapter 3) showing that the broad signal centred at g=2 was the result of weakly interacting Mn₂^{II,II} dimers (Figure 4.12c). The spectra of the heterodimers revealed less of the dimeric Mn₂^{II,II} structure

than found in the BFR¹-ZM sample. This coincides with the ITC data which shows weaker binding of the second metal ion in the heterodimers versus BFR¹-ZM. There is also a larger signal centred at 150 mT ($g=4$) associated with the quino-protein which is due to monomeric Mn^{II} bound within a highly structured, weak ligand field.

Figure 4.12: EPR spectra of a.) BFR-H2 ZM CoQ₀, b.) BFR-H1 ZM and c.) BFR¹-ZM. All samples were in tricine 40mM KCl 100mM pH7.7 buffer. Microwave frequency 9.4 GHz Microwave power 10mW, modulation amplitude 10G, temperature 5K.

Iron Oxidation Rates – UV-Vis

The enzymatic activity of all of the BFR mutants was assessed using a well established iron oxidation assay (Andrews *et al.* 1995). Bacterioferritin catalyses the oxidation of Fe^{II} to Fe^{III} with the formation of oxidised iron producing absorbance at 340nm which enables the rate of oxidation to be monitored (Figure 4.13). The rate of uncatalysed iron oxidation, without the protein present, was very low for buffers with a pH less than 6 but above this pH the rate increased significantly. The assay was therefore carried out at pH5.9.

Figure 4.13: Iron oxidation monitored by absorbance at 340nm. Protein concentrations 5μM, iron 2000μM, in 40mM MES buffer pH 5.9. Iron was injected at time zero with rapid stirring.

The initial rates of iron oxidation, given as the change in absorbance at 340nm per minute, are very similar for the wild-type and BFR¹ mutant (Table 4.3). The heterodimers on the other hand show much increased initial rates of oxidation. BFR-H1 which from the CD spectra appears to be more structured than BFR-H2 shows slower iron oxidation. BFR-H2 shows very fast iron oxidation but the rate tails off significantly over time. The uncatalysed rate is negligible and is not significant when compared to the catalysed rates.

Table 4.3: Iron oxidation rates as monitored by UV-visible spectroscopy

Sample	Rate ($\Delta \epsilon_{340} / \text{min}$)
Iron	0.01
wt-BFR	0.08
BFR ¹	0.07
BFR-H1	0.18
BFR-H2	0.38

4.2.2. Quinone Binding

The covalent binding of CoQ₀ to a protein or other molecule via a sulphur substitution reaction has a major influence on the UV-visible spectra of the quinone. The significant changes in the blue region of the spectra enable the substitution reaction to be monitored. Binding a glutathione molecule (γ -L-Glutamyl-L-cysteinylglycine (2S)-2-Amino-), which is a cysteine containing tri-peptide, to the quinone reveals the changes in absorbance which can be expected upon covalently attaching the quinone to a protein. Figure 4.14 shows the UV-visible absorbance spectra of CoQ₀, glutathione and the CoQ₀-Glutathione complex in aqueous buffer pH7.7. CoQ₀ initially has an absorbance band at 268nm which red shifts upon sulphur substitution to 276nm. The new CoQ₀-glutathione adduct produces intense π - π^* transition bands at 276 nm and 343 nm

Figure 4.14: Absorbance spectra of CoQ₀ (black), glutathione (red) and the product of reacting CoQ₀ with reduced glutathione producing the CoQ₀-Glutathione complex (blue). All samples were in tricine 40mM KCl 100mM pH7.7 buffer.

Monitoring the absorbance spectra for the binding reaction of CoQ₀ with BFR-H2 produces a spectrum which is a composite of the protein signal and the substituted quinone (Figure 4.15). The spectra revealed maxima at 276 and 343 nm characteristic for the quinone with thioether linkage to the protein. The relative amount of CoQ₀ bound to BFR-H2 equates to ~1:1.

Figure 4.15: UV-visible absorbance spectra of BFR-H2 CoQ₀ (black), BFR-H2 (red) and CoQ₀-Glutathione complex (blue). The concentration of all protein and quino-adducts was 17μM in tricine 40mM, KCl 100mM pH7.7 buffer.

Binding ZnCe₆ to the BFR-H2 CoQ₀ protein reveals a spectrum very similar to that of BFR¹ with ZnCe₆ bound (Table 4.4). Binding produces a red shift in the absorbance maxima of the ZnCe₆ Soret band from 411 to 422nm, whilst the Q_x band shifts from 517 to 521nm and the Q_y band from 632 to 637nm (Figure 4.16). The absorbance spectra of ZnCe₆ bound to BFR-H2 ZM and to BFR-H2 ZM CoQ₀ are almost identical, which would suggest that there is no ground-state coupling between the chlorin and quinone.

Figure 4.16: Absorbance spectra of ZnCe₆ free in solution (black), and bound to BFR-H2 CoQ₀ (red). The inset shows the absorbance maxima of the Q_x band magnified eight times. Samples were prepared in tricine 40mM KCl 100mM pH7.7 buffer.

Table 4.4: ZnCe₆ absorbance maxima when bound to BFR proteins

Binding Partner	Absorbance Maxima (nm) ⁴
Free in solution	411, 517, 632
BFR ¹	423, 522, 639
BFR-H2 CoQ ₀	422, 521, 637

EPR Dark Spectra of Quinones

EPR spectroscopy was utilised to monitor quinone bound to the BFR protein constructs. Unbound or bound quinone could be visualised with this technique. The singly reduced quinone radical anion is readily picked up in the region of $g=2$. By increasing the pH the semi quinone radical is generated and can be observed at room temperature. Figure 4.17 shows the spectra of the semi quinone form for CoQ₀ and also 1,2-naphthoquinone (NQ).

Figure 4.17: Room temperature EPR spectra of a.) CoQ₀ 6.6mM (spectra multiplied by 5) microwave power 20mW and b.) 1,2-naphthoquinone 10mM, microwave power 5mW. Samples made up in 80mM CAPS pH 9.5 buffer. Microwave frequency 9.8 GHz, modulation amplitude 0.1mW.

⁴ The optical properties of a porphyrin can be accounted for by electronic transitions from the two highest occupied molecular orbital's to the two lowest unoccupied molecular orbital's. The interaction of these orbital's gives rise to the relatively weak Q band (~500-650 nm) and the intense Soret band in the UV region. The porphyrin ring system of the chlorin is perturbed, splitting the nominally degenerate Q band into a lower energy Q_y band and a higher energy Q_x band Gouterman M (1961) Spectra of porphyrins. Journal of Molecular Spectroscopy 6: 138-163. Ligation of the porphyrins changes the energy levels of these orbital's producing shifts in the absorbance spectra.

CoQ₀ was covalently bound to BFR² producing the homodimer with two copies of the quinone bound. The generation of the semi quinone radical at room temperature using high pH reveals a much broadened and less featured signal than the free quinone (Figure 4.18). The semi-quinone is only formed with a low yield and as such the signal is quite noisy.

Figure 4.18: Room temperature EPR spectra of a.) CoQ₀ free in solution and b.) CoQ₀ bound to BFR² 170μM. Samples in 80mM CAPS pH9.5 buffer. Microwave frequency 9.8 GHz, modulation amplitude 2G, microwave power 20mW.

4.2.3. Light Induced Changes in BFR Heterodimers

Light Induced Mn²⁺ Oxidation as Measured by EPR

The BFR-H1 ZM complex was prepared containing stoichiometric amounts of ZnCe₆ and Mn (i.e. 1 ZnCe₆ and 4 Mn²⁺ per BFR heterodimer unit) in the dark. EPR was used to monitor changes in the Mn₂^{II,II} dimer signal centred at g=2 upon illumination. The sample was illuminated for 2 minutes at 200K with white light then quick frozen, and the structured Mn₂^{II,II} dimer signal was found to decrease by a small amount (Figure 4.19). No new Mn signal was observed in perpendicular or parallel mode EPR. It would be beneficial to use longer illumination times at room temperature to increase the amount of manganese which is oxidised (see Chapter 3).

Figure 4.19: EPR spectrum of BFR-H1 ZM before and after illumination at 200K for 2 minutes. Microwave frequency 9.4 GHz Microwave power 10mW, modulation amplitude 20G, temperature 5K.

Light Induced Chlorin Oxidation / Tyrosine Oxidation as Measured by EPR.

As was found for BFR^I, light induced oxidation of ZnCe₆ bound to the BFR heterodimers was readily observed by EPR (Chapter 3). Upon illumination of the BFR-H1 ZM complex at 200K a broad radical signal was generated (Figure 4.20a). The same sample with no manganese present (i.e. BFR-H1 Z) produced a much narrower 0.75 mT wide radical centred at $g = 2.0022$ (Figure 4.20b). Subtraction of the spectra of the narrow radical signal from the broad signal reveals a structured underlying signal (2.0 mT) centred at $g=2.0031$, which appears very similar to a tyrosine radical species (Figure 4.20c) (Un *et al.* 1995; Bleifuss *et al.* 2001; Hofbauer *et al.* 2001). This broad structured radical signal appears to be the same signal as that found in the BFR^I protein though the turning points are not as well defined. Again the formation of the tyrosine radical was dependent on Mn being bound to the metal centre.

Figure 4.20: EPR spectra of a.) BFR-H1 ZM, b.) BFR-H1 Z and c.) a. minus b. The samples were illuminated for 2 minutes at 200K before measurements were taken. Microwave frequency 9.4 GHz Microwave power 0.01mW, modulation amplitude 5G, temperature 5K

Light Induced O_2 Consumption of the BFR Metal Site

BFR^I has been shown to be capable of binding manganese at the ferroxidase site and when ZnCe₆ is bound oxidising the bound Mn^{II} to Mn^{III} (see Chapter 3). This occurred concomitant with the uptake of molecular oxygen from solution and this O_2 consumption can be directly monitored using a Clarke type oxygen electrode. The analysis of BFR^I, BFR-H1 and BFR-H2 constructs found them to all consume oxygen, but the rate of consumption and amount taken up varied according to the construct. The BFR-H1 ZM, BFR-H2 ZM and BFR-H2 ZM CoQ₀ complexes did not auto-oxidise Mn in the dark. Though subsequent illumination of these complexes did lead to oxygen uptake (Figure 4.21).

Control measurements demonstrated that minimal O_2 was consumed upon illuminating quinone free in solution (Figure 4.21a.). However, illumination of the heterodimer constructs with ZnCe₆ and Mn bound revealed high levels of O_2 consumption. The BFR-H1 heterodimer took up the least amount of O_2 (Figure 4.21b.) consuming slightly less than that found for BFR^I (Table 4.5). BFR-H2 ZM consumed the same amount of O_2 as BFR^I but interestingly BFR-H2 ZM CoQ₀ took up considerably more oxygen than any of the other constructs (Figure 4.21c. and d.).

Figure 4.21: Oxygen electrode traces showing light induced oxygen uptake. a.) 10µM quinone b.) BFR-H1 ZM c.) BFR-H2 ZM d.) BFR-H2 ZM CoQ₀. Protein concentration 15µM, ZnCe₆ 15µM, Mn 60µM in 50mM MES buffer pH6.5

BFR²-ZM (homodimer with E51C mutation) was shown to bind very little Mn and then only weakly (see section 4.2.2), yet it still consumes almost as much oxygen as BFR¹-ZM (Table 4.5). This protein also had the fastest initial rate of oxygen uptake. The BFR-H2 ZM CoQ₀ protein had the next highest rate of oxygen uptake, consuming almost half again the amount of oxygen found for the other constructs, which suggests more Mn centres were oxidised.

Table 4.5: Oxygen uptake data

Sample	Total O ₂ uptake (µM)	O ₂ uptake / metal ion	Max Rate (µMol/s)*
10µM CoQ ₀ control	16	-	0.16
BFR ¹ ZM	127	2.12	0.58
BFR ² ZM	118	1.97	1.03
BFR-H1 ZM	107	1.78	0.63
BFR-H2 ZM	128	2.13	0.63
BFR-H2 ZM CoQ ₀	200	3.33	0.78

Note: BFR 30µM, ZnCe₆ 15µM, Mn 60µM.

BFR-H 15µM, ZnCe₆ 15µM, Mn 60µM

*Maximum rate was determined over an average of the initial 10 seconds.

Steady State Fluorescence Properties

The steady state fluorescence of ZnCe_6 when bound to oxidised forms of both the BFR-H2 ZM and quinone bound form of this same protein (BFR-H2 ZM CoQ_0) produced very different yields of fluorescence. The fluorescence emission of ZnCe_6 is decreased by 21.7% ($\pm 1.2\%$) when bound to the quino-protein (Figure 4.22). The only difference between the two proteins is the presence of the quinone within its binding pocket and as such any changes in yield are likely due to energy dissipation to the quinone via a route not involving fluorescence i.e. electron transfer (Hay *et al.* 2004).

Figure 4.22: Fluorescence spectra of a.) BFR-H2 ZM b.) BFR-H2 ZM CoQ_0 c.) spectra a. minus spectra b. BFR-H2 proteins 2.5 μM , ZnCe_6 2.5 μM , MnCl_2 66 μM in 50mM MES pH6.5.

When the proteins are reduced with sodium dithionite the overall fluorescence increases significantly for ZnCe_6 bound to both the quino-protein and the non-quinone protein (see Figure 4.23). The fluorescence levels of the reduced forms of both proteins are identical ($\pm 1.9\%$). The reduced forms show an almost flat difference spectrum as both proteins now have no electron acceptors. The fact that the spectra are so similar when reduced suggests that the binding of the quinone to the protein complex does not change the binding environment of the ZnCe_6 .

Figure 4.23: Fluorescence spectra of sodium dithionite reduced BFR heterodimer proteins. a.) BFR-H2 ZM b.) BFR-H2 ZM CoQ₀ c.) resultant spectra of a. minus b. BFR-H2 proteins 2.5μM, ZnCe₆ 2.5μM, MnCl₂ 66μM in 50mM MES pH6.5.

EPR Spectra of BFR-H2 Quino-Protein

In order to try and monitor light induced changes in the BFR-H2 quino-protein EPR spectroscopy was again utilised. For these experiments 1,2-naphthoquinone (NQ) was bound covalently to the BFR-H2 protein to produce the BFR-H2 NQ protein. Samples were monitored at room temperature in the dark and then subsequent spectra were taken under the same conditions but with constant illumination. All spectra shown are light minus dark spectra.

It was found that under constant illumination at room temperature the BFR-H2 Z quino-protein (no Mn) produced a radical signal which was broader than a simple ZnCe₆ cation radical (Figure 4.24a). The same protein, without NQ bound, produced a signal which was somewhat narrower, centred at $g=2.0022$, and characteristic of the ZnCe₆ radical (Figure 4.24b). The BFR-H2 protein without NQ produced a much larger ZnCe₆ signal than the quino-protein. Subtraction of a portion of the ZnCe₆ signal from the quino-protein spectra reveals a radical which is broader and with a higher g value centred at $g=2.0032$. This signal appears to be the singly reduced semi-quinone radical (Figure 4.24c). The formation of the semi-quinone radical could only be due to electron transfer from the ZnCe₆.

Figure 4.24: Illuminated room temperature EPR spectra of a.) BFR-H2 Z NQ, b.) BFR-H2 Z, and c.) a. minus b. Samples were under constant illumination during measurement. Microwave frequency 9.8 GHz, modulation amplitude 4G, microwave power 20mW

Looking at the same quino-protein with manganese bound revealed some significant differences. BFR-H2 ZM produced a broad radical signal as expected for this sample which is a combination of the ZnCe_6 cation radical and the tyrosine radical (Figure 4.25a). The same protein, with NQ covalently bound, produced a radical signal which was much narrower (Figure 4.25b). Subtraction of the quino-protein spectra from the original protein spectra reveals a radical signal which is remarkably similar to the tyrosine radical signal (Figure 4.25c). Thus it appears that the binding of NQ to BFR-H2 results in the loss of a large percentage of the tyrosine radical species. Fifty percent of the tyrosine radical is expected to be lost due to quinone binding preventing light and manganese dependent tyrosine oxidation in one half of the protein. The binding of NQ which is larger than CoQ_0 may also have detrimental effects on the other Mn centre inhibiting tyrosine oxidation.

Figure 4.25: Illuminated room temperature EPR spectra of a.) BFR-H2 ZM, b.) BFR-H2 ZM NQ, c.) a. minus b. Samples were under constant illumination during measurement. Microwave frequency 9.8 GHz, modulation amplitude 4G, microwave power 20mW

4.3. Discussion

I have taken a bioengineering approach to study the light-activated electron transport of the water oxidation reactions during oxygenic photosynthesis. The bacterioferritin protein utilised in Chapter 3 was modified to produce a minimalist model of PSII in which not only donor but also acceptor cofactors were bound. These included a photo-oxidisable chlorin, a quinone electron acceptor, a redox active tyrosine, and a di-nuclear manganese centre (Figure 4.26). This model system demonstrates porphyrin excitation, quinone reduction and metal centre oxidation.

Figure 4.26: A cartoon model of the BFR heterodimer which combines a manganese metal centre as electron donor (D), a redox active tyrosine, a light activated pigment (P), and a covalently bound quinone electron acceptor (A).

4.3.1. Cofactor-Protein Assembly

Linker Effects

The BFR-H1 and BFR-H2 heterodimer constructs were generated by linking the C-terminus of subunit A to the N terminus of subunit B using a 22 amino acid flexible linker. BFR-H2 incorporated a mutation to allow a quinone to be bound within the

metal binding site of subunit B. Upon expression of these constructs they were found to bind heme as per the wild-type and there were no shifts in the absorbance spectrum of the bound heme for any of the mutant forms.

In order to bind the heme it is expected that the dimer must form properly and the structure of the proteins remain intact. Even upon removal of the heme and any bound metal ions the BFR heterodimer proteins still retained the majority of secondary structure as determined by CD spectroscopy. This was supported by the retained ability of the mutant BFR proteins to spontaneously bind porphyrins (heme or ZnCe_6) and metal ions (Fe or Mn).

The CD spectra revealed that the heterodimer proteins were less structured than the BFR¹ mutant. The decrease in secondary structure was in part expected as the addition of a completely unstructured 22 amino acid loop region would increase the random coil component by at least 7%. The addition of metal ions was found to increase the alpha-helical constant of both heterodimer forms. This was particularly evident in the BFR-H2 construct which was less structured than the BFR-H1 construct due to the extra mutation of Glu51 to a cysteine (Table 4.1).

It is not certain but appears likely that the BFR heterodimer proteins do not form the large 24mer sphere structure. This indicated that the Gly-Gly-Ser linker was either inhibiting the 24mer forming due to the fact that it extends from a region which is inherently involved in the interactions needed to form the sphere; or that the reduction in alpha-helical structure destabilizes the ionic interactions linking the components. Addition of the linker region is estimated to increase the iso-electric point (pI) of the protein from the literature value of 4.72 up to 4.79, due to the basic nature of the linker. This is a small amount but the BFR complex relies on negative interactions to form the dodecamer of dimers.

Isothermal Titration Calorimetry

Manganese ligates to the di-nuclear metal binding site of bacterioferritin (Conlan *et al.* 2007) via four glutamate residues (E18, E51, E94, E127) one from each helix and two histidine residues (H54, H130) (Frolow *et al.* 1994). E51 and E127 form di- μ -1,3-

caboxylato bridges, where each oxygen atom is coordinated to a different metal ion, removal of E51 was expected to cancel metal binding within the site.

ITC analysis of Mn binding to BFR² revealed that there was no longer a distinct binding site within the protein and the small amount of manganese which did bind was not ligated very strongly (Table 4.2). The large and negative ΔH_{b1} value found for Mn binding to BFR¹ was significantly less in BFR² which reflects the lack of structural changes normally associated with metal binding at the ferroxidase site (LeBrun *et al.* 1996).

BFR-H1 and BFR-H2 both bind manganese ions with similar association constants but BFR-H1 binds more Mn as it contains two di-metal binding centres whereas the BFR-H2 mutant only contains one di-metal binding centre. BFR-H1 and BFR-H2 when binding the first manganese ion have a relatively small and positive ΔH_b but a large ΔS_b indicating that binding Mn to site 1 is entropically driven (Table 4.2). The most likely contributions to this positive entropy are changes in the hydration of the protein and of the metal ion upon binding to the protein (Bou-Abdallah *et al.* 2003). The first Mn ion binds to the heterodimers relatively strongly but the second ion binds much more weakly. Both the association constants are significantly smaller than that found for BFR¹. This may be due to strains placed on the metal binding site due to the large linker and the associated reduction in secondary structure. Ligands are required from all four helices of the 4-helix bundle to bind the metal ions so changes in just one helix could destabilise metal binding. Though the binding is not as tight for the heterodimers there is still definite ligation of Mn ions within these proteins.

The BFR-H1 heterodimer appeared to bind ZnCe₆ quite differently to that found for BFR¹. The heterodimer appeared to have two binding events taking place. The initial binding event which was quite strong and proceeded until 0.47 BFR-H1 molecules were titrated in per ZnCe₆ molecule. The next binding event was not as strong and may have to do with multimerisation of the dimer subunits as they become more structured upon binding to ZnCe₆. This second binding event may also reflect heterogeneity in the sample; i.e. some of the protein aggregating at high concentration. ITC analysis revealed that the second binding event of ZnCe₆ to the hydrophobic heme pocket has a large enthalpy term (Table 4.2). This large and negative ΔH_b value probably reflects

strong non-covalent interactions (van der Waals and hydrogen bonds) between ZnCe_6 and BFR^1 relative to their interaction with the solvent. The driving force for binding is likely the hydrophobic interactions between the porphyrin and the hydrophobic amino acid residues lining the binding pocket (Zhang *et al.* 2007).

Iron Oxidation Rates

Analysis of iron oxidation rates is a good method to determine if the native activity of the BFR protein is retained upon production of different mutants. All forms of the BFR protein tested retained iron oxidation activity (Figure 4.13). The wild-type and BFR^1 forms had very similar rates of oxidation, though the heterodimers appear to oxidise iron at a much faster rate (Table 4.3). The initial rate of oxidation for BFR-H2 is very fast but this rate tails off significantly over time. This change in apparent rate may be due to precipitation of the oxidised iron which is no longer being stored within the sphere structure of the BFR protein shell.

The rate of oxidation seems to increase as the number of mutations and changes in secondary structure increase. The slight unwinding the BFR protein when expressed as the heterodimeric form appears to speed up iron oxidation in solution. It is suggested here that loosening of the structure opens the ferroxidase site slightly which allows faster access and exit of reactants and products to this site.

Quinone Binding

The covalent binding of quinones to proteins or other molecules produces distinctive shifts in the absorbance spectrum (Figure 4.14) (Hay *et al.* 2004; Li *et al.* 2005; Li *et al.* 2006). Binding CoQ_0 to the reduced forms of BFR^2 and BFR-H2, produced absorbance bands at 276 nm and 343 nm due to the covalent linkage to the cysteine sulphur. This binding was further validated by EPR analysis of the singly reduced semi-quinone form of the quino-proteins. The semi-quinone radical can be detected by EPR at high pH (~9).

For para-benzoquinone in water at pH 7.0, $E(\text{Q}/\text{Q}^{\bullet-}) = 99\text{mV}$, and $E(\text{Q}/\text{Q}^{2-}) = 286\text{mV}$ (Depew and Wan 1988) and no semi-quinone is observed. As the pH is increased above pH 7, $E(\text{Q}/\text{Q}^{\bullet-})$ doesn't change but $E(\text{Q}/\text{Q}^{2-})$ decreases by 120mV per pH unit above pH

7 up until the pK (QH/QH_2) which is approximately 10 (Chambers 1988). This decreases $[E(Q/Q^{\bullet-}) - E(Q/Q^{2-})]$ and stabilises the semi-quinone form. The yield of semi-quinone upon increasing the pH is predicted to be very low (Hay 2005).

The solution EPR spectra of CoQ_0 and NQ reveal complex hyperfine splitting centred around $g=2$. CoQ_0 did not resolve as many peaks as NQ and the yield of semi-quinone when placed into high pH buffer was much lower for CoQ_0 than NQ. The level of semi-quinone formed is greater for the NQ under the conditions used (i.e pH 9.5) due to the pK of NQ being higher than CoQ_0 . CoQ_0 is either forming the dianion (QH_2) or degrading at the high pH used.

4.3.2. Light Induced Effects

Mn^{2+} Oxidation State as Measured by EPR

The EPR spectra of the heterodimer with Mn bound showed less of the dimeric $Mn_2^{II,II}$ structure than did the BFR¹ homodimer. When compared to the homodimer the spectra was narrower suggesting less interaction of the Mn ions due to an increase in the distance separating them. This same phenomenon was found in ~50% of the homodimer metal centres when $ZnCe_6$ was bound but appears to occur in a greater number of centres in the heterodimer construct. The amount of bound monomeric Mn^{II} signal centred at $g=4$ is increased in the heterodimer again pointing to a decoupling of the manganese ions in the metal centre.

The BFR-H1 ZM complex showed some changes in the amount of $Mn_2^{II,II}$ dimer signal centred at $g=2$ upon illumination. Illumination produced a small decrease in the structured $Mn_2^{II,II}$ dimer signal, though no new signals were observed. It would be beneficial to use longer illumination times at room temperature to increase the amount of manganese oxidised (see Chapter 3). Longer illumination times should also produce new EPR signals as a result of Mn^{III} species forming.

Light Induced Chlorin Oxidation / Tyrosine Oxidation as Measured by EPR.

Illumination of the BFR-H1 heterodimer with no metal bound produced a large pronounced ZnCe_6 radical signal. When manganese was bound to the same protein then a broader signal was produced. Subtraction of the narrow ZnCe_6 radical from the broader signal reveals a structured radical signal (2.0 mT) centred at $g=2.0031$ (Figure 4.20). This broad radical signal appears to be the same signal as that found in the BFR^1 protein though the g value is slightly shifted and the turning points are not quite as well defined.

The protein environment is known to affect the g -value of tyrosine radicals (Un *et al.* 1995; Bleifuss *et al.* 2001; Hofbauer *et al.* 2001). There are some slight structural changes in the heterodimer protein as compared to BFR^1 so it is not surprising that the g -value of the tyrosine signal is slightly shifted. The presence of the tyrosine radical does confirm that the system is still intact and capable of carrying out light induced electron transfer.

Light Induced Oxygen Consumption of the BFR Metal Site

BFR functions in biology to bind iron and oxidize Fe_2^{II} to Fe_2^{III} forming a stabilizing μ -oxo bond between the two metal ions. This reaction consumes dissolved O_2 (see Chapter 3) (Liu *et al.* 2004). BFR^1 has been shown to be capable of binding manganese at the ferroxidase site and when ZnCe_6 is bound oxidising the bound Mn from 2+ to 3+. BFR^1 , BFR-H1 and BFR-H2 all consume significant amounts of oxygen upon illumination (Figure 4.21).

BFR-H1 and BFR-H2 consumed similar amounts of oxygen as found for the BFR^1 homodimer (Table 4.5). The addition of a quinone electron acceptor was found to significantly increase both the rate of uptake and the total oxygen consumed. A control of quinone in solution proved that the increase was not due to light effects on the quinone itself. The total amount of oxygen consumed far exceeded the other constructs indicating that more manganese centres were oxidised than in the other constructs. The increased oxygen uptake is likely through improvements in the efficiency of transferring the electron away from the ZnCe_6 pigment resulting in a longer lived charge-separated state. As suggested in Chapter 3 it is possible that coinciding with the extra oxygen

consumption more than one μ -oxo bond is formed. This hypothesis could in future be tested using EXAF measurements to look at bond lengths of the metal complex.

Steady State Fluorescence Properties

Comparison of the steady-state fluorescence spectra of BFR-H2 ZM CoQ₀ with the reference protein BFR-H2 ZM indicates significant fluorescence quenching due to the presence of the CoQ₀ moiety (Figure 4.22). The 22% quenching of the chlorin fluorescence by the quinone in its oxidized state is either due to non-radiative Forster energy transfer (the absence of ground-state coupling rules-out a Dexter mechanism) or electron transfer. Forster energy transfer is a mechanism of excitation transfer which can occur between molecular entities separated by distances considerably exceeding the sum of their van der Waals radii. It involves a dipole-dipole coupling mechanism between donor and acceptor. This form of energy transfer requires spectral overlap of the donor emission spectrum and the acceptor absorption spectrum. There is no detectable spectral overlap between the ZnCe₆ emission and quinone absorbance and as the quenching is only observed in the presence of the oxidised quinone, which is a renowned electron acceptor, the later explanation of electron transfer is favoured.

It follows from this that ~78% of the excited-state population decays through channels other than photo-induced electron transfer, such as internal conversion to the ground state and intersystem crossing to the triplet state whose lower energy compared to that of the singlet excited state may not provide sufficient driving force for the photo-induced electron transfer reaction.

The quenching of fluorescence as a result of electron transfer to an oxidised quinone was reported by Hay et al (2004). The Cyt *b*₅₆₂ quino-protein with a ZnCe₆ bound at the heme binding site showed a similar reduction in steady state fluorescence yield. Measurements of the fluorescence lifetimes quenching gave very similar efficiency values to that found by steady state fluorescence quenching (i.e. ~20%).

Having manganese bound to the BFR-H2 quino-protein was found to have very little effect on fluorescence yield. There was a very slight quenching when Mn was bound but this was hardly significant. This lack of effect was expected due to the fast back

reaction of the electron from the quinone to the ZnCe_6 and the slow movement of the electron hole to the metal centre (Rappaport *et al.* 1994).

EPR Spectra of BFR-H2 Quinone

Illumination of the BFR-H2 Z naphthoquinone protein produced a radical signal which was broader than a simple ZnCe_6 cation radical. Subtraction of some of the simple ZnCe_6 signal revealed a radical which was broader and with a higher g value centred at $g=2.0032$. This signal appeared to be the singly reduced semi-quinone radical (Figure 4.24).

This protein has no metal ions bound and as such we do not expect to see any tyrosine radical. This simplifies analysis of the EPR spectra as the only two possible radical signals are the ZnCe_6 cation radical and the semi-quinone anion radical signal. Subtraction of the ZnCe_6 portion of the spectra reveals a signal which could only be due to the singly reduced semi-quinone.

Looking at the solution EPR spectra of the same quino-protein but with manganese bound revealed some significant differences. The BFR-H2 ZM sample produces the expected mix of ZnCe_6 radical signal and an oxidised tyrosine radical. (Figure 4.25). When the NQ was bound the protein a much narrower $g=2$ EPR signal, remarkably similar to the BFR-H2 Z spectra, was produced with little tyrosine species detectable. Thus it appears that the binding of NQ to BFR-H2 results in the loss of a large percentage of the tyrosine radical species. It is expected that at least 50% of the tyrosine radical signal would be removed due to no metal ions binding in one half of the protein. There may also be changes in the metal binding site which could prevent the formation of the tyrosine radical which is dependent on Mn being bound and the subsequent deprotonation of the metal amino acid ligands (Conlan *et al.* 2007).

The complexity of the sample makes it very difficult to assign the radicals to one particular species. There are three potential radical signals all overlaid within these spectra. There is the ZnCe_6 cation radical, the tyrosine radical and the semi-quinone radical. All of these are centred around $g=2$ with only slight shifts due to the nature of the radicals. Further studies of the Mn containing heterodimer need to be carried out to visualise electron transfer to the quinone by EPR.

4.3.3. Electron Pathways Modelling

The optimal electron tunnelling pathway (Beratan *et al.* 1992) from ZnCe₆ to the bound quinone was calculated using the HARLEM program (Kurnikov and Beratan 1996). This program uses a modified superexchange coupling model developed by Beratan, Onuchic, and co-workers which accommodates the structural complexity of a protein matrix in electron transfer reactions (Onuchic and Beratan 1990; Beratan *et al.* 1991, 1992a; Onuchic *et al.* 1992)

In this tunnelling-pathway model, the medium between donor and acceptor is broken down into small subunits linked by covalent bonds (ϵ_C), hydrogen bonds (ϵ_H), or through-space jumps (ϵ_S). A coupling decay is assigned to each link, and a structure-dependent search algorithm is used to identify the optimum coupling pathway between the two redox sites. The electronic coupling matrix element (H_{AB}) reflects the strength of the interaction between reactants and products. The total coupling is given as a product of the couplings for the individual links (Equation 4-1):

$$H_{AB} \propto \prod \epsilon_C \prod \epsilon_H \prod \epsilon_S$$

Equation 4-1

The tunnelling-pathway model has proved to be a useful method for estimating distant electronic coupling and as such was used to compare the bacterioferritin system and the Cyt *b*₅₆₂ system of Hay and co-workers (Beratan *et al.* 1987; Onuchic and Beratan 1990; Onuchic *et al.* 1992; Hay *et al.* 2004; Hay 2005). The optimal electron tunnelling pathway from ZnCe₆ to the bound quinone was predicted to proceed via the axial ligand sulphur of Met51 and transfer along the protein backbone through Cys51 which is covalently attached to the quinone (Figure 4.27). Altogether the path includes 9 covalent bonds. This corresponds to an effective tunnelling length $\sigma_t = 15.8 \text{ \AA}$ and an electronic coupling decay factor, $H_{AB} = 6.05 \times 10^{-3}$. The direct ZnCe₆ to quinone distance $r = 10.6 \text{ \AA}$.

Figure 4.27: The calculated best pathway for electron transfer from the ZnCe_6 to the quinone in BFR-H2 ZM CoQ_0 . The electron likely travels along the axial ligand Met51 and then along Cys52 which is covalently linked to CoQ_0 .

When the previously characterised quino-protein of Hay *et al* (2004) was modelled (a protein proven to have high electron transfer efficiency), the coupling is theoretically less favourable than that of the BFR quino-protein. The best pathway coupling between donor and acceptor $H_{AB} = 7.84 \times 10^{-5}$ is much smaller than that of the BFR heterodimer system. In the Cyt b_{562} model the electron has an effective tunnelling length $\sigma_t = 11.3 \text{ \AA}$ and the straight ZnCe_6 to quinone distance $r = 9.5 \text{ \AA}$. Even though this quinone is placed $\sim 1.1 \text{ \AA}$ closer the smaller coupling is due to the electron having to transfer through space as opposed to along covalently bonded atoms (Figure 4.28). Tunnelling is much more efficient (decays more slowly) through bonded orbitals than through space because the effective potential barrier is lower.

Figure 4.28: The calculated best pathway for electron transfer from the ZnCe_6 to the quinone in the Cyt b_{562} quino-protein of Hay *et al* (2004). The electron likely travels from the ZnCe_6 outer carbon atoms through space to Met34 and then again through space to the oxygen of the methoxy group of the covalently linked to CoQ_0 .

Assuming that CoQ_0 is buried equally within the protein framework for both Cyt b_{562} and BFR-H2 then we could expect similar reorganisation energies for the system. The same porphyrin is utilised and the driving force of the system is likely to be similar. Under such conditions, we would expect better yields of electron transfer in the BFR quino-protein than that found in the Cyt b_{562} protein due to the electronic coupling being higher in the BFR system. Then again the BFR system may be limited to a single dominant pathway of electron transfer via the axial ligand which could generate a single-pathway limit and slower rates. (Prytkova *et al.* 2007). There remain many further measurements to be carried out on this system to determine the efficiency of electron transfer.

4.4. Conclusion

In conclusion, engineering a functional heterodimer was successful and analysis suggests that a linear electron transport chain was instigated. This is the first generation of a synthetic protein with photochemical charge separation involving a bound donor and acceptor. This scheme was borrowed from design principles of cofactors found in PSII. The quino-protein form of the heterodimer appears to carry out light induced electron transfer from the chlorin to the quinone with approximately 20% efficiency as suggested by fluorescence and EPR. The oxygen measurements combined with EPR suggest that the heterodimer quino-protein behaves similarly to the homodimer and should oxidise the manganese centre in the process.

Further work is needed on these constructs in order to fully characterise them. This will include spectroscopy of donor and acceptor function and will be continued in the future in the Photobioenergetics lab. It is hoped that these studies will be able to answer some of the many questions that have been raised in the study of these engineered proteins and refine the heterodimer reaction centre mimic of PSII.

CHAPTER 5

5 Conclusions and Future Work.

The work carried out in this thesis was aimed at studying light activated electron transfer within model protein systems. One of the goals was to capture light energy as stored chemical oxidation states with a long term view to use these stored oxidation states to oxidise water as occurs in Photosystem II. Using the electron donor side of Photosystem II as a guide, a homodimer of bacterioferritin from *E. coli* was bio-engineered to carry out light-induced electron transfer from a di-nuclear Mn complex to a photo-oxidized pigment. The new construct (BFR¹-ZM) was capable of using light energy to drive photochemical charge separation and couple the oxidising potential to multiple oxidation events at the bound Mn₂^{II} complex. This work is the first bioengineered protein capable of photo-catalytic, multi-electron oxidation of a di-manganese centre that has been reported in the literature.

Having shown that a protein can be modified to undergo a photochemical reaction, the next step was to design and construct a bacterioferritin heterodimer to mimic the PSII reaction centre. PSII is made up of a heterodimeric protein core which binds multiple cofactors necessary to create a light dependent charge separated state. The modified bacterioferritin is also a heterodimer with multiple cofactors capable of forming such a state. Both heterodimers bind light active pigments which transfer an electron to a quinone electron acceptor bound to the protein complex. Also both proteins have multiple symmetrically positioned tyrosine residues which are selectively oxidised during light activation. These tyrosines likely mediate transfer of electrons from the manganese centre.

The heterodimer proteins are a successful attempt at engineering a linear electron transport chain, which contains a number of the cofactors found in PSII. The quino-protein form of the heterodimer appears to carry out light induced electron transfer from the chlorin to the quinone with approximately 20% efficiency as suggested by fluorescence and EPR. The oxygen measurements combined with EPR suggest that the heterodimer quino-protein behaves similarly to the homodimer and should oxidise the manganese centre in the process.

More work needs to be completed on these constructs to fully characterise them. Nevertheless, the modified heterodimer of BFR promises to be a good model system for testing out the properties of PSII reactions, e.g.

- (1) If there are two nearby tyrosine donors to the oxidised pigment, is electron transfer purely distance dependent or controlled by intervening chemical bonds?
- (2) If a H-bond can be engineered to the tyrosine, how does this affect the redox potentials of the oxidation? Can a H-acceptor histidine be engineered.
- (3) If the di-metal site can be modified to bind more than two Mn ions within the complex, how does this affect the oxidation sequence?
- (4) What is the affect if two quinone electron acceptors are incorporated into the heterodimer.

The most immediate question to be answered is the nature of the tyrosine radical found in illuminated samples of the bacterioferritin complex with chlorin and manganese bound. Mutation of the three potentially redox active tyrosine's, found within close proximity of the chlorin and metal centre, should reveal which is oxidised during illumination. This is currently being studied in the Photobioenergetics lab with some interesting observations on the tyrosine story already emerging.

Limited attempts have been made in the generation of a protein-bound manganese centre that acts as an electron donor species. However numerous synthetic di-manganese compounds have been produced as models of the OEC, of which several are able to catalyse the oxidation of water to molecular oxygen (Limburg *et al.* 1999; Limburg *et al.* 2001; Poulsen *et al.* 2005; Tagore *et al.* 2008). The use of a di-manganese centre in our protein based system could therefore potentially facilitate the oxidation of water. Theoretically introduction of mutations to the BFR system that increases the number of hydrogen bonds to the $ZnCe_6$ could substantially increase the oxidation potential (Kalman *et al.* 2005). This could promote oxidation of the di-manganese centre to the higher oxidation states that are necessary for water splitting (Kurz *et al.* 2007).

The unique spherical dodecamer of dimers of the BFR model system has a large internal cavity of $\sim 8 \text{ \AA}$, which could potentially be used to compartmentalize reactions within the same molecular assembly. This could allow separation of oxidative and reductive processes across a protein interface. It is proposed that the internal cavity of the BFR sphere structure is anaerobic (Baaghil *et al.* 2003) in which case the addition of a hydrogenase site within the sphere would limit O_2 poisoning the hydrogenase catalyst. If for instance the Mn centre was capable of photo-catalytic oxidation of water this oxidative reaction could be coupled to reduction of protons to hydrogen gas on the inside of the complex.

The ultimate aim of this ongoing research is to produce an artificial catalyst whereby water is efficiently split into molecular oxygen, protons and electrons. To my knowledge this work is novel and the first example of a bioengineered protein capable of photo-catalytic, multi-electron oxidation of a dimanganese centre. This system should enable theoretical PSII problems to be tested on a much simpler protein complex with fewer cofactors to confuse analysis. This work represents one of the first steps in developing synthetic photo-catalytic, 'green enzymes' that utilize light energy to catalyse oxygen evolution and hydrogen production.

References

- Abragam A and Bleaney BI (1970) *Electron Paramagnetic Resonance of Transition Ions*. Oxford University Press, Oxford
- Ahmad I, Cusanovich MA and Tollin G (1981) Laser Flash-Photolysis Studies of Electron-Transfer between Semi-Quinone and Fully Reduced Free Flavins and Horse Heart Cytochrome-C. *Proc Natl Acad Sci USA* 78: 6724-6728
- Al'tshuler SA and Kozyrev BM (1974) *Electron Paramagnetic Resonance in Compounds of Transition Elements*. Halsted-Wiley, New York, NY, USA
- Alfonta L, Zhang ZW, Uryu S, Loo JA and Schultz PG (2003) Site-specific incorporation of a redox-active amino acid into proteins. *J Am Chem Soc* 125: 14662-14663
- Allen JP and Williams JC (1995) Relationship between the Oxidation Potential of the Bacteriochlorophyll Dimer and Electron-Transfer in Photosynthetic Reaction Centers. *J Bioenerg Biomembr* 27: 275-283
- Allen JP and Williams JC (1998) Photosynthetic reaction centers. *Febs Letters* 438: 5-9
- Andrews SC, Harrison PM and Guest JR (1989) Cloning, Sequencing, and Mapping of the Bacterioferritin Gene (bfr) of *Escherichia coli* K-12. *Journal of Bacteriology* 171: 3940-3947
- Andrews SC, Lebrun NE, Barynin V, Thomson AJ, Moore GR, Guest JR and Harrison PM (1995) Site-Directed Replacement of the Coaxial Heme Ligands of Bacterioferritin Generates Heme-Free Variants. *J Biol Chem* 270: 23268-23274
- Anthony C (2004) The quinoprotein dehydrogenases for methanol and glucose. *Arch Biochem Biophys* 428: 2-9
- Armstrong FA (2008) Why did nature choose manganese to make oxygen? *Philos Trans R Soc Lond B Biol Sci* 363: 1263-1270
- Arnesano F, Banci L, Bertini I, Faraone-Mennella J, Rosato A, Barker PD and Fersht AR (1999) The solution structure of oxidized *Escherichia coli* cytochrome b(562). *Biochemistry* 38: 8657-8670
- Baaghil S, Lewin A, Moore GR and Le Brun NE (2003) Core formation in *Escherichia coli* bacterioferritin requires a functional. *Biochemistry* 42: 14047-14056
- Baaghil S, Thomson AJ, Moore GR and Le Brun NE (2002) Studies of copper(II)-binding to bacterioferritin and its effect on iron(II) oxidation. *J Chem Soc Dalton*: 811-818
- Barker PD, Nerou EP, Cheesman MR, Thomson AJ, deOliveira P and Hill HAO (1996) Bis-methionine ligation to heme iron in mutants of cytochrome b(562) .1.

- Spectroscopic and electrochemical characterization of the electronic properties'. *Biochemistry* 35: 13618-13626
- Barynin VV, Whittaker MM, Antonyuk SV, Lamzin VS, Harrison PM, Artymiuk PJ and Whittaker JW (2001) Crystal structure of manganese catalase from *Lactobacillus plantarum*. *Structure* 9: 725-738
- Bauminger ER, Harrison PM, Hechel D, Nowik I and Treffry A (1991) Mossbauer Spectroscopic Investigation of Structure-Function Relations in Ferritins. *Biochim Biophys Acta* 1118: 48-58
- Bellelli A, Brzezinski P, Arese M, Cutruzzola F, Silvestrini MC and Brunori M (1996) Electron transfer in zinc-reconstituted nitrite reductase from *Pseudomonas aeruginosa*. *Biochem J* 319: 407-410
- Beratan DN, Onuchic JN, Winkler JR and Gray HB (1992) Electron-Tunneling Pathways in Proteins. *Science* 258: 1740-1741
- Blankenship RE (2002) *Molecular Mechanisms of Photosynthesis*. Blackwell Science Ltd
- Bleifuss G, Kolberg M, Potsch S, Hofbauer W, Bittl R, Lubitz W, Graslund A, Lassmann G and Lendzian F (2001) Tryptophan and tyrosine radicals in ribonucleotide reductase: A comparative high-field EPR study at 94 GHz. *Biochemistry* 40: 15362-15368
- Bou-Abdallah F, Arosio P, Levi S, Janus-Chandler C and Chasteen ND (2003) Defining metal ion inhibitor interactions with recombinant human H- and L-chain ferritins and site-directed variants: an isothermal titration calorimetry study. *J Biol Inorg Chem* 8: 489-497
- Boussac A, Rappaport F, Carrier P, Verbavatz JM, Gobin R, Kirilovsky D, Rutherford AW and Sugiura M (2004) Biosynthetic $\text{Ca}^{2+}/\text{Sr}^{2+}$ exchange in the photosystem II oxygen-evolving enzyme of *Thermosynechococcus elongatus*. *J Biol Chem* 279: 22809-22819
- Campbell KA, Force DA, Nixon PJ, Dole F, Diner BA and Britt RD (2000) Dual-mode EPR detects the initial intermediate in photoassembly of the photosystem II Mn cluster: The influence of amino acid residue 170 of the D1 polypeptide on Mn coordination. *J Am Chem Soc* 122: 3754-3761
- Campbell KA, Lashley MR, Wyatt JK, Nantz MH and Britt RD (2001) Dual-mode EPR study of Mn(III) salen and the Mn(III) salen-catalyzed epoxidation of cis-beta-methylstyrene. *J Am Chem Soc* 123: 5710-5719
- Campbell KA, Yikilmaz E, Grant CV, Gregor W, Miller AF and Britt RD (1999) Parallel polarization EPR characterization of the Mn(III) center of oxidized manganese superoxide dismutase. *J Am Chem Soc* 121: 4714-4715
- Carrondo MA (2003) Ferritins, iron uptake and storage from the bacterioferritin viewpoint. *The EMBO Journal* 22: 1959-1968

- Catanzariti A-M, Soboleva TA, Jans DA, Board PG and Baker RT (2004) An efficient system for high-level expression and easy purification of authentic recombinant proteins. *Protein Sci* 13: 1331-1339
- Cecchini G, Maklashina E, Yankovskaya V, Iverson TM and Iwata S (2003) Variation in proton donor/acceptor pathways in succinate : quinone oxidoreductases. *Febs Letters* 545: 31-38
- Chambers J (1988) The Chemistry of The Quinoid Compounds. In: Patai S and Rappaport Z (eds), pp 719-759. John Wiley and Sons
- Chang IJ, Gray HB and Winkler JR (1991) High-Driving-Force Electron-Transfer in Metalloproteins - Intramolecular Oxidation of Ferrocycytochrome-C by Ru(2,2'-Bpy)2(Im)(His-33)3+. *J Am Chem Soc* 113: 7056-7057
- Chen M, Eggink LL, Hooper JK and Larkum AWD (2005) Influence of structure on binding of chlorophylls to peptide ligands. *J Am Chem Soc* 127: 2052-2053
- Cheung E, Taylor K, Kornblatt JA, English AM, McLendon G and Miller JR (1986) Direct Measurements of Intramolecular Electron-Transfer Rates between Cytochrome-C and Cytochrome-C Peroxidase - Effects of Exothermicity and Primary Sequence on Rate. *Proc Natl Acad Sci USA* 83: 1330-1333
- Clausen J, Debus RJ and Junge W (2004) Time-resolved oxygen production by PSII: chasing chemical intermediates. *Biochim Biophys Acta* 1655: 184-194
- Conlan B (2008) Designing photosystem II: molecular engineering of photo-catalytic proteins. *Photosyn Res* 98: 687-700
- Conlan B, Wydrzynski T and Hillier W (2007) Designing artificial photosynthesis: Engineering bacterioferritin for light-activated electron transfer. *Photosyn Res* 91: 281-281
- Connick WB, DiBilio AJ, Hill MG, Winkler JR and Gray HB (1995) Tricarbonyl(1,10-phenanthroline)(imidazole)rhenium(I): A powerful photooxidant for investigations of electron tunneling in proteins. *Inorg Chim Acta* 240: 169-173
- Cowan JA (1996) *Inorganic Biochemistry - an introduction*. Wiley-VCH
- Cowan JA and Gray HB (1989) Synthesis and Properties of Metal-Substituted Myoglobins. *Inorg Chem* 28: 2074-2078
- Dasgupta J, Ananyev GM and Dismukes GC (2008) Photoassembly of the water-oxidizing complex in photosystem II. *Coord Chem Rev* 252: 347-360
- Dau H and Haumann M (2008) The manganese complex of photosystem II in its reaction cycle - Basic framework and possible realization at the atomic level. *Coord Chem Rev* 252: 273-295

- Dautant A, Meyer JB, Yariv J, Precigoux G, Sweet RM, Kalb AJ and Frolow F (1998) Structure of a monoclinic crystal form of cytochrome b1 (bacterioferritin) from E-coli. *Acta Crystallogr D* 54: 16-24
- Depew M and Wan J (1988) The Chemistry of the Quinoid Compounds. In: Patai S and Rappaport Z (eds), pp 899-962. John Wiley and Sons
- Dexheimer SL, Gohdes JW, Chan MK, Hagen KS, Armstrong WH and Klein MP (1989) Detection of Epr-Spectra in S = 2 States of Trivalent Manganese Complexes. *J Am Chem Soc* 111: 8923-8925
- Di Bilio AJ, Crane BR, Wehbi WA, Kiser CN, Abu-Omar MM, Carlos RM, Richards JH, Winkler JR and Gray HB (2001) Properties of photogenerated tryptophan and tyrosyl radicals in structurally characterized proteins containing rhenium(I) tricarbonyl diimines. *J Am Chem Soc* 123: 3181-3182
- Di Bilio AJ, Dennison C, Gray HB, Ramirez BE, Sykes AG and Winkler JR (1998) Electron transfer in ruthenium-modified plastocyanin. *J Am Chem Soc* 120: 7551-7556
- Diner BA and Britt RD (2005) The Redox-Active Tyrosines Y_Z and Y_D. In: Wydrzynski T and Satoh K (eds) *Photosystem II - The Light Driven Water:Plastoquinone Oxidoreductase*, pp 207-233. Springer, Dordrecht
- Dismukes GC (1996) Manganese enzymes with binuclear active sites. *Chem Rev* 96: 2909-2926
- Eggink LL and Hooper JK (2000) Chlorophyll binding to peptide maquettes containing a retention motif. *J Biol Chem* 275: 9087-9090
- Epel B, Schafer KO, Quentmeier A, Friedrich C and Lubitz W (2005) Multifrequency EPR analysis of the dimanganese cluster of the putative sulfate thiohydrolase SoxB of *Paracoccus pantotrophus*. *J Biol Inorg Chem* 10: 636-642
- Ferreira KN, Iverson TM, Maghlaoui K, Barber J and Iwata S (2004) Architecture of the photosynthetic oxygen-evolving center. *Science* 303: 1831-1838
- Fisher N and Rich PR (2000) A motif for quinone binding sites in respiratory and photosynthetic systems. *J Mol Biol* 296: 1153-1162
- Freire E, Mayorga OL and Straume M (1990) Isothermal Titration Calorimetry. *Analytical Chemistry* 62: A950-A959
- Frolow F, Kalb AJ and Yariv J (1994) Structure of a Unique Twofold Symmetrical Heme-Binding Site. *Nature Structural Biology* 1: 453-460
- Frolow FK, A.J. (2001) Handbook of metalloproteins. In: A. M (ed) *Handbook of metalloproteins*, pp 782-790. Wiley, New York
- Furukawa Y, Ishimori K and Morishima I (2000) Electron transfer reactions in Zn-substituted cytochrome P450cam. *Biochemistry* 39: 10996-11004

- Gerritsen HJ and Sabisky ES (1963) Paramagnetic Resonance of Trivalent Manganese in Rutile (TiO₂). *Physical Review* 132: 1507-&
- Golombek AP and Hendrich MP (2003) Quantitative analysis of dinuclear manganese(II) EPR spectra. *J Magn Reson* 165: 33-48
- Gouterman M (1961) Spectra of porphyrins. *Journal of Molecular Spectroscopy* 6: 138-163
- Grabolle M and Dau H (2005) Energetics of primary and secondary electron transfer in Photosystem II membrane particles of spinach revisited on basis of recombination-fluorescence measurements. *Biochim Biophys Acta* 1708: 209-218
- Gray HB and Winkler JR (1996) Electron transfer in proteins. *Annu Rev Biochem* 65: 537-561
- Gray HB and Winkler JR (2003) Electron tunneling through proteins. *Q Rev Biophys* 36: 341-372
- Griffith JS (1971) *Theory of Transition: Metal Ions*. Cambridge University Press
- Groni S, Dorlet P, Blain G, Bourcier S, Guillot R and Anxolabehere-Mallart E (2008) Reactivity of an aminopyridine [LMnII](2+) complex with H₂O₂. Detection of intermediates at low temperature. *Inorg Chem* 47: 3166-3172
- Gunner MR (1991) The Reaction Center Protein from Purple Bacteria - Structure and Function *Curr Top Bioenerg*, pp 319-367.
- Hamada K, Bethge PH and Mathews FS (1995) Refined Structure of Cytochrome B(562) from Escherichia-Coli at 1.4 Angstrom Resolution. *J Mol Biol* 247: 947-962
- Harrison PM and Arosio P (1996) Ferritins: Molecular properties, iron storage function and cellular regulation. *Biochim Biophys Acta* 1275: 161-203
- Haumann M, Liebisch P, Muller C, Barra M, Grabolle M and Dau H (2005) Photosynthetic O₂ formation tracked by time-resolved X-ray experiments. *Science* 310: 1019-1021
- Hay S (2005) Thesis - Protein Engineering of Novel Porphyrin/Quinone-Binding Proteins for Light-Induced Electron Transfer, Australian National University, Canberra
- Hay S, Wallace BB, Smith TA, Ghiggino KP and Wydrzynski T (2004) Protein engineering of cytochrome *b*₅₆₂ for quinone binding and light-induced electrons transfer. *Proc Natl Acad Sci USA* 101: 17675-17680
- Heathcote P, Fyfe PK and Jones MR (2002) Reaction centres: the structure and evolution of biological solar power. *Trends in Biochemical Sciences* 27: 79-87

- Hendrich MP and Debrunner PG (1988) Electron-Paramagnetic-Res Spectra of Quintet Ferrous Myoglobin and a Model Heme Compound. *J Magn Reson* 78: 133-141
- Hendrich MP and Debrunner PG (1989) Integer-Spin Electron-Paramagnetic Resonance of Iron Proteins. *Biophys J* 56: 489-506
- Hendry G and Wydrzynski T (2003) O-18 isotope exchange measurements reveal that calcium is involved in the binding of one substrate-water molecule to the oxygen-evolving complex in photosystem II. *Biochemistry* 42: 6209-6217
- Hersleth HP, Ryde U, Rydberg P, Gorbitz CH and Andersson KK (2006) Structures of the high-valent metal-ion haem-oxygen intermediates in peroxidases, oxygenases and catalases. *J Inorg Biochem* 100: 460-476
- Hillier W and Messinger J (2005) Mechanism of Photosynthetic Oxygen Production. In: Wydrzynski T and Satoh K (eds) *Photosystem II: The light-driven water:plastoquinone oxidoreductase.*, pp 567-608. Springer, Dordrecht
- Ho FM and Styring S (2008) Access channels and methanol binding site to the CaMn₄ cluster in Photosystem II based on solvent accessibility simulations, with implications for substrate water access. *Biochim Biophys Acta* 1777: 140-153
- Ho TF, McIntosh AR and Bolton JR (1980) Intramolecular Photochemical Electron-Transfer in a Linked Porphyrin-Quinone Molecule as a Model for the Primary Step of Photosynthesis. *Nature* 286: 254-256
- Hofbauer W, Zouni A, Bittl R, Kern J, Orth P, Lendzian F, Fromme P, Witt HT and Lubitz W (2001) Photosystem II single crystals studied by EPR spectroscopy at 94 GHz: The tyrosine radical Y-D(center dot). *Proc Natl Acad Sci USA* 98: 6623-6628
- Hoganson CW, Ghanotakis DF, Babcock GT and Yocum CF (1989) Mn-2+ Reduces Y_Z⁺ in Manganese-Depleted Photosystem-II Preparations. *Photosyn Res* 22: 285-293
- Hughes JL, Razeghifard R, Logue M, Oakley A, Wydrzynski T and Krausz E (2006) Magneto-optic spectroscopy of a protein tetramer binding two exciton-coupled chlorophylls. *J Am Chem Soc* 128: 3649-3658
- Iverson TM, Luna-Chavez C, Cecchini G and Rees DC (1999) Structure of the Escherichia coli fumarate reductase respiratory complex. *Science* 284: 1961-1966
- Jiang W, Yun D, Saleh L, Barr EW, Xing G, Hoffart LM, Maslak M-A, Krebs C and Bollinger JM, Jr. (2007) A Manganese(IV)/Iron(III) Cofactor in Chlamydia trachomatis Ribonucleotide Reductase. *Science* 316: 1188-1191
- Johnson ET and Parson WW (2002) Electrostatic interactions in an integral membrane protein. *Biochemistry* 41: 6483-6494

- Kalman L, LoBrutto R, Allen JP and Williams JC (1999) Modified reaction centres oxidize tyrosine in reactions that mirror photosystem II. *Nature* 402: 696-699
- Kalman L, LoBrutto R, Williams JC and Allen JP (2006) Iron as a bound secondary electron donor in modified bacterial reaction centers. *Biochemistry* 45: 13869-13874
- Kalman L, Williams J and Allen J (2005) Mimicking the properties of photosystem II in bacterial reaction centers. In: Wydrzynski T and Satoh K (eds) *Photosystem II: The light-driven water:plastoquinone oxidoreductase.* , pp 715-727. Springer, Dordrecht
- Kaneko Y, Tamura M and Yamazaki I (1980) Formation of Porphyrin-Pi Cation Radical in Zinc-Substituted Horseradish-Peroxidase. *Biochemistry* 19: 5795-5799
- Kehoe JW, Meadows KA, Parkes-Loach PS and Loach PA (1998) Reconstitution of core light-harvesting complexes of photosynthetic bacteria using chemically synthesized polypeptides. 2. Determination of structural features that stabilize complex formation and their implications for the structure of the subunit complex. *Biochemistry* 37: 3418-3428
- Klinman JP (2001) Commentary - How many ways to craft a cofactor? *Proc Natl Acad Sci USA* 98: 14766-14768
- Koder RL and Dutton PL (2006) Intelligent design: the de novo engineering of proteins with specified functions. *Dalton Trans*: 3045-3051
- Krieger-Liszka A and Rutherford AW (1998) Influence of Herbicide Binding on the Redox Potential of the Quinone Acceptor in Photosystem II: Relevance to Photodamage and Phytotoxicity. *Biochemistry* 37: 17339-17344
- Krieger A, Rutherford AW and Johnson GN (1995) On the determination of redox midpoint potential of the primary quinone electron acceptor, QA, in Photosystem II. *Biochim Biophys Acta* 1229: 193-201
- Krishtalik LI (1986) Energetics of Multielectron Reactions - Photosynthetic Oxygen Evolution. *Biochim Biophys Acta* 849: 162-171
- Kruse O, Rupprecht J, Mussgnug JR, Dismukes GC and Hankamer B (2005) Photosynthesis: a blueprint for solar energy capture and biohydrogen production technologies. *Photoch Photobio Sci* 4: 957-970
- Kulik LV, Epel B, Lubitz W and Messinger J (2007) Electronic Structure of the Mn4OxCa Cluster in the S0 and S2 States of the Oxygen-Evolving Complex of Photosystem II Based on Pulse 55Mn-ENDOR and EPR Spectroscopy. *J. Am. Chem. Soc.* 129: 13421-13435
- Kurnikov IV and Beratan DN (1996) Ab initio based effective Hamiltonians for long-range electron transfer: Hartree-Fock analysis. *Journal of Chemical Physics* 105: 9561-9573

- Kurz P, Berggren G, Anderlund MF and Styring S (2007) Oxygen evolving reactions catalysed by synthetic manganese complexes: A systematic screening. *Dalton Trans*: 4258-4261
- Ladbury JE and Chowdhry BZ (1996) Sensing the heat: The application of isothermal titration calorimetry to thermodynamic studies of biomolecular interactions. *Chem Biol* 3: 791-801
- Leavitt S and Freire E (2001) Direct measurement of protein binding energetics by isothermal titration calorimetry. *Curr Opin Struct Biol* 11: 560-566
- LeBrun NE, Andrews SC, Guest JR, Harrison PM, Moore GR and Thomson AJ (1995a) Identification of the Ferroxidase Center of Escherichia-Coli Bacterioferritin. *Biochem J* 312: 385-392
- LeBrun NE, Keech AM, Mauk MR, Mauk AG, Andrews SC, Thomson AJ and Moore GR (1996) Charge compensated binding of divalent metals to bacterioferritin: H⁺ release associated with cobalt(II) and zinc(II) binding at dinuclear metal sites. *Febs Letters* 397: 159-163
- LeBrun NE, Moore GR and Thomson AJ (1995b) Magnetic Circular-Dichroism Spectroscopy of the Iron Cores of Ferritin and Bacterioferritin. *Molecular Physics* 85: 1061-1068
- Lewis NS and Nocera DG (2006) Powering the planet: Chemical challenges in solar energy utilization. *Proc Natl Acad Sci USA* 103: 15729-15735
- Li WW, Heinze J and Haehnel W (2005) Site-specific binding of quinones to proteins through thiol addition and addition-elimination reactions. *J Am Chem Soc* 127: 6140-6141
- Li WW, Hellwig P, Ritter M and Haehnel W (2006) De novo design, synthesis, and characterization of quinoproteins. *Chem-Eur J* 12: 7236-7245
- Limburg J, Vrettos JS, Chen HY, de Paula JC, Crabtree RH and Brudvig GW (2001) Characterization of the O₂-evolving reaction catalyzed by [(terpy)(H₂O)Mn-III(O)(2)Mn-IV(OH₂)(terpy)](NO₃) (terpy=2,2':6,2''-terpyridine). *J Am Chem Soc* 123: 423-430
- Limburg J, Vrettos JS, Liable-Sands LM, Rheingold AL, Crabtree RH and Brudvig GW (1999) A functional model for O-O bond formation by the O₂-evolving complex in photosystem II. *Science* 283: 1524-1527
- Lin RJ, Lin KS and Chang IJ (1996) Photophysical properties of tricarbonyl(histidine)(diimine)rhenium(I) complexes in aqueous solution. *Inorg Chim Acta* 242: 179-183
- Lin X, Murchison HA, Nagarajan V, Parson WW, Allen JP and Williams JC (1994) Specific Alteration of the Oxidation Potential of the Electron-Donor in Reaction

- Centers from Rhodobacter-Sphaeroides. *Proc Natl Acad Sci USA* 91: 10265-10269
- Liu HL, Zhou HN, Xing WM, Zhao HF, Li SX, Huang JF and Bi RC (2004) 2.6 angstrom resolution crystal structure of the bacterioferritin from *Azotobacter vinelandii*. *Febs Letters* 573: 93-98
- Loll B, Kern J, Saenger W, Zouni A and Biesiadka J (2005) Towards complete cofactor arrangement in the 3.0 angstrom resolution structure of photosystem II. *Nature* 438: 1040-1044
- Magner E and McLendon G (1989) Ground-State and Excited-State Electron-Transfer Reactions of Zinc Cytochrome-C. *J Phys Chem* 93: 7130-7134
- Marcus RA (1956) On the Theory of Oxidation-Reduction Reactions Involving Electron Transfer. I. *J Chem Phys* 24: 966-978
- Marcus RA and Sutin N (1985) Electron transfers in chemistry and biology. *Biochim Biophys Acta, Rev Bioenerg* 811: 265-322
- Martin SR and Schilstra MJ (2008) Circular dichroism and its application to the study of biomolecules. *Biophysical Tools for Biologists: Vol 1 In Vitro Techniques*, pp 263-293. Elsevier Academic Press Inc, San Diego
- McEvoy JP and Brudvig GW (2006) Water-splitting chemistry of photosystem II. *Chem Rev* 106: 4455-4483
- McEvoy JP, Gascon JA, Batista VS and Brudvig GW (2005) The mechanism of photosynthetic water splitting. *Photoch Photobio Sci* 4: 940-949
- McGourty JL, Petersonkennedy SE, Ruo WY and Hoffman BM (1987) Characterization of Long-Range Electron-Transfer in Mixed-Metal [Zinc,Iron] Hybrid Hemoglobins. *Biochemistry* 26: 8302-8312
- McGregor WC, Swierczek SI, Bennett B and Holz RC (2007) Characterization of the catalytically active Mn(II)-loaded argE-encoded N-acetyl-L-ornithine deacetylase from *Escherichia coli*. *J Biol Inorg Chem* 12: 603-613
- McLendon GL, Winkler JR, Nocera DG, Mauk MR, Mauk AG and Gray HB (1985) Quenching of Zinc-Substituted Cytochrome-C Excited-States by Cytochrome-B5. *J Am Chem Soc* 107: 739-740
- Meadows KA, Parkes-Loach PS, Kehoe JW and Loach PA (1998) Reconstitution of core light-harvesting complexes of photosynthetic bacteria using chemically synthesized polypeptides. 1. Minimal requirements for subunit formation. *Biochemistry* 37: 3411-3417
- Mennenga A, Gartner W, Lubitz W and Gerner H (2006) Effects of noncovalently bound quinones on the ground and triplet states of zinc chlorins in solution and bound to de novo synthesized peptides. *Phys Chem Chem Phys* 8: 5444-5453

- Messinger J, Badger M and Wydrzynski T (1995) Detection of One Slowly Exchanging Substrate Water Molecule in the S-3 State of Photosystem-II. *Proc Natl Acad Sci USA* 92: 3209-3213
- Messinger J, Robblee JH, Bergmann U, Fernandez C, Glatzel P, Visser H, Cinco RM, McFarlane KL, Bellacchio E, Pizarro SA, Cramer SP, Sauer K, Klein MP and Yachandra VK (2001) Absence of Mn-centered oxidation in the S-2 → S-3 Transition: Implications for the mechanism of photosynthetic water oxidation. *J Am Chem Soc* 123: 7804-7820
- Miller JE, Di Bilio AJ, Wehbi WA, Green MT, Museth AK, Richards JR, Winkler JR and Gray HB (2004) Electron tunneling in rhenium-modified *Pseudomonas aeruginosa* azurins. *Biochim Biophys Acta* 1655: 59-63
- Miloslavina Y, Szczepaniak M, Muller MG, Sander J, Nowaczyk M, Rogner M and Holzwarth AR (2006) Charge separation kinetics in intact photosystem II core particles is trap-limited. A picosecond fluorescence study. *Biochemistry* 45: 2436-2442
- Moore GR (1991) Bacterial 4-Alpha-Helical Bundle Cytochromes. *Biochim Biophys Acta* 1058: 38-41
- Moore GR, Williams RJP, Peterson J, Thomson AJ and Mathews FS (1985) A Spectroscopic Investigation of the Structure and Redox Properties of *Escherichia-Coli* Cytochrome B-562. *Biochim Biophys Acta* 829: 83-96
- Moser C and Dutton L (1996) Outline of theory of protein electron transfer. In: Bendall DS (ed) *Protein Electron Transfer*, pp 300. Bios Scientific Publishers, Oxford, UK
- Moser C and Dutton PL (1992) Engineering protein structure for electron transfer function in photosynthetic reaction centers. *Biochim Biophys Acta* 1101: 171-176
- Mure M (2004) Tyrosine-derived quinone cofactors. *Acc Chem Res* 37: 131-139
- Murray JW and Barber J (2007) Structural characteristics of channels and pathways in photosystem II including the identification of an oxygen channel. *J Struct Biol* 159: 228-237
- Narvaez AJ, Kalman L, LoBrutto R, Allen JP and Williams JC (2002) Influence of the protein environment on the properties of a tyrosyl radical in reaction centers from *Rhodobacter sphaeroides*. *Biochemistry* 41: 15253-15258
- Noy D, Discher BM, Rubtsov IV, Hochstrasser RA and Dutton PL (2005) Design of amphiphilic protein maquettes: Enhancing maquette functionality through binding of extremely hydrophobic cofactors to lipophilic domains. *Biochemistry* 44: 12344-12354

- Noy D and Dutton PL (2006) Design of a minimal polypeptide unit for bacteriochlorophyll binding and self-assembly based on photosynthetic bacterial light-harvesting proteins. *Biochemistry* 45: 2103-2113
- Petersonkennedy SE, McGourty JL and Hoffman BM (1984) Temperature-Dependence of Long-Range Electron-Transfer in [Zn,Fe-III] Hybrid Hemoglobin. *J Am Chem Soc* 106: 5010-5012
- Pierce MM, Raman CS and Nall BT (1999) Isothermal titration calorimetry of protein-protein interactions. *Methods* 19: 213-221
- Polak E and Ribiere G (1969) Note on Convergence of Conjugate Direction Methods. *Revue Francaise D Informatique De Recherche Operationnelle* 3: 35-&
- Poulsen AK, Rompel A and McKenzie CJ (2005) Water oxidation catalyzed by a dinuclear Mn complex: A functional model for the oxygen-evolving center of photosystem II. *Angew Chem Int Edit* 44: 6916-6920
- Prytkova TR, Beratan DN and Skourtis SS (2007) Photoselected electron transfer pathways in DNA photolyase. *Proc Natl Acad Sci USA* 104: 802-807
- Qin L and Kostic NM (1994) Photoinduced Electron-Transfer from the Triplet-State of Zinc Cytochrome-C to Ferricytochrome B5 Is Gated by Configurational Fluctuations of the Diprotein Complex. *Biochemistry* 33: 12592-12599
- Rappaport F, Blancharddesce M and Lavergne J (1994) Kinetics of Electron-Transfer and Electrochromic Change During the Redox Transitions of the Photosynthetic Oxygen-Evolving Complex. *Biochim Biophys Acta* 1184: 178-192
- Rappaport F and Diner BA (2008) Primary photochemistry and energetics leading to the oxidation of the (Mn)₄Ca cluster and to the evolution of molecular oxygen in Photosystem II. *Coord Chem Rev* 252: 259-272
- Rau HK, Snigula H, Struck A, Robert B, Scheer H and Hachnel W (2001) Design, synthesis and properties of synthetic chlorophyll proteins. *Eur J Biochem* 268: 3284-3295
- Razeghifard AR and Wydrzynski T (2003) Binding of Zn-chlorin to a synthetic four-helix bundle peptide through histidine ligation. *Biochemistry* 42: 1024-1030
- Redfearn E (1965) *Biochemistry of Quinones*. Academic Press, New York
- Reed G and Poyner R (2000) Chapter 7: Mn²⁺ as a Probe of Divalent Metal Ion Binding and Function in Enzymes and Other Proteins. In: Sigel HS, A. (ed) *Manganese and its role in biological processes*, pp 183-207. Marcel Dekker, Inc., New York
- Rehm D and Weller A (1970) Kinetics of Fluorescence Quenching by Electron and H-Atom Transfer. *Isr J Chem* 8: 259
- Rex JH (2000) Purification of genomic DNA from Gram-Negative bacteria. *Focus* 22: 26-27

- Rich PR (2004) The quinone chemistry of bc complexes 13th European Bioenergetics Conference (EBEC 2004). Elsevier Science Bv, Pisa, ITALY
- Robinson CR, Liu YF, Thomson JA, Sturtevant JM and Sligar SG (1997) Energetics of heme binding to native and denatured states of cytochrome b(562). *Biochemistry* 36: 16141-16146
- Sato H, Uchida A, Nakayama K and Okada M (2001) Water-soluble chlorophyll protein in Brassicaceae plants is a stress-induced chlorophyll-binding protein. *Plant Cell Physiol* 42: 906-911
- Sauer K, Yano J and Yachandra VK (2005) X-ray spectroscopy of the Mn₄Ca cluster in the water-oxidation complex of Photosystem II. *Photosyn Res* 85: 73-86
- Schaab MR, Barney BM and Francisco WA (2006) Kinetic and spectroscopic studies on the quercetin 2,3-dioxygenase from *Bacillus subtilis*. *Biochemistry* 45: 1009-1016
- Schafer KO, Bittl R, Lendzian F, Barynin V, Weyhermuller T, Wieghardt K and Lubitz W (2003) Multifrequency EPR investigation of dimanganese catalase and related Mn(III)Mn(IV) complexes. *J Phys Chem B* 107: 1242-1250
- Sharp RE, Moser CC, Rabanal F and Dutton PL (1998) Design, synthesis, and characterization of a photoactivatable flavocytochrome molecular maquette. *Proc Natl Acad Sci USA* 95: 10465-10470
- Siegbahn PEM (2006) O-O bond formation in the S-4 state of the oxygen-evolving complex in photosystem II. *Chem-Eur J* 12: 9217-9227
- Siegbahn PEM (2008) Mechanism and energy diagram for O-O bond formation in the oxygen-evolving complex in photosystem II. *Philos Trans R Soc Lond B Biol Sci* 363: 1221-1228
- Simolo KP, McLendon GL, Mauk MR and Mauk AG (1984) Photoinduced Electron-Transfer within a Protein Protein Complex Formed between Physiological Redox Partners - Reduction of Ferricytochrome-B5 by the Hemoglobin Derivative Alpha-2zn-Beta-2fe-Iiicn. *J Am Chem Soc* 106: 5012-5013
- Snell J and Weissberger A (1939) The reaction of thiol compounds with quinones. *J Am Chem Soc*: 450-453
- Springs SL, Bass SE, Bowman G, Nodelman I, Schutt CE and McLendon GL (2002) A multigeneration analysis of cytochrome b(562) redox variants: Evolutionary strategies for modulating redox potential revealed using a library approach. *Biochemistry* 41: 4321-4328
- Su JH, Lubitz W and Messinger J (2008) Probing mode and site of substrate water binding to the oxygen-evolving complex in the S-2 state of photosystem II by O-17-HYSCORE Spectroscopy. *J Am Chem Soc* 130: 786-787

- Tagore R, Crabtree RH and Brudvig GW (2008) Oxygen Evolution Catalysis by a Dimanganese Complex and Its Relation to Photosynthetic Water Oxidation. *Inorg. Chem.* 47: 1815-1823
- Tang XS, Zheng M, Chisholm DA, Dismukes GC and Diner BA (1996) Investigation of the differences in the local protein environments surrounding tyrosine radicals Y-Z(center dot) and Y-D(center dot) in photosystem II using wild-type and the D2-Tyr160Phe mutant of *Synechocystis* 6803. *Biochemistry* 35: 1475-1484
- Teale FWJ (1959) Cleavage of the Haem-Protein Link by Acid Methylethylketone. *Biochim Biophys Acta* 35: 543-543
- Teutloff C, Schafer KO, Sinnecker S, Barynin V, Bittl R, Wieghardt K, Lendzian F and Lubitz W (2005) High-field EPR investigations of Mn(III)Mn(IV) and Mn(II)Mn(III) states of dimanganese catalase and related model systems. *Magn Reson Chem* 43 Spec no.: S51-64
- Thielges M, Uyeda G, Camara-Artigas A, Kalman L, Williams JC and Allen JP (2005) Design of a redox-linked active metal site: Manganese bound to bacterial reaction centers at a site resembling that of photosystem II. *Biochemistry* 44: 7389-7394
- Tommos C, Skalicky JJ, Pilloud DL, Wand AJ and Dutton PL (1999) De novo proteins as models of radical enzymes. *Biochemistry* 38: 9495-9507
- Twitchett MB, Ferrer JC, Siddarth P and Mauk AG (1997) Intramolecular Electron Transfer Kinetics of a Synthetic Flavocytochrome c. *J. Am. Chem. Soc.* 119: 435-436
- Tyryshkin AM, Watt RK, Baranov SV, Dasgupta J, Hendrich MP and Dismukes GC (2006) Spectroscopic evidence for Ca²⁺ involvement in the assembly of the Mn₄Ca cluster in the photosynthetic water-oxidizing complex. *Biochemistry* 45: 12876-12889
- Un S, Atta M, Fontecave M and Rutherford AW (1995) G-Values as a Probe of the Local Protein Environment - High-Field Epr of Tyrosyl Radicals in Ribonucleotide Reductase and Photosystem-II. *J Am Chem Soc* 117: 10713-10719
- Vanderkooi JM, Adar F and Erecinska M (1976) Metallocytochromes c : Characterization of Electronic Absorption and Emission Spectra of Sn⁴⁺ and Zn²⁺ Cytochromes c. *Eur J Biochem* 64: 381-387
- Warncke K, Gunner MR, Braun BS, Gu LQ, Yu CA, Bruce JM and Dutton PL (1994) Influence of Hyrdocarbon Tail Structure on Quinone Binding and Electron-Transfer Performance at the Q_A and Q_B Sites of the Photosynthetic Reaction-Center Protein. *Biochemistry* 33: 7830-7841
- Weng TC, Hsieh WY, Uffelman ES, Gordon-Wylie SW, Collins TJ, Pecoraro VL and Penner-Hahn JE (2004) XANES evidence against a manganyl species in the S-3 state of the oxygen-evolving complex. *J Am Chem Soc* 126: 8070-8071

- Whiting AK, Boldt YR, Hendrich MP, Wackett LP and Que L (1996) Manganese(II)-dependent extradiol-cleaving catechol dioxygenase from *Archrobacter globiformis* CM-2. *Biochemistry* 35: 160-170
- Whitney SM and Sharwood RE (2006) Linked rubisco subunits can assemble into functional oligomers without impeding catalytic performance. *J. Biol. Chem.*: M610479200
- Whittaker JW and Whittaker MM (1991) Active-Site Spectral Studies on Manganese Superoxide-Dismutase. *J Am Chem Soc* 113: 5528-5540
- Whittaker MM, Barynin VV, Igarashi T and Whittaker JW (2003) Outer sphere mutagenesis of *Lactobacillus plantarum* manganese catalase disrupts the cluster core - Mechanistic implications. *Eur J Biochem* 270: 1102-1116
- Williams JC, Haffa ALM, McCulley JL, Woodbury NW and Allen JP (2001) Electrostatic interactions between charged amino acid residues and the bacteriochlorophyll dimer in reaction centers from *Rhodobacter sphaeroides*. *Biochemistry* 40: 15403-15407
- Winkler JR, Di Bilio AJ, Farrow NA, Richards JH and Gray HB (1999) Electron tunneling in biological molecules. *Pure Appl Chem* 71: 1753-1764
- Winkler JR and Gray HB (1997) Electron tunneling in proteins: Role of the intervening medium. *J Biol Inorg Chem* 2: 399-404
- Wraight CA (2004) Proton and electron transfer in the acceptor quinone complex of photosynthetic reaction centers from *Rhodobacter sphaeroides*. *Front. Biosci.* 9: 309-337
- Wu AJ, Penner-Hahn JE and Pecoraro VL (2004) Structural, spectroscopic, and reactivity models for the manganese catalases. *Chem Rev* 104: 903-938
- Wydrzynski T, Hillier W and Conlan B (2007) Engineering model proteins for Photosystem II function. *Photosyn Res* 94: 225-233
- Wydrzynski T, Hillier W and Messinger J (1996) On the functional significance of substrate accessibility in the photosynthetic water oxidation mechanism. *Physiol Plant* 96: 342-350
- Wydrzynski T and Satoh K (2005) Photosystem II : the light-driven water:plastoquinone oxidoreductase. Springer, Dordrecht
- Yachandra VK (2005) The Catalytic Manganese Cluster: Organization of the Metal Ions. In: Satoh K and Wydrzynski T (eds) Photosystem II - The light driven water:plastoquinone oxidoreductase, pp 235-260. Springer, Dordrecht
- Yang X, Le Brun NE, Thomson AJ, Moore GR and Chasteen ND (2000) The Iron Oxidation and Hydrolysis Chemistry of *Escherichia coli* Bacterioferritin. *Biochemistry* 39: 4915-4923

- Yariv J, Kalb AJ, Sperling R, Bauminger ER, Cohen SG and Ofer S (1981) The Composition and the Structure of Bacterioferritin of *Escherichia-Coli*. *Biochem J* 197: 171-175
- Yoder DW, Hwang J and Penner-Hahn JE (2000) Chapter 16: Manganese Catalases. In: Sigel HS, A. (ed) *Manganese and Its Role in Biological Processes* pp 527-557. Marcell Dekker, Inc., New York
- Zhang B, Harb JN, Davis RC, Kim JW, Chu SH, Choi S, Miller T and Watt GD (2005) Kinetic and thermodynamic characterization of the cobalt and manganese oxyhydroxide cores formed in horse spleen ferritin. *Inorg Chem* 44: 3738-3745
- Zhang LY, Wang LJ, Kao YT, Qiu WH, Yang Y, Okobiah O and Zhong DP (2007) Mapping hydration dynamics around a protein surface. *Proc Natl Acad Sci USA* 104: 18461-18466