ANALYSIS & PDE

Volume 9

No. 6

2016

ELINA ANDRIYANOVA AND SHIBING CHEN

**BOUNDARY $C^{1,\alpha}$ REGULARITY OF POTENTIAL FUNCTIONS
IN OPTIMAL TRANSPORTATION WITH QUADRATIC COST**

BOUNDARY $C^{1,\alpha}$ REGULARITY OF POTENTIAL FUNCTIONS IN OPTIMAL TRANSPORTATION WITH QUADRATIC COST

ELINA ANDRIYANOVA AND SHIBING CHEN

We provide a different proof for the global $C^{1,\alpha}$ regularity of potential functions in the optimal transport problem, which was originally proved by Caffarelli. Our method applies to a more general class of domains.

1. Introduction

We study the global $C^{1,\alpha}$ regularity of potential functions in optimal transportation with quadratic cost. Let Ω and Ω^* be the source and target domains associated with densities $1/C < f < C$ and $1/C < g < C$, respectively, where C is a positive constant. The optimal transport problem with quadratic cost is about finding a map $T : \Omega \rightarrow \Omega^*$ among all measure-preserving maps minimizing the transportation cost

$$\int_{\Omega} |x - Tx|^2 dx.$$

Here the term “measure-preserving” means that $\int_{T^{-1}(B)} f = \int_B g$ for any Borel set $B \subset \Omega^*$. Brenier [1991] proved that one can find a convex function u such that

$$T(x) = Du(x) \quad \text{for a.e. } x \in \Omega.$$

Indeed, the convex function u satisfies $\int_{(\partial u)^{-1}B} f = \int_B g$ for any Borel set $B \subset \Omega$, where ∂u is the standard subgradient map of the convex function u . We call u a *Brenier solution* of the optimal transport problem if it satisfies the property above. When the target domain Ω^* is convex, Caffarelli proved that $\partial u(\Omega) = \Omega^*$ and that u is an *Alexandrov solution*, namely u satisfies $(1/C)|A \cap \Omega| \leq |\partial u(A)| \leq C|A \cap \Omega|$ for any Borel set $A \subset \Omega$. Moreover, if we extend u to $\mathbb{R}^n$ via

$$\tilde{u} := \sup\{L \mid L \text{ is linear, } L|_{\Omega} \leq u, L(z) = u(z) \text{ for some } z \in \Omega\},$$

then $\tilde{u}$ is a globally Lipschitz convex solution of

$$C^{-1}\chi_{\Omega} \leq \det \tilde{u}_{ij} \leq C\chi_{\Omega}.$$

We will still use u to denote this extended function. Caffarelli [1992b] proved interior $C^{1,\alpha}$ regularity by using his techniques for studying the standard Monge–Ampère-type equation; see [Caffarelli 1990a; 1990b; 1991].

This work is supported by the Australian Research Council. The authors would like thank Professor Xu-Jia Wang for sharing his proof of Lemma 2.5.

MSC2010: 35B45, 35J60, 49Q20, 35J96.

Keywords: optimal transport, quadratic cost, boundary regularity.

Then, Caffarelli [1992a] proved the boundary $C^{1,\alpha}$ regularity result under the condition that both Ω and Ω^* are convex. Below we will briefly discuss the main ideas involved in his proof. First, Caffarelli established a fundamental property of convex functions, namely the existence of sections centred at a given point (see the statement of Lemma 2.5). Then, he proved that such sections are decaying geometrically, namely there exists a constant δ such that

$$S_{\delta h}(y) \subset \frac{3}{4}S_h(x) \quad \text{for any } y \in \frac{1}{2}S_h(x). \quad (1-1)$$

Here $S_h(x)$ denotes the section of u centred at x with height h . From (1-1) we obtain the quantitative strict convexity estimate

$$u(z) \geq u(x) + Du(x) \cdot (z - x) + C|z - x|^\beta \quad \text{for any } x, z \in \bar{\Omega}, \quad (1-2)$$

for some $\beta > 1$. From (1-2), it is easy to check that u^* , the standard Legendre transform of u , is $C^{1,\alpha}$ on $\bar{\Omega}^*$. Recall the well-known fact that u^* is indeed the potential function of the optimal transport problem from Ω^* to Ω . Therefore, by switching the role of u and u^* one can show the global $C^{1,\alpha}$ regularity of u .

The convexity of domains is crucial in Caffarelli's approach. Indeed, the convexity of Ω ensures that u^* is an Alexandrov solution, while the convexity of Ω^* ensures that the sections of u^* , centred at some point in $\bar{\Omega}^*$, have some doubling property. Here we provide a different proof of the global $C^{1,\alpha}$ result. Instead of deducing the $C^{1,\alpha}$ regularity of u from the strict convexity of u^* , we prove the $C^{1,\alpha}$ regularity of u directly. Moreover, our method works for a slightly more general class of domains, namely we allow the source to be a domain obtained by removing finitely many disjoint convex subsets from a convex domain.

We would like to mention that in recent years the regularity of optimal transport maps has attracted much interest and there are many important works related to it; to cite a few, see [Figalli and Loeper 2009; Liu 2009; Trudinger and Wang 2009b; 2009a; Figalli and Rifford 2009; Loeper 2011; Loeper and Villani 2010; Liu et al. 2010; Kim and McCann 2010; Figalli et al. 2010; 2011; 2012; 2013a; 2013b].

The rest of the paper is organized as follows. In Section 2 we introduce some notations and preliminaries, and state the main results. Section 3 is devoted to the proof of global C^1 regularity. In the last section we complete the proof of the main results.

2. Preliminaries and main result

The main result of this paper is the following theorem:

Theorem 2.1. *Let Ω and Ω^* be two bounded domains in $\mathbb{R}^n$, $n \geq 2$, and f and g be densities of two positive probability measures defined in Ω and Ω^* , respectively, satisfying $C^{-1} \leq f, g \leq C$ for a positive constant C . Assume that Ω^* is convex and Ω is Lipschitz.*

- (i) *If, for any given $x \in \bar{\Omega}$, there exists a small ball $B_{r_x}(x)$ such that, for any convex set $\omega \subset B_{r_x}(x)$ centred in Ω , we have $\int_\omega f \leq C \int_{\omega/2} f$ for some constant C independent of ω , then the potential function u is $C^1(\bar{\Omega})$. (Here f is defined to be 0 outside Ω .)*
- (ii) *If Ω is a domain obtained by removing finitely many disjoint convex subsets from a convex set, then the potential function u is $C^{1,\alpha}(\bar{\Omega})$ for some $\alpha \in (0, 1)$.*

Remark 2.2. (a) It is easy to see that in [Theorem 2.1\(i\)](#) we allow Ω to be any polytope (not necessarily convex). We also note that the C^1 regularity always holds in dimension two without any condition on Ω . This is a classical result of Alexandrov; see also [\[Figalli and Loeper 2009\]](#).

(b) One may want to prove higher regularity when the densities are smooth; however, in view of the following simple example we see that this is impossible. Let the dimension be $n = 2$. Let $\Omega := B_2 - B_1$, with uniform probability density, and let $\Omega^* := B_{\sqrt{3}}$, with uniform probability density. Then by symmetry it is easy to compute that the optimal transport map is $T(x) = \sqrt{|x|^2 - 1} x/|x|$, which is only $C^{1/2}$ on $\partial B_1 \subset \partial \Omega$.

In the following we will use $S_h(x_0)$ to denote a section of u with height h , namely

$$S_h(x_0) := \{x \mid u < p \cdot (x - x_0) + h\},$$

where p is chosen so that x_0 is the centre of mass of $S_h(x_0)$. We say a point $x_0 \in \bar{\Omega}$ is localized (with respect to u) if, for any sequences $h_k \rightarrow 0$ and $x_k \rightarrow x_0$ satisfying $x_0 \in S_{h_k}(x_k)$, we have that $S_{h_k}(x_k)$ shrinks to the point $x_0 \in \bar{\Omega}$.

Now we record a fundamental property of convex sets.

Lemma 2.3 (John's lemma). *Let $U \subset \mathbb{R}^n$ be a bounded, convex domain with its centre of mass at the origin. There exists an ellipsoid E , also centred at the origin, such that*

$$E \subset U \subset n^{3/2} E.$$

The original John's lemma does not require that the ellipsoid is centered at the origin, and the constant $n^{3/2}$ can be replaced by n . We refer the reader to [\[Liu and Wang 2015\]](#) for a simple proof of the existence and uniqueness of such an ellipsoid.

By John's lemma we can show the following property of convex functions:

Lemma 2.4. *Let $u : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function. Let L be a supporting function of u . Then any extreme point of $\{u = L\}$ is localized.*

Proof. Suppose to the contrary that there exists an extreme point x_0 of $\{u = L\}$ which is not localized. Then there exist sequences $x_k \rightarrow x_0$ and $h_k \rightarrow 0$ such that $x_0 \in S_{h_k}(x_k)$, and that $S_{h_k}(x_k)$ contains a segment of length greater than or equal to some positive constant δ . Since $S_{h_k}(x_k)$ is convex and centred at x_k , by John's lemma there exists a unit vector ξ_k such that I_k , the segment connecting $x_k - \delta/(2n^{3/2}) \xi_k$ and $x_k + \delta/(2n^{3/2}) \xi_k$, is contained in $S_{h_k}(x_k)$. Denote by L_k the defining function of $S_{h_k}(x_k)$, namely $S_{h_k}(x_k) = \{u \leq L_k\}$. Then it is easy to see that DL_k is bounded; hence, by passing to a subsequence, $L_k \rightarrow L_\infty$ for some linear function L_∞ . Also by passing to a subsequence we may assume $\xi_k \rightarrow \xi_\infty$ for some unit vector ξ_∞ . Then u is linear on I_∞ , which is the segment connecting $x_0 - \delta/(2n^{3/2}) \xi_\infty$ and $x_0 + \delta/(2n^{3/2}) \xi_\infty$. Hence $I_\infty \subset \{u = L\}$, which contradicts the assumption that x_0 is an extreme point of $\{u = L\}$. $\square$

The following property of sections of convex functions was proved by Caffarelli [\[1992a\]](#). Here we provide a different proof by using a well-known fact that if a continuous map from a ball to itself fixes the boundary then it must be surjective. We learned this method from Wang; see [\[Sheng et al. 2004, Section 4\]](#).

Lemma 2.5. *Let $u : \mathbb{R}^n \rightarrow [0, \infty]$ be a convex function. Assume that:*

- (1) $u(0) = 0$, $u \geq 0$.
- (2) u is finite in a neighbourhood of 0.
- (3) The graph of u contains no complete lines.

Then for $h > 0$ there exists a slope p such that the centre of mass of the section

$$S_{h,p} := \{x \mid u \leq x \cdot p + h\}$$

is defined and equal to 0.

Proof. Let

$$\begin{cases} u_k(x) = u(x) & \text{in } B_k, \\ u_k = \infty & \text{in } \mathbb{R}^n - B_k. \end{cases} \quad (2-1)$$

We only need to show the existence of sections $S^k := \{x \mid u_k \leq x \cdot p_k + h\}$ centred at 0 with bounded p_k . Then $S_{h,p} = \lim_{k \rightarrow \infty} S^k$ is the desired section in the lemma.

Take a large ball B_r . For any $p \in B_r$, let z_p be the centre of mass of the section $S_p := \{x \mid u_k(x) \leq x \cdot p + h\}$. Then we obtain a mapping $M_1 : p \rightarrow z_p$ from B_r to $\mathbb{R}^n$. If $p \in \partial B_r$, it is easy to see that $p \cdot z_p > 0$ provided r is sufficiently large.

If there is no $p \in B_r$ such that $z_p = 0$, then we can define a mapping $M_2 : z_p \rightarrow t_p z_p$, where $t_p > 0$ is a constant such that $t_p z_p \in \partial B_r$. We then obtain a continuous mapping $M = M_2 \circ M_1$ from B_r to ∂B_r with the property that

$$p \cdot M(p) > 0 \quad \text{on } \partial B_r. \quad (2-2)$$

To get a contradiction, we extend the mapping M to B_{2r} as follows. For any point $p \in \partial B_{2r}$, let $p_1 = p$, $p_0 = \frac{1}{2}p \in \partial B_r$ and $p_t = (1-t)p_0 + p_1$. We extend the mapping M to B_{2r} by letting $M(p_t) = (1-t)M(p_0) + tp_1$. Then, by (2-2), $M(p) \neq 0$ on B_{2r} and M is the identity mapping on ∂B_{2r} . This is a contradiction.

Hence, for each $k > 0$, there exists a $p_k \in \mathbb{R}^n$ such that $S^k := \{x \mid u_k \leq x \cdot p_k + h\}$ is centred at 0. Moreover, $|p_k| \leq C$ for some constant independent of k . Indeed, we can argue as follows: By rotating the coordinates we may assume $p_k = (a, 0, \dots, 0)$ with $a > 0$. Let $\alpha^+ = \sup\{x_1 \mid (x_1, 0, \dots, 0) \in S^k\}$ and $\alpha^- = -\inf\{x_1 \mid (x_1, 0, \dots, 0) \in S^k\}$. Then $\alpha^+/\alpha^- \rightarrow \infty$ as $a \rightarrow \infty$. Since S_k is centred at 0, a cannot be too large. $\square$

The following Alexandrov-type estimates were proved by Caffarelli [1996]:

Lemma 2.6. *Let u be a convex solution of*

$$\det D^2 u = d\mu$$

in the convex domain S with $u = 0$ on ∂S . Assume S is normalized, namely $B_1 \subset S \subset n^{3/2}B_1$. Assume $d\mu(S) \leq \theta d\mu(\frac{1}{2}S)$ for some constant θ , where $\frac{1}{2}S$ is a dilation of S with respect to the origin.

- (a) $(1/C)|\inf_S u|^n \leq d\mu(S) \leq C|\inf_S u|^n$, where C is a constant depending only on θ .
- (b) $|u(x)|^n \leq C d\mu(S) d(x, \partial S)$.

3. Global C^1 regularity

In this section, we prove [Theorem 2.1\(i\)](#).

Lemma 3.1. *Suppose u is a globally Lipschitz convex function. Assume that u is C^1 at all of the extreme points of a convex set $K = \{u = L\}$, where L is a linear function satisfying $u \geq L$ and $u(y) = L(y)$ for some $y \in \mathbb{R}^n$. Then u is C^1 on K .*

Proof. By subtracting L we may assume $K = \{u = 0\}$. If K is a bounded convex set, then for any $x \in K$ we have

$$x = \sum_{i=1}^k \lambda_i x_i,$$

where $x_i, i = 1, \dots, k$, are extreme points of K , $\lambda_i \geq 0$ and $\sum_{i=1}^k \lambda_i = 1$. Since u is C^1 at $x_i, i = 1, \dots, k$, we have $0 \leq u(z) = o(z - x_i), i = 1, \dots, k$. Now, by convexity we have

$$0 \leq u(z) = u\left(\sum_{i=1}^k \lambda_i (z - x + x_i)\right) \leq \sum_{i=1}^k \lambda_i u(z - x + x_i) = \sum_{i=1}^k \lambda_i o(z - x) = o(z - x).$$

Hence, u is C^1 at x .

If K is unbounded, it is well-known that $K = \text{convex}[K] + \text{rc}[K]$, where $\text{convex}[K]$ is the convex hull of the extreme points of K , and $\text{rc}[K] := \lim_{\lambda \downarrow 0} \lambda K$ is the recession cone of K . Hence we need only to show that u is C^1 at points represented by $x = x_0 + q$, where x_0 is an extreme point of K and $q \in \text{rc}[K]$. For any $M \geq 0$, by using the facts that u is Lipschitz and $x_1 := x_0 + Mq \in K$ we have that $u(z - x + x_1) \leq C|z - x|$. By convexity we have

$$u(z) = u\left(\frac{M-1}{M}(z - x + x_0) + \frac{1}{M}(z - x + x_1)\right) \leq \frac{M-1}{M}o(|z - x|) + \frac{C}{M}|z - x|.$$

By letting $M \rightarrow \infty$ we have $0 \leq u(z) \leq o(|z - x|)$. Hence u is C^1 at x . $\square$

Since u is convex, for any unit vector γ the lateral derivatives

$$\partial_\gamma^+ u(x) =: \lim_{t \searrow 0} t^{-1}(u(x + t\gamma) - u(x)) \quad \text{and} \quad \partial_\gamma^- u(x) =: \lim_{t \searrow 0} t^{-1}(u(x) - u(x - t\gamma))$$

exist. To prove that $u \in C^1(\bar{\Omega})$, it suffices to prove that

$$\partial_\gamma^+ u(x_0) = \partial_\gamma^- u(x_0) \tag{3-1}$$

at any point $x_0 \in \partial\Omega$ for any unit vector γ . By convexity, it suffices to prove this for $\xi = \xi_k$ for all $k = 1, 2, \dots, n$, where $\xi_k, k = 1, \dots, n$, are any n linearly independent unit vectors.

Proof of Theorem 2.1(i). By Lemmas 3.1 and 2.4 we only need to show that u is C^1 at localized points. Assume to the contrary that u is not C^1 at $x_0 \in \partial\Omega$. Let us assume that $x_0 = 0$, $u(0) = 0$, $u \geq 0$ and $\partial_1^+ u(0) > \partial_1^- u(0) = 0$. Since $\partial\Omega$ is Lipschitz, we may also assume that $-te_1 \in \Omega$ for $t \in (0, 1)$, where e_1 is the first coordinate direction.

Figure 1. Centred section.

Now we consider a section $S_h(x')$, where $x' = (-a', 0, \dots, 0)$ for some small constant $0 < a' < \frac{1}{2}r_0$, where $r_0 := r_{x_0}$ is the radius in the condition of [Theorem 2.1\(i\)](#). Note that by John's lemma there exists an ellipsoid E with centre x' such that $E \subset S_h(x') \subset n^{3/2}E$. Since u is Lipschitz and $\partial_1^+ u(0) > 0$, we have that $C^{-1}\varepsilon \leq u(\varepsilon e_1) \leq C\varepsilon$ for any small positive ε , where C is a positive constant. Since $\partial_1^- u(0) = 0$, we have $u(-Ma'e_1) = o(a')$, where $M = 2n^{3/2}$. Hence, we can choose small ε and a' so that the following properties hold:

- (1) $o(a') = u(-Ma'e_1) \leq C^{-1}\varepsilon \ll a'$,
- (2) εe_1 is on the boundary of some section $S_h(x')$, and
- (3) $S_h(x') \subset B_{r_0}(0)$.

The existence of such a section $S_h(x')$ in (2) follows from the property that a centred section, say $S_h(x)$, varies continuously with respect to the height h ; see [\[Caffarelli and McCann 2010, Lemma A.8\]](#), and (3) follows from the assumption that $x_0 = 0$ is localized.

Let L be the defining linear function of $S_h(x')$; by (1) it is easy to see that L is increasing in the e_1 direction (see [Figure 1](#)); hence,

$$(L - u)(0) \geq (L - u)(x') = h. \quad (3-2)$$

Since $\int_{S_h(x')} f \leq C \int_{\frac{1}{2}S_h(x')} f$, we have that

$$(L - u)(0) \leq C \left(\frac{\varepsilon}{a'} \right)^{\frac{1}{n}} h, \quad (3-3)$$

contradicting (3-2), since $a' \gg \varepsilon$. Here we have followed the argument of [\[Caffarelli 1996\]](#). Indeed, let A be an affine transform normalizing $S_h(x')$; then $v := (u - L)(A^{-1}x)/h$ satisfies $\det D^2 v = f(A^{-1}x)/h^n$ in $A(S_h(x'))$ and $v = 0$ on $\partial S_h(x')$. Hence, by applying [Lemma 2.6](#) to v and translating back to u we get (3-3).

Hence, u must be C^1 at any localized point x_0 . Therefore $u \in C^1(\mathbb{R}^n)$. □

Remark 3.2. The proof of [Theorem 2.1\(i\)](#) shares some similarities with the proof of C^1 regularity for the obstacle problem in [\[Savin 2005\]](#) (see Proposition 2.8 in that paper).

4. Global $C^{1,\alpha}$ regularity

In this section, we prove [Theorem 2.1\(ii\)](#). First we point out that to prove $u \in C^{1,\alpha}(\bar{\Omega})$, it suffices to prove that there exist constants $C > 0$, $\alpha \in (0, 1)$ and $r > 0$ such that, for any point $x_0 \in \bar{\Omega}$,

$$u(x) - \ell_{x_0}(x) \leq C|x - x_0|^{1+\alpha} \quad (4-1)$$

for every $x \in B_r(x_0) \cap \bar{\Omega}$. From (4-1) one can prove that $u \in C^{1,\alpha}(\bar{\Omega})$, using the convexity of u . In the following we will show that a relaxed version of (4-1) is enough to show $u \in C^{1,\alpha}(\bar{\Omega})$, and it has the advantage of avoiding some annoying limiting picture.

By the assumption of [Theorem 2.1\(ii\)](#) we write $\Omega = U - \sum_{i=1}^k C_i$, where U is an open convex set, and C_i , $i = 1, \dots, k$, are closed disjoint convex subsets of U ; see [Figure 2](#). Given any $x \in \bar{\Omega}$, we introduce the function

$$\rho_x(t) := \sup \left\{ u(z) - u(x) - Du(x) \cdot (z - x) \mid |z - x| = t, x + s \frac{z - x}{|z - x|} \in \bar{\Omega} \text{ for any } s \in [0, r_0] \right\}, \quad (4-2)$$

where r_0 is a fixed small positive constant depending on Ω , and its smallness will be clear in the proof of [Lemma 4.1](#). Indeed, we need to take r_0 small enough that $B_{r_0}(x) \cap \partial U$ can be represented as the graph of some Lipschitz function for any $x \in \partial U$ with the Lipschitz constant independent of x , and that

$$r_0 \ll \min \{ \text{dist}(\partial U, \partial C_i), \text{dist}(\partial C_j, \partial C_l) \mid i = 1, \dots, k, 1 \leq j \neq l \leq k \}.$$

Lemma 4.1. *Suppose that there exist $r > 0$ and $\delta \in (0, 1)$ such that for any $x \in \bar{\Omega}$ we have*

$$\rho_x\left(\frac{1}{2}t\right) \leq \frac{1}{2}(1 - \delta)\rho_x(t) \quad (4-3)$$

whenever $t \leq r$. Then $u \in C^{1,\alpha}(\bar{\Omega})$ for some $\alpha \in (0, 1)$.

Figure 2. Domain Ω .

Proof. For $t = r/2^k$, we have

$$\rho_x(t) \leq \frac{(1-\delta)^k}{2^k} \rho_x(r) \leq \frac{t}{r} (1-\delta)^{\log(r/t)/\log 2} \rho_x(r) \leq C t^{1+\alpha}, \quad (4-4)$$

where C depends on r , δ and $\rho_x(r)$, and $\alpha = -\log(1-\delta)/\log 2$.

Suppose $x, y \in \bar{\Omega}$ and $|x - y| \ll r \ll r_0$. We need to consider two cases:

- (a) x, y are close to ∂U .
- (b) x, y are close to ∂C_i for some $1 \leq i \leq k$.

We will deal with case (a) first; case (b) follows from a similar argument. Without loss of generality we may assume that $B_{3r_1} \subset U$ for some small fixed r_1 , that $r_0 \ll r_1$, and that $\text{dist}(\partial B_{3r_1}, \partial U) \gg r_1$. Denote by $\mathcal{C}_{x,r_1}$ the convex hull of x and B_{r_1} . By convexity, $\mathcal{C}_{x,3r_1} \subset U$. Then we prove the following claim:

Claim 1. For any $z \in B_{r/2}(x) \cap \mathcal{C}_{x,2r_1}$, we have $|Du(x) - Du(z)| \leq C|x - z|^\alpha$.

Proof of Claim 1. Observe that $\text{dist}(z, \partial \mathcal{C}_{x,3r_1}) \geq (1/C)|x - z|$ for some large constant C . Hence, $B_{(1/C)|x-z|}(z) \subset B_r \cap \mathcal{C}_{x,3r_1}$. Now, for any $\tilde{z} \in \partial B_{(1/C)|x-z|}(z)$, by (4-4) we have that

$$u(\tilde{z}) \leq u(x) + Du(x) \cdot (\tilde{z} - x) + C|\tilde{z} - x|^{1+\alpha}. \quad (4-5)$$

By convexity we also have

$$u(\tilde{z}) \geq u(z) + Du(z) \cdot (\tilde{z} - z) \quad (4-6)$$

and

$$u(z) \geq u(x) + Du(x) \cdot (z - x). \quad (4-7)$$

By (4-5), (4-6) and (4-7) we have

$$(Du(z) - Du(x)) \cdot (\tilde{z} - z) \leq C|\tilde{z} - x|^{1+\alpha}. \quad (4-8)$$

Note that $|\tilde{z} - z| \approx |\tilde{z} - x| \approx |z - x|$ provided $\tilde{z} \in \partial B_{(1/C)|x-z|}(z)$ and C is sufficiently large. Since (4-8) holds for any $\tilde{z} \in \partial B_{(1/C)|x-z|}(z)$, it follows that $|Du(x) - Du(z)| \leq C|x - z|^\alpha$. $\square$

Now suppose $|x - y| \ll r$. If either $y \in \mathcal{C}_{x,2r_1}$ or $x \in \mathcal{C}_{y,2r_1}$ holds, then by Claim 1 we have $|Du(x) - Du(y)| \leq C|x - y|^\alpha$. Otherwise one may find a point $z \in \mathcal{C}_{x,r_1} \cap \mathcal{C}_{y,r_1}$ such that $|z - x| \approx |z - y| \approx |x - y|$. Then by applying the estimate in Claim 1 we have

$$|Du(x) - Du(y)| \leq |Du(x) - Du(z)| + |Du(y) - Du(z)| \leq C(|x - z|^\alpha + |y - z|^\alpha) \leq C|x - y|^\alpha.$$

We can prove case (b) by a similar argument. Indeed, $\partial C_1 \cap B_r(x)$ can be represented as the graph of some Lipschitz function for any fixed $x \in \partial C_1$ provided $r \ll r_0$. Then, by the assumption that the C_i are disjoint, it is easy to find a small ball $B_{3r_1} \subset \Omega$ such that $\mathcal{C}_{z,3r_1} \subset \Omega$ for any $z \in B_r(x) \cap \bar{\Omega}$. Then, by a similar argument to the proof of case (a), we can show that $|Du(x) - Du(y)| \leq C|x - y|^\alpha$ provided $|x - y| \ll r$. $\square$

The following lemma shows that the centred sections are well-localized provided the heights are sufficiently small.

Lemma 4.2. *There exists a height $h_0 > 0$ such that, for any $x \in \bar{\Omega}$, the section $S_h(x)$ intersects at most one of $\partial U, \partial C_i, i = 1, \dots, m$, provided $h \leq h_0$.*

Proof. Suppose to the contrary there exist sequences $x_k \in \bar{\Omega}$ and $h_k \rightarrow 0$, such that $S_{h_k}(x_k)$ intersects at least two of $\partial U, \partial C_i, i = 1, \dots, m$. Passing to a subsequence we may assume $x_k \rightarrow y \in \bar{\Omega}$. Since u is strictly convex in the interior of Ω , we have either $y \in \partial U$ or $y \in \partial C_i$ for some i . Denote by L_k the defining function of $S_{h_k}(x_k)$, namely $S_{h_k}(x_k) = \{u \leq L_k\}$. Then, passing to a subsequence we may assume $L_k \rightarrow L$ for some affine function L , and $S_{h_k}(x_k) \rightarrow S \subset \{u \leq L\}$. It follows from the properties of $S_{h_k}(x_k)$ that:

- (i) S is centred at y .
- (ii) S intersects at least two of $\partial U, \partial C_i, i = 1, \dots, m$.
- (iii) $L(y) = \lim_{k \rightarrow \infty} L_k(x_k) = \lim_{k \rightarrow \infty} u(x_k) + h_k = u(y)$.

By (i) and (iii) we have that $S \subset \{u = L\}$. Then by (ii) we see that S passes through the interior of Ω , which contradicts the fact that u is strictly convex in the interior of Ω . $\square$

Proof of Theorem 2.1(ii). Step 1. The main observation in this step is that if (4-3) is violated for small δ , then u is close to a linear function on a segment connecting x and some point $z_\delta \in \bar{\Omega}$. Hence, if (4-3) is violated for arbitrary r, δ , then one can find a sequence of points x_k such that u is more and more linear around x_k in some direction as $k \rightarrow \infty$. The “almost linearity” will be clear if we perform blow-up and an affine transform on u properly restricted to some carefully chosen section around x_k , and a line segment will appear on the graph of the limiting function. The detailed argument goes as follows.

To prove $\rho_x(t) \leq Ct^{1+\alpha}$ for any $x \in \bar{\Omega}$ and any $t \leq r$, by Lemma 4.1 we assume to the contrary that there exist sequences $t_k \leq 1/k, \delta_k = 1/k$ and $x_k \in \bar{\Omega}$ such that

$$\rho_{x_k}(\tfrac{1}{2}t_k) \geq \tfrac{1}{2}(1 - 1/k)\rho_{x_k}(t_k). \quad (4-9)$$

Suppose the supremum in (4-2) (when $x = x_k$ and $t = \tfrac{1}{2}t_k$) is attained at $\tfrac{1}{2}(x_k + z_k) \in \bar{\Omega}$; by the definition of ρ_x we see that $\overline{z_k x_k} \subset \bar{\Omega}$, where $\overline{z_k x_k}$ denotes the segment connecting z_k and x_k . By passing to a subsequence, we may assume $x_k \rightarrow x_\infty \in \partial\Omega$.

Choosing sections. For each k , let $S_{h_k}(x_k)$ be a section of u with centre x_k , where h_k is chosen so that $z_k \in \partial S_{h_k}(x_k)$. Similar to the proof of Theorem 2.1(i), the existence of such a section follows from the property that a centred section, say $S_h(x)$, varies continuously with respect to the height h ; see [Caffarelli and McCann 2010, Lemma A.8] for a proof. It is easy to see that $h_k \rightarrow 0$.

Normalization. Let L_k be the defining function of $S_{h_k}(x_k)$. We normalize the section $S_{h_k}(x_k)$ by a linear transformation T_k , and let $S_k = T_k(S_{h_k}(x_k))$. Note that $T_k(x_k) = 0$ and $B_1 \subset S_k \subset n^{3/2}B_1$. Also we let $u_k = (u - L_k)(T_k^{-1}x)/h_k$. Then u_k solves

$$\begin{cases} \det D^2 u_k = f_k & \text{in } S_k, \\ u_k = 0 & \text{on } \partial S_k, \end{cases} \quad (4-10)$$

where $f_k = h_k^{-n}(\det T_k)^{-1}f(T_k^{-1}x)/g(Du(T_k^{-1}x))$. After a rotation of coordinates, we may assume $T_k(z_k)$ is on the x_1 -axis.

Linearity estimate. Let

$$v_k(x) := u(x) - Du(x_k) \cdot (x - x_k) - u(x_k);$$

from (4-9) we have that $v_k(\frac{1}{2}(x_k + z_k)) \geq \frac{1}{2}(1 - 1/k)v_k(z_k)$. Let

$$\tilde{L}_k(x) := L_k(x) - Du(x_k) \cdot (x - x_k) - u(x_k).$$

Then we have that $S_{h_k}(x_k) = \{v_k \leq \tilde{L}_k\}$. Since $S_{h_k}(x_k)$ is centred at x_k , $z_k \in \partial S_{h_k}(x_k)$, $v_k \geq 0$ and $\tilde{L}_k(x_k) = h_k$, by John's lemma we have that $0 \leq \tilde{L}_k(z_k) \leq 2n^{3/2}h_k$. Now,

$$(v_k - \tilde{L}_k)(\frac{1}{2}(x_k + z_k)) - \frac{1}{2}\left(1 - \frac{1}{k}\right)((v_k - \tilde{L}_k)(x_k) + (v_k - \tilde{L}_k)(z_k)) \geq -\frac{1}{2k}(\tilde{L}_k(x_k) + \tilde{L}_k(z_k)) \geq -\frac{3n^{3/2}}{2k}h_k.$$

Since $v_k - \tilde{L}_k = u - L_k$, from the above estimate and the definition of u_k we have

$$u_k(\frac{1}{2}T_k z_k) \geq \frac{1}{2}\left(1 - \frac{1}{k}\right)(u_k(0) + u_k(T_k z_k)) - \frac{3n^{3/2}}{2k}. \quad (4-11)$$

Limiting problem. Now, by convexity we may take limits $S_k \rightarrow S_\infty$ and $u_k \rightarrow u_\infty$. Let f_∞ be the weak limit of f_k . Then u_∞ satisfies $\det D^2 u_\infty = f_\infty$ in the Alexandrov sense. Let $z_\infty := \lim_{k \rightarrow \infty} T_k(z_k)$. By (4-11) we have

$$u_\infty = L \text{ on the segment connecting } 0 \text{ and } z_\infty, \quad (4-12)$$

where L is a supporting function of u_∞ at 0.

Step 2. In this step, we need to consider two situations:

- (a) $x_\infty \in \partial C_i$ for some $1 \leq i \leq k$.
- (b) $x_\infty \in \partial U$.

In each case, a contradiction is obtained at some carefully chosen extreme point (denoted by y) of $\{u_\infty = L\}$. Heuristically, we can choose a section of u_∞ (denoted by S) around y such that y is much closer to ∂S in one direction than in the opposite direction. Hence, on one hand the Alexandrov-type estimate Lemma 2.6(a) shows that h , the height of the section S , should not be too small. On the other hand, Lemma 2.6(b) shows that h is very small, which is a contradiction.

We deal with case (a) first.

Proof in case (a). Note that since $x_\infty \in \partial C_i$ for some $1 \leq i \leq k$ and $h_k \rightarrow 0$ as $k \rightarrow \infty$, by Lemma 4.2 we have that the support of f_k can be represented by $S_k - A_k$ when k is large, where A_k is an open convex subset of S_k . Let the convex set A_∞ be the limit of the A_k . Then $S_\infty - A_\infty$ is the support of f . Since the centre of mass of S_∞ is 0 and $0 \in S_\infty - A_\infty$, we have that the volume of $S_\infty - A_\infty$ is positive. Hence, it is easy to see that there exists a constant C such that $C^{-1}\chi_{S_\infty - A_\infty} \leq f_\infty \leq C\chi_{S_\infty - A_\infty}$.

Since $\overline{z_k x_k} \subset \bar{\Omega}$, we have $\overline{0 z_\infty} \cap A_\infty = \emptyset$.

Subcase 1: $\{u_\infty = L\}$ contains an interior point of $S_\infty - \bar{A}_\infty$.

Subcase 2: $\{u_\infty = L\} \cap S_\infty \subset \bar{A}_\infty$.

Figure 3. Two related sections.

For subcase 1, take $x_0 \in (S_\infty - \bar{A}_\infty) \cap \{u = L\}$. Take δ sufficiently small that $B_\delta(x_0) \subset S_\infty - \bar{A}_\infty$.

Choosing an extreme point. Let $y \in \{u = L\}$ be the point such that:

- (1) $u_\infty(y) = \inf_{\{u=L\}} u_\infty$.
- (2) y is an extreme point of the convex set $\{u_\infty = L\} \cap \{u_\infty = u(y)\}$.

It is easy to see that y is an extreme point of $\{u_\infty = L\}$.

Cutting a suitable section. By rotating the coordinates we may assume that $\{u_\infty = L\} \subset \{x_1 \leq b\}$ for some constant $b > 0$, and that $\{u_\infty = L\} \cap \{x_1 = b\} = \{y\}$. Then we consider the section $S = \{u_\infty < L + \varepsilon(x_1 - b + a)\}$ (see Figure 3), where we fix a sufficiently small and then take $\varepsilon \ll a$, so that $S \Subset S_\infty$ and $a \gg d := \max\{x_1 \mid (x_1, 0, \dots, 0) \in S\} - b$.

Using Alexandrov estimates to obtain a contradiction. On one hand, by the Alexandrov estimate we have

$$|S|^2 > C \frac{a}{d} \varepsilon^n. \quad (4-13)$$

On the other hand, we consider another section $\tilde{S} = \{u_\infty < L + C\varepsilon\}$. Since u is Lipschitz, it is easy to see that $S \subset \tilde{S}$ provided C (independent of ε) is sufficiently large. By convexity we have $|B_\delta(x_0) \cap \tilde{S}| \geq C|\tilde{S}|$

for some constant C . We claim

$$|S|^2 \leq C\varepsilon^n, \quad (4-14)$$

where the constant C is independent of d . The claim follows from the following argument. Let $v = u_\infty - L - C\varepsilon$. Let $G := \tilde{S} \cap B_\delta(x_0)$. By John's lemma, there exists an affine transformation A with $\det A = 1$ such that

$$B_{\bar{r}} \subset A(G) \subset n^{3/2} B_{\bar{r}}$$

for some $\bar{r}$. Now $\bar{v} = v(A^{-1}x)$ satisfies $\det D^2 \bar{v} = f_\infty(A^{-1}x) \geq C^{-1}$ in $A(G)$ and $|\bar{v}| \leq C\varepsilon$ in $A(G)$. Then we have

$$C^{-1}|G| \leq \int_{G/2} f_\infty = |\partial \bar{v}(A(\tfrac{1}{2}G))| \leq C \frac{\varepsilon^n}{\bar{r}^n}. \quad (4-15)$$

Equation (4-14) follows from (4-15) and the fact that $|\tilde{S}| \approx |G| \approx \bar{r}^n$. Since $d \ll a$, it is easy to see that (4-14) contradicts (4-13).

For subcase 2, we need to choose the extreme point more carefully.

Choosing an extreme point. Let $\tilde{K} \subset \mathbb{R}^n$ be a supporting plane of the convex set A_∞ at 0. If A_∞ is not C^1 at 0 we choose $\tilde{K}$ to be the one containing $\overline{z_\infty 0}$. Let y' be the point where u_∞ attains its minimum on $D := \{u = L\} \cap \tilde{K} \cap \bar{S}_\infty$. It is easy to check that D is a convex set, and the set $D \cap \{x \mid u(x) = u(y')\}$ is also convex. Let y be an extreme point of $D \cap \{x \mid u(x) = u(y')\}$. We claim that y is an extreme point of $\{u = L\}$. Indeed, suppose not; then there exist $y_1, y_2 \in \{u = L\} \cap S_\infty \subset \bar{A}_\infty$ such that $y = \frac{1}{2}(y_1 + y_2)$. Since $\tilde{K}$ is a supporting plane of A_∞ and $y \in \bar{A}_\infty$, we have that $y_1, y_2 \in D$. However, since $u(y) = \min\{u(x) \mid x \in D\}$, we have $y_1, y_2 \in D \cap \{x \mid u(x) = u(y')\}$, which contradicts the choice of y as an extreme point of $D \cap \{x \mid u(x) = u(y')\}$.

Cutting a suitable section. By subtracting L and translating the coordinates we may assume that $y = 0$, that $u_\infty \geq 0$, that $u_\infty(te_1) = 0$ for $t \in (0, 1)$, and that $u_\infty(te_1) > 0$ for $t < 0$. Let $0 < \varepsilon \ll a$ be small positive numbers. Let $S_h(ae_1)$ be a section of u_∞ with centre ae_1 , where h is chosen so that $-\varepsilon e_1 \in \partial S_h(ae_1)$. Since y is an extreme point of $\{u = L\}$, we have that $S_h(ae_1) \Subset S_\infty$ provided h is sufficiently small. Note that $h \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Using Alexandrov estimates to obtain a contradiction. Since A_∞ is convex, it is easy to see that

$$\int_{S_h(ae_1)} f_\infty \leq C \int_{\frac{1}{2}S_h(ae_1)} f_\infty$$

for some constant C . Let L_1 be the defining function of the section $S_h(ae_1)$, which is obviously decreasing in the e_1 direction. Hence $(L_1 - u_\infty)(0) \geq h$. Then by Lemma 2.6 we also have

$$(L_1 - u_\infty)(0) \leq C \left(\frac{\varepsilon}{a}\right)^{1/n} h,$$

which contradicts the previous estimate. $\square$

Proof in case (b). The proof in case (b) follows from a similar argument to the proof of [Caffarelli 1992a, Lemma 4]; we sketch the argument here. Note that f_k is now supported in a convex domain $D_k \subset \bar{S}_k$.

Let $D_\infty := \lim_{k \rightarrow \infty} D_k$. We have $z_\infty \in D_\infty$. Let L be the supporting function of u_∞ at 0 such that $\overline{0z_\infty} \subset \{u_\infty = L\}$. Similarly to the proof of subcase 1 of case (a), let $y \in \{u_\infty = L\}$ be the point such that:

- (1) $u_\infty(y) = \inf_{\{u_\infty = L\}} u_\infty$.
- (2) y is an extreme point of the convex set $\{u_\infty = L\} \cap \{u_\infty = u(y)\}$.

It is easy to see that y is an extreme point of $\{u_\infty = L\}$. Observe that $y \in D_\infty$, since otherwise u_k has positive Monge–Ampère measure outside D_k for large k . Let $z = (1 - \sigma)y + \sigma z_\infty$ for some small positive σ ; we may also find a section satisfying $S_h(z) := \{u_\infty < L\} \Subset S_\infty$ and $y + \varepsilon(y - z_\infty)/|y - z_\infty| \in \partial S_h(z)$ for small $\varepsilon \ll \sigma$. Since $y \in D_\infty$, there exists a sequence $y_k \in D_k$ such that $y_k \rightarrow y$ as $k \rightarrow \infty$. Let

$$\tilde{z}_k := (1 - \sigma)y_k + \sigma T(z_k);$$

it is easy to see that $\tilde{z}_k \rightarrow z$ as $k \rightarrow \infty$. Recall that $z_\infty := \lim_{k \rightarrow \infty} T(z_k)$ with $T(z_k) \in D_k$. Let $\tilde{S}_k := \{u_k \leq L_k\}$ be a section of u_k centred at $\tilde{z}_k$ with height h . Then, passing to a subsequence, $\tilde{S}_k \rightarrow S_h(z)$ in Hausdorff distance. In particular, $\tilde{S}_k \Subset S_k$ provided k is sufficiently large. Then, by [Lemma 2.6](#), we have that

$$Ch \leq (L_k - u_k)(y_k) \leq \left(\frac{\varepsilon}{\sigma}\right)^{1/n} h$$

for large k , which is a contradiction because $\varepsilon \ll \sigma$. □

[Theorem 2.1\(ii\)](#) follows from the above discussions. □

References

- [Brenier 1991] Y. Brenier, “Polar factorization and monotone rearrangement of vector-valued functions”, *Comm. Pure Appl. Math.* **44**:4 (1991), 375–417. [MR 1100809](#) [Zbl 0738.46011](#)
- [Caffarelli 1990a] L. A. Caffarelli, “A localization property of viscosity solutions to the Monge–Ampère equation and their strict convexity”, *Ann. of Math. (2)* **131**:1 (1990), 129–134. [MR 1038359](#) [Zbl 0704.35045](#)
- [Caffarelli 1990b] L. A. Caffarelli, “Interior $W^{2,p}$ estimates for solutions of the Monge–Ampère equation”, *Ann. of Math. (2)* **131**:1 (1990), 135–150. [MR 1038360](#) [Zbl 0704.35044](#)
- [Caffarelli 1991] L. A. Caffarelli, “Some regularity properties of solutions of Monge–Ampère equation”, *Comm. Pure Appl. Math.* **44**:8–9 (1991), 965–969. [MR 1127042](#) [Zbl 0761.35028](#)
- [Caffarelli 1992a] L. A. Caffarelli, “Boundary regularity of maps with convex potentials”, *Comm. Pure Appl. Math.* **45**:9 (1992), 1141–1151. [MR 1177479](#) [Zbl 0778.35015](#)
- [Caffarelli 1992b] L. A. Caffarelli, “The regularity of mappings with a convex potential”, *J. Amer. Math. Soc.* **5**:1 (1992), 99–104. [MR 1124980](#) [Zbl 0753.35031](#)
- [Caffarelli 1996] L. A. Caffarelli, “Boundary regularity of maps with convex potentials, II”, *Ann. of Math. (2)* **144**:3 (1996), 453–496. [MR 1426885](#) [Zbl 0916.35016](#)
- [Caffarelli and McCann 2010] L. A. Caffarelli and R. J. McCann, “Free boundaries in optimal transport and Monge–Ampère obstacle problems”, *Ann. of Math. (2)* **171**:2 (2010), 673–730. [MR 2630054](#) [Zbl 1196.35231](#)
- [Figalli and Loeper 2009] A. Figalli and G. Loeper, “ C^1 regularity of solutions of the Monge–Ampère equation for optimal transport in dimension two”, *Calc. Var. Partial Differential Equations* **35**:4 (2009), 537–550. [MR 2496656](#) [Zbl 1170.35400](#)
- [Figalli and Rifford 2009] A. Figalli and L. Rifford, “Continuity of optimal transport maps and convexity of injectivity domains on small deformations of $\mathbb{S}^2$ ”, *Comm. Pure Appl. Math.* **62**:12 (2009), 1670–1706. [MR 2569074](#) [Zbl 1175.49040](#)
- [Figalli et al. 2010] A. Figalli, L. Rifford, and C. Villani, “On the Ma–Trudinger–Wang curvature on surfaces”, *Calc. Var. Partial Differential Equations* **39**:3–4 (2010), 307–332. [MR 2729302](#) [Zbl 1203.53034](#)

- [Figalli et al. 2011] A. Figalli, L. Rifford, and C. Villani, “Necessary and sufficient conditions for continuity of optimal transport maps on Riemannian manifolds”, *Tohoku Math. J. (2)* **63**:4 (2011), 855–876. MR 2872966 Zbl 1262.58013
- [Figalli et al. 2012] A. Figalli, L. Rifford, and C. Villani, “Nearly round spheres look convex”, *Amer. J. Math.* **134**:1 (2012), 109–139. MR 2876141 Zbl 1241.53031
- [Figalli et al. 2013a] A. Figalli, Y.-H. Kim, and R. J. McCann, “Hölder continuity and injectivity of optimal maps”, *Arch. Ration. Mech. Anal.* **209**:3 (2013), 747–795. MR 3067826 Zbl 1281.49037
- [Figalli et al. 2013b] A. Figalli, Y.-H. Kim, and R. J. McCann, “Regularity of optimal transport maps on multiple products of spheres”, *J. Eur. Math. Soc. (JEMS)* **15**:4 (2013), 1131–1166. MR 3055758 Zbl 1268.49053
- [Kim and McCann 2010] Y.-H. Kim and R. J. McCann, “Continuity, curvature, and the general covariance of optimal transportation”, *J. Eur. Math. Soc. (JEMS)* **12**:4 (2010), 1009–1040. MR 2654086 Zbl 1191.49046
- [Liu 2009] J. Liu, “Hölder regularity of optimal mappings in optimal transportation”, *Calc. Var. Partial Differential Equations* **34**:4 (2009), 435–451. MR 2476419 Zbl 1166.35331
- [Liu and Wang 2015] J. Liu and X.-J. Wang, “Interior a priori estimates for the Monge–Ampère equation”, pp. 151–177 in *Surveys in differential geometry 2014: regularity and evolution of nonlinear equations*, edited by H.-D. Cao et al., Surv. Differ. Geom. **19**, International Press, Somerville, MA, 2015. MR 3381500 Zbl 1326.35168
- [Liu et al. 2010] J. Liu, N. S. Trudinger, and X.-J. Wang, “Interior $C^{2,\alpha}$ regularity for potential functions in optimal transportation”, *Comm. Partial Differential Equations* **35**:1 (2010), 165–184. MR 2748621 Zbl 1189.35142
- [Loeper 2011] G. Loeper, “Regularity of optimal maps on the sphere: the quadratic cost and the reflector antenna”, *Arch. Ration. Mech. Anal.* **199**:1 (2011), 269–289. MR 2754343 Zbl 1231.35280
- [Loeper and Villani 2010] G. Loeper and C. Villani, “Regularity of optimal transport in curved geometry: the nonfocal case”, *Duke Math. J.* **151**:3 (2010), 431–485. MR 2605867 Zbl 1192.53041
- [Savin 2005] O. Savin, “The obstacle problem for Monge Ampere equation”, *Calc. Var. Partial Differential Equations* **22**:3 (2005), 303–320. MR 2118901 Zbl 1084.35026
- [Sheng et al. 2004] W. Sheng, N. Trudinger, and X.-J. Wang, “Convex hypersurfaces of prescribed Weingarten curvatures”, *Comm. Anal. Geom.* **12**:1-2 (2004), 213–232. MR 2074877 Zbl 1065.53008
- [Trudinger and Wang 2009a] N. S. Trudinger and X.-J. Wang, “On strict convexity and continuous differentiability of potential functions in optimal transportation”, *Arch. Ration. Mech. Anal.* **192**:3 (2009), 403–418. MR 2505359 Zbl 1214.49038
- [Trudinger and Wang 2009b] N. S. Trudinger and X.-J. Wang, “On the second boundary value problem for Monge–Ampère type equations and optimal transportation”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **8**:1 (2009), 143–174. MR 2512204 Zbl 1182.35134

Received 20 Nov 2015. Revised 8 Mar 2016. Accepted 29 Apr 2016.

ELINA ANDRIYANOVA: elina.andriyanova@anu.edu.au

Mathematical Science Institute, The Australian National University, Canberra ACT 2601, Australia

SHIBING CHEN: shibing.chen@anu.edu.au

Mathematical Science Institute, The Australian National University, Canberra ACT 2601, Australia

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard
patrick.gerard@math.u-psud.fr
Université Paris Sud XI
Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Yuval Peres	University of California, Berkeley, USA peres@stat.berkeley.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	András Vasy	Stanford University, USA andras@math.stanford.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpmmms.cam.ac.uk	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2016 is US \$235/year for the electronic version, and \$430/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

mathematical sciences publishers
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 9 No. 6 2016

- A complete study of the lack of compactness and existence results of a fractional Nirenberg equation via a flatness hypothesis, I 1285
Wael Abdelhedi, Hichem Chtioui and Hichem Hajaiej
- On positive solutions of the (p, A) -Laplacian with potential in Morrey space 1317
Yehuda Pinchover and Georgios Psaridakis
- Geometric optics expansions for hyperbolic corner problems, I: Self-interaction phenomenon 1359
Antoine Benoit
- The interior C^2 estimate for the Monge–Ampère equation in dimension $n = 2$ 1419
Chuanqiang Chen, Fei Han and Qianzhong Ou
- Bounded solutions to the Allen–Cahn equation with level sets of any compact topology 1433
Alberto Enciso and Daniel Peralta-Salas
- Hölder estimates and large time behavior for a nonlocal doubly nonlinear evolution 1447
Ryan Hynd and Erik Lindgren
- Boundary $C^{1,\alpha}$ regularity of potential functions in optimal transportation with quadratic cost 1483
Elina Andriyanova and Shibing Chen
- Commutators with fractional differentiation and new characterizations of BMO-Sobolev spaces 1497
Yanping Chen, Yong Ding and Guixiang Hong