

Measurements and Theoretical Predictions of Charge Exchange Cross Sections and Emission Spectra for O^{6+} with H_2O , CO , CO_2 , CH_4 , N_2 , NO , N_2O and Ar

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 022076

(<http://iopscience.iop.org/1742-6596/635/2/022076>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 130.56.97.48

This content was downloaded on 11/11/2016 at 10:05

Please note that [terms and conditions apply](#).

You may also be interested in:

[N₂O Decomposed by Discharge Plasma with Catalysts](#)

Hu Hui, Huang Hao, Xu Jie et al.

[The ionization and dissociation of nitrous oxide studied by fast electron impact](#)

Y W Liu, M Lin and L F Zhu

Measurements and Theoretical Predictions of Charge Exchange Cross Sections and Emission Spectra for O^{6+} with H_2O , CO , CO_2 , CH_4 , N_2 , NO , N_2O and Ar

J. R. Machacek¹, D. P. Mahapatra^{2,3}, D. R. Schultz^{4*}, Yu. Ralchenko⁵, A. Moradmand²,
 M. O. A. El Ghazaly², and A. Chutjian^{2†}

¹ Positron Physics Group, The Australian National University, ACT 2601, Australia

² Astrophysics and Space Sciences Section, Jet Propulsion Laboratory/Caltech, Pasadena CA 91109

³ Department of Physics, Utkal University, Bhubaneswar-751004, India

⁴ Department of Physics, Univ. of North Texas, Denton, TX 76203

⁵ Atomic Spectroscopy Group, National Institute of Standards and Technology, Gaithersburg, MD 20899-8422

Synopsis Relevant to modeling and understanding X-ray emission from cometary and planetary atmospheres, total charge-exchange cross sections for 1.17 and 2.33 keV/u O^{6+} ions colliding with H_2O , CO , CO_2 , CH_4 , N_2 , NO , N_2O , and Ar have been measured and calculated for the processes of single, double, and triple exchanges. Synthetic emission spectra spanning the X-ray, UV, and visible ranges have also been calculated, based on theoretical treatment of the transfer of between one and six electrons from the target neutral to the projectile ion, followed by radiative and non-radiative decay of the resulting highly-excited projectile states.

The phenomenon of charge exchange (CE) between highly-charged ions (HCIs) and neutral molecules is observed throughout our Solar System, and is very likely occurring in more-distant stellar atmospheres. In such exchanges an HCI in (for example) the solar wind (SW) captures an electron from the neutral gas into a state with high principal quantum number n that subsequently de-excites by emitting one or more photons at visible-to-X-ray wavelengths [1]. The gaseous targets can be comet or planetary atmospheres, circumstellar clouds, and the interplanetary medium; while the HCIs arise from the SW, the Jovian magnetosphere and cosmic rays [2]. Since O^{6+} is the most abundant heavy ion in the SW, its interactions can be used to probe objects as distant as Pluto.

Reported herein are absolute CE cross sections for up to three exchanges of the O^{6+} ion with electrons from the comet/planetary gases H_2O , CO , CO_2 , CH_4 , N_2 , NO , N_2O and Ar. The measurements are compared with results in calculations using the classical trajectory Monte Carlo (CTMC) method. With the measurements as benchmarks, synthetic emission spectra spanning the X-ray, UV, and visible range have been calculated based on theoretical treatment of the transfer of between one and six electrons from the target neutrals to O^{6+} , followed by radiative and non-radiative decay of the highly-excited ion states.

Experimental measurements were carried out using the HCI source at JPL. Details are given in [3] for results with He and H_2 targets; as well as details of the CTMC calculational approach.

Extension of the theory for calculating cross sections and spectra for 1-6 captures using an *ab initio* formulation of the atomic structure and decay parameters will be described in [4, 5].

Shown in Table 1 are results of measurements and calculations for CO and H_2O , for three exchanges, at the lower energy of 1.17 keV/u [4]. Results will be presented for three exchanges in all the target gases at 1.17 and 2.33 keV/u, and simulated emission spectra for relaxation of the $O^{(5-3)+}$ ions to their lower electronic states.

Target	CE Channel	CE Cross Section (Exp.)	CE Cross Section (Theory)
CO	σ_{6-5}	5.41 ± 0.35	4.51
	σ_{6-4}	1.10 ± 0.08	0.58
	σ_{6-3}	0.37 ± 0.03	0.05
H_2O	σ_{6-5}	4.73 ± 0.31	5.50
	σ_{6-4}	0.83 ± 0.06	0.63
	σ_{6-3}	0.37 ± 0.03	0.06

Table 1. Experimental and CTMC results for CE of O^{6+} with CO and H_2O at an energy of 1.17 keV/u. Units are 10^{-15} cm^2 , and errors are at the 2σ level.

References

- [1] N. A. Schwadron, T. E. Cravens 2000 *ApJ* **554** 558
- [2] A. Bhardwaj *et al* 2007 *Planet. Sp. Sci.* **55** 1135
- [3] J. R. Machacek *et al* 2014 *Phys. Rev. A* **90** 052708
- [4] J. R. Machacek *et al.* 2015 *ApJ*, in preparation
- [5] D. R. Schultz and Yu. Ralchenko, in preparation

* E-mail: david.schultz@unt.edu † E-mail: ara.chutjian@jpl.nasa.gov

