



# Regulation of *Arabidopsis* root development by small signaling peptides

Christina Delay, Nijat Imin and Michael A. Djordjevic\*

Division of Plant Sciences, Research School of Biology, College of Medicine, Biology and Environment, The Australian National University, Canberra, ACT, Australia

## Edited by:

Wolfgang Schmidt, Academia Sinica, Taiwan

## Reviewed by:

Ive De Smet, Vlaams Instituut voor Biotechnologie, Belgium  
Reidunn Birgitta Aalen, University of Oslo, Norway

## \*Correspondence:

Michael A. Djordjevic, Division of Plant Sciences, Research School of Biology, College of Medicine, Biology and Environment, The Australian National University, Linnaeus Bld. 134, Linnaeus Way, Canberra, ACT 0200, Australia  
e-mail: michael.djordjevic@anu.edu.au

Plant root systems arise *de novo* from a single embryonic root. Complex and highly coordinated developmental networks are required to ensure the formation of lateral organs maximizes plant fitness. The *Arabidopsis* root is well-suited to dissection of regulatory and developmental networks due to its highly ordered, predictable structure. A myriad of regulatory signaling networks control the development of plant roots, from the classical hormones such as auxin and cytokinin to short-range positional signaling molecules that relay information between neighboring cells. Small signaling peptides are a growing class of regulatory molecules involved in many aspects of root development including meristem maintenance, the gravitropic response, lateral root development, and vascular formation. Here, recent findings on the roles of regulatory peptides in these aspects of root development are discussed.

**Keywords:** small signaling peptides, small post-translationally modified peptides, regulatory peptides, root development, CLE, RGF/CLEL/GLV, IDA, CEP

## INTRODUCTION

The entire plant root system is formed post-embryonically from a single primary root. Growth and development of this system requires coordinated regulation of hardwired developmental programs together with input from environmental signals (Casimiro et al., 2003; Malamy, 2005). While the classical phytohormones are key players in many aspects of root development, cell-to-cell communication is a vital component of most developmental processes. Regulatory peptides are one class of small signaling molecules that mediate intercellular communication. Roles for small signaling peptides in shoot development have been elucidated and are relatively well-characterized (Fukuda and Higashiyama, 2011). Recently, there has been a leap in our understanding of the roles of regulatory peptides in root development.

Small signaling peptides arise from genes that typically encode an N-terminal signal peptide region, one or more conserved peptide domains and variable regions that flank one or both sides of the discrete peptide domains. There are two structural classes of signaling peptides: small (5–20 amino acids) post-translationally modified peptides, which are the focus of this review; and larger cysteine-rich peptides (approximately 50 amino acids) that undergo disulfide bond formation as part of the maturation process. Common post-translational modifications to the smaller peptides class include hydroxylation, sulfation, and arabinosylation. These modifications may increase peptide stability, assist with receptor interactions and provide a further degree of regulation. The precursor proteins undergo processing to form the mature peptide product. While the maturation process is poorly understood, it was recently shown that four residues upstream of the peptide domain are required for CLE (CLAVATA3/ESR-related) peptide endoproteolytic processing (Ni et al., 2011). Most peptides are thought to act as extracellular signaling molecules that are ligands for membrane bound receptors although few


ligand/receptor interactions have been validated (Matsubayashi et al., 2002; Hirakawa et al., 2008; Ogawa et al., 2008).

Several families of regulatory peptides, defined by homology of the peptide domain, have been implicated in various developmental processes in the roots. The CLE peptide family has a conserved 12–14 amino acid CLE motif at or near the C-terminus (Cock and McCormick, 2001). The RGF (ROOT GROWTH FACTOR) family of peptides, also known as GLV (GOLVEN) and CLEL (CLE-Like), has a conserved 14 amino acid peptide domain containing the tyrosine sulfation motif Asp-Tyr (Matsuzaki et al., 2010; Meng et al., 2012b; Whitford et al., 2012). The IDA (INFLORESCENCE DEFICIENT IN ABCISSION) peptide family has conserved, functionally active 20 amino acid motif (EPIP; Butenko et al., 2003; Stenvik et al., 2008). The CEP (C-TERMINALLY ENCODED PEP-TIDE) family has a conserved 15 amino acid peptide domain with two proline residues that may be hydroxylated (Ohyama et al., 2008).

Recent work implicates regulatory peptides in many aspects of root growth and development, including meristem maintenance, gravitropism, lateral root (LR) development, and protoxylem differentiation (Figures 1A–G). This review outlines the recently discovered roles of these peptide families in root development.

## ROOT MERISTEM MAINTENANCE

For continual growth and development, roots require a source of new cells that are able to differentiate into various tissue types. The root meristem contains a population of stem cells maintained by the quiescent center (QC). An extraordinary amount of precise regulation is required to ensure that the identity of the QC cells, the surrounding stem cells and their daughters is maintained. Several transcription factors are essential for this. *WOX5* (*WUSCHEL*-related homeobox 5) is expressed in the QC to maintain stem cell identity (Haecker et al., 2004). *SCR* (*SCARECROW*)



and *SHR* (*SHORTROOT*) are expressed along the radial axis to provide positional information and also maintain QC identity (Di Laurenzio et al., 1996; Helariutta et al., 2000; Sabatini et al., 2003). *PLT1* (*PLETHORA1*) and *PLT2* (*PLETHORA2*) are auxin

responsive genes that form a gradient distribution to maintain stem cell identity when expressed at high levels and promote differentiation when present at low levels (Aida et al., 2004; Galinha et al., 2007). An auxin maximum is also required in the stem cell niche (Sabatini et al., 1999; Friml et al., 2003).

A recent report implicated the CLE40 peptide in maintaining QC and columella stem cell identity (Figure 1D; Stahl et al., 2009). *CLE40* is expressed in the stele and in differentiating columella cells. *cle40* loss of function mutants had short roots with irregular tips caused by delayed differentiation of columella stem cells. Excessive CLE40 led to columella stem cell differentiation. The expression of *WOX5* was perturbed when CLE40 was deregulated, extending beyond the QC in the *cle40* mutant and being reduced in the QC and shifted toward the proximal meristem upon CLE40 peptide addition to WT plants. CLE40, together with ACR4 (*Arabidopsis* CRINKLY 4), a receptor-like kinase, regulated the spatial expression of *WOX5* to maintain columella stem cell identity. Additionally, the *CLV2* (*CLAVATA 2*) leucine-rich repeat receptor was found to be necessary for CLE40-dependent root growth suppression, but not for differentiation of the daughters of columella stem cells (Stahl et al., 2009), indicating that the same peptide may interact with more than one receptor to regulate different developmental processes. Indeed, further investigation showed that subcellular localization, environment, and concentration affect the formation of receptor complexes (Stahl et al., 2013). The leucine-rich repeat receptor kinase *CLV1* was shown to be involved in the signaling pathway of CLE40 and ACR4 (Figure 1D). *ACR4* and *CLV1* have similar expression patterns in the root meristem. CLE40 regulates *ACR4* expression but not *CLV1* expression. Using Förster resonance energy transfer and multiparameter fluorescence image spectroscopy, the transmembrane domains of ACR4 and CLV1 were shown to interact and form homomeric and heteromeric complexes depending on localization (plasma membrane or plasmodesmata). Both the *clv1* and *acr4* single mutants were impaired in aspects of meristem maintenance and responded differently to CLE40 peptide application. This indicated that the homo- and heteromeric complexes differentially regulate root meristem maintenance.

In another recent report, the sulfated RGF1 peptide was found to regulate meristem maintenance by providing a link between *PLT* expression and auxin through sulfation by TPST1 (TYROSYL-PROTEIN SULFOTRANSFERASE 1; Figure 1G; Matsuzaki et al., 2010; Zhou et al., 2010). The *tpst1* mutant displayed pleiotropic effects including decreased meristem size and cell division activity, QC misregulation and starch granule accumulation. This mutant lacked the only copy of tyrosylprotein sulfotransferase found in the *Arabidopsis* genome (Matsuzaki et al., 2010; Zhou et al., 2010). *TPST1* was induced by auxin and helped to maintain auxin distribution by regulating expression of *PIN-FORMED* (*PIN*) and auxin biosynthetic genes (Zhou et al., 2010).

The RGF peptide family was found by a bioinformatic search for putatively sulfated peptide-encoding genes (Matsuzaki et al., 2010). Applying synthetic, sulfated RGF1 peptide to *tpst1* roots ameliorated the stem cell misregulation phenotype and increased cell division activity, indicating that sulfated RGF1 was required for these processes. Further investigation showed the sulfated peptides PSK (PHYTOSULFOKINE) and PSY1 (PLANT PEPTIDE

CONTAINING SULFATED TYROSINE 1) were also required for full restoration of cell division activity. Eight of the nine sulfated RGF peptides complemented the *tpst1* mutant in this way, however, non-sulfated peptides were unable to do so. It was found that RGF1 acts by regulating spatial expression and expression levels of *PLT* transcription factors. This work provided a link between auxin levels, the RGF peptide and *PLT* expression through TPST1. These studies also exemplified the extra layer of regulation that can be provided by post-translational modification of signaling peptides.

The *CEP* family has also been shown to affect root meristem development (Ohyama et al., 2008). Several forms of cleaved and modified peptides were confirmed using mass spectrometry on *CEP1* overexpression lines. Synthetic 15 amino acid CEP1 peptide application or *CEP1* overexpression arrested root growth. Meristematic cell division and expansion was repressed by CEP1, although QC specification was not affected. A more detailed analysis of the CEP family and the mode of action of individual CEPs is required.

### VASCULAR DEVELOPMENT

Once cells leave the meristematic zone of the root, they go through phases of elongation and differentiation into specific tissue types. Procambial and cambial cells differentiate into xylem and phloem, the vascular tissues required for long-distance transport of water and nutrients through the plant. Several regulatory peptides are known to play important roles in vascular development, including the CLE-family member TDIF (TRACHEARY ELEMENT DIFFERENTIATION INHIBITORY FACTOR; Ito et al., 2006). More recently, another member of this family, CLE10, was found to inhibit protoxylem formation through interaction with cytokinin signaling (Figure 1F; Kondo et al., 2011).

A screen for *CLE* genes up-regulated during vascular development together with synthetic peptide assays implicated *CLE10* in early xylem differentiation (Kondo et al., 2011). *CLE10* peptide application rapidly down-regulated the transcription of *ARR5* and *ARR6*, two type A *Arabidopsis* RESPONSE REGULATORS that negatively regulate cytokinin signaling. This indicated that cytokinin signaling was increased by *CLE10*, leading to decreased protoxylem development, one of the documented roles of cytokinin. The leucine-rich repeat receptor-like protein CLV2 and protein kinase CRN (CORYN)/SOL2 (SUPPRESSOR OF LLP2) were both required for *CLE10* suppression of *ARR5* and *ARR6* transcription, indicating they may be the receptor module for the *CLE10* peptide.

### GRAVITROPIC RESPONSE

Response to gravity stimulus is an essential component of root development. This process relies on gravity sensing in the root tip together with manipulation of auxin gradients, mediated by the localization of the *AUX1/LIKE-AUX1* family of auxin influx carriers and the *PIN* family of auxin efflux carriers (Vieten et al., 2007). The GLV/RGF peptide family has been implicated in this process (Figure 1C). Using a reverse genetic screen, Whitford et al. (2012) found that *GLV1/RGF6*, *GLV2/RGF9*, and *GLV3/RGF4* overexpression (or sulfated peptide addition) caused an altered gravitropic response in roots. *GLV1/RGF6* and *GLV3/RGF4*

post-transcriptionally regulated *PIN2* within minutes of peptide application, presenting evidence that GLV/RGF signaling controls the trafficking and stability of *PIN2*. A perturbation in the *PIN2* pool would abolish the finely tuned auxin gradient necessary for normal gravitropic response. A loss of function *glv3/rgf4* mutant displayed gravitropic defects (Fernandez et al., 2013) implying *GLV3/RGF4* is required for correct gravitropic response.

In a separate study, *CLEL/RGF* genes were found by homology with *CLE18*, a peptide precursor gene with a CLE motif in the variable region and a CLEL/RGF peptide domain at the C-terminus (Meng et al., 2012a). Unmodified *CLEL6/RGF6* and *CLEL8/RGF1* peptide application and *CLEL6/RGF6* and *CLEL7/RGF5* overexpression induced long roots on WT plants as well as an additional “wavy” root phenotype. This wavy root phenotype was reported to be independent of thigmotropism and phototropism. It was also reported to be independent of gravitropism as the phenotype persisted when unmodified *CLEL6/RGF6* peptide was applied to *eir1-1* and *aux1-7* mutants, which are impaired in the gravitropic response. This is in disagreement with the aforementioned study which implicates *CLEL6/GLV1/RGF6* in gravitropic response.

Interestingly, the peptide-encoding gene at the center of this debate, *CLEL6/GLV1/RGF6*, is reportedly not expressed in any part of the root (Whitford et al., 2012; Fernandez et al., 2013). Furthermore, a *GLV1/RGF6* knockdown line did not show impaired gravitropic response (Fernandez et al., 2013). This indicates regulation of peptide gene expression is paramount to its *in planta* function. Excessive peptide treatment by overexpression or peptide application should be interpreted in a biologically relevant context and should be supported by data from loss of function mutants.

### LATERAL ROOT DEVELOPMENT

The formation of LR allows the plant to exploit water and nutrients in the surrounding soil. LRs arise from a repetitive process which begins in the root meristem region. The earliest described event in LR development is priming or pre-branch site formation. This process requires auxin (De Smet et al., 2007) and the oscillation of over 3,000 genes (Moreno-Risueno et al., 2010). As the cells move into the differentiation zone, a site of relatively low auxin levels, LR founder cells are specified from xylem pole pericycle cells (Dubrovsky et al., 2008, 2011). The founder cells are initiated in an auxin-dependent manner (Dubrovsky et al., 2011) and begin to undergo a series of divisions. Eight stages of LR development have been defined (Malamy and Benfey, 1997), from the first asymmetric division (stage I) through many subsequent rounds of cell division (stages II–VII), to LR emergence (stage VIII). In order for the nascent LR to pass through the cortical, endodermal, and epidermal cell layers before emerging, auxin-dependent degradation and remodeling of cell walls is required (Swarup et al., 2008).

Peptides from the RGF family were reported to regulate LR development in an auxin-independent manner (Figure 1A; Meng et al., 2012a). Overexpression of *CLEL6/RGF6* or *CLEL7/RGF5* resulted in a significant reduction in LR number, due to abnormal cell divisions at stage I of LR development. The authors speculated that *CLEL6/RGF6* and *CLEL7/RGF5* may interact with the

receptor kinase ACR4, as it too plays an important role in the early stages on LR development. Other RGF genes were found to be specifically expressed during different stages of LR development (Fernandez et al., 2013). Promoter reporter constructs showed that *GLV6/RGF8* was active from stage I of LR development, *GLV5/RGF2* and *GLV10/RGF5* at stage II, *GLV7/RGF3* and *GLV11/RGF1* at stage IV, *GLV3/RGF4* at stage V, and *GLV9* and *GLV2/RGF9* after emergence. Overexpression of these genes resulted in decreased LR number as well as root waving and enlarged root meristems. These data indicate that RGF peptides may act at different stages of LR development, however, specific mechanisms remain to be elucidated.

The IDA peptide and its receptors, HAE (HAESA) and HSL2 (HAESA-Like 2) were recently shown to play a role in LR emergence (Figure 1E; Kumpf et al., 2013) in addition to their known roles in floral abscission (Butenko et al., 2003; Stenvik et al., 2008). The *ida*, *hae*, *hsl2* single mutants, and *hae hsl2* double mutant showed a significant reduction in the number of LR (Kumpf et al., 2013). LR primordia in these mutants encountered difficulties in penetrating the cortical, endodermis, and epidermal cell layers and often displayed irregular flattened shapes. IDA was strongly induced by auxin, whereas the two receptors were only transiently induced by auxin, indicating the receptors are used to limit IDA function. Two stages for IDA-mediated cell wall remodeling (CWR) were identified. During early primordia development (stage I and II), auxin from the LR primordium induced IDA expression in the endodermal cells, where HAE and HSL2 were already present. IDA signaling led to the expression of CWR enzymes, which allowed the nascent LR to pass through the endodermal cell layer. At a later stage in primordia development (stage V), auxin was derived from the auxin influx carrier LIKE AUX1-3 (LAX3), expressed in the neighboring cortical and epidermal cells. This induced the degradation of SOLITARY ROOT1 which in turn released the transcription factors AUXIN RESPONSE FACTOR (ARF) 7 and 19. ARF7 was required for the subsequent induction of IDA that triggered the expression of CWR genes through HAE/HSL2 signaling to allow the primordia to emerge from the parent root.

## OTHER ROLES OF SMALL SIGNALING PEPTIDES IN ROOT DEVELOPMENT

A recent report described *CLELn* (CLE-Like protein in the nucleus), a RGF family gene that lacked the archetypal N-terminal signal sequence (Meng et al., 2012b). A green fluorescent protein (GFP)-fusion assay suggested this peptide localizes to the nucleus and western blots indicated it is specifically processed from the precursor. Overexpression of this gene gave a long root phenotype similar to *GLV2/RGF9* overexpression, however, synthetic peptide application elicited the long and wavy phenotype seen when *GLV1/RGF6* and *GLV11/RGF1* synthetic peptides were assayed. These data raise the possibility that regulatory peptides do not act solely as intercellular signals and may play roles in nuclear signaling or are secreted by non-conventional routes.

Regulatory peptides also play important roles in nodule organogenesis in legumes. Nodules form in response to infection by symbiotic bacteria called rhizobia. This process requires systemic regulation (autoregulation of nodulation; Caetano-Anolles

and Gresshoff, 1991), a process that has parallels with shoot meristem regulation in *Arabidopsis* by *CLV3* and *CLV1* and their downstream processes (Mayer et al., 1998; Brand et al., 2000; Schoof et al., 2000; Leibfried et al., 2005; Osipova et al., 2012). Two root-specific pathways that involve Nod factors and cytokinin signaling are also required for nodulation. *Medicago truncatula* *CLE12* and *CLE13* are up-regulated in nodulating roots (Mortier et al., 2010). Upon overexpression, wild type plants do not form nodules. Suppression of nodulation by *MtCLE12* and *MtCLE13* is dependent upon *SUNN* (*SUPER NUMERIC NODULES*, orthologous to *CLAVATA1* in *Arabidopsis*) and induces type A response regulators, leading to cytokinin signaling (Mortier et al., 2010; Saur et al., 2011). Similar results were found in *Glycine max* (Reid et al., 2011) and *Lotus japonicus* (Okamoto et al., 2009), indicating at least two CLE peptides play an essential role in the autoregulation of nodulation. Recently a nodule-specific CLE in *L. japonicus* (CLE-RS2) was shown to be a root produced arabinosylated peptide (Okamoto et al., 2013). It interacted with HAR1, which shares functional similarity with *SUNN*, in an arabinose chain and sequence-dependent manner. CLE-RS2 was found in shoot-collected xylem sap, indicating that the CLE-RS glycopeptide is a long distance, root-to-shoot signal that controls autoregulation. These exciting results add a new dimension to how regulatory peptides control root development.

## FUTURE PERSPECTIVES

Although a number of peptides from the CLE and RGF families have been implicated in different aspects of root development, the specific function and mechanistic action of other family members remains to be elucidated (Figure 1B). In a step toward this, the expression patterns and effects of misregulation of all RGF and CLE peptide family members have been assessed (Jun et al., 2010; Fernandez et al., 2013). Further work is required to elucidate the functions of the CEP family in root development.

While genetic redundancy has hindered elucidation of peptide function by loss of function mutants, a recently reported technology may assist in overcoming this. It was shown that by substituting Gly6 in the *CLV3* peptide domain for Ala or Thr, dominant-negative *clv3* phenotypes were obtained (Song et al., 2013). When applied in combination with the unsubstituted *CLV3* peptide, it was shown that the antagonistic effect was a result of competition between the two peptides. It was hypothesized that *CLV3*<sup>Thr6</sup> was able to bind the *CLV3* receptor without eliciting a response. This concept was also used to make antagonists for *CLE8* and *CLE22* peptides. This technology may assist in elucidating the function of specific regulatory peptides.

As highlighted in this review, small signaling peptides play important roles in root development. There are several layers of regulation which serve to add specificity to the roles of individual peptides. These include post-translational modification, tissue-specific expression, regulation of receptor expression, the subcellular localization of the receptor and the potential for long-distance movement. Recent bioinformatic approaches have indicated that over 7,000 small, unannotated open reading frames exist in the *A. thaliana* genome (Hanada et al., 2013). As proportion of these are likely to be regulatory peptide-encoding genes,

there is still a long way to go in fully exploring the extent of peptide-mediated developmental regulation.

## ACKNOWLEDGMENTS

Christina Delay was supported by an Australia Postgraduate Award and GRDC Grains Industry Research Scholarship (GRS10329).

## REFERENCES

- Aida, M., Beis, D., Heidstra, R., Willemssen, V., Blilou, I., Galinha, C., et al. (2004). The PLETHORA genes mediate patterning of the *Arabidopsis* root stem cell niche. *Cell* 119, 109–120. doi: 10.1016/j.cell.2004.09.018
- Brand, U., Fletcher, J. C., Hobe, M., Meyerowitz, E. M., and Simon, R. (2000). Dependence of stem cell fate in *Arabidopsis* on a feedback loop regulated by CLV3 activity. *Science* 289, 617–619. doi: 10.1126/science.289.5479.617
- Butenko, M. A., Patterson, S. E., Grini, P. E., Stenvik, G.-E., Amundsen, S. S., Mandal, A., et al. (2003). INFLORESCENCE DEFICIENT IN ABSCISSION controls floral organ abscission in *Arabidopsis* and identifies a novel family of putative ligands in plants. *Plant Cell* 15, 2296–2307. doi: 10.1105/tpc.014365
- Caetano-Anolles, G., and Gresshoff, P. M. (1991). Plant genetic control of nodulation. *Annu. Rev. Microbiol.* 45, 345–382. doi: 10.1146/annurev.mi.45.100191.002021
- Casimiro, I., Beeckman, T., Graham, N., Bhalerao, R., Zhang, H., Casero, P., et al. (2003). Dissecting *Arabidopsis* lateral root development. *Trends Plant Sci.* 8, 165–171. doi: 10.1016/S1360-1385(03)00051-7
- Cock, J. M., and McCormick, S. (2001). A large family of genes that share homology with CLAVATA3. *Plant Physiol.* 126, 939–942. doi: 10.1104/pp.126.3.939
- De Smet, I., Tetsumura, T., De Rybel, B., Frey, N. F. D., Laplace, L., Casimiro, I., et al. (2007). Auxin-dependent regulation of lateral root positioning in the basal meristem of *Arabidopsis*. *Development* 134, 681–690. doi: 10.1242/dev.02753
- Di Lorenzo, L., Wysocka-Diller, J., Malamy, J. E., Pysh, L., Helariutta, Y., Freshour, G., et al. (1996). The SCARECROW gene regulates an asymmetric cell division that is essential for generating the radial organization of the *Arabidopsis* root. *Cell* 86, 423–433. doi: 10.1016/S0092-8674(00)80115-4
- Dubrovsky, J. G., Napsucialy-Mendivil, S., Duclercq, J., Cheng, Y., Shishkova, S., Ivanchenko, M. G., et al. (2011). Auxin minimum defines a developmental window for lateral root initiation. *New Phytol.* 191, 970–983. doi: 10.1111/j.1469-8137.2011.03757.x
- Dubrovsky, J. G., Sauer, M., Napsucialy-Mendivil, S., Ivanchenko, M. G., Friml, J., Shishkova, S., et al. (2008). Auxin acts as a local morphogenetic trigger to specify lateral root founder cells. *Proc. Natl. Acad. Sci. U.S.A.* 105, 8790–8794. doi: 10.1073/pnas.0712307105
- Fernandez, A., Drozdzecki, A., Hoogewijs, K., Nguyen, A., Beeckman, T., Madder, A., et al. (2013). Transcriptional and functional classification of the GLV/RGF/CLEL signaling peptides reveals their role in lateral root and hair formation. *Plant Physiol.* 161, 954–970. doi: 10.1104/pp.112.206029
- Friml, J., Vieten, A., Sauer, M., Weijers, D., Schwarz, H., Hamann, T., et al. (2003). Efflux-dependent auxin gradients establish the apical-basal axis of *Arabidopsis*. *Nature* 426, 147–153. doi: 10.1038/nature02085
- Fukuda, H., and Higashiyama, T. (2011). Diverse functions of plant peptides: entering a new phase. *Plant Cell Physiol.* 52, 1–4. doi: 10.1093/pcp/pcq193
- Galinha, C., Hofhuis, H., Luijten, M., Willemsen, V., Blilou, I., Heidstra, R., et al. (2007). PLETHORA proteins as dose-dependent master regulators of *Arabidopsis* root development. *Nature* 449, 1053–1057. doi: 10.1038/nature06206
- Haecker, A., Groß-Hardt, R., Geiges, B., Sarkar, A., Breuninger, H., Herrmann, M., et al. (2004). Expression dynamics of WOX genes mark cell fate decisions during early embryonic patterning in *Arabidopsis thaliana*. *Development* 131, 657–668. doi: 10.1242/dev.00963
- Hanada, K., Higuchi-Takeuchi, M., Okamoto, M., Yoshizumi, T., Shimizu, M., Nakaminami, K., et al. (2013). Small open reading frames associated with morphogenesis are hidden in plant genomes. *Proc. Natl. Acad. Sci. U.S.A.* 110, 2395–2400. doi: 10.1073/pnas.1213958110
- Helariutta, Y., Fukaki, H., Wysocka-Diller, J., Nakajima, K., Jung, J., Sena, G., et al. (2000). The SHORT-ROOT gene controls radial patterning of the *Arabidopsis* root through radial signaling. *Cell* 101, 555–567. doi: 10.1016/S0092-8674(00)80865-X
- Hirakawa, Y., Shinohara, H., Kondo, Y., Inoue, A., Nakanomyo, I., Ogawa, M., et al. (2008). Non-cell-autonomous control of vascular stem cell fate by a CLE peptide/receptor system. *Proc. Natl. Acad. Sci. U.S.A.* 105, 15208–15213. doi: 10.1073/pnas.0808444105
- Ito, Y., Nakanomyo, I., Motose, H., Iwamoto, K., Sawa, S., Dohmae, N., et al. (2006). Dodeca-CLE peptides as suppressors of plant stem cell differentiation. *Science* 313, 842–845. doi: 10.1126/science.1128436
- Jun, J., Fiume, E., Roeder, A. H. K., Meng, L., Sharma, V. K., Osmont, K. S., et al. (2010). Comprehensive analysis of CLE polypeptide signaling gene expression and over-expression activity in *Arabidopsis*. *Plant Physiol.* 154, 1721–1736. doi: 10.1104/pp.110.163683
- Kondo, Y., Hirakawa, Y., Kieber, J. J., and Fukuda, H. (2011). CLE peptides can negatively regulate protoxylem vessel formation via cytokinin signaling. *Plant Cell Physiol.* 52, 37–48. doi: 10.1093/pcp/pcq129
- Kumpf, R. P., Shi, C.-L., Larrieu, A., Stø, I. M., Butenko, M. A., Péret, B., et al. (2013). Floral organ abscission peptide IDA and its HAE/HSL2 receptors control cell separation during lateral root emergence. *Proc. Natl. Acad. Sci. U.S.A.* 110, 5235–5240. doi: 10.1073/pnas.1210835110
- Leibfried, A., To, J. P. C., Busch, W., Stehling, S., Kehle, A., Demar, M., et al. (2005). WUSCHEL controls meristem function by direct regulation of cytokinin-inducible response regulators. *Nature* 438, 1172–1175. doi: 10.1038/nature04270
- Malamy, J. E. (2005). Intrinsic and environmental response pathways that regulate root system architecture. *Plant Cell Environ.* 28, 67–77. doi: 10.1111/j.1365-3040.2005.01306.x
- Malamy, J. E., and Benfey, P. N. (1997). Organization and cell differentiation in lateral roots of *Arabidopsis thaliana*. *Development* 124, 33–44.
- Matsubayashi, Y., Ogawa, M., Morita, A., and Sakagami, Y. (2002). An LRR receptor kinase involved in perception of a peptide plant hormone, phytosulfokine. *Science* 296, 1470–1472. doi: 10.1126/science.1069607
- Matsuzaki, Y., Ogawa-Ohnishi, M., Mori, A., and Matsubayashi, Y. (2010). Secreted peptide signals required for maintenance of root stem cell niche in *Arabidopsis*. *Science* 329, 1065–1067. doi: 10.1126/science.1191132
- Mayer, K. F. X., Schoof, H., Haecker, A., Lenhard, M., Jürgens, G., and Laux, T. (1998). Role of WUSCHEL in regulating stem cell fate in the *Arabidopsis* shoot meristem. *Cell* 95, 805–815. doi: 10.1016/S0092-8674(00)81703-1
- Meng, L., Buchanan, B. B., Feldman, L. J., and Luan, S. (2012a). CLE-like (CLEL) peptides control the pattern of root growth and lateral root development in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 109, 1760–1765. doi: 10.1073/pnas.1119864109
- Meng, L., Buchanan, B. B., Feldman, L. J., and Luan, S. (2012b). A putative nuclear CLE-Like (CLEL) peptide precursor regulates root growth in *Arabidopsis*. *Mol. Plant* 5, 955–957. doi: 10.1093/mp/sss060
- Moreno-Risueno, M. A., Van Norman, J. M., Moreno, A., Zhang, J., Ahnert, S. E., and Benfey, P. N. (2010). Oscillating gene expression determines competence for periodic *Arabidopsis* root branching. *Science* 329, 1306–1311. doi: 10.1126/science.1191937
- Mortier, V., Den Herder, G., Whitford, R., Van De Velde, W., Rombauts, S., D'haeseleer, K., et al. (2010). CLE peptides control *Medicago truncatula* nodulation locally and systemically. *Plant Physiol.* 153, 222–237. doi: 10.1104/pp.110.153718
- Ni, J., Guo, Y., Jin, H., Hartsell, J., and Clark, S. (2011). Characterization of a CLE processing activity. *Plant Mol. Biol.* 75, 67–75. doi: 10.1007/s11103-010-9708-2
- Ogawa, M., Shinohara, H., Sakagami, Y., and Matsubayashi, Y. (2008). *Arabidopsis* CLV3 peptide directly binds CLV1 ectodomain. *Science* 319, 294. doi: 10.1126/science.1150083
- Ohyama, K., Ogawa, M., and Matsubayashi, Y. (2008). Identification of a biologically active, small, secreted peptide in *Arabidopsis* by in silico gene screening, followed by LC-MS-based structure analysis. *Plant J.*

This work was supported by an Australian Research Council grant to Michael A. Djordjeric and Nijat Imin (DP140103714).

## CONTRIBUTION

Christina Delay wrote the paper. Nijat Imin and Michael A. Djordjeric critically read and edited the paper.

- 55, 152–160. doi: 10.1111/j.1365-313X.2008.03464.x
- Okamoto, S., Ohnishi, E., Sato, S., Takahashi, H., Nakazono, M., Tabata, S., et al. (2009). Nod factor/nitrate-induced CLE genes that drive HARI-mediated systemic regulation of nodulation. *Plant Cell Physiol.* 50, 67–77. doi: 10.1093/pcp/pcn194
- Okamoto, S., Shinohara, H., Mori, T., Matsubayashi, Y., and Kawaguchi, M. (2013). Root-derived CLE glycopeptides control nodulation by direct binding to HARI receptor kinase. *Nat. Commun.* 4, article number 2191. doi: 10.1038/ncomms3191
- Osipova, M. A., Mortier, V., Demchenko, K. N., Tsyganov, V. E., Tikhonovich, I. A., Lutova, L. A., et al. (2012). WUSCHEL-RELATED HOMEODOMAIN5 gene expression and interaction of CLE peptides with components of the systemic control add two pieces to the puzzle of autoregulation of nodulation. *Plant Physiol.* 158, 1329–1341. doi: 10.1104/pp.111.188078
- Reid, D. E., Ferguson, B. J., and Gresshoff, P. M. (2011). Inoculation- and nitrate-induced CLE peptides of Soybean control NARK-dependent nodule formation. *Mol. Plant Microbe Interact.* 24, 606–618. doi: 10.1094/MPMI-09-10-0207
- Sabatini, S., Beis, D., Wolkenfelt, H., Murfett, J., Guilfoyle, T., Malamy, J., et al. (1999). An auxin-dependent distal organizer of pattern and polarity in the *Arabidopsis* root. *Cell* 99, 463–472. doi: 10.1016/S0092-8674(00)81535-4
- Sabatini, S., Heidstra, R., Wildwater, M., and Scheres, B. (2003). SCARECROW is involved in positioning the stem cell niche in the *Arabidopsis* root meristem. *Genes Dev.* 17, 354–358. doi: 10.1101/gad.252503
- Saur, I. M. L., Oakes, M., Djordjevic, M. A., and Imin, N. (2011). Crosstalk between the nodulation signaling pathway and the autoregulation of nodulation in *Medicago truncatula*. *New Phytol.* 190, 865–874. doi: 10.1111/j.1469-8137.2011.03738.x
- Schoof, H., Lenhard, M., Haecker, A., Mayer, K. F. X., Jürgens, G., and Laux, T. (2000). The stem cell population of *Arabidopsis* shoot meristems is maintained by a regulatory loop between the CLAVATA and WUSCHEL genes. *Cell* 100, 635–644. doi: 10.1016/S0092-8674(00)80700-X
- Song, X.-F., Guo, P., Ren, S.-C., Xu, T.-T., and Liu, C.-M. (2013). Antagonistic peptide technology for functional dissection of CLV3/ESR genes in *Arabidopsis*. *Plant Physiol.* 161, 1076–1085. doi: 10.1104/pp.112.211029
- Stahl, Y., Grabowski, S., Bleckmann, A., Kühnemuth, R., Weidtkamp-Peters, S., Pinto, K. G., et al. (2013). Moderation of *Arabidopsis* root stemness by CLAVATA1 and ARABIDOPSIS CRINKLY4 receptor kinase complexes. *Curr. Biol.* 23, 362–371. doi: 10.1016/j.cub.2013.01.045
- Stahl, Y., Wink, R. H., Ingram, G. C., and Simon, R. (2009). A signaling module controlling the stem cell niche in *Arabidopsis* root meristems. *Curr. Biol.* 19, 909–914. doi: 10.1016/j.cub.2009.03.060
- Stenvik, G.-E., Tandstad, N. M., Guo, Y., Shi, C.-L., Kristiansen, W., Holmgren, A., et al. (2008). The EPIP peptide of INFLORESCENCE DEFICIENT IN ABSCISSION is sufficient to induce abscission in *Arabidopsis* through the receptor-like kinases HAESA and HAESA-LIKE2. *Plant Cell* 20, 1805–1817. doi: 10.1105/tpc.108.059139
- Swarup, K., Benkova, E., Swarup, R., Casimiro, I., Peret, B., Yang, Y., et al. (2008). The auxin influx carrier LAX3 promotes lateral root emergence. *Nat. Cell Biol.* 10, 946–954. doi: 10.1038/ncb1754
- Vieten, A., Sauer, M., Brewer, P. B., and Friml, J. (2007). Molecular and cellular aspects of auxin-transport-mediated development. *Trends Plant Sci.* 12, 160–168. doi: 10.1016/j.tplants.2007.03.006
- Whitford, R., Fernandez, A., Tejos, R., Pérez, A. C., Kleine-Vehn, J., Vanneste, S., et al. (2012). GOLVEN secretory peptides regulate auxin carrier turnover during plant gravitropic responses. *Dev. Cell* 22, 678–685. doi: 10.1016/j.devcel.2012.02.002
- Zhou, W., Wei, L., Xu, J., Zhai, Q., Jiang, H., Chen, R., et al. (2010). *Arabidopsis* tyrosylprotein sulfotransferase acts in the auxin/PLETHORA pathway in regulating postembryonic maintenance of the root stem cell niche. *Plant Cell* 22, 3692–3709. doi: 10.1105/tpc.110.075721

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 04 July 2013; accepted: 21 August 2013; published online: 06 September 2013.

Citation: Delay C, Imin N and Djordjevic MA (2013) Regulation of *Arabidopsis* root development by small signaling peptides. *Front. Plant Sci.* 4:352. doi: 10.3389/fpls.2013.00352

This article was submitted to *Plant Systems Biology*, a section of the journal *Frontiers in Plant Science*.

Copyright © 2013 Delay, Imin and Djordjevic. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.