Correlation energy of two electrons in a ball

Pierre-François Loos^{a)} and Peter M. W. Gill^{b)}

Research School of Chemistry, Australian National University, Canberra, Australian Capital Territory 0200, Australia

(Received 7 April 2010; accepted 28 May 2010; published online 17 June 2010)

We study the ground-state correlation energy E_c of two electrons of opposite spin confined within a D -dimensional ball ($D \geq 2$) of radius R . In the high-density regime, we report accurate results for the exact and restricted Hartree–Fock energy, using a Hylleraas-type expansion for the former and a simple polynomial basis set for the latter. By investigating the exact limiting correlation energy $E_c^{(0)} = \lim_{R \rightarrow 0} E_c$ for various values of D , we test our recent conjecture [P.-F. Loos and P. M. W. Gill, *J. Chem. Phys.* **131**, 241101 (2009)] that in the large- D limit, $E_c^{(0)} \sim -\delta^2/8$ for any spherically symmetric confining external potential, where $\delta = 1/(D-1)$. © 2010 American Institute of Physics. [doi:10.1063/1.3455706]

I. INTRODUCTION

In the early days of quantum chemistry, there was considerable interest in cavity-confined atoms as a model for high-density atomic gas^{1–3} and extrapolation of high-density results provides a convenient but powerful route to understanding the intermediate-density regime.⁴ Thanks to Hylleraas' work,⁵ the compressed heliumlike ions have been widely studied^{6,7} and interest in these continues unabated.^{8–12} Other confined systems such as electrons in square,^{13,14} cylindrical,¹⁵ and spherical^{16–21} boxes have also attracted attention. The last of these has been extensively used for the assessment of density-functional approximations^{16–18} and the study of Wigner molecules²² at low densities.^{19–21}

In a previous article,²³ we studied the high-density correlation energy $E_c^{(0)}$ for various two-electron systems confined to a D -dimensional space ($D \geq 2$) by an external potential $V(r) \propto r^m$. As the high-density limit sheds light on intermediate densities, the large-dimension limit provides useful insights into the $D=3$ case.^{24,25} For the heliumlike ions ($m=-1$), spherium atoms^{26–34} ($m=0$), and Hooke's law atoms^{35–38} ($m=2$), we found that in the large- D limit,

$$E_c^{(0)} \sim -\delta^2/8 - C\delta^3, \quad (1)$$

where $\delta = 1/(D-1)$ and the coefficient $C \approx 1/6$ varies slowly with m . On this basis, we conjectured that Eq. (1) is true for any spherically symmetric confining external potential.

At the end of our previous work,²³ we observed that it would be highly desirable to consider D -ballium, the system in which the two electrons are trapped in a D -dimensional ball of radius R . This model is a severe test of our conjecture because it corresponds to $m=\infty$.

The present study focuses mainly on the high-density regime ($R > 0$ but small) and the corresponding limiting case ($R=0$). We report accurate results for the restricted Hartree–Fock (HF) and exact energies (Secs. II and III, respectively). For the limiting case (Sec. IV), perturbation theory is used to

expand both the HF and exact energies and this allows us to determine the limiting correlation energy in D -ballium. We use atomic units throughout.

The Hamiltonian of D -ballium is

$$\hat{H} = -\frac{\nabla_1^2}{2} - \frac{\nabla_2^2}{2} + V(r_1) + V(r_2) + \frac{1}{r_{12}}, \quad (2)$$

where $r_{12} = |\mathbf{r}_1 - \mathbf{r}_2|$ is the interelectronic distance and the external potential is defined by

$$V(r) = \begin{cases} 0, & \text{if } r < R, \\ \infty, & \text{otherwise.} \end{cases} \quad (3)$$

Any physically acceptable eigenfunction of Eq. (2) must satisfy the Dirichlet boundary condition

$$\Psi(r_1 = R) = \Psi(r_2 = R) = 0. \quad (4)$$

II. RESTRICTED HF APPROXIMATION

The spin-restricted HF solution,³⁹ which is the only HF solution in the high-density regime, is given by

$$\Psi_{\text{HF}}(r_1, r_2) = \phi(r_1)\phi(r_2). \quad (5)$$

If we introduce the scaled coordinate $t = r/R$, the HF orbital $\phi(t)$ is an eigenfunction of the Fock operator

$$\hat{F} = -\frac{1}{2R^2}\nabla_t^2 + \frac{1}{R}J_\phi(t). \quad (6)$$

For S states in a D -dimensional space,^{23,40} we have

$$\nabla_t^2 = \frac{d^2}{dt^2} + \frac{D-1}{t} \frac{d}{dt}, \quad (7)$$

$$J_\phi(t) = \int_0^1 \frac{\phi(x)^2}{\max(t, x)} F\left[\frac{3-D}{2}, \frac{1}{2}, \frac{D}{2}, \alpha^2\right] x^{D-1} dx, \quad (8)$$

where $\alpha = \min(t, x)/\max(t, x)$ and F is the hypergeometric function.

^{a)}Electronic mail: loos@rsc.anu.edu.au.

^{b)}Electronic mail: peter.gill@anu.edu.au.

Unlike Thompson and Alavi,²¹ who expanded the HF orbital in a basis of spherical Bessel functions,⁴¹ we chose to explore an even-degree polynomial basis, writing

$$\phi(t) = (1 - t^2) \sum_{k=0}^{N-1} c_k t^{2k}. \quad (9)$$

Any such orbital is smooth at the center of the ball, i.e.,

$$\phi'(0) = 0, \quad (10)$$

and is cusped and vanishes at the boundary, i.e.,

$$\phi'(1) < 0, \quad \phi(1) = 0. \quad (11)$$

It can be shown that the resulting HF energy is

$$E_{\text{HF}} = \frac{1}{R^2} \frac{T}{S} + \frac{1}{R} \frac{U}{S^2}, \quad (12)$$

with

$$S = \sum_{ij} \frac{c_i c_j}{\left(i + j + \frac{D}{2}\right)_3}, \quad (13)$$

$$T = \sum_{ij} c_i c_j \left[\frac{D}{\left(i + j + \frac{D}{2}\right)_2} + \frac{4ij}{\left(i + j + \frac{D}{2} - 1\right)_3} \right], \quad (14)$$

$$U = \sum_{ijkl} c_i c_j c_k c_l [\beta_{i+j+k+l-1/2} \gamma_{k+l} - 2\beta_{i+j+k+l+1/2} \gamma_{k+l+1} + \beta_{i+j+k+l+3/2} \gamma_{k+l+2}], \quad (15)$$

where

$$(a)_b = \frac{\Gamma(a+b)}{\Gamma(a)} \quad (16)$$

is the Pochhammer symbol and Γ is the Gamma function.⁴¹ The coefficients β_n and γ_n are given by

$$\beta_n = \frac{1}{(n+D)_3}, \quad (17)$$

$$\gamma_n = \frac{{}_3F_2\left(\frac{3-D}{2}, \frac{1}{2}, n + \frac{D}{2}; \frac{D}{2}, n + 1 + \frac{D}{2}; 1\right)}{n + \frac{D}{2}}, \quad (18)$$

where ${}_pF_q$ is the generalized hypergeometric function.⁴¹ For $D=3$, the hypergeometric functions reduce to unity.

The energy (12) can be minimized with respect to the coefficients c_k using a numerical solver,⁴² thus avoiding the self-consistent field procedure usually needed for this kind of calculation.⁴³ Henceforth, we define the accuracy of an energy E as

$$\mathcal{A} = -\log_{10}[(E - E^*)/E^*], \quad (19)$$

where E^* is our best estimate of the exact energy. In loose terms, $\mathcal{A}$ is the number of correct decimal digits.

Figure 1 shows how the accuracy of the HF energy of 3-ballium ($R=1$) improves as N increases. For very small N ,

FIG. 1. Accuracy $\mathcal{A}$ of the HF energy of 3-ballium ($R=1$) with respect to basis set size N . TA results taken from Ref. 21 and LG from the present study.

the spherical Bessel basis²¹ (TA) is more accurate than the polynomial (LG) basis. However, although both the TA and LG bases seem to exhibit exponential convergence as N increases, the TA energy improves by roughly one order of magnitude and the LG energy by roughly two orders of magnitude as each basis function is added. As a result, one obtains the HF energy to 20 digits using Eq. (9) with $N=10$. However, the origin of the superiority of the polynomial basis is not clear. We find that the resulting expansion coefficients c_k decay roughly exponentially and the convergence behavior for other D is similar.

Numerical results for 3-ballium ($R=1$) are shown in Table I. For $N=7$ basis functions, the Bessel and polynomial bases yield HF energies that lie 13 nE_h and 2.8 pE_h above the HF limit, respectively. Analogous behavior is observed for the larger values of R , including those that lie in the low-density regime where a lower-energy Unrestricted Hartree-Fock (UHF) solution exists ($R \geq 6$ for $D=3$).²¹

III. EXPLICITLY CORRELATED CALCULATIONS

We write the correlated wave function of D -ballium as

$$\Psi = \sum_{n=0}^{\omega} \sum_{l=0}^n \sum_{m=0}^{\omega} c_{nlm} (1 + \hat{P}_{12}) \psi_{nlm}, \quad (20)$$

where $\hat{P}_{12}$ is the permutation operator between electron 1 and 2, which ensures the correct symmetry for the 1S ground state, and the basis functions are

$$\psi_{nlm} = (1 - x^2)(1 - y^2)x^{2n}y^{2l}z^m, \quad (21)$$

the scaled coordinates are

$$x = \frac{r_1}{R}, \quad y = \frac{r_2}{R}, \quad z = \frac{r_{12}}{R}, \quad (22)$$

and n , l , and m are non-negative integers. Such functions ensure that Ψ is smooth at $r_1=0$ and $r_2=0$, i.e.,

$$\left. \frac{\partial \Psi}{\partial r_1} \right|_{r_1=0} = \left. \frac{\partial \Psi}{\partial r_2} \right|_{r_2=0} = 0, \quad (23)$$

and that Ψ is cusped at the boundary and satisfies Eq. (4). The total number of basis functions in Eq. (21) is

TABLE I. HF and exact energies of 3-ballium for $R=1, 5$, and 20 .

	Basis set size	$R=1$	$R=5$	$R=20$
HF energy				
Thompson and Alavi ^a	7	11.641 747 645	0.739 761 807	0.105 378 511
Present work using Eq. (9)	7	11.641 747 631 859	0.739 761 794 626	0.105 378 488 0
	10	11.641 747 631 855 851 828	0.739 761 794 625 138	0.105 378 488 024
Exact energy				
Thompson and Alavi ^b	210	11.591 380 285	0.701 706 934	0.086 577 117
	Extrap.	11.590 81(4)	0.701 606 (2)	0.086 577 0(0)
Jung and Alvarellos ^c	6296	11.590 906	...	...
Present work using Eq. (20)	196	11.590 838 69	0.701 613 820	0.086 576 568 4
	726	11.590 838 689 02	0.701 613 820 002	0.086 576 568 358 529

^aReference 21.^bReferences 16 and 44.^cReference 17.

$$N = \frac{(\omega + 1)^2(\omega + 2)}{2}. \quad (24)$$

The ground-state energy is the lowest eigenvalue of

$$\mathbf{S}^{-1/2}(\mathbf{T} + \mathbf{U})\mathbf{S}^{-1/2}, \quad (25)$$

where $\mathbf{S}$, $\mathbf{T}$, and $\mathbf{U}$ are the overlap, kinetic, and repulsion matrices, respectively.⁴⁵

Although our (x, y, z) coordinates are equivalent to the (s, t, u) coordinates of Hylleraas,^{5,45} ours lead to simpler closed-form expressions for the required integrals. All the required matrix elements can be found in closed form using the general formula in the Appendix.

Figure 2 shows how the accuracy of the exact energy of 3-ballium improves as the number N of terms in the expansion increases and Table I reports numerical values of the exact energy for various $R=1, 5$, and 20 . Our explicitly correlated results are compared with the Configuration Interaction energies of Thompson and Alavi^{16,44} and of Jung and Alvarellos.¹⁷ Convergence for other values of D is similar.

Explicitly correlated calculations converge much faster than CI calculations because the former include terms (21) with $m=1$ satisfying the Kato cusp condition.^{46–48} For example, for the unit ball, Thompson and Alavi^{16,44} obtained

$E=11.591\,380\,285$ using 210 basis functions, Jung and Alvarellos¹⁷ subsequently found $E=11.590\,906$ using 6296 functions, but we obtain $E=11.590\,838\,689$ using only 196 functions ($\omega=6$). Our energy is consistent with the extrapolated⁴⁹ estimate $E=11.590\,81(4)$ of Thompson and Alavi.^{16,44} Likewise, using 6296 basis functions, Jung and Alvarellos¹⁷ found $E=22.033\,71$ for the first excited S state and, using 196 explicitly correlated functions, we obtain $E=22.033\,562\,4E_h$.

Figure 2 reveals that, for $R=1$ and $R=5$, the rate of convergence of the Hylleraas basis set is very similar. When R exceeds the Wigner–Seitz critical value ($r_s \approx 6$ for $D=3$), a Wigner molecule is formed, characterized by a minimum of the electron density at the center of the box.^{16,44} Although the CI calculations of Thompson and Alavi show only small variations of the rate of convergence, the Wigner molecule formation dramatically modifies the energy convergence of our explicitly correlated calculations. For $R=20$ and short expansions ($N < 50$), the CI and Hylleraas calculations lead to similar results but, for larger basis sets, the Hylleraas scheme is superior and the accuracy rapidly improves.

Correlation energies E_c for $D=2, 3, \dots, 7$ and $R=1, 5$, and 20 are given in Table II. For fixed D and increasing R , E_c decreases. For fixed R and increasing D , although both the exact and HF energies increase, E_c decreases.^{50–52}

FIG. 2. Accuracy $\mathcal{A}$ of the exact energy of 3-ballium with respect to the basis set size N for various R . TA results taken from Ref. 16, JA from Ref. 17 and LG from the present study.

IV. LIMITING CORRELATION ENERGY

Following Hylleraas perturbation theory,⁵³ we expand both the exact and HF energies as series in R , yielding

$$E = \frac{E^{(0)}}{R^2} + \frac{E^{(1)}}{R} + E^{(2)} + O(R), \quad (26)$$

$$E_{\text{HF}} = \frac{E^{(0)}}{R^2} + \frac{E^{(1)}}{R} + E_{\text{HF}}^{(2)} + O(R). \quad (27)$$

The limiting correlation energy is then given by

$$E_c^{(0)} = \lim_{R \rightarrow 0} E_c = E^{(2)} - E_{\text{HF}}^{(2)}. \quad (28)$$

The one-electron Hamiltonian for D -ballium is

TABLE II. Exact, HF, and correlation energies for various finite R . Zeroth-order, first-order, and second-order energies, and limiting correlation energies.

D	2	3	4	5	6	7
Exact, HF, and correlation energies of ballium for $R=1$						
E	8.104 931	11.590 839	16.151 742	21.519 813	27.612 654	34.391 191
E_{HF}	8.326 496	11.641 748	16.172 654	21.530 902	27.619 443	34.395 746
E_{c}	-0.221 565	-0.050 909	-0.020 912	-0.011 089	-0.006 789	-0.004 555
Exact, HF, and correlation energies of ballium for $R=5$						
E	0.586 796	0.701 614	0.863 437	1.063 334	1.296 170	1.559 045
E_{HF}	0.711 077	0.739 762	0.880 997	1.073 192	1.302 407	1.563 317
E_{c}	-0.124 281	-0.038 148	-0.017 560	-0.009 858	-0.006 237	-0.004 272
Exact, HF, and correlation energies of ballium for $R=20$						
E	0.078 628	0.086 577	0.096 381	0.108 129	0.121 815	0.137 388
E_{HF}	0.123 044	0.105 378	0.107 060	0.114 985	0.126 552	0.140 835
E_{c}	-0.044 416	-0.018 801	-0.010 679	-0.006 856	-0.004 737	-0.003 447
Zeroth-order and first-order energies of ballium, from Eq. (32)						
$E^{(0)}$	5.783 186	9.869 604	14.681 971	20.190 729	26.374 616	33.217 462
$E^{(1)}$	2.596 157	1.786 073	1.496 754	1.343 463	1.2468 45	1.179 626
Second-order energies of ballium, from Eqs. (34) and (36)						
$E^{(2)}$	-0.324 120	-0.069 618	-0.028 107	-0.014 770	-0.008 977	-0.005 983
$E_{\text{HF}}^{(2)}$	-0.057 959	-0.014 442	-0.006 194	-0.003 333	-0.002 037	-0.001 352
Limiting correlation energies $E_{\text{c}}^{(0)}$, from Ref. 23 and Eq. (28)						
Helium	-0.220 133	-0.046 663	-0.018 933	-0.010 057	-0.006 188	-0.004 176
Spherium	-0.227 411	-0.047 637	-0.019 181	-0.010 139	-0.006 220	-0.004 189
Hookium	-0.239 641	-0.049 703	-0.019 860	-0.010 439	-0.006 376	-0.004 280
Ballium	-0.266 161	-0.055 176	-0.021 913	-0.011 437	-0.006 940	-0.004 631

$$\hat{H}_0 = -\frac{1}{2} \left[\frac{d^2}{dr^2} + \frac{D-1}{r} \frac{d}{dr} \right] + V(r), \quad (29)$$

$$E_{\text{HF}}^{(2)} = - \int_0^1 \frac{W(r)^2}{r^{D-1} \psi_0(r)^2} dr, \quad (34)$$

and the associated zeroth-order wave function is

$$\Psi_0(r_1, r_2) = \psi_0(r_1) \psi_0(r_2), \quad (30)$$

$$W(r) = 2 \int_0^r [J_{\psi_0}(x) - E^{(1)}] \psi_0(x)^2 x^{D-1} dx, \quad (35)$$

where

$$\psi_0(r) = \frac{\sqrt{2}}{J_{D/2}(\kappa)} \frac{J_{D/2-1}(\kappa r)}{r^{D/2-1}}, \quad (31)$$

In Eq. (31), $\kappa = j_{D/2-1,1}$ and $j_{\mu,k}$ is the k th zero of the Bessel function of the first kind $J_{D/2-1}$.⁴¹ The $E^{(0)}$ and $E^{(1)}$ values are easily obtained from the relations

$$E^{(0)} = \kappa^2, \quad E^{(1)} = \langle \Psi_0 | r_{12}^{-1} | \Psi_0 \rangle, \quad (32)$$

and are reported in Table II. For odd D , $E^{(1)}$ can be found in closed form. For example, for $D=3$,

$$E^{(1)} = 2 \left[1 - \frac{\text{Si}(2\pi)}{2\pi} + \frac{\text{Si}(4\pi)}{4\pi} \right], \quad (33)$$

where Si is the sine integral function.⁴¹

A. HF energy

Values of $E_{\text{HF}}^{(2)}$ have been determined using the generalization of the Byers-Brown–Hirschfelder equations⁵⁴

where $J_{\psi_0}(x)$ is given by Eq. (8).

B. Exact energy

The second-order energy $E^{(2)}$, which minimizes the Hylleraas functional,⁵³ is given by

$$E^{(2)} = -\mathbf{b}^T \mathbf{A}^{-1} \mathbf{b}, \quad (36)$$

where

$$\mathbf{A} = \mathbf{T} - E^{(0)} \mathbf{S}, \quad (37)$$

$$\mathbf{b} = \mathbf{C}^T [E^{(1)} \mathbf{S} - \mathbf{U}]. \quad (38)$$

The matrices $\mathbf{S}$, $\mathbf{T}$, and $\mathbf{U}$ have been defined in Sec. III. The vector $\mathbf{C}$ contains the coefficients of the zeroth-order wave function (30) expanded in the basis set (21). The basis set has been enlarged by progressively increasing the maximum value of ω .

C. Correlation energy

The exact and HF second-order energies, as well as the limiting correlation energy $E_c^{(0)}$, of D -ballium are reported in Table II. The latter is compared with previously reported results²³ for related two-electron systems (helium, spherium, and hookium).

The first observation is the tendency of the limiting correlation energies to decrease with increasing dimensionality.⁵⁵ As D increases, all of the energies decrease dramatically and the correlation energies fall by almost two orders of magnitude between $D=2$ and $D=7$. This point has been already discussed and explained in previous works.^{55,23}

We used the method developed by Herschbach and collaborators^{40,56–59} to find that the large- D expansion of $E_c^{(0)}$ in D -ballium is

$$E_c^{(0)} \sim -\frac{1}{8}\delta^2 - \frac{53}{128}\delta^3 + \dots, \quad (39)$$

where $\delta=1/(D-1)$.²⁵ This supports our recent conjecture (1) that the leading term $-\delta^2/8$ is universal and independent of the radial external potential $V(r)$. We note that the coefficient of δ^3 is larger than those in the other two-electron systems and this explains why the limiting correlation energy of D -ballium is always larger than those in helium, spherium, and hookium.

V. CONCLUSION

In this article, we reported accurate results for the exact and HF ground-state energies of two electrons of opposite spin confined within a ball of radius R in a D -dimensional space. We call this system D -ballium.

Our results, focusing mainly on the high-density regime (small- R) and the limit where $R=0$, extend and complete earlier studies on 3-ballium.^{16,17,21} The exact results have been obtained using a Hylleraas-type expansion, while the HF limit has been reached using a simple polynomial basis set.

We have also shown that in the large-dimension limit, the limiting correlation energy behaves as $-\delta^2/8 - C\delta^3$, in agreement with our recent conjecture,²³ and is larger than the limiting correlation energy in other two-electron systems. A rigorous proof of the conjecture will be reported elsewhere,⁶⁰ where we will show that this result is related to the Kato cusp factor. Using continuity arguments, it seems clear that the conjecture does not apply to nonspherical external potentials.

ACKNOWLEDGMENTS

P.M.W.G. thanks the NCI National Facility for a generous grant of supercomputer time and the Australian Research Council (Grant No. DP0984806) for funding.

APPENDIX: HYLLERAAS-TYPE INTEGRALS

The integrals needed to compute the different matrix elements are of the form

$$\mathcal{I}_{nlm} = \int x^n y^l z^m d\tau, \quad (A1)$$

with the volume element

$$d\tau = xyz \mathcal{J}^{D-3/2} dx dy dz, \quad (A2)$$

$$\mathcal{J} = (x+y+z)(x-y+z)(x+y-z)(x-y-z), \quad (A3)$$

and domain of integration

$$\int d\tau = \int_0^1 dx \int_0^1 dy \int_{|x-y|}^{x+y} dz. \quad (A4)$$

One eventually finds

$$\mathcal{I}_{nlm} = \sqrt{\pi} \frac{\Gamma\left(\frac{D-1}{2}\right)}{\Gamma\left(\frac{D}{2}\right)} \frac{R^{n+l+m+2D}}{n+l+m+2D} (I_n^m + I_l^m), \quad (A5)$$

and

$$I_a^b = \frac{{}_3F_2\left(\frac{a+D}{2}, -\frac{b}{2}, -\frac{b+D-2}{2}; \frac{a+D+2}{2}, \frac{D}{2}; 1\right)}{a+D}. \quad (A6)$$

¹ A. Michels, J. de Boer, and A. Bijl, *Physica (Amsterdam)* **4**, 981 (1937).

² A. Sommerfeld and H. Welker, *Ann. Phys.* **424**, 56 (1938).

³ S. R. De Groot and C. A. Seldam, *Physica (Amsterdam)* **12**, 669 (1946).

⁴ M. Gell-Mann and K. A. Brueckner, *Phys. Rev.* **106**, 364 (1957).

⁵ E. A. Hylleraas, *Z. Phys.* **54**, 347 (1929).

⁶ C. A. Ten Seldam and S. R. De Groot, *Physica (Amsterdam)* **18**, 891 (1952).

⁷ B. M. Gimarc, *J. Chem. Phys.* **47**, 5110 (1967).

⁸ N. Aquino, A. Flores-Riveros, and J. F. Rivas-Silva, *Phys. Lett. A* **307**, 326 (2003).

⁹ N. Aquino, J. Garza, A. Flores-Riveros, J. F. Rivas-Silva, and K. D. Sen, *J. Chem. Phys.* **124**, 054311 (2006).

¹⁰ N. Aquino, *Adv. Quantum Chem.* **57**, 123 (2009).

¹¹ A. Flores-Riveros, N. Aquino, and H. E. Montgomery, Jr., *Phys. Lett. A* **374**, 1246 (2010).

¹² A. Flores-Riveros, N. Aquino, and H. E. Montgomery, Jr., *Phys. Lett. A* (in press).

¹³ A. Alavi, *J. Chem. Phys.* **113**, 7735 (2000).

¹⁴ S. Ghosh and P. M. W. Gill, *J. Chem. Phys.* **122**, 154108 (2005).

¹⁵ I. G. Ryabinkin and V. N. Staroverov, *Phys. Rev. A* **81**, 032509 (2010).

¹⁶ D. C. Thompson and A. Alavi, *Phys. Rev. B* **66**, 235118 (2002).

¹⁷ J. Jung and J. E. Alvarez, *J. Chem. Phys.* **118**, 10825 (2003).

¹⁸ J. Jung, P. Garcia-Gonzalez, J. E. Alvarez, and R. W. Godby, *Phys. Rev. A* **69**, 052501 (2004).

¹⁹ D. C. Thompson and A. Alavi, *Phys. Rev. B* **69**, 201302 (2004).

²⁰ D. C. Thompson and A. Alavi, *J. Phys.: Condens. Matter* **16**, 7979 (2004).

²¹ D. C. Thompson and A. Alavi, *J. Chem. Phys.* **122**, 124107 (2005).

²² E. Wigner, *Phys. Rev.* **46**, 1002 (1934).

²³ P.-F. Loos and P. M. W. Gill, *J. Chem. Phys.* **131**, 241101 (2009).

²⁴ E. Witten, *Phys. Today* **33** (7), 38 (1980).

²⁵ L. G. Yaffe, *Phys. Today* **36** (8), 50 (1983).

²⁶ G. S. Ezra and R. S. Berry, *Phys. Rev. A* **25**, 1513 (1982).

²⁷ G. S. Ezra and R. S. Berry, *Phys. Rev. A* **28**, 1989 (1983).

²⁸ P. C. Ojha and R. S. Berry, *Phys. Rev. A* **36**, 1575 (1987).

²⁹ R. J. Hinde and R. S. Berry, *Phys. Rev. A* **42**, 2259 (1990).

³⁰ J. W. Warner and R. S. Berry, *Nature (London)* **313**, 160 (1985).

³¹ M. Seidl, *Phys. Rev. A* **75**, 062506 (2007).

³² P.-F. Loos and P. M. W. Gill, *Phys. Rev. A* **79**, 062517 (2009).

³³ P.-F. Loos and P. M. W. Gill, *Phys. Rev. Lett.* **103**, 123008 (2009).

³⁴ P.-F. Loos, *Phys. Rev. A* **81**, 032510 (2010).

³⁵ N. R. Kestner and O. Sinanoglu, *Phys. Rev.* **128**, 2687 (1962).

³⁶ S. Kais, D. R. Herschbach, and R. D. Levine, *J. Chem. Phys.* **91**, 7791 (1989).

³⁷ M. Taut, *Phys. Rev. A* **48**, 3561 (1993).

³⁸ J. Cioslowski and K. Penal, *J. Chem. Phys.* **113**, 8434 (2000).

- ³⁹T. Helgaker, P. Jørgensen, and J. Olsen, *Molecular Electronic-Structure Theory* (Wiley, Chichester, 2000).
- ⁴⁰D. R. Herschbach, *J. Chem. Phys.* **84**, 838 (1986).
- ⁴¹M. Abramowitz and I. E. Stegun, *Handbook of Mathematical Functions* (Dover, New York, 1972).
- ⁴²Wolfram Research, Inc., MATHEMATICA 7, 2008.
- ⁴³S. Ragot, *J. Chem. Phys.* **128**, 164104 (2008).
- ⁴⁴D. C. Thompson and A. Alavi, *Phys. Rev. B* **68**, 039901 (2003).
- ⁴⁵E. A. Hylleraas, *Adv. Quantum Chem.* **1**, 1 (1964).
- ⁴⁶T. Kato, *Commun. Pure Appl. Math.* **10**, 151 (1957).
- ⁴⁷W. Kutzelnigg, *Theor. Chim. Acta* **68**, 445 (1985).
- ⁴⁸W. Kutzelnigg and W. Klopper, *J. Chem. Phys.* **94**, 1985 (1991).
- ⁴⁹A. Halkier, T. Helgaker, P. Jørgensen, W. Klopper, H. Koch, J. Olsen, and A. K. Wilson, *Chem. Phys. Lett.* **286**, 243 (1998).
- ⁵⁰J. G. Loeser and D. R. Herschbach, *J. Chem. Phys.* **84**, 3882 (1986).
- ⁵¹J. G. Loeser and D. R. Herschbach, *J. Chem. Phys.* **84**, 3893 (1986).
- ⁵²J. G. Loeser and D. R. Herschbach, *J. Chem. Phys.* **86**, 3512 (1987).
- ⁵³E. A. Hylleraas, *Z. Phys.* **65**, 209 (1930).
- ⁵⁴W. Byers Brown and J. O. Hirschfelder, *Proc. Natl. Acad. Sci. U.S.A.* **50**, 399 (1963).
- ⁵⁵D. R. Herrick and F. H. Stillinger, *Phys. Rev. A* **11**, 42 (1975).
- ⁵⁶J. G. Loeser and D. R. Herschbach, *J. Phys. Chem.* **89**, 3444 (1985).
- ⁵⁷D. J. Doren and D. R. Herschbach, *Chem. Phys. Lett.* **118**, 115 (1985).
- ⁵⁸J. G. Loeser and D. R. Herschbach, *J. Chem. Phys.* **86**, 2114 (1987).
- ⁵⁹D. Z. Goodson and D. R. Herschbach, *J. Chem. Phys.* **86**, 4997 (1987).
- ⁶⁰P.-F. Loos and P. M. W. Gill, "Invariance of the high-density correlation energy in two-electron systems," *Phys. Rev. Lett.* (submitted); arXiv:1005.0676v1.