Health Change in the Asia-Pacific Region

The Asia-Pacific region has seen great social, environmental and economic change across the past century, with great acceleration of change in the last 20 years or so, leading to dramatic changes in the health profiles of all populations represented in South East Asia, East Asia, Pacific Islands and the islands of Melanesia. This volume will consider recent evidence concerning prehistoric migration, and colonial, regional and global processes in the production of health change in the Asia-Pacific region. Notably, it will examine ways in which a health pattern dominated by undernutrition and infection has been displaced in many ways, and is being displaced elsewhere, by over-nutrition and the degenerative diseases associated with it. This book will present a cohesive view of the ways in which exchange relationships, economic modernization, migration and transnational linkages interact with changing rural subsistence ecologies to influence health patterns in this region.

RYUTARO OHTSUKA is President of the National Institute for Environmental Studies, Japan. Until 2005, he was Professor of Human Ecology in the University of Tokyo and conducted research among various populations in the Asia-Pacific region.

STANLEY J. ULIJASZEK is Professor of Human Ecology in the Institute of Social and Cultural Anthropology, University of Oxford. His research is focused on the biocultural ecology of human populations, past and present, and nutritional anthropology.

Cambridge Studies in Biological and Evolutionary Anthropology

Series editors

HUMAN ECOLOGY

C. G. Nicholas Mascie-Taylor, University of Cambridge

Michael A. Little, State University of New York, Binghamton

GENETICS

Kenneth M. Weiss, Pennsylvania State University

HUMAN EVOLUTION

Robert A. Foley, University of Cambridge

Nina G. Jablonski, California Academy of Science

PRIMATOLOGY

Karen B. Strier, University of Wisconsin, Madison

Also available in the series

39 *Methods in Human Growth Research* Roland C. Hauspie, Noel Cameron & Luciano Molinari (eds.) 0 521 82050 2

40 *Shaping Primate Evolution* Fred Anapol, Rebecca L. German & Nina G. Jablonski (eds.) 0 521 81107 4

41 *Macaque Societies – A Model for the Study of Social Organization* Bernard Thierry, Mewa Singh & Werner Kaumanns (eds.) 0 521 81847 8

42 *Simulating Human Origins and Evolution* Ken Wessen 0 521 84399 5

43 *Bioarchaeology of Southeast Asia* Marc Oxenham & Nancy Tayles (eds.) 0 521 82580 6

44 *Seasonality in Primates* Diane K. Brockman & Carel P. van Schaik 0 521 82069 3

45 *Human Biology of Afro-Caribbean Populations* Lorena Madrigal 0 521 81931 8

46 *Primate and Human Evolution* Susan Cachel 0 521 82942 9

47 *The First Boat People* Steve Webb 0 521 85656 6

48 *Feeding Ecology in Apes and Other Primates* Gottfried Hohmann, Martha Robbins & Christophe Boesch (eds.) 0 521 85837 2

49 *Measuring Stress in Humans: A Practical Guide for the Field* Gillian Ice & Gary James (eds.) 0 521 84479 7

50 *The Bioarchaeology of Children: Perspectives from Biological and Forensic Anthropology* Mary Lewis 0 521 83602 6

51 *Monkeys of the Tai Forest* W. Scott McGraw, Klaus Zuberbühler & Ronald Noë (eds.) 0 521 81633 5

52 *Health Change in the Asia-Pacific Region: Biocultural and Epidemiological Approaches* Ryutaro Ohtsuka & Stanley J. Ulijaszek (eds.) 978 0 521 83792 7

Health Change in the Asia-Pacific Region

Biocultural and Epidemiological Approaches

EDITED BY

RYUTARO OHTSUKA

National Institute for Environmental Studies, Tsukuba City, Japan

STANLEY J. ULIJASZEK

Institute of Social and Cultural Anthropology, University of Oxford, UK

CAMBRIDGE
UNIVERSITY PRESS

CAMBRIDGE UNIVERSITY PRESS
Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore, São Paulo

Cambridge University Press
The Edinburgh Building, Cambridge CB2 2RU, UK

Published in the United States of America by Cambridge University Press,
New York

www.cambridge.org
Information on this title: www.cambridge.org/9780521837927

© Cambridge University Press 2007

This publication is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

First published 2007

Printed in the United Kingdom at the University Press, Cambridge

A catalogue record for this publication is available from the British Library

Library of Congress Cataloguing in Publication data

Health change in the Asia-Pacific region: biocultural and epidemiological
approaches / edited by Ryutaro Ohtsuka and Stanley J. Ulijaszek.

p. ; cm. — (Cambridge studies in biological and evolutionary anthropology ; 52)

Includes bibliographical references.

ISBN-13: 978 0 521 83792 7 (hardback)

1. Health transition – Southeast Asia. 2. Health transition – East Asia. 3. Health
transition – Islands of the Pacific. 4. Nutrition – Southeast Asia. 5. Nutrition – East
Asia. 6. Nutrition – Islands of the Pacific. I. Ohtsuka, Ryutaro. II. Ulijaszek,
Stanley J. III. Series.

[DNLM: 1. Health Transition – Asia, Southeastern. 2. Health Transition – Far
East. 3. Health Transition – Pacific Islands. 4. Anthropology, Cultural – Asia,
Southeastern. 5. Anthropology, Cultural – Far East. 6. Anthropology, Cultural –
Pacific Islands. 7. Nutritional Status – Asia, Southeastern. 8. Nutritional Status –
Far East. 9. Nutritional Status – Pacific Islands. 10. Socioeconomic Factors – Asia,
Southeastern. 11. Socioeconomic Factors – Far East. 12. Socioeconomic Factors –
Pacific Islands. WA 105 H4342 2007] I. Title. II. Series.

RA650.7.S68H43 2007
362.10959–dc22 2006036793

ISBN-13 978 0 521 83792 7 hardback

Cambridge University Press has no responsibility for
the persistence or accuracy of URLs for external or
third-party internet websites referred to in this publication,
and does not guarantee that any content on such
websites is, or will remain, accurate or appropriate.

Contents

List of contributors

vii

Acknowledgements

ix

- 1 Health change in the Asia-Pacific region: disparate
end-points?
STANLEY J. ULIJASZEK AND RYUTARO OHTSUKA 1
- 2 Interactions of nutrition, genetics and infectious disease in the
Pacific: implications for prehistoric migrations
STEPHEN OPPENHEIMER 21
- 3 Biocultural adaptation and population connectedness
in the Asia-Pacific region
RYUTARO OHTSUKA 44
- 4 Changing nutritional health in South East Asia
GEOFFREY C. MARKS 64
- 5 Obesity and nutritional health in Hong Kong
Chinese people
GARY T. C. KO 88
- 6 Modernization, nutritional adaptability and health in Papua
New Guinea Highlanders and Solomon Islanders
TARO YAMAUCHI 101
- 7 Tongan obesity: causes and consequences
TSUKASA INAOKA, YASUHIRO MATSUMURA
AND KAZUHIRO SUDA 127
- 8 Nutrition and health in modernizing Samoans: temporal
trends and adaptive perspectives
EMBER D. KEIGHLEY, STEPHEN T. MCGARVEY, CHRISTINE
QUESTED, CHARLES MCCUDDIN, SATUPAITEA VIALI AND
UTO'OFILI A. MAGA 147
- 9 Health patterns of Pacific Islanders and Asians in the
United States
W. PARKER FRISBIE, ROBERT A. HUMMER, T. ELIZABETH
DURDEN AND YOUNGTAE CHO 192

10	Impacts of modernization and transnationalism on nutritional health of Cook Islanders	219
	STANLEY J. ULIJASZEK	
11	Mortality decline in the Pacific: economic development and other explanations	234
	ALISTAIR WOODWARD AND TONY BLAKELY	
12	Health changes in Papua New Guinea: from adaptation to double jeopardy?	254
	ROBERT ATTENBOROUGH	
	<i>Index</i>	303

Contributors

Robert Attenborough

The School of Archaeology and Anthropology, A. D. Hope Building, The Australian National University, Canberra, ACT 0200, Australia

Tony Blakely

Department of Public Health, Wellington School of Medicine and Health Sciences, University of Otago, 23A Mein Street, Newtown, Wellington, New Zealand

Youngtae Cho

School of Public Health Seoul, National University Seoul, Korea

T. Elizabeth Durden

Department of Sociology and Anthropology, Bucknell University, Lewisburg, PA 17837

W. Parker Frisbie

Population Research Center and Department of Sociology, College of Liberal Arts, University of Texas at Austin, 1 University Station A1700, Austin, TX 78712-0543, USA

Robert A. Hummer

The University of Texas at Austin, Population Research Center, 1 University Station G1800, Austin, TX 78712-0543, USA

Tsukasa Inaoka

Department of Environmental Sociology, Saga University, 1 Honjo-machi, Saga City, Japan

Ember D. Keighley

International Health Institute, Brown University, Box G-B495, Providence, RI 02912, USA

Gary T. C. Ko

Hong Kong Alice Ho Miu Ling Nethersole Hospital, 11 Chuen On Road, Tai Po, New Territory, Hong Kong

Uto'ofili A. Maga

Department of Health, Pago Pago, American Samoa 96799

Geoffrey C. Marks

Australian Centre for International and Tropical Health and Nutrition, University of Queensland, Level 4, Public Health Building, Herston Road, Herston, Brisbane, Queensland 4006, Australia

Yasuhiro Matsumura

National Institute of Health and Nutrition, Toyama 1-23-1 Shinjuku-ku, Tokyo 162-8636, Japan

Charles McCuddin

Tafuna Family Health Center, Department of Health, Pago Pago, American Samoa 9679

Stephen T. McGarvey

International Health Institute, Brown University, Box G-B497, Providence RI 02912, USA

Ryutaro Ohtsuka

National Institute for Environmental Studies, 16-2 Onogawa, Tsukuba City, Ibaraki 305-8506, Japan

Stephen Oppenheimer

Green College, Woodstock Road, Oxford OX2 6HG, UK

Christine Quested

Nutrition Centre, Ministry of Health, Government of Samoa, Apia, Samoa

Kazuhiro Suda

Hokkai-Gakuen University, Asahi-machi 4-1-40 Toyohira-ku, Sapporo 062-8605, Japan

Stanley J. Ulijaszek

Institute of Social and Cultural Anthropology, University of Oxford, 51 Banbury Road Oxford OX2 6PF, UK

Satupaitea Viali

Oceania University of Medicine Samoa, Apia Campus and Tupua Tamasese Meaole Hospital, Ministry of Health, Government of Samoa, Apia, Samoa

Alistair Woodward

School of Population Health, University of Auckland, Private Bag 92019, Auckland, New Zealand

Taro Yamauchi

Department of Human Ecology, School of International Health, Graduate School of Medicine, University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

Acknowledgements

The meeting which formed a basis for this volume took place in Tokyo, as part of the International Union of Anthropological and Ethnographic Sciences Congress, held in 2002. We thank the Japan Society for the Promotion of Science for helping to support this symposium financially. We also thank Robin Hide, Tony McMichael, Vicki Luker, Don Gardner and Donald Denoon for their comments and suggestions at various stages of this project.

12 *Health changes in Papua New Guinea: from adaptation to double jeopardy?*

ROBERT ATTENBOROUGH

Introduction

Are the health patterns of Papua New Guinea (PNG) changing? If so, in what ways are they doing so? Specifically, are they changing in line with theoretical models which invoke modernization, development and globalization to account for the decline of infections and the rise of lifestyle- and ageing-related health problems? Or does communicable disease – persistent, recrudescing or emergent – still predominate? In this review, I aim to discuss selected evidence on the health of Papua New Guineans, with a particular focus on examining past trends and on considering, so far as possible, future prospects – whether by projection from the past or on some other rationale. In the process, I shall consider the adequacy of conventional health modernization models to describe PNG's present situation.

To be modern is to have characteristics typical of recent times and the present day, by contrast with the more distant past. This sounds open-ended; but, in the health sciences as in the arts and social sciences, modernity is (explicitly or implicitly) expected to bring changes of a specific character. This follows usually from the concept that socioeconomic modernization, discussed by Ulijaszek (1995), is accompanied, anywhere in the world and whenever it may take place, by a demographic, health or epidemiological transition (Caldwell *et al.* 1990; Riley 2001). In such transitions, not only are death rates alleviated and life expectancies increased, but also one set of causes of ill-health (life-threatening and otherwise) declines in importance, to be replaced by a different set. In broad terms, infectious causes, often acting early in life and sometimes ecologically sensitive, decline, while non-infectious causes, usually late-acting and often related to diet or other aspects of lifestyle, become the leading health problems.

Thirty years ago, a sceptic could tenably have argued that the demographic transition was a unique phenomenon embedded in the culture and

historical experience of industrialization in Europe, and in settler nations predominantly populated from Europe (Teitelbaum 1975). The corollary of this was that attempts to extrapolate key features of transition theory to other parts of the world undergoing a later and faster economic development were inadequately grounded and perhaps wrong. Nowadays, such scepticism has largely evaporated. Debate continues as to both the theoretical standing of the central propositions and the details of the transition process in particular populations. But in broad outline it is now widely accepted that health and mortality transitions as just characterized (and most often transitions from high to low fertility as well) do regularly, albeit with variations, accompany socioeconomic modernization almost anywhere. Among the variations, however, improvements in morbidity and mortality are currently slowing down in most Pacific countries, even though developed-country levels have not yet been reached (McMurray 2004). Other chapters in this book provide more detailed evidence from the Asia-Pacific region, much of which essentially bears out the established expectations of declining infection and rising lifestyle-related health problems with modernization. How well do these expectations hold for PNG?

Papua New Guinea

The eastern portion of the large island of New Guinea, the Bismarck archipelago and many other smaller islands nearby together make up the modern nation of PNG. Now celebrating over 30 years of independence, PNG emerged in 1975 from a century or less of colonialism – effectively much less in the remotest areas – under Germany and Britain initially, and later Australia (Alpers and Attenborough 1992; Ulijaszek 1995). It retains strong economic and other links with Australia as well as other Asia-Pacific nations.

With a land area of about 463,000 km² (Hanson *et al.* 2001) and a population of about 5.7 million, PNG is not very populous by Asian standards, but is easily the largest of the Pacific Community nations and territories, with 65% of their total population and one of their shorter population doubling times, currently 32 years (Secretariat of the Pacific Community 2004). It also has a larger population than New Zealand or Hawaii. In its famous diversity and complexity – linguistic and cultural, topographic and climatic – PNG bears comparison with any nation in the Asia-Pacific region. The circumstances as well as the timing of socioeconomic modernization in PNG are about as different as can be imagined from mid-eighteenth-century France, Sweden or Britain, where the earliest signs of a health transition can be traced (Riley 2001).

It would be a mistake to consider PNG's undocumented pre-colonial past as timeless or changeless. Two particularly salient transitions for which we have evidence are the development of taro and banana agriculture in the highlands, and the later advent of the South American sweet potato as an alternative basis for root crop agriculture (Denham *et al.* 2003; Ballard *et al.* 2005; Pawley *et al.* 2005). The possible implications of these for diet and lifestyle are discussed below. Nonetheless, the starting point for PNG's modern transition might be taken either at the earliest effective exertions of European colonial control (from the 1870s in some coastal regions to the 1950s in parts of the interior) or at the Second World War, during and following which major social and economic changes, with lasting impacts, took place in most parts of it. Since that war, and with gathering pace and increasing complexity since independence, modernization has brought to PNG astonishing changes within the lifetimes of individuals. Some people still alive now grew up outside the ambit of colonial control. Their parents may have witnessed wartime aerial dog-fights without any prior awareness of foreign armies on New Guinea soil: many others, of course, were far more involved. Post-war development has proceeded in different ways and at different speeds in different regions, accentuating PNG's pre-existing socioeconomic diversity and resulting in what many have regarded as increasingly unbalanced development (Allen 1992; Hanson *et al.* 2001).

The starkest manifestation of that uneven development, arguably, is urbanization. Table 12.1 provides some urban settlement figures from the year 2000 census (National Statistical Office 2002). Port Moresby, the capital of PNG and of Central Province, has over a quarter of a million inhabitants. Other larger urban settlements (generally provincial capitals) range from nearly 80,000 people in Lae, through Arawa, Madang and Mount Hagen, down to approximately 20,000 each in Wewak, Popondetta, Goroka and Kokopo/Vunamami (which, following the September 1994 volcanic eruptions, has taken over from Rabaul as provincial capital). Nine of the other eleven provincial capitals, plus five non-capital centres, also have populations now over 5,000. Many of these centres have shown substantial recent population growth: Brinkhoff (2003) conveniently tabulates comparative figures for 1980, 1990 and the year 2000. The total urban population of PNG now stands at 686,000 (National Statistical Office 2003a), of whom all but 38,000 live in one of the 26 settlements listed in Table 12.1. This is a substantial urban population, though well short of the one million anticipated for the year 2000 by Allen (1992); and it is growing absolutely and as a proportion of the country's total population (from 10% in 1980 to 13% in the year 2000).

Table 12.1. *Populations of urban settlements, Papua New Guinea, Census of 9–15 July 2000*

Settlement	Province	Population
Port Moresby ^a	National Capital District	254,158
Lae ^a	Morobe	78,692
Arawa ^a	North Solomons (Bougainville Autonomous Region)	31,462
Madang ^a	Madang	28,547
Mount Hagen ^a	Western Highlands	27,877
Wewak ^a	East Sepik	20,257
Kokopo/Vunamami	East New Britain	19,933
Popondetta ^a	Northern (Oro)	19,866
Goroka ^a	Eastern Highlands	19,523
Mendi ^a	Southern Highlands	17,128
Kimbe ^a	West New Britain	14,184
Wau/Bulolo	Morobe	13,037
Daru ^a	Western	12,935
Kavieng ^a	New Ireland	11,274
Alotau ^a	Milne Bay	9,888
Vanimo ^a	West Sepik (Sandaun)	9,778
Tari	Southern Highlands	8,824
Kiunga	Western	8,295
Kundiawa ^a	Chimbu (Simbu)	8,087
Kainantu	Eastern Highlands	6,788
Lorengau ^a	Manus	5,874
Ialibu	Southern Highlands	5,479
Kerema ^a	Gulf	5,124
Wabag ^a	Enga	4,208
Rabaul ^a	East New Britain	3,907
Balimo	Western	3,143

Note:

^a Denotes provincial capitals.

Source: 2000 census, National Statistical Office (2002).

Urban dwellers range from the more educated élite to the squatter settlements where often people have migrated from the villages to start a new life with their city *wantoks* (speakers of the same indigenous language). The larger of PNG's cities and towns have hospitals, university campuses and many of the same institutions and facilities as are found in urban centres worldwide.

The other 87% of the total population of PNG still live in rural areas, where, despite networks of government stations, aid posts, health centres, agricultural extension centres and schools, facilities are much more restricted. Agriculture, especially of root crops, and husbandry, especially of pigs, supplemented in some places by hunting and collecting, still supply

rural people with their staple foods. Many use earnings from cash-cropping or plantation labour among other occupations to supplement what they can grow or acquire directly, or rely principally on these sources of income. Modern life in PNG contains many elements and contrasts which foreigners are apt to find arresting, even paradoxical, in their compression or bypassing of the earlier stages of technological modernity as experienced elsewhere. There is, for example, familiarity with air transport even in areas where there is no overland transport except on foot and in canoes; and for some, there is access to video and electronic technology even in areas where there are no cinemas or televisions and perhaps not even reticulated electricity.

Are modernization and globalization, at the scale and pace experienced in PNG, at risk of overwhelming people's capacity to cope, whether in urban or rural areas? Do the health deficits they bring outweigh the benefits? Do other extrinsic shocks, coming perhaps from the worlds of infectious microorganisms or of climate change, bring similar risks? Or do the assumptions behind such questions underestimate people's capacity to adapt, both biologically and socially, as the recent rise of interest in resilience (e.g., Barnett 2002) might lead us to think?

Perhaps this last question cannot be directly answered from conventional population health data. But a sampling of recent comparative statistics certainly offers little cause for complacency over PNG's present situation. At 93 per 1,000, PNG's rate of child mortality (under five years) is currently the 49th highest in the world, the 11th highest among non-African nations (United Nations Children's Fund 2004a: 105). Within the East Asia-Pacific region, only Cambodia and the Democratic People's Republic of Korea have seen slower progress than PNG towards the Millennium Development Goal of reducing under-five mortality by two-thirds between 1990 and 2015 (United Nations Children's Fund 2004b: 20). Among Pacific Community nations and territories, PNG has, if the data can be accepted, the highest infant mortality rate (IMR; death rate among those below one year of age: 64 per 1,000 live births) except for the Solomon Islands; a very high maternal mortality rate of 370 per 100,000 live births, only 45% of which are attended by trained personnel; the highest rate of HIV/AIDS diagnosis; and the lowest life expectancy (54 years for males, 55 for females) with the striking exception of Nauruan males (McMurray 2004; Secretariat of the Pacific Community 2004). Of the World Health Organization's 191 member states in 1997, PNG ranked 150th in overall health system attainment, a composite measure covering average health, health equality, system responsiveness, responsiveness equality and financial fairness; and 148th in overall health system performance, as measured

Table 12.2. Overall health system attainment* and performance[†] rankings of 22 countries, selected from 191 World Health Organization member states, 1997

Attainment		Performance	
Rank	Country	Rank	Country
1	Japan	1	France
6	France	10	Japan
9	UK	18	UK
12	Australia	32	Australia
15	USA	37	USA
26	New Zealand	41	New Zealand
54	Philippines	60	Philippines
78	Fiji	76	Sri Lanka
80	Sri Lanka	80	Solomon Islands
82	Samoa	92	Indonesia
85	Tonga	96	Fiji
88	Cook Islands	107	Cook Islands
100	Russian Federation	112	India
106	Indonesia	116	Tonga
108	Solomon Islands	119	Samoa
121	India	125	Brazil
125	Brazil	127	Vanuatu
132	China	130	Russian Federation
134	Vanuatu	144	China
150	Papua New Guinea	148	Papua New Guinea
175	Myanmar	190	Myanmar
191	Sierra Leone	191	Sierra Leone

Source: World Health Organization 2000, Annex Tables 9* and 10[†], which also present uncertainties around these rankings. See explanation and discussion in text.

by the efficiency with which health expenditure is translated into health outcomes (World Health Organization 2000). Table 12.2 presents selected international, including Asia-Pacific, comparisons on these rankings. Of course some of the comparison countries have *a priori* advantages; but whatever the reasons, PNG is not fortunately placed. While much remains uncertain, therefore, it seems clear that concerns as to PNG's health prospects raised by previous authors (e.g., Inaoka 1995 and n.d.; Ulijaszek 1995; Connell 1997; Naraqi *et al.* 2003) should remain a focus of attention.

In this chapter, then, my aim is to examine some (necessarily selected) evidence on health trends in PNG. I shall ask how far the changes evidenced, and those that one can project into the future, correspond to the picture, established from other countries' experience, of the health trends

to be expected from modernization; and, on the other hand, how far other processes, such as those that underlie the persistence, emergence or re-emergence of infections, might blow the evolution of PNG's health trends off their expected course.

I shall begin at the beginning, with the earliest known human presence in New Guinea and the concept of human adaptation to the region's environment. In view of the paucity of available evidence, I shall be brief on the long unwritten pre-colonial history of human health in PNG. Even the literature from the earlier part of the colonial period, though informative in some ways (Attenborough and Alpers 1995), has severe limitations as a source of population health information. Frequently, as elsewhere in colonialism's history, it was marked by the power asymmetries and the now discarded beliefs about human difference characteristic of that period; by closer attention to the health of the colonizers than of the colonized; by conceptions of infectious disease processes and of epidemiological quantification then still nascent; and above all by biomedical problems of a variety and complexity disproportionate (even more than nowadays) to the research capacity available on the ground. In most contexts, therefore, my timescale will be little more than six decades into the past, often less, according to the availability since the Second World War of literature in which such problems are better controlled; and perhaps a decade or two into the future. At times I shall draw on national statistics, but for detail and time depth I shall sometimes opt instead for local studies, often better controlled than national ones can be, and sometimes (too seldom) repeated locally so as to allow trends to emerge, not obscured by extreme regional diversity.

It is not by chance that this chapter's title refers to health changes in the plural. There are of course many health changes at work – changes in different parts of the country, at different periods and on different time-scales, and affecting different aspects of health. Some changes may be aptly described as quantitative trends, in that they are gradual shifts over a considerable time in the burden of mortality and morbidity in a consistent direction, often but not always the direction predicted by health transition theory and experience. Other changes have different characters. In addition to infections which may have reached New Guinea with the earliest people, PNG's medical history includes infections which arrived later than that but before European colonialism: Wigley (1977, 1991) has argued that this is true of tuberculosis, although also that it became a major health problem only under colonial conditions. There have also been introductions during the colonial period or after it – from the late nineteenth century in the case of some sexually transmitted infections for example (Attenborough and Alpers 1995; Hughes 1997), up to at least the

late 1980s when PNG's first AIDS case was reported. There has been disappearance under colonialism followed by re-emergence, as in the case of yaws – apparently widely endemic at outsider contact and responsive to intervention (Gajdusek 1990), but later recrudescence (Garner *et al.* 1986; Manning and Ogle 2002). There have been introduced epidemic infections which never became recurrent or endemic, for example smallpox (Fenner *et al.* 1988; Allen 1989); and others which did, for example measles (Adels and Gajdusek 1963; Benjamin and Dramoi 2002). There have been major but finite epidemics under specific historical circumstances, such as the bacillary dysentery epidemic during the Second World War (Allen 1983; Burton 1983). Pigbel, until recently a major endemic cause of child death in the PNG highlands but responsive to vaccination campaigns, has been enormously reduced, although it has not disappeared entirely (Lawrence 1992; Poka and Duke 2003). And of course PNG's health changes are not confined to infectious disease. A range of non-infectious conditions, including but not solely the classical diseases of modernization, were essentially new in PNG in the later twentieth century: for example, asthma (Anderson and Woolcock 1992). Accidents, violence (including the Second World War) and sudden cataclysmic events have also taken a major toll, episodically in some cases. As an example of the last, the Sissano (Sandaun Province) tsunami of 17 July 1998, though less catastrophic than the Indian Ocean one of 26 December 2004, killed over 2,100 people and led to the hospitalization of at least 800 people, the permanent displacement of over 10,000 people and the abandonment of ten villages, all in one quite restricted area (Dengler and Preuss 2003).

In the space available here, I naturally cannot review PNG's health changes comprehensively, nor even cover adequately all the examples listed above. I shall review trends in overall mortality, as one indicator of health change; examine the rise of some diseases of modernization with attention to diabetes in particular; and finally consider, selectively, the question of infectious disease, old and new, persistent and emergent, with special focus on malaria and to a lesser extent HIV/AIDS. But first, I turn to early human adaptations in the New Guinea region.

Pre-colonial adaptation and health

By 40,000 years ago, people had reached a New Guinea then connected by land bridge to Australia and Tasmania (Groube *et al.* 1986), and not much later, via a further sea voyage, the Bismarck Archipelago (Specht 2005). By 30,000 years ago, the valleys of the central highlands were also reached

(Hope and Haberle 2005). For millennia, therefore, people have lived – for most of that time as hunter-gatherers – in the lands which now constitute PNG (Lilley 1992). With some riders, one must assume, then, that there has been sufficient time, on any plausible model, for them to adapt to specific critical features of Melanesia's humid tropical environment.

Riders to this statement must cover complications affecting both people's exposure to environmental pressures and those pressures themselves. On the one hand, some present-day Papua New Guineans will have some ancestors, including speakers of Austronesian languages, who arrived there more recently than the earliest peopling (Foley 1992; Kirk 1992; Spriggs 1997). Also, changes in ways of life (including agriculture), diet and rising population density will have exposed people to pressures additional to those intrinsic to the natural environment. On the other hand, the environment itself is, as already discussed, far from homogeneous; and it has changed significantly since earliest times (e.g., the infilling of the inland sea that once occupied the lower Sepik-Ramu basin: Swadling and Hope 1992, Swadling and Hide 2005).

No ecological change in the New Guinea region since the first peopling is likely to have been more important, or more devastating, for human adaptation and health than the arrival of malaria. All of the four species of the protozoon *Plasmodium* which infect humans occur in New Guinea. The global prehistory of *Plasmodium falciparum*, which as the cause of the most dangerous malaria will receive most attention here, remains under debate. While one school of thought proposes a 'Malaria's Eve' hypothesis whereby extant *P. falciparum* variants have evolved from a quite recent (3,000 to 8,000 years) common ancestor, another school argues for a much older one (100,000 to 200,000 years): see review by Hume *et al.* (2003). Recently, Joy *et al.* (2003) have presented evidence that while the origins of *P. falciparum* appear very ancient, much of its numerical expansion, including a second dispersion beyond Africa, appears to have occurred within the last 10,000 years. For PNG, their evidence suggests a rapid expansion event of the order of 13,000 to 19,000 years ago (but with a small sample and large uncertainties, so PNG data are hard to differentiate unequivocally from the African data). Thus *P. falciparum* may be a relatively recent – or if we follow Groube's (1993, 2000) speculation, very recent (within the last 1,000 to 2,000 years) – arrival in Melanesia. None of this necessarily holds true for the other three *Plasmodium* species which cause less severe malarias in humans; though in fact Clark and Kelly (1993) took seriously the possibility of malaria (species not stated) arriving with Austronesian-speakers in the fourth millennium before the present, and Leclerc *et al.* (2004) concluded that the worldwide expansion of *P. vivax* too was recent, perhaps within the last 10,000 years.

But despite indications of the relative recency of malaria, especially that caused by *P. falciparum*, it seems plausible that there has been sufficient time for substantial genetic adaptation to intense selection pressure, such as falciparum malaria at least does exert (Riley 1983). The sickle-cell gene, though not one of the suite of malaria adaptations indigenous to Melanesia (Lavu *et al.* 2002), supplies the main, albeit approximate, model for others. Wiesenfeld (1967) concluded that even 1,000–1,500 years would be enough for such adaptation; Livingstone (1989) found that present distributions were best approximated by 3,000–4,500 years of simulated malaria selection and diffusion; and Fix (2003) too was impressed by the potential rapidity of genetic change, with substantial change occurring even within 300 years. On balance then – and taking malaria as the classical exemplar of the environmental and biotic pressures at work – even if we cannot entirely exclude Groube's (1993) suggestion that accommodation to falciparum malaria is still under way, it seems reasonable to work with the proposition that by the beginning of the colonial era New Guineans had reached a high level of adaptive equilibrium. At any rate, a very effective functional adaptability has been well demonstrated (Lourie *et al.* 1992; Norgan 1997), even if specific genetic adaptations, other than to malaria, are hard to identify (Serjeantson *et al.* 1992). And if low population growth suggests some sort of adaptive equilibrium, that too is evidenced, at least in genealogical data covering recent centuries for one case, the Gidra of Western Province (Ohtsuka 1986).

A high level of adaptive equilibrium should not, of course, be equated with good population health and low mortality, as we would rate them on modern expectations. Indeed, it is one of New Guinea's conundrums that population density and cultural efflorescence such as those witnessed in the Sepik basin (Forge 1972, 1990) should have emerged and been sustained under conditions of holoendemic malaria, severe child growth deficit (Tyson 1987) and atrocious infant mortality (to cite an extreme example of the latter, Forge's estimate of 570 per 1,000 for the Abelam: Peters 1960). It is not particularly reassuring to reflect that, theoretically, some mortality régimes that strike us as appalling can be compensated by quite realistically attainable fertility régimes: for example, under definable conditions, a total fertility rate of 6.6 children is sufficient to offset a life expectancy as low as 20 years (Attenborough 2002).

The obstacle to assessing how good or bad population health was in New Guinea during the pre-colonial period (or any period not covered by systematic data collection) is self-evident. Our only direct evidence is that of palaeopathology, and for New Guinea this is limited by the paucity of studies as well as methodological constraints. In a skeletal sample of

42 individuals from Motupore Island near Port Moresby, dated at between AD 1200 and AD 1700, Webb (1995) diagnosed pathology of some kind in 22 individuals (both sexes, all but two of them adults), representing a higher prevalence of infection particularly than in his comparative Australian samples. The most common infectious disease diagnosis was treponematosi (11 individuals), with no definite signs of tuberculosis, in contrast to the nearby and roughly contemporary site of Nebira (Hope *et al.* 1983). Haematological disease was diagnosed in three cases, including one child considered to be a thalassaemia sufferer, suggestive of endemic malaria in the region by that time. Arthritis, of the cervical and lumbar vertebrae especially, was a very common diagnosis (16 individuals); trauma less so (2 individuals). Dental caries, attributed to a diet high in sago, was substantially more common than at Nebira. Greenburg-Hudd (1998), in a survey of protohistoric and historic dental palaeopathology museum specimens originating in PNG, found periodontal disease to be the most prevalent pathology: attrition was only moderate, tooth loss low to moderate and caries low, perhaps owing to the cariostatic properties of betel nut.

Beyond palaeopathology, we can only resort to indirect evidence and broad inferences. Caution is required if we are to assume that the long hunter-gatherer phase of New Guinea prehistory (and some of its history) resembles that of recent low-latitude hunter-gatherers, not in close interaction with agriculturalists, elsewhere in the world. But if we do make that assumption, it is likely (Froment 2001) that population density was low, and that the main causes of morbidity and mortality were certain parasites and infections (though others would have been absent or at low level, owing to the protections afforded by low population density and semi-isolation), plus accidental trauma, violence, snakebite and so forth, perhaps overt malnutrition on occasion, with no more than a minor role for degenerative disease except perhaps arthritis and poor dental health. Psychiatric health would be hardest of all to assess in such contexts.

Subsequent developments may all have brought impacts on human adaptation and health. These include (1) the emergence of taro agriculture and banana cultivation from at least 6,500 to 7,000 years ago (Denham *et al.* 2003); (2) the arrival of outsiders, including Austronesian-speakers in many coastal areas from the fourth millennium before the present (Lilley 1992; Spriggs 1997); (3) faunal introductions such as the pig at a much debated date not more recent than 1,900 years ago (Hide 2003); and (4) the introduction of the sweet potato from the Americas probably only a few hundred years ago, displacing taro in many places (Lilley 1992; Yen 2005). There is clearly great potential for health problems related to heavy

reliance on staple crops, where these do not provide adequate protein and micronutrients in proportion to energy, are too bulky to be satisfactory foods for younger children, and/or bring disaster if they fail; and related also to village living conditions, somewhat denser and more sedentary. Sweet potato, given its productivity and quantitative dependence of many populations upon it, has major dietary implications and was no doubt involved in the build-up of denser, more settled populations in the highlands, certainly relative to hunter-gatherer subsistence and probably also relative to taro agriculture. The dietary role of pork even nowadays is much debated and regionally variable: probably its contribution was modest but real – smaller than an outsider might expect, on account of the social and symbolic importance and only occasional slaughter of domestic pigs (Hide 2003). Both sweet potato and pork are involved in the complex of factors that can precipitate pigbel amongst PNG highlanders (Lawrence 1992), although the antiquity of this disease is unclear. Pigs are potential, and in some cases actual, sources of parasitic zoonoses in New Guinea; but on the whole, relatively few zoonoses are recorded as causing human disease in PNG even today, a fact for which it can thank in large measure its relative isolation historically and the absence until recently of large placental mammals other than pig and dog (Owen 2005).

Even after the establishment of agriculture, population densities must be presumed to have remained moderate at most, as they were historically (only exceeding 30 persons per km² in parts of the highlands, the Sepik River basin and the Gazelle peninsula of East New Britain: Ward and Lea 1970). Settlements would have been relatively small: Forge (1972), in his review, found settlements to have fewer than 300 residents each historically in most areas, and somewhere over 1,000 in a few areas including the Sepik. Mobility between settlements increases their effective overall size from the viewpoint of sustaining infection, but even the largest language group in PNG would not have been large enough in earlier times to sustain acute viral infections with properties like those of measles, for example (Attenborough and Alpers 1995). So, even within a health profile dominated by infection, pre-colonial New Guinea would have been free of certain infections which have become a problem more recently, as measles now is. Furthermore, biogeographic isolation would have protected New Guinea for a long time, perhaps up to the colonial period, from the arrival of certain other pathogenic microorganisms, including a number of them causing sexually transmitted infections (Attenborough and Alpers 1995; Hughes 1997). Finally, the presence in many lowland New Guinea populations, both north and south of the main cordillera, of genetic polymorphisms believed to confer some malaria-protective benefit on carriers

suggests the presence of malaria in New Guinea on a millennial timescale (Attenborough and Alpers 1995). These polymorphisms number at least six – five of them affecting *P. falciparum* (α -thalassaemia, β -thalassaemia, glucose-6-phosphate dehydrogenase deficiency, ovalocytosis, and Gerbich negativity) and a sixth affecting *P. vivax* (Duffy antigen variation) (Serjeantson *et al.* 1992; Genton *et al.* 1995a; Mgone *et al.* 1996; Allen *et al.* 1997; Zimmerman *et al.* 1999, 2003; Cortés *et al.* 2004). The presence of so many such polymorphisms, in some cases clearly the product of multiple mutations, is hard to interpret in other than as adaptive to the presence of malaria in the ancestral populations for an evolutionarily significant period of time.

Putting together hints such as these, then, we have a few pencil marks towards a sketch of pre-colonial health in New Guinea: a set of populations with long exposure and adaptation to most facets of their complex tropical environment, but certainly not leading idyllically healthy lives, subject to many infections including malaria and yaws, though protected from some others by isolation and small social scale, and no doubt living with frequent trauma and arthritis. It is difficult to go beyond these broad qualitative indications.

Health impacts of colonization

How much had things changed by the time of the first systematic health surveys conducted under colonial régimes, which must form the baseline for the examination of quantitative trends to which I shall shortly turn? The earliest (late-nineteenth-century) written records, such as the diaries of Mikloucho-Maclay (1975) and the first Annual Reports of the Territories of Papua and New Guinea (discussed by Attenborough and Alpers 1995), do not capture the earliest contacts; but even they tend, in the earlier years at least, to be anecdotal, at best semi-quantitative. Early contacts with outsiders from colonizing powers, whether their formal representatives or not, had a potential to change health, deliberately and otherwise, in many ways. They not only introduced novel infections: they also introduced new foods and religions; forbade warfare; directed employment and labour migration; undertook vaccination; regulated disposal of the dead; implemented sanitation measures; conducted mass treatment campaigns; made dietary recommendations; instituted constraints on sexual behaviour; inspected premises for slaughtering and food preparation; and undertook environmental measures against malaria (Attenborough and Alpers 1995).

Public health in the initially distinct colonies of Papua and New Guinea has attracted a number of detailed historical accounts (Denoon *et al.* 1989; Burton-Bradley 1990; Spencer 1999; Davies 2002). In one famous instance, colonial prohibitions on consumption of human flesh unexpectedly interrupted the transmission of kuru in the Eastern Highlands during the 1950s (Alpers 1992). The overall impact of colonial activities on indigenous health was clearly mixed and is hard to assess. Some activities, including the introduction of novel infections, clearly had profound population health impacts, sometimes running ahead of the frontier of direct colonial contact. Other activities seem likely to have been quite local and/or temporary in any effects they may have had on health, bringing little lasting alteration of the disease environment of the country overall. In the gap, then – longest in the areas contacted earliest – between the pre-colonial era and the baseline studies for quantitative population health research, health patterns would have changed significantly in some respects, certainly including novel infections; probably not greatly in many others.

Mortality trends

In reviewing more recent trends, I start with the overview of health offered by mortality and life expectancy data. At a national level, the principal data sources are the two stratified sampling censuses of 1967 and 1971 and the three full censuses of 1980, 1990 and 2000. These can be supplemented with survey data which lack national coverage but are indicative and sometimes offer greater detail. There is not space here to discuss data problems, estimation methods and other complexities in depth as Riley and Lehmann (1992) do, but it is now possible to update the picture they drew.

Table 12.3 amplifies the findings collated by Riley and Lehmann (1992: Table 3.2), with other findings, mostly later ones reported by the PNG National Statistics Office (National Statistical Office 2003b: Tables III-1–III-4 and text). Mortality clearly has been and remains high by modern international standards. Nonetheless, although aspects of these findings are open to debate, it is clear that, on any indicator, a substantial mortality decline did take place in the second half of the twentieth century. Whatever the negative health impacts of modernization and other contemporary changes, they have yet to outweigh its benefits, at least at the level of mortality (unlike certain African countries since the advent of HIV/AIDS). Infant mortality, as the most sensitive of these indicators to socio-economic and public health changes, reflects most strongly PNG's last

Table 12.3. *Estimates of mortality indicators in Papua New Guinea, 1946–2000*

	Crude mortality/ population	Infant mortality/1000 live births		Mortality 1–4 years/1000 live births		Life expectancy at birth		Life expectancy at age 25	
		Males	Females	Males	Females	Males (years)	Females (years)	Males (years)	Females (years)
1946	30–35	252	251			31	32		
1966	21	161	157			44	43		
1960s ^a						53	51		
1971	20	130	120			49	50	36	36
1971		142	125	92	90	40	41		
1980	13	78	66	47	43	49	51	31	32
1991		88	76	57	55	52	51	34	35
1996	12		73	24		55	54	39	37
2000	11	67	61	27	23	54	55	38	38

Note:

^aCombining four studies of small groups from highlands, lowland foothills and islands, published 1967–69; van de Kaa (1971).
Source: Data mostly collated from various sources by Riley and Lehmann (1992) and National Statistical Office (2003b).

half-century of such changes, and they do cover the greater part of that period. Mortality improvements are inherently hard to sustain, however, and these figures, taken at face value, appear to show that in recent times improvements have become less consistent as well as more slight.

A similar picture, of improvement slowing at some point in the 1990s, appears to emerge from the other indicators also; and, of most concern, it is a slowing at a level short of the improvements one might hope for on the development experience of, for example, Kerala, Sri Lanka, Costa Rica and China (Caldwell 1986). Riley and Lehmann (1992) attribute the driving roles in the earlier improvement to health service provision, especially the availability of penicillin for pneumonia and chloroquine for malaria, together with other anti-malaria measures, and pertussis and tetanus immunization. Immunization campaigns are likely to have played a particularly large part in accounting for such improvements as have been seen in both infant and one to four years mortality. As Ulijaszek (1995) has pointed out, however, while immunization has brought demonstrable improvements which will have contributed to the alleviation of mortality (e.g., for measles; but see also Duke 2003), there are signs of trends in the opposite direction, even before 1990, in conditions not targeted by immunization (e.g., diarrhoea and malnutrition). This among other factors may have contributed to the unevenness of mortality trends.

Consistently over time (and consistently across all provinces in the year 2000 except Manus), infant and child mortality rates are higher for males than females. In keeping with so many dimensions of PNG life, however, mortality and mortality trends are highly variable regionally, as shown in Table 12.4. Here, for the 1980 and 2000 censuses only, key mortality indicators are shown province by province (updating Allen 1992: Table 2.2; on the basis of National Statistical Office 2003b: Tables III-1–III-4). The provinces are arranged not in the conventional order but so as to highlight trends: on the left, in ascending order of the 1980 to 2000 change in life expectancy (in years), and on the right, in descending order of the change in infant mortality across the same period (in deaths/1,000; the third numerical column in each case). These orders are highly correlated though not identical. Mortality at one to four years follows a very similar ordering to infant mortality, suggesting similarity in the underlying factors. Life expectancy at 25 years of age (e_{25}) also follows a similar ordering to life expectancy at birth (LEB; e_0), but the similarity is less close. In all cases, there is a weak negative correlation, never accounting for more than half the variation, between the 1980 value and the subsequent change: thus there is an unsurprising tendency, most noticeable for life expectancy at 25 years, for initially better results to show less

Table 12.4. Mortality indicator changes between 1980 and 2000 by province: life expectancy at birth (e_0) and at 25 years (e_{25}); infant mortality rate (IMR) and 1-4 years mortality rate (1-4MR). Under each heading the third column is the difference between the first two. Females and males are combined throughout.

	e_0			e_{25}			IMR			1-4MR		
	1980	2000	'00-'80	1980	2000	'00-'80	1980	2000	'00-'80	1980	2000	'00-'80
PNG	49.6	54.2	4.6	34.6	37.9	3.3	72	64	-8	45	25	-20
Milne Bay	57.1	54.1	-3.0	39.2	38.3	-0.9	71	103	32	48	57	9
Gulf	47.3	46.4	-0.9	32.3	34.5	2.2	50	69	19	27	28	1
N. Solomons	59.6	59.6	0.0	39.6	41.1	1.5	62	80	18	39	38	-1
Madang	50.7	51.1	0.4	34.6	36.5	1.9	62	78	16	39	35	-4
Morobe	50.9	51.7	0.8	34.8	37.3	2.5	33	47	14	17	15	-2
E. Highlands	53.1	55.4	2.3	36.1	38.0	1.9	104	105	1	74	58	-16
NCD	56.7	59.2	2.5	37.2	37.9	0.7	55	54	-1	33	19	-14
E. Sepik	49.3	52.2	2.9	36.4	37.4	1.0	57	54	-3	34	19	-15
W. Sepik	42.1	46.0	3.9	30.6	34.4	3.8	60	55	-5	37	19	-18
E. New Britain	52.8	57.1	4.3	36.0	39.5	3.5	67	59	-8	43	22	-21
W. Highlands	51.9	56.2	4.3	37.6	38.0	0.4	62	52	-10	36	17	-19
Central	51.3	56.4	5.1	34.8	38.1	3.3	55	45	-10	34	14	-20
New Ireland	52.7	57.9	5.2	36.4	40.0	3.6	59	47	-12	36	15	-21
Northern	49.2	54.5	5.3	33.7	37.6	3.9	35	22	-13	20	5	-15
Enga	47.1	52.5	5.4	34.1	36.8	2.7	94	79	-15	56	36	-20
W. New Britain	51.3	56.7	5.4	34.9	39.2	4.3	83	66	-17	53	26	-27
Chimbu	50.2	56.8	6.6	36.6	39.2	2.6	91	69	-22	58	28	-30
Western	47.7	54.3	6.6	33.8	38.2	4.4	87	54	-33	51	19	-32
Manus	51.8	58.6	6.8	34.8	39.9	5.1	81	48	-33	46	15	-31
S. Highlands	43.8	55.2	11.4	33.2	38.4	5.2	116	61	-55	75	23	-52

Note:

To the left, provinces are arranged in ascending order of change in life expectancy at birth; to the right, they are arranged in descending order of change in infant mortality rate.

Source: Data from National Statistical Office (2003b).

improvement than initially poorer ones. The weakness of this trend, however, and counter-examples visible in the table confirm that in no sense have inherent obstacles to improvement been encountered yet. The absolute mortality levels observed convey the same message.

In Gulf, Milne Bay, Morobe, Madang and North Solomons Provinces, there have been large rises in infant mortality (whether viewed absolutely or proportionally), and these provinces have also seen rises or very small declines in one to four years mortality. Accordingly, LEB has declined in Milne Bay and Gulf, and remained essentially static in the other three. All five provinces are below the national figure for improvement in life expectancy at age 25 years also, but only in Milne Bay has this indicator actually declined: in Gulf and Morobe especially, adults appear relatively protected from the adverse changes affecting children. With the civil conflicts there during this time, it is not surprising to see North Solomons among these five provinces, as a result of rising infant mortality. What is perhaps surprising is that overall life expectancy there has remained static: North Solomons had the highest life expectancy of all provinces in 1980, and perhaps the conditions which permitted that outcome, whatever they may have been, still had sufficient effect in the year 2000 to protect its life expectancy from actually declining. There is no space here to discuss possible method problems or data errors, but demographic data collected in a conflict zone should clearly be interpreted cautiously.

At the other end of the spectrum, Southern Highlands Province has the largest improvement recorded in PNG on all these indicators. With the worst infant mortality of all in 1980, Southern Highlands had the most scope for improvements, and the data suggest that these have ensued. But with civil strife there too, this province's demographic results also require particularly cautious interpretation. Other highlands provinces, Western Highlands, Chimbu and Enga, also registered large improvements over initially high levels of child mortality; but Eastern Highlands levels were already lower, and its mortality indicators have improved less, in the case of infant mortality hardly at all. Western Province and East Sepik have also shown marked alleviations of initially high levels of infant and child mortality. Manus and West New Britain have shown conspicuous rises in life expectancy despite only moderate falls in infant and child mortality, bringing Manus to the third position on both life expectancy measures, behind only North Solomons and National Capital District on e_0 , North Solomons and New Ireland on e_{25} , and almost to the point where a 25-year-old can 'expect' to live to 65 years of age.

While the right-hand half of Table 12.4 explicitly focuses on measures of pre-adult death, and LEB is also most substantially affected by that, life

expectancy at 25 years focuses attention on premature adult death. Not surprisingly, this parameter is least closely tied to the other indicators listed here. Thus Gulf, West Sepik, Morobe and East New Britain Provinces all have more favourable trends in e_{25} than one might expect from their trends in e_0 ; while in East Sepik, Enga, Chimbu and most dramatically Western Highlands, significant benefits in child survival seem not to have flowed through to adult survival as well as one might have hoped. On the figures, National Capital District also belongs in this latter category, but conditions are probably unique there – on the one hand, relatively good and improving child survival patterns set a high benchmark; and on the other, an e_{25} no higher than the national average, exhibiting only marginal net improvement in 20 years despite the capital's facilities, probably provides the clearest signal apparent in this table of encroaching health problems (infectious and non-infectious) of modernization.

Table 12.4 arranges provinces according to absolute mortality trends rather than relative trends or absolute outcomes. Some features of mortality variation not highlighted by this procedure nonetheless deserve attention. For example, West Sepik (Sandaun) Province sits inconspicuously near the middle of both listings, being at neither extreme in its mortality trends; but its absolute mortality levels are extreme. Already one of the more disadvantaged provinces in 1971, by 1980 it had clearly the shortest life expectancies of any province, and the highest pre-adult mortality rates except for the Southern Highlands Province. By the year 2000, strikingly unlike Southern Highlands, infant mortality had not improved at all; and although other measures had improved modestly, West Sepik was the worst placed province on all measures, with Gulf now not far behind. The deterioration in Gulf Province has been more dramatic but has not resulted in a worse outcome in absolute terms. Worse figures than this are reported from specific localized populations in West and East Sepik Provinces; for example, IMRs among the Abelam (cited above), the Saniyo-Hiowe (400 per 1000: Townsend 1985) and the Mianmin (200 per 1000: Attenborough *et al.* in prep.).

It would be very interesting to renew the question (cf. Caldwell 1986; Inaoka n.d.) where PNG fits in terms of the contemporary relationship between economic variables and mortality improvement. Has PNG been able, as Sri Lanka was, for example (Caldwell 1986; Caldwell *et al.* 1989), to achieve good health and mortality outcomes relative to other nations at a similar economic level; or similar, or worse ones? Unfortunately there are data obstacles to answering this question rigorously. Diversity of income is as striking as many of PNG's other dimensions of diversity (Hanson *et al.*

Table 12.5. *Life expectancies at birth and trends in life expectancy at birth (e_0) for countries with gross national income (GNI) between US\$ 400 and US\$ 600 per caput, 2003*

	GNI/caput US\$ 2003	e_0 2003	Years added to e_0 since 1970
Zimbabwe	480	33	– 22
Lesotho	590	35	– 14
Moldova	590	69	4
Uzbekistan	420	70	7
Benin	440	51	9
Mauritania	530	53	11
Mongolia	480	64	11
Timor-Leste	430	50	11
Guinea	430	49	12
Senegal	550	53	12
Comoros	450	61	13
Pakistan	470	61	13
Sudan	460	56	13
India	530	64	15
PNG	510	58	15
Solomon Islands	600	69	15
Bangladesh	400	62	18
Vietnam	480	69	20
Yemen	520	60	22

Note:

Countries are arranged in ascending order of life expectancy trend.

Source: Data from UNICEF (2004a).

2001), so much so that an average should not attract too much attention. Furthermore, with 87% of the population of PNG still living in rural areas and predominantly still covered by traditional means of access to land, food, shelter and community support, resource measures based only on the formal money economy are far from satisfactory for this purpose. Nevertheless, for what they may be worth, some selected international comparison figures are presented in Table 12.5, where absolute and trend life expectancy figures are shown for countries (many with significant traditional resource access also) with annual per caput incomes of US\$400 to US\$600. On the estimates of the United Nations Children's Fund (UNICEF), PNG emerges neither as well placed as Bangladesh, Vietnam and Yemen, which have both higher life expectancies and larger gains in life expectancy, nor as unfortunate as those southern African

countries worst afflicted by AIDS, or even, in terms of the trend rather than the absolute figures, the former Soviet republics included in the table. Rather, PNG is near the upper end of an intermediate group, with gains in life expectancy similar to India and Solomon Islands, but absolute life expectancy for 2003 (though higher than the NSO figure for 2000 cited above) below theirs. Within this income range, no clear relationship between income and life expectancy emerges.

Leading causes of death are hard to identify in the absence of comprehensive vital registration systems. Figures at a national level tend to come from hospitals and health centres, and since many deaths take place away from such centres, there is an obvious potential for distortion in the data available. Local surveys, such as those collated by Riley and Lehmann (1992) for the 1960s to 1980s, confirm, however, that the pattern of the past has been one dominated by infectious causes of death: particularly respiratory infections and, especially (but not exclusively) in the lowlands, malaria. Diarrhoeal disease is important but less so than in many developing countries. Neonatal and congenital causes are important in early childhood, as are accidents and violence in adults. On World Health Organization summary data (World Health Organization Regional Office for the Western Pacific 2005), the leading causes of mortality in the year 2000 were, in descending order, pneumonia, perinatal conditions, malaria, tuberculosis and meningitis; and of morbidity, obstetric causes, pneumonia, malaria and perinatal conditions. Thus the recent picture has much in common with the earlier one. This chapter cannot be comprehensive in examining trends in these and other conditions; but trends in selected causes of mortality and morbidity will be examined more closely in what follows.

Health problems of modernization

First, I shall consider some evidence on trends in health problems associated with modernization. There can be no doubt but that significant health and nutritional benefits have accompanied modernization in PNG. At Ok Tedi, where mine employment and elements of a 'modern' lifestyle were introduced over very few years in the 1980s to a previously very remote region in Western Province, and also in Simbu Province, where data extend back more than four decades documenting somewhat less dramatic socioeconomic changes associated with cash-cropping for example, improvements in nutritional health were evident for at least a large part of that time: see review by Ulijaszek (1995). Whether trends in

single locations or contemporaneous contrasts between more and less economically developed locations are considered, modernization appears linked, usually, with faster physical growth of children (Ulijaszek 1995; Müller *et al.* 2001) and higher birth weights (Ulijaszek 2001; Müller *et al.* 2002a; but see also Lourie 1986; Tracer *et al.* 1998).

A review of the impact of cash-cropping on nutritional health (Heywood and Hide 1994) found cash cropping to be associated with increased child growth, especially in the highlands from where the best data are available, but probably also in the lowlands where infant mortality declined over the same period. The evidence did not bear out pessimistic anticipations of rising malnutrition. As one might expect, there was a concomitant secular increase not only in adult weight and height, but also in the prevalence of various syndromes and diseases of modernization. While it is difficult to establish thresholds for individuals beyond which growth and anthropometric increases bring more cost than benefit, two broad conclusions seem clear: (a) that modernization has brought substantial nutritional health improvements to many PNG communities; and (b) that there have also been health costs at a population level to the nutritional changes that accompany modernization.

No cases of myocardial infarction were found in a large series of autopsies in Rabaul, New Britain, in the 1920s and 1930s (T. C. Backhouse, reported Riley and Lehmann 1992). In 1966 a survey of the Enga community of Murapin in the PNG highlands found no evidence of coronary artery disease, cerebrovascular disease, hypertensive disease or diabetes (Sinnott 1977). Serum cholesterol, serum triglycerides and blood pressure were all low, and showed no increase with advancing age, or even fell. While few studies from that period are so comprehensive, it seems reasonable to accept that these results are quite representative of PNG communities under conditions little influenced by modernization. Those cardiovascular diseases which are associated with modernization, and many of their biological risk factors, are essentially recent phenomena in PNG.

It has been proposed, on the basis of African experience, that in general hypertension is the first disorder of modernization to emerge as the process gets under way, followed by obesity, type 2 diabetes mellitus (or non-insulin-dependent diabetes, NIDDM, usually mature-onset, of which impaired glucose tolerance (IGT) is a precursor), cerebrovascular disease and finally ischaemic heart disease (Trowell 1981). By the 1960s certain more urbanized communities, for example in the Port Moresby region, were providing evidence of hypertension, with blood pressures that had begun to rise with age, weight increases and higher, though still not high, prevalence of type 2 diabetes (Sinnott *et al.* 1992). At the Ok Tedi

mine site, blood pressure rose significantly within three years in the mid 1980s, among a range of other human biological changes (Lourie 1987; Ulijaszek 1995). Hypertension has now been described in a rural Purari delta population (Ulijaszek 1997). The first description of a myocardial infarction in an indigenous Papua New Guinean was that of Conyers (1971). More cases were identified through the 1970s, and by the 1980s death was attributable to ischaemic heart disease in 11% of autopsy samples in Port Moresby, but only in 2% of similar samples in the smaller town of Goroka (Misch 1988). A further series of cases with high case fatality in Port Moresby hospital confirmed the trend (Kevau 1990).

Type 2 diabetes, similarly, was a very rare diagnosis in the 1960s (Ogle 2001). A study of five PNG communities in 1991, urban, peri-urban and rural, lowland and highland, used an existing 'modernity score' to track the development of diabetes and other heart disease risk factors (Hodge *et al.* 1996). This score was based on education, employment, access to and experience of urban centres, type of housing and so on (King and Collins 1989). A subsequent study has found distance from urban centre to be a useful prime index of modernization (Ulijaszek *et al.* 2005). Hodge *et al.* (1996) found 'modernity score' to be positively associated with overweight and type 2 diabetes. Furthermore, they found cholesterol concentrations to be higher than previously recorded in PNG and to show age-related increases; the same was true of triglyceride concentrations in women. In an urban community, total cholesterol was similar to Australian levels while HDL cholesterol was lower. The authors point out that PNG's earlier freedom from heart disease by no means signals that Papua New Guineans are protected from developing blood lipid levels that bring risk in other populations – no evidence of special adaptation here – and that without preventive strategies the incidence of coronary heart disease can be expected to increase further.

For the three coastal communities included in the study of Hodge *et al.* (1996), there are also diabetes data from earlier studies (Dowse *et al.* 1994). The majority of the results on both NIDDM IGT exhibit an upward trend (Table 12.6), especially those from Koki; but these results are also diverse. Koki is a squatter settlement near the centre of Port Moresby. Most residents of Koki originate from Wanigela, about 200 km southeast of Port Moresby. As we might expect, it has both the highest modernity score and the highest IGT and NIDDM prevalences, very high on a global scale. Wanigela has medium modernity scores, very similar to Kalo, which geographically is situated between the other two. But in frank diabetes mellitus particularly, Kalo is much less severely affected. The authors could not explain this last contrast in lifestyle terms, and proposed that the two

Table 12.6. Trends in age-standardized prevalence of impaired glucose tolerance (IGT) and non-insulin-dependent diabetes (NIDDM) in three coastal communities in Papua New Guinea

	1977/1983	1991
<i>IGT (%)</i>		
Koki	18.0	34.2
Wanigela	6.7	14.1
Kalo	14.1	10.3
<i>NIDDM (%)</i>		
Koki	16.7	27.5
Wanigela	9.8	11.1
Kalo	1.2	1.5

Source: Dowse *et al.* (1994), where sample sizes, confidence intervals, age standardization method and diagnostic criteria are detailed.

(largely endogamous) Austronesian-speaking groups of Wanigela origin may have a high frequency of a specific genotype, perhaps more susceptible to the diabetogenic effects of a modernizing life style than any other known. Neither this lifestyle nor the hypothesized genotype currently affect the majority of the populations of PNG to the extent seen in Wanigelans: on the other hand, it has become clear that any genetic protection non-Austronesian-speakers may have is only relative (King *et al.* 1989). The contrast between Koki and Wanigela underlines the potency of lifestyle factors, without denying that genetic factors may be important too. Prevalence in the country as a whole is rising but not yet high, and above all diverse (Ogle 2001).

Modernization does not only take place *in situ*: people migrate towards it, on a large scale in the case of PNG's larger towns. Studies among migrants to Port Moresby have drawn attention to the emergence of cardiovascular and diabetes risk factors in these groups also, and to how this varies markedly according to their geographical origin within PNG, length of stay in Port Moresby, occupation and sex (Natsuhara *et al.* 2000).

The role of a modern lifestyle in increasing the prevalence of cardiovascular disease, in which NIDDM is a major risk factor, has received enormous attention and other widely accepted risk factors include (1) high dietary intake of fats, especially saturated fats; (2) high dietary intake of sodium, in connexion with hypertension and haemorrhagic

stroke; (3) excessive alcohol consumption; (4) low levels of physical activity; (5) obesity, especially with centripetal fat distribution; and (6) psychosocial stress. These classical risk factors account for some but not all of the observed variation in cardiovascular disease within- and between-populations around the world. Newer additional hypotheses implicate early-life undernutrition, marked catch-up growth and/or infectious agents such as the common bacteria *Chlamydia pneumoniae* and *Helicobacter pylori* in the disease processes leading to elevated risk of cardiovascular disease (Pollard 1997). These agents, perhaps potentiated only in the presence of the classical risk factors already discussed, could, if either or both of these hypotheses are correct, be particularly devastating for populations currently experiencing a rapid transition from under-development to modernization. PNG in the near future may perhaps represent just such a case.

Discussions of modernization and health frequently select, as I have here, hypertension, ischaemic heart disease, type 2 diabetes and their risk factors for special attention; and there are good reasons for this. But these are by no means the only non-infectious diseases that show a rise associated with modernization: dozens have been listed (Trowell and Burkitt 1981), including, for example, asthma (Anderson and Woolcock 1992).

Infections, old and new

Is the rise of diseases of modernization the main feature of population health in PNG to emerge as the nation modernizes? If we extrapolate existing trends, can we foresee an automatic dwindling of the diseases of underdevelopment, as the classic diseases of modernization rise?

Other processes, especially affecting communicable disease, ensure that the prospects are not so simple. Despite some declines in particular infections in conformity with the epidemiological transition model, it is infectious diseases collectively that still represent the country's main health problem, causing large numbers of deaths, especially deaths of children and younger adults, as well as many non-fatal episodes of ill-health (Carrad 1987; Riley and Lehmann 1992). Pneumonia, due to a variety of bacterial and viral pathogens, and malaria, due to four *Plasmodium* species as discussed above, predominate among these (Cattani 1992; Riley *et al.* 1992).

There is not enough space here to explore all the important themes, so despite their prominence in the statistics, I shall comment only briefly on respiratory infections. The continuing importance of pneumonia (acute lower respiratory infections, ALRI), and of specific acute (e.g., influenza)

and chronic (e.g., tuberculosis) respiratory infections is clear. Pneumonia is especially salient in the highlands, where malaria is less prevalent, but important everywhere. At Tari in the Southern Highlands Province, in the period 1977 to 1983, ALRI accounted for 53% of infant deaths, 34% of child deaths (one to four years), 18% of deaths in people over 60 years of age and 23% of deaths overall (Lehmann 2002). By comparison, neonatal/congenital abnormalities accounted for 21% of infant deaths, other febrile illness (presumably including malaria in lower-lying parts of the study area) for 8% of them, gastroenteritis for 5% and measles for 4%. Mortality due to ALRI had actually risen since 1972 to 1974, for which there are several possible reasons including penicillin resistance. However, after a pneumococcal vaccine trial in the area in 1981 it fell significantly, even though half the participants had received only a placebo vaccine. The vaccine has been shown to be effective (Riley 2002), and this development appears to offer the best way forward for public health approaches to ALRI. Unfortunately the benefits it offers have yet to be translated into lasting benefits for large numbers of Papua New Guineans, because the vaccine has not become available at a price that PNG can afford.

Since the Second World War and into the period of national independence, PNG has developed ways of bringing the benefits of its health care system beyond the cities and towns to its predominantly rural population, notably a network of aid posts and health centres linked to the urban hospitals. At a minimum, government and non-government (e.g., mission) facilities alike have been able to offer first aid, antibiotics (e.g., procaine-penicillin injections) and malaria medications (e.g., chloroquine/amodiaquine), and have run maternal and child health clinics. Immunization programmes have been a particularly important part of PNG's public health system in both rural and urban areas. Indeed, as already discussed, PNG's successes in countering infectious disease at a national level have essentially been with those infections against which an immunization programme has been widely implemented – and not, to date, with pneumonia, influenza, diarrhoea or malaria.

A further concern arose during the 1990s and remains current, to the effect that even those public health gains which had been made by then are not secure; that health systems are experiencing some deterioration owing to funding and organizational problems; and that population health improvements are stalling as a result. For example, real per caput expenditure on health is declining, geographical coverage is uneven, many aid posts have closed, and harassment of staff, theft of equipment and vandalism of facilities are reported to be frequent (Ministry of Health 2001). In these circumstances it is not surprising that existing immunization

programmes only reach half of all children, or that, as discussed elsewhere, more than half of women giving birth do so without medical or paramedical assistance, and overall mortality and morbidity outcomes in the past 15 years have tended to be disappointing. Connell (1997) argues strongly that both health systems and population health status have deteriorated substantially since the 1980s and that the prospects for improvement are poor; see also Edwards (1994).

Malaria

Among PNG's myriad health problems, only malaria – the most frequent outpatient diagnosis and the second most frequent diagnosis amongst health facility admissions in many endemic areas – rivals pneumonia in importance at the present time (Ministry of Health 2001). In some provinces and age categories, it equals pneumonia as a primary cause of death. At its maximum, malaria transmission in PNG occurs on a holoendemic scale comparable with equatorial Africa.

Malaria is an environmental disease among other things, dependent on warmth (minimum ambient temperatures of 18 °C for *Plasmodium falciparum* and 16 °C for *P. vivax* for completion of the parasite's life cycle in the mosquito) and aquatic breeding places for *Anopheles* mosquitos. It is thus not surprising that its transmission in humid equatorial PNG is difficult to disrupt (Cattani 1992). In PNG, environmental temperature depends mainly on altitude, and so malaria is highly endemic in the lowlands (except during the marked dry season found in some southern lowland areas). Transmission decreases with elevation above sea level, becoming unstable and more epidemic in character around 1,300 to 1,600 m, and ceasing above 1,700 to 1,800 m (Müller *et al.* 2003). With that said, malaria prevalence and transmission are highly variable geographically, even on a local (village to village) scale, with mosquito biting rates of up to 500 bites/person/night in some places, much less in others a few kilometres or less away (Cattani *et al.* 1986a; Burkot *et al.* 1988; Attenborough *et al.* 1997).

Plasmodium infections do not always result in clinical illness, but most frequently do so in those with the least effective immune protection, especially children under ten years of age, pregnant women, and residents of non-endemic areas. While its seriousness is not in question, severe and lethal malaria is less frequent in PNG than in areas of comparable endemicity in Africa, perhaps because of the multiple genetic protections against malaria already discussed (more numerous and varied than

Africa's principal genetic protection, the sickle-cell gene), and/or because of the higher prevalence than in most parts of Africa of vivax malaria, which may confer some cross-protection to the more dangerous falciparum malaria (Maitland *et al.* 1996; Smith *et al.* 2001; Bruce and Day 2003; Müller *et al.* 2003; Mayxay *et al.* 2004). High rates of spleen enlargement in adults as well as children are another distinctive feature of malaria in PNG, especially in the highland fringes and associated with high risk of premature death from indeterminate causes (Crane 1986).

Malaria's prehistory in New Guinea has been discussed above. The beginning of systematic data collection there came with the work of Robert Koch and his colleague Otto Dempwolff in Kaiser-Wilhelmsland (German New Guinea), just over a century ago (Ewers 1972). Malaria was a major cause of mortality among Europeans, Javanese and Chinese in the newly founded German colony and the high death rate caused survivors to abandon some of their settlements. Thirteen out of 45 Europeans died of malaria during one three-month period of 1891 at Finschhafen (now in Madang Province). Of 273 Chinese at Stephansort, near the present town of Madang, 125 died during 1898, mostly of malaria.

Koch and Dempwolff also studied malaria among the indigenous people between 1899 and 1903 and found it to be highly endemic through much of the area, though they also considered some villages and some off-shore islands to be malaria-free or to be experiencing malaria as a new disease. Koch was impressed with the importance of malaria as a cause of illness and death; and, we may now wryly read in the light of the subsequent retreat from such ambitions, optimistically instructed Dempwolff not only to study malaria amongst village people but also to attempt its eradication. Both around the Madang coast and on the Bismarck Islands, they found the prevalence of malaria (as indicated by parasitaemia at all ages, with all species) to be generally between 15% and 25%. These are low figures by comparison with more recent ones (e.g., approximately 35% to 45% for Madang area communities in 1981 to 1982, reported by Cattani *et al.* 1986b). With geographical variations, seasonal fluctuation and the possibility of different or less sensitive methods all confounding the comparison, however, we cannot be sure that malaria was less prevalent in 1900 around Madang than it is now. At any rate, despite all efforts made in terms of malaria treatment and control campaigns, there is not even a hint here of any downward trend over the twentieth century.

The locations which now have New Guinea's highest recorded endemicities of malaria, in the Sepik basin, were not covered in this early German colonial research. But one such location, Maprik District, provides one of the best data sets in New Guinea for malaria, from the 1950s, during the

Table 12.7. Malaria prevalence (%) in Maprik, East Sepik Province: proportion of positive blood slides

	Date	P.f.	P.m.	P.v.	c.r.
All ages	1957	16.1	17.0	29.5	62.6
	1963	13.6	21.2	9.7	44.5
	1984	22.6	19.5	7.7	49.8
1-6 years	1957	24.1	25.6	38.5	90.3 ^a
	1963	19.8	19.8	16.0	55.6
	1984	50.0	37.5	0.0	87.5
7-15 years	1957	11.7	13.0	28.2	62.3 ^a
	1963	13.2	25.3	8.4	46.9
	1984	24.3	21.6	21.6	67.5
>15 years	1957	7.1	7.3	18.7	38.2 ^a
	1963	10.4	20.4	7.0	37.8
	1984	14.0	11.3	4.7	30.0

Notes:

P.f. = *Plasmodium falciparum*; P.m. = *Plasmodium malariae*; P.v. = *Plasmodium vivax*; c.r. = crude parasitaemia rate, i.e., % of blood slides positive for any *Plasmodium* species.

^a Usually, crude rates may be lower than the sum of the species-specific rates, owing to infections with multiple species (not apparent here), but are not higher. Desowitz and Spark do not comment on why they are 2-10% higher for the 1957 age-specific data from Peters (1960). Explanations invoking the possibility that in some cases the species was indeterminate or was the rare *P. ovale* would have to cover the fact that the all-ages data do not have a higher total crude rate.

Source: Desowitz and Spark (1987).

Australian colonial period, to quite recent times. The available data (Table 12.7) allow us to look at trends from the 1950s to 1980s, in the main *Plasmodium* species as well as in the crude rate of infection (Desowitz and Spark 1987). The crude rate went down and up again, but what should perhaps strike and concern us most is the changing species balance. While *P. malariae* remained more or less constant (and higher at the location studied than at most others), *P. vivax* declined; and *P. falciparum*, the most dangerous of the species, increased in prevalence, both relatively and absolutely. In 1990 to 1992, surveys in another part of the Maprik District, Wosera, undertaken in preparation for vaccine trials, again found high prevalence of malaria at 60% (at all ages, with all species, ranging from 49% to 70% between villages: Genton *et al.* 1995b), a similar level overall to those of the 1950s shown in Table 12.7, and up from those of the 1960s and 1980s. *Plasmodium falciparum* was the predominant species (observed in 55% of all positive blood slides: Genton *et al.*

Table 12.8. Malaria prevalence (%) in Karimui, South Simbu: proportion of positive blood slides

	P.f.	P.m.	P.v.	c.r.
1965	7.1	5.6	6.7	19.4
1971	4.0	0.7	2.4	7.1
1981	12.3	3.9	14.1	30.3
2001/02	21.3	3.3	6.6	31.2

Notes:

P.f. = *Plasmodium falciparum*; P.m. = *Plasmodium malariae*; P.v. = *Plasmodium vivax*; c.r. = crude parasitaemia rate, i.e., % of blood slides positive for any *Plasmodium* species.

Source: Müller *et al.* (2005).

1995b), a position it had approached in the 1980s (45%) but not in the 1960s (31%) or the 1950s (25%) (Desowitz and Spark 1987). There is little seasonality to confound this picture, although local differences within Maprik may do so to some extent. Most recently, a significant drop in the prevalence of *P. falciparum* but not *P. vivax* has been observed in the Maprik area (Kasehagen *et al.* 2006). Still, the overall impression remains that the problem is not declining.

The shift in species balance is not a trend purely local to Maprik or the Sepik basin. The 1980s Madang figures show a predominance of *P. falciparum*, although we do not know the earlier species balance there. A local survey in the highlands fringes of West Sepik Province (Attenborough *et al.* 1996) shows a predominance of *P. falciparum* also, though again we have no time series data. On the Karimui plateau, Simbu province, at 900 to 1,200 m, malaria is more prevalent than at similar altitudes in some other locations, but here we do have another good time series (Table 12.8; Müller *et al.* 2005). The crude parasitaemia rate rose, later than in the Sepik lowlands, and most strikingly *P. falciparum* prevalence rose both absolutely and relatively.

Are the shifts in species balance that we have seen in several locations linked to economic modernization and globalization; or do they merely coincide with these developments? The changes matter either way: perhaps it is more important to identify specific factors that might be responsible than attempt to answer that question in broad terms. In reality, both overall prevalence and species balance have changed in ways too complex and too frequently reversed to be plausibly tied to a single process, whether we label it modernization or not. No single factor explains all aspects of the pattern, but ones which may be relevant to variations in malaria epidemiology both in time and space include (Müller *et al.* 2003): (1) vector control programmes,

for example, DDT and dieldrin residual spraying; (2) large-scale drug administration (in the past mainly with chloroquine); (3) use of bednets, especially when insecticide-impregnated; (4) drug resistance evolved by *Plasmodium*, especially *P. falciparum*; (5) variations in the ecology, behaviour and vectorial capacity of the mosquitos which act as vectors in PNG (principally the *Anopheles punctulatus* group, now known to number at least 10 species in PNG, rather than the three recognized before 1993: Beebe and Cooper 2002); and (6) relevant variations in human numbers, behaviour and genetics.

In Maprik, in 1957, no mosquito-spraying operations had yet taken place and anti-malarial drugs were available only for the treatment of acute cases. By 1963, there was still no spraying locally (it started shortly afterwards); but anti-malarial drugs, especially chloroquine, were more readily available, and this may have led to a reduction in the prevalence of *P. vivax*. In 1984, spraying had virtually ceased, but chloroquine was still more accessible and this appears to account for a further reduction in *P. vivax* rates. Mechanisms for these trends and especially for the rise of *P. falciparum* remain to be clarified. The clinical drug-resistance of some *P. falciparum* strains, especially to chloroquine, was not in evidence until the 1970s; but by the 1980s it was highly prevalent (Müller *et al.* 2003), so clearly it could be important in the rise of *falciparum* apparent in the 1984 and 1990 to 1992 Maprik data. The recent drop in *P. falciparum* prevalence in Maprik may be related to the introduction in the year 2000 of a new standard first-line treatment (chloroquine/amodiaquine plus sulphadoxine-pyrimethamine).

In Karimui, the 1965 survey preceded control measures. The 1971 survey, immediately following a mass drug (chloroquine and pyrimethamine) administration campaign, and early in the period of indoor residual DDT spraying to control vectors, found substantially reduced overall malaria rates but also a species balance apparently tipped towards *P. falciparum*. Vector control measures became less effective from 1978 onwards, and were abandoned in 1984, in line with similar trends elsewhere as it became apparent that eradication would not be achieved; thus the 1981 survey took place at a time of failing control. This is no doubt linked to the large increase in both *P. vivax* and *P. falciparum* prevalence found in this survey, relative to both 1971 and 1965 levels. In the following 20 years from 1981, not only did control measures cease but also chloroquine-resistant *P. falciparum* became abundant, which may play a key part in explaining why by 2001 *P. falciparum* had become much more prevalent while *vivax* had declined. Population growth, a shift from dispersed long-houses to more centralized villages, and correspondingly larger, more clustered land cultivation patterns may also have contributed (Müller *et al.* 2005).

The rise of chloroquine-resistant *falciparum* malaria seems to have a clear role in the shifting species balance and the absolute rise of

P. falciparum. The timing of the DDT indoor residual house spraying programme, relative to the rise of *P. falciparum* and the decline of *P. vivax*, suggests that it may have a role, though again the mechanism is unclear. One suggestion is that it led to a shift from indoor late-night biting to outdoor dawn and dusk biting (Spencer *et al.* 1974), perhaps by younger mosquitoes or different species, with different propensities to transmit *P. falciparum* and *P. vivax* (Bockarie *et al.* 1996). However, this remains to be resolved. Also some evidence (discussed above) might support hypotheses that reduction in *vivax* prevalence is conducive to an absolute as well as relative rise of *P. falciparum*. Among possible candidate factors suggested as having favoured *P. vivax* initially, related to its long-lasting liver stages and other biological characteristics, are insecticide spraying (irregular or otherwise) and mass drug administration. Vines (1970) found *P. vivax* to be more prevalent in sprayed areas, and more recent results elsewhere have also shown a shift to *P. vivax* in the course of effective vector-control programmes. Indiscriminate anti-malarial drug use patterns and the rise of drug resistance in *P. falciparum* particularly in the post-vector-control period appear to have reversed any such initial advantage to *P. vivax*, however.

It can plausibly be argued that before the Australian colonial intrusions that began in 1930, the central highlands of PNG were generally free of endemic malaria, except in low-lying and swampy areas. Before mid-century, legislation regulated recruitment of plantation labourers from above 1,200 m, which delayed the spread of malaria into the highlands (Radford *et al.* 1976). Eventually, however, under the influence of post-war economic development, and with relaxation of controls on recruitment and of compulsory malaria prophylaxis for highlanders working at the coast, epidemic malaria outbreaks in the highlands became a more significant phenomenon than before, with a more fluctuating and epidemic pattern than in the lowlands, and with marked seasonality. Between the late 1940s and the early 1960s, these outbreaks were – successfully, at the time – counteracted by means of a DDT-spraying programme in the most densely populated areas, including Eastern Highlands, Western Highlands and Simbu Provinces. By 1974, the health authorities were no longer attempting to control the spread of infection through controls on the mass movement of people, or through vector control. Sharp (1982) reported a parasite rate in non-epidemic situations in western Enga Province of 12% in 1979, all due to *P. vivax*; but a parasite rate of 75% (in surveys angled towards febrile individuals) below 1,600 m during epidemics there in 1980, nearly all due to *P. falciparum*. Recent surveys have shown increased malaria in outlying highlands areas over 1,200 m, though

not in all areas of this altitude (I. Müller, personal communication). Malaria is now a troublesome health problem throughout the highlands provinces (Müller *et al.* 2002b).

Two further factors not mentioned so far promise, or threaten, to exert impacts on trends in malaria prevalence. On the one hand, we still have the hope of the vaccine, on which much has been achieved in research terms, although there are still obstacles to its general implementation to bring about a major reduction in the population health burden imposed by malaria (Brown and Reeder 2002; Genton *et al.* 2003). Even when it is available, the malaria vaccine is expected to bring a reduction in frequency and severity rather than a prospect of eradication. On the other hand, there is at least one further reason to predict an upward trend in malaria, unless the vaccine or other control measures can offset it – climate change. While the smaller Pacific nations have other reasons to be concerned about global climate change, a major consideration for PNG is the possibility of its effects on vector-borne disease including malaria – even the possibility that such effects are already under way, contributing to the trend in highlands malaria just discussed. Available data for New Guinea are insufficient to say at present, but projected climate change would be sufficient to raise the altitude cut-offs for the transmission of malaria by an amount that could be significant in terms of the location and movements of human populations. In the Ethiopian highlands, monthly night-time temperatures are reported to have risen by about 1.5 °C from 1980 to 1993, and with them – though there can be debate about what this means – has risen the incidence of falciparum malaria and the altitude cut-off for malaria there (Tulu 1996; Kovats *et al.* 2001).

One of the main points of the chapter is then, using malaria as a prime example, to underline the continuing salience of communicable disease in PNG's overall health profile. It is not that some primaevial pattern simply continues unchanged: but rather that malaria is still there, continuing unabated, but shifting in its expression as its prevalence rises, its species balance changes and its geographical territory expands, under the influence of evolving drug resistance, changing strategies of treatment and prevention and their implementation, modernizing ways of life and patterns of movement, perhaps dietary trends, possibly climate change and no doubt other factors yet to be definitely identified.

HIV/AIDS

While malaria is my prime example of the continuing prominence of communicable disease, some counterbalance is required to illustrate the variety of

those communicable disease problems which show little sign of spontaneous decline as modernization proceeds. Other vector-borne diseases could be chosen, especially since many of them might be expected to spread or intensify under climate change: for example, a number of arboviruses; or Bancroftian filariasis (Kazura and Bockarie 2003). Given the practical reality in rural PNG that acute febrile illnesses are often treated presumptively as malaria until there are indications otherwise (Müller *et al.* 2003), it is likely that the incidence of such illnesses tends to be underestimated. Similarly, tuberculosis remains an important problem, undefeated and likely to be exacerbated as AIDS increasingly takes hold. But it is AIDS itself, strongly contrasting with malaria in several respects, including the historical recency of its presence in PNG, which will constitute my main counterbalancing example.

HIV/AIDS was first diagnosed in PNG in 1987 (Centre for International Economics 2002; National AIDS Council Secretariat and Department of Health 2002). Initially thereafter, numbers of cases reported remained small; but from the mid 1990s the upward trend became steeper, more than trebling between 1996 and 1998, and continuing to rise thereafter without evidence of plateauing in the data available so far. By March 2002, 5,239 people (2,665 male, 2,391 female, 183 sex not recorded) had been diagnosed as infected, more of them through presenting clinically with an AIDS-defining illness than through a positive HIV test; and 273 deaths had been reported (National AIDS Council Secretariat and Department of Health 2002). Seventy-one percent of these cases were detected in the National Capital District, 12% in Western Highlands Province, and 5% each in Eastern Highlands and Morobe Provinces, although no province was case-free. The manner of transmission was unrecorded for most cases, but where recorded, exposure was predominantly by heterosexual sex (72% for males, 75% for females), with vertical transmission the only other frequently recorded exposure category. Age at diagnosis, where recorded, was most commonly in the 20 to 34 years age range for both sexes: males outnumbered females in all adult age categories except those of 15 to 29 years.

These figures, based as they are on reported cases, need to be interpreted cautiously. Numbers of AIDS diagnoses and HIV-positive test results naturally depend on the availability and uptake of diagnostic and testing facilities. The higher female numbers during childbearing years, for example, probably reflect routine testing at the Port Moresby General Hospital antenatal clinic. Therefore we cannot be confident that the available data mirror the historical course of the epidemic in PNG to date or give an accurate measure of current incidence or prevalence, especially beyond the major urban centres where facilities are concentrated. In fact, we can be confident that the available data seriously underestimate the true extent of

HIV infection, although the degree of underestimation is hard to gauge: the reported cases may represent 5% to 25% of the true numbers according to some estimates made (Centre for International Economics 2002). It is unclear whether the true trend since 1987 or since 1996 is more steep, or less so, than appears from the available figures; whether the age/sex and transmission category figures of the recorded cases are representative of all cases; and whether, when the urban bias due to the availability of facilities is allowed for, HIV/AIDS is still a problem much more of Port Moresby and other urban centres than of the rural population.

Nonetheless, it is difficult to doubt that HIV infection is rising in prevalence, spreading geographically, and likely to continue both trends. The National AIDS Council has run a vigorous public campaign, including promotion of condom use; but a number of factors militate against its message. These include (Lemeki 2003): (1) sexual mores and attitudes to fertility such that one or both sexes are often reluctant to use condoms; (2) contexts for sexual interaction which can include, on a significant scale, multiple sexual partnerships, sexual violence and exchange of sex for cash, whether commercial or simply occasional; (3) much poverty, economic inequality and urban-rural migration; (4) high levels of other sexually transmitted infections in some areas and of tuberculosis; and (5) impoverished facilities for HIV/AIDS treatment as well as diagnosis. Comparisons with the situation in the worst affected southern African and South East Asian countries are being made.

The Centre for International Economics (2002), in a report for the Australian government aid agency AusAID, reviewed many of the points just summarized, and concluded, while acknowledging many uncertainties, that PNG appears poised on the brink of a major epidemic. It raised the prospects that by the year 2020 there could be (1) a decline in PNG's already low life expectancy, by up to 20% to 25%; (2) a national population between 13% and 38% smaller than it would have been without AIDS; and (3) negative economic growth. Concern as to the likely future impact of HIV/AIDS on PNG's population health is supported by subsequent reports that HIV/AIDS has become the second most common admission diagnosis at Port Moresby General Hospital after tuberculosis, and its most common cause of death (G. Tau, reported McBride 2005); and that the prevalence of HIV antibody in 300 patients (both sexes, ages where known, ranging from 10 to 69 years) presenting to the emergency department at Port Moresby General Hospital was 18% (Curry *et al.* 2005). HIV-positive individuals in the study of Curry *et al.* were of both sexes, and ranged in age from 18 to 60 years: their presenting illnesses were predominantly respiratory tract infections (most often pulmonary tuberculosis) or gastrointestinal tract infections.

Conclusion: double jeopardy?

In this chapter, with a view to exploring the health patterns of PNG as it makes its complex transitions from tradition to modernity, I have traversed a very wide range of topics, from prehistoric genetic adaptations to the nutritional and lifestyle diseases and emergent infections associated with modernization. How do the various themes link with each other?

There is some validity in evolutionary arguments (e.g., Boyden 2004) according to which a high level of adaptedness can be expected to build up in humans (or any other organisms) exposed for many millennia to a constant or only slowly changing environment. Rapid and radical change in environment or way of life may cause adaptedness to break down in some respects, and with this breakdown in adaptedness, quite plausibly, comes the emergence of new health problems. Thus adaptation may help explain the relative success with which Papua New Guinean populations have accommodated to malaria: its breakdown may help explain the rise of circulatory disease, diabetes and asthma there (and elsewhere). But this approach can only take us so far. An adaptive equilibrium does not guarantee good health and long life. As we have seen, modernization has been accompanied by rising life expectancy and the alleviation of some old health problems, even while new ones have emerged.

One intellectual move which may be appropriate here is to expand the concept of adaptation beyond the strictly evolutionary model which I have followed thus far. Perhaps vaccination programmes, for example, can be viewed as cultural adaptations – so long as this does not lead us to count on them as automatic. A 'people will adapt' attitude, sometimes encountered in the climate change literature for example, risks complacency. If we are to think in terms of successful biocultural adaptation, then surely thought-through responses must be part of that. It will take more than the slow ticking of genetic evolution, the exercise of physiological homeostasis and the unplanned accretion of historical and cultural change to meet the complex challenges of current health trends in PNG.

There is also, unquestionably, considerable validity in health transition models. It is clear that in some of PNG's highly diverse communities, more of them than in the past, and with more reach from the urban into the rural areas, the syndromes and diseases associated with modernization are becoming epidemiologically important. That is a trend that can be expected to continue. Some communities may be genetically predisposed, and/or predisposed by the coexistence of early undernutrition and highly prevalent bacterial infection, to the development of type 2 diabetes and ischaemic heart disease, although the role of modern diet and activity patterns is better

established. Non-infectious disease is now a matter for real and rising concern (Temu 1991; Tefuarani *et al.* 2002). But, here too, in the PNG context, reality is more complex than many of the models, and it would be dangerously misleading to ascribe a kind of automatism to the health transition process, important as it is. It is by no means clear yet that time and economic development will be sufficient to ensure that lifestyle diseases eventually overtake infections, or even that gains so far in the control of infection and increase of life expectancy are secure for the future.

By reviewing trends in selected aspects of PNG's population health with special focus – mortality, type 2 diabetes, malaria, HIV/AIDS – and touching briefly on many others – including pneumonia, influenza, tuberculosis, measles, smallpox, sexually transmitted infections, yaws, diarrhoea, dysentery, pigbel, arboviruses, filariasis, maternal mortality, heart disease, asthma, arthritis, trauma – I have aimed to illustrate empirically some of PNG's many simultaneous and successive health trends, too complex to be drawn into a single overarching model unless that itself is very complex. Some observed trends bear out the expected declines of early-acting infections, increases in life expectancy and rises of late-acting non-infectious disease. Others, however, follow their own dynamic: malnutrition, diarrhoea and even measles continue to cause uncontrolled problems; tuberculosis is increasing; and most saliently, the emergence of HIV/AIDS threatens a large increase in the infectious death toll and even a reduction in life expectancy; while malaria shows no sign of conforming to health transition expectations, rather developing in increasingly dangerous ways, with a rise in the proportion due to *P. falciparum*, increasing drug resistance in *P. falciparum*, territorial expansion across the highlands, and the possibility of aggravation under climate change.

The PNG Ministry of Health (2001: 3) itself acknowledges in its current National Health Plan that 'The poor status of the health of the people, in particular rural women and children, is not improving.' The Ministry has numerous proposals to address this situation, though as it documents, there are funding and organizational hurdles to overcome in implementing them. It is beyond the scope of this chapter to review such proposals. Perhaps the strongest avenue in conventional public health terms would lie in redoubled immunization programmes – the improved coverage of vaccines already offered, the implementation of vaccines already feasible and tested, and the development and implementation of ones still at the research stage; of course all this too would present funding problems.

The search for low-cost alternative ways to improve population health is not easy. Maybe one possibility lies in regulation: PNG has a history of

innovation in public health regulation, as witnessed by its ban on the advertisement of baby feeding bottles and breast-milk substitutes, and requirement of a prescription for the sale of feeding bottles (Biddulph 1983). Perhaps it is time for another imaginative step in this direction – if the right step can be found. Any suggestion to follow Bhutan's lead by banning the sale of tobacco products and public smoking (BBC News 2004) may attract scepticism in PNG for some time to come, despite the near universal prevalence of cough.

Since some of the problems are social – for example in the integration of health services into communities – maybe some social solutions, beyond purely technological and public health ones, are needed too. Perhaps a hint in this direction is given by Macintyre's (2004) case study from the Lihir islands, New Ireland Province. Contrary to the argument sometimes made that traditional ideas and cultural practices tend to be at odds with modern medical services and impede their uptake, she found, encouragingly, that women there have embraced new practices in relation to medical treatment and hospital birth where they saw benefits in them, and have experienced dramatic health benefits in consequence.

To conclude, then, despite such encouraging signs as those provided by Macintyre's study, there remain major population health problems for PNG. On the one hand, for a complex of environmental, biological, economic and sociocultural reasons, communicable diseases still play an extremely strong part in PNG's health profile, and the immediate outlook for most Papua New Guineans is not currently one dominated by the expected 'diseases of modernization'. On the other hand, the rise of those mainly non-infectious diseases and their risk factors, especially in the urban centres and the areas most influenced by them, is well attested and substantial. There is thus, as my title suggests, a kind of double jeopardy now at work – as for indigenous minorities in developed countries (Kunitz 1994), though with differences of scale and pattern – whereby both old causes and new, infectious and non-infectious, diminish the prospects of Papua New Guineans for long and healthy lives.

Acknowledgments

I thank Professor Ryutaro Ohtsuka for his invitation to participate in the Tokyo International Union of Anthropological and Ethnographic Sciences meeting on which this volume is based; Professor Tai-Ichiro Takemoto for his invitation to give a subsequent presentation at Nagasaki University Medical School; audiences in Tokyo and Nagasaki

for their comments and questions; Drs Michael Alpers, Robin Hide, Ivo Müller and Stanley Uliaszek for their critical readings of part or all of this chapter in draft; Drs Nigel Beebe, Colin Filer, Deirdre Joy, Judith Littleton, Christine McMurray, Kim Streatfield and Zhongwei Zhao for drawing my attention to particular points and sources; and the editors of this volume for their assistance and forbearance. All remaining errors and omissions are my own responsibility.

References

- Adels, B. R. and Gajdusek, D. C. (1963). Survey of measles patterns in New Guinea, Micronesia and Australia. *American Journal of Hygiene* **77**, 317–43.
- Allen, B. J. (1983). A bomb or a bullet or the bloody flux? Population change in the Aitape inland, Papua New Guinea, 1941–1945. *Journal of Pacific History* **18**, 218–35.
- (1989). Infection, innovation and residence: illness and misfortune in the Torricelli foothills from 1800. In *A Continuing Trial of Treatment: Medical Pluralism in Papua New Guinea*, ed. S. Frankel and G. Lewis. Dordrecht: Kluwer, pp. 35–68.
- (1992). The geography of Papua New Guinea. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 36–66.
- Allen, S. J., O'Donnell, A., Alexander, N. D. E., *et al.* (1997). α + -thalassemia protects children against disease caused by other infections as well as malaria. *Proceedings of the National Academy of Sciences of the USA* **94**, 14736–41.
- Alpers, M. P. (1992). Kuru. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 313–34.
- Alpers, M. P. and Attenborough, R. D. (1992). Human biology in a small cosmos. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 1–35.
- Anderson, H. R. and Woolcock, A. J. (1992). Chronic lung disease and asthma in Papua New Guinea. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 289–301.
- Attenborough, R. D. (2002). Ecology, homeostasis and survival in human population dynamics. In *Human Population Dynamics: Cross-Disciplinary Perspectives*, ed. H. Macbeth and P. Collinson. Cambridge: Cambridge University Press, pp. 186–208.
- Attenborough, R. D. and Alpers, M. P. (1995). Change and variability in Papua New Guinea's patterns of disease. In *Human Populations: Diversity and Adaptation*, ed. A. J. Boyce and V. Reynolds. Oxford: Oxford University Press, pp. 189–216.

- Attenborough, R. D., Gardner, D. S. and Gibson, F. D. (1996). *Malaria and Filariasis amongst the Mianmin of the Highland Fringes of Sandaun Province, Papua New Guinea: A Report*. Canberra: Department of Archaeology and Anthropology, Australian National University.
- Attenborough, R. D., Burkot, T. R. and Gardner, D. S. (1997). Altitude and the risk of bites from mosquitoes infected with malaria and filariasis among the Mianmin people of Papua New Guinea. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **91**, 8–10.
- Ballard, C., Brown, P., Bourke, R. M. and Harwood, T. (eds.) (2005). *The Sweet Potato in Oceania: A Reappraisal*. Pittsburgh & Sydney: Ethnology Monographs and Oceania Monographs.
- Barnett, J. (2002). Resilience and adaptation. In *Asia-Pacific Network Workshop on Ethnographic Perspectives on Resilience to Climate Variability in Pacific Island Countries*, ed. J. Barnett and M. Busse. Christchurch, New Zealand: Macmillan Brown Centre for Pacific Studies, University of Canterbury, pp. 5–11.
- BBC News (2004). Bhutan forbids all tobacco sales. BBC. http://news.bbc.co.uk/2/hi/south_asia/4012639.stm (accessed December 2005).
- Beebe, N. W. and Cooper, R. D. (2002). Distribution and evolution of the *Anopheles punctulatus* group (Diptera: Culicidae) in Australia and Papua New Guinea. *International Journal for Parasitology* **32**, 563–74.
- Benjamin, A. L. and Dramoi, V. (2002). Outbreak of measles in the National Capital District, Papua New Guinea in 2001. *Papua New Guinea Medical Journal* **45**, 178–84.
- Biddulph, J. (1983). Legislation to protect breastfeeding. *Papua New Guinea Medical Journal* **26**, 9–12.
- Bockarie, M. J., Alexander, N., Bockarie, F., *et al.* (1996). The late biting habit of parous *Anopheles* mosquitoes and pre-bedtime exposure of humans to infective female mosquitoes. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **90**, 23–5.
- Boyden, S. (2004). *The Biology of Civilisation: Understanding Human Culture as a Force in Nature*. Sydney: University of New South Wales Press.
- Brinkhoff, T. (2003). City population. <http://www.citypopulation.de> (accessed 25 November 2005).
- Brown, G. V. and Reeder, J. C. (2002). Malaria vaccines. *Medical Journal of Australia* **177**, 230–1.
- Bruce, M. C. and Day, K. P. (2003). Cross-species regulation of *Plasmodium* parasitemia in semi-immune children from Papua New Guinea. *Trends in Parasitology* **19**, 271–7.
- Burkot, T. R., Graves, P. M., Paru, R., Wirtz, R. A. and Heywood, P. F. (1988). Human malaria transmission studies in the *Anopheles punctulatus* complex: sporozoite rates, inoculation rates and sporozoite densities. *American Journal of Tropical Medicine and Hygiene* **39**, 135–44.
- Burton, J. (1983). A dysentery epidemic in New Guinea and its mortality. *Journal of Pacific History* **18**, 236–61.

- Burton-Bradley, B. G. (ed.) (1990). *A History of Medicine in Papua New Guinea: Vignettes of an Earlier Period*. Kingsgrove, New South Wales: Australasian Medical Publishing Company.
- Caldwell, J. C. (1986). Routes to low mortality in poor countries. *Population and Development Review* **12**, 171–220.
- Caldwell, J. C., Gajanayake, I., Caldwell, P. and Peiris, I. (1989). Sensitization to illness and the risk of death: an explanation for Sri Lanka's approach to good health for all. *Social Science and Medicine* **28**, 365–79.
- Caldwell, J. C., Findley, S., Caldwell, P., et al. (eds.) (1990). *What We Know about Health Transition: The Cultural, Social and Behavioural Determinants of Health*. Canberra: Health Transition Centre, Australian National University.
- Carrad, E. V. (1987). *Review of Disease Patterns in Papua New Guinea*. Port Moresby: University of Papua New Guinea Press.
- Cattani, J. A. (1992). The epidemiology of malaria in Papua New Guinea. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 302–12.
- Cattani, J. A., Moir, J. S., Gibson, F. D., et al. (1986a). Small-area variations in the epidemiology of malaria in Madang Province. *Papua New Guinea Medical Journal* **29**, 11–17.
- Cattani, J. A., Tulloch, J. L., Vrbova, H., et al. (1986b). The epidemiology of malaria in a population surrounding Madang, Papua New Guinea. *American Journal of Tropical Medicine and Hygiene* **35**, 3–15.
- Centre for International Economics (2002). *Potential Economic Impacts of an HIV/AIDS Epidemic in Papua New Guinea*. Canberra: AusAID.
- Clark, J. T. and Kelly, K. M. (1993). Human genetics, paleoenvironments and malaria: relationships and implications for the settlement of Oceania. *American Anthropologist* **95**, 612–30.
- Connell, J. (1997). Health in Papua New Guinea: a decline in development. *Australian Geographical Studies* **35**, 271–93.
- Conyers, R. A. J. (1971). Myocardial infarction in a New Guinean. *Medical Journal of Australia* **ii**, 412–17.
- Cortés, A., Benet, A., Cooke, B. M., Barnwell, J. W. and Reeder, J. C. (2004). Ability of *Plasmodium falciparum* to invade Southeast Asian ovalocytes varies between parasite lines. *Blood* **104**, 2961–6.
- Crane, G. G. (1986). Recent studies of hyperreactive malarious splenomegaly (tropical splenomegaly syndrome) in Papua New Guinea. *Papua New Guinea Medical Journal* **29**, 35–40.
- Curry, C., Bunungam, P., Annerud, C. and Babona, D. (2005). HIV antibody seroprevalence in the emergency department at Port Moresby General Hospital, Papua New Guinea. *Emergency Medicine Australasia* **17**, 359–62.
- Davies, M. (2002). *Public Health and Colonialism: The Case of German New Guinea 1884–1914*. Wiesbaden: Harrassowitz Verlag.
- Dengler, L. and Preuss, J. (2003). Mitigation lessons from the July 17, 1998 Papua New Guinea tsunami. *Pure and Applied Geophysics* **160**, 2001–31.
- Denham, T. P., Haberle, S. G., Lentfer, C., et al. (2003). Origins of agriculture at Kuk swamp in the highlands of New Guinea. *Science* **301**, 189–93.
- Denoon, D., Dugan, K. and Marshall, L. (1989). *Public Health in Papua New Guinea: Medical Possibility and Social Constraint, 1884–1984*. Cambridge: Cambridge University Press.
- Desowitz, R. S. and Spark, R. A. (1987). Malaria in the Maprik area of the Sepik region, Papua New Guinea: 1957–1984. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **81**, 175–6.
- Dowse, G. K., Spark, R. A., Mavo, B., et al. (1994). Extraordinary prevalence of non-insulin-dependent diabetes mellitus and bimodal plasma glucose distribution in the Wanigela people of Papua New Guinea. *Medical Journal of Australia* **160**, 767–74.
- Duke, T. (2003). The crisis of measles and the need to expand the ways of delivering vaccines in Papua New Guinea. *Papua New Guinea Medical Journal* **46**, 1–7.
- Edwards, K. N. (1994). Rural health service crisis in Papua New Guinea: causes, implications and possible solutions. *Papua New Guinea Medical Journal* **37**, 145–51.
- Ewers, W. H. (1972). Malaria in the early years of German New Guinea. *Journal of the Papua and New Guinea Society* **6**, 3–30.
- Fenner, F., Henderson, D. A., Arita, I., Jezek, Z. and Ladnyi, I. D. (1988). *Smallpox and Its Eradication*. Geneva: World Health Organization.
- Fix, A. G. (2003). Simulating hemoglobin history. *Human Biology* **75**, 607–18.
- Foley, W. A. (1992). Language and identity in Papua New Guinea. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 136–49.
- Forge, A. (1972). Normative factors in the settlement size of neolithic cultivators (New Guinea). In *Man, Settlement and Urbanism*, ed. P. J. Ucko, R. Tringham and G. W. Dimbleby. London: Duckworth, pp. 363–76.
- (1990). The power of culture and the culture of power. In *Sepik Heritage: Tradition and Change in Papua New Guinea*, ed. N. Lutkehaus, C. Kaufmann, W. E. Mitchell, L. Osmundsen and M. Schister. Durham, North Carolina: Carolina Academic Press, pp. 160–70.
- Froment, A. (2001). Evolutionary biology and health of hunter-gatherer populations. In *Hunter-Gatherers: An Interdisciplinary Perspective*, ed. C. Panter-Brick, R. H. Layton and P. Rowley-Conwy. Cambridge: Cambridge University Press, pp. 239–66.
- Gajdusek, D. C. (1990). Raymond Pearl Memorial Lecture, 1989: Cultural practices as determinants of clinical pathology and epidemiology of venereal infections: implications for predictions about the AIDS epidemic. *American Journal of Human Biology* **2**, 347–51.
- Garner, P. A., Talwat, E. N., Hill, G., Reid, M. S. and Garner, M. F. (1986). Yaws reappears. *Papua New Guinea Medical Journal* **29**, 247–52.
- Genton, B., Al-Yaman, F., Mgone, C. S., et al. (1995a). Ovalocytosis and cerebral malaria. *Nature* **378**, 564–5.
- Genton, B., Al-Yaman, F., Beck, H.-P., et al. (1995b). The epidemiology of malaria in the Wosera area, East Sepik Province, Papua New Guinea, in preparation

- for vaccine trials. I. Malariometric indices and immunity. *Annals of Tropical Medicine and Parasitology* **89**, 359–76.
- Genton, B., Anders, R., Alpers, M. P. and Reeder, J. C. (2003). The malaria vaccine development program in Papua New Guinea. *Trends in Parasitology* **19**, 264–70.
- Greenburg-Hudd, S. (1998). *Dental palaeopathology and subsistence in protohistoric and historic skeletal remains from Papua New Guinea*. M.Litt. thesis, Archaeology & Anthropology, Australian National University, Canberra.
- Groube, L. M. (1993). Contradictions and malaria in Melanesian and Australian prehistory. In *A Community of Culture: The People and Prehistory of the Pacific*, ed. M. Spriggs, D. E. Yen, W. Ambrose, R. Jones, A. Thorne and A. Andrews. Canberra: Department of Prehistory, Research School of Pacific Studies, Australian National University, pp. 164–86.
- (2000). *Plasmodium falciparum*: the African genesis. In *Australian Archaeologist: Collected Papers in Honour of Jim Allen*, ed. A. Anderson and T. Murray. Canberra: Coombs Academic Publishing, Australian National University, pp. 131–44.
- Groube, L. M., Chappell, J., Muke, J. and Price, D. (1986). A 40 000-year-old occupation site at Huon Peninsula, Papua New Guinea. *Nature* **324**, 453–5.
- Hanson, L. W., Allen, B. J., Bourke, R. M. and McCarthy, T. J. (2001). *Papua New Guinea Rural Development Handbook*. Canberra: The Australian National University.
- Heywood, P. F. and Hide, R. L. (1994). Nutritional effects of export-crop production in Papua New Guinea: a review of the evidence. *Food and Nutrition Bulletin* **15**, 233–49.
- Hide, R. (2003). *Pig Husbandry in New Guinea: A Literature Review and Bibliography*. Canberra: Australian Centre for International Agricultural Research.
- Hodge, A. M., Dowse, G. K., Erasmus, R. T., et al. (1996). Serum lipids and modernization in coastal and highland Papua New Guinea. *American Journal of Epidemiology* **144**, 1129–42.
- Hope, G. S. and Haberle, S. (2005). The history of the human landscapes of New Guinea. In *Papuan Pasts: Cultural, Linguistic and Biological Histories of Papuan-Speaking Peoples*, ed. A. K. Pawley, R. D. Attenborough, J. Golson and R. L. Hide. Canberra: Pacific Linguistics, The Australian National University, pp. 541–54.
- Hope, G. S., Golson, J. and Allen, J. (1983). Palaeoecology and prehistory in New Guinea. *Journal of Human Evolution* **12**, 37–60.
- Hughes, J. (1997). A history of sexually transmitted diseases in Papua New Guinea. In *Sex, Disease and Society: A Comparative History of Sexually Transmitted Diseases and HIV/AIDS in Asia and the Pacific*, ed. M. Lewis, S. Bamber and M. Waugh. Westport, CT: Greenwood Press, pp. 231–47.
- Hume, J. C. C., Lyons, E. J. and Day, K. P. (2003). Malaria in antiquity: a genetics perspective. *World Archaeology* **35**, 180–92.

- Inaoka, T. (1995). Health and survival in modernizing Papua New Guinea societies. *Anthropological Science* **103**, 339–47.
- (n.d.). The modernization of Papua New Guinea and the changes in disease patterns – recent health transition and emerging/re-emerging infectious diseases. *JCAS Area Studies Research Reports* **2**, 201–17 [In Japanese: transl. A. Callaway].
- Joy, D. A., Feng, X., Mu, J., et al. (2003). Early origin and recent expansion of *Plasmodium falciparum*. *Science* **300**, 318–21.
- Kasehagen, L., Müller, I., McNamara, D. T. et al. (2006). Changing patterns of *Plasmodium* blood stage infections in the Wosera region of Papua New Guinea as monitored by light microscopy and high through-put PCR. *American Journal of Tropical Medicine & Hygiene* **67**, 588–96.
- Kazura, J. W., and Bockarie, M. J. (2003). Lymphatic filariasis in Papua New Guinea: interdisciplinary research on a national health problem. *Trends in Parasitology* **19**, 260–3.
- Kevau, I. H. (1990). Clinical documentation of twenty cases of acute myocardial infarction in Papua New Guineans. *Papua New Guinea Medical Journal* **33**, 275–80.
- King, H. and Collins, A. M. (1989). A modernity score for individuals in Melanesian society. *Papua New Guinea Medical Journal* **32**, 11–22.
- King, H., Finch, C., Collins, A. M., et al. (1989). Glucose tolerance in Papua New Guinea: ethnic differences, association with environmental and behavioural factors and the possible emergence of glucose intolerance in a highland community. *Medical Journal of Australia* **151**, 204–10.
- Kirk, R. L. (1992). Population origins in Papua New Guinea – a human biological overview. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 172–97.
- Kovats, R. S., Campbell-Lendrum, D. H., McMichael, A. J., Woodward, A. and Cox, J. S. (2001). Early effects of climate change: do they include changes in vector-borne disease? *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences* **356**, 1057–68.
- Kunitz, S. J. (1994). *Disease and Social Diversity: The European Impact on the Health of Non-Europeans*. New York: Oxford University Press.
- Lavu, E. K., Oswyn, G. and Vince, J. D. (2002). Sick cell/ β +thalassaemia in a Papua New Guinean: the first reported case of the sickle gene in Papua New Guinea. *Medical Journal of Australia* **176**, 70–1.
- Lawrence, G. (1992). Pigbel. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 335–44.
- Leclerc, M. C., Durand, P., Gauthier, C., et al. (2004). Meager genetic variability of the human malaria agent *Plasmodium vivax*. *Proceedings of the National Academy of Sciences of the USA* **101**, 14455–60.
- Lehmann, D. (2002). Demography and causes of death among the Huli in the Tari basin. *Papua New Guinea Medical Journal* **45**, 51–62.
- Lemeki, M. (2003). *Sexually transmitted diseases and HIV/AIDS: knowledge and sexual behaviour among Eastern Highlands youth, Papua New Guinea*. MA thesis, Australian National University, Canberra.

- Lilley, I. (1992). Papua New Guinea's human past: the evidence of archaeology. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 150–71.
- Livingstone, F. B. (1989). Simulation of the diffusion of the β -globin variants in the Old World. *Human Biology* **61**, 297–309.
- Lourie, J. A. (1986). Trends in birthweight over 43 years at Kwato, Milne Bay Province. *Papua New Guinea Medical Journal* **29**, 337–43.
- (ed.) (1987). *Ok Tedi Health and Nutrition Project Papua New Guinea 1982–1986: Final Report*. Port Moresby and Tabubil: University of Papua New Guinea and Ok Tedi Mining Limited.
- Lourie, J. A., Budd, G. and Anderson, H. R. (1992). Physiological adaptability in Papua New Guinea. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 268–80.
- Macintyre, M. (2004). 'Thoroughly modern mothers': maternal aspirations and declining mortality on the Lihir islands, Papua. *Health Sociology Review* **13**, 43–53.
- Maitland, K., Williams, A. I., Bennett, N. M., *et al.* (1996). The interaction *Plasmodium falciparum* and *P. vivax* in children on Espiritu Santo island, Vanuatu. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **90**, 614–20.
- Manning, L. A. and Ogle, G. D. (2002). Yaws in the periurban settlements of Port Moresby, Papua New Guinea. *Papua New Guinea Medical Journal* **45**, 206–12.
- Mayxay, M., Pukrittayakamee, S., Newton, P. N. and White, N. J. (2004). Mixed-species malaria infections in humans. *Trends in Parasitology* **20**, 233–40.
- McBride, W. J. (2005). HIV/AIDS in Papua New Guinea: an unfolding disaster? *Emergency Medicine Australasia* **17**, 304–6.
- McMurray, C. (2004). *Morbidity and Mortality Patterns in the Pacific*. United Nations Economic and Social Commission for Asia and the Pacific. <http://www.unescap.org/esid/psis/population/popseries/apss163/index.asp> (accessed 12 September 2005).
- Mgone, C. S., Koki, G., Paniu, M. M., *et al.* (1996). Occurrence of the erythrocyte band 3 (AE1) gene deletion in relation to malaria endemicity in Papua New Guinea. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **90**, 228–31.
- Mikloucho-Maclay, N. N. (1975). *New Guinea Diaries, 1871–1883*. Madang (PNG): Kristen Pres.
- Ministry of Health (2001). *National Health Plan 2001–2010: Health Vision 2010*. Port Moresby: Ministry of Health, Independent State of Papua New Guinea.
- Misch, K. A. (1988). Ischaemic heart disease in urbanized Papua New Guinea: an autopsy study. *Cardiology* **75**, 71–5.
- Müller, I., Vounatsu, P., Allen, B. J. and Smith, T. (2001). Spatial patterns of child growth in Papua New Guinea and their relation to environment, diet, socio-economic status and subsistence activities. *Annals of Human Biology* **28**, 263–80.
- Müller, I., Betuela, I. and Hide, R. L. (2002a). Regional patterns of birthweights in Papua New Guinea in relation to diet, environment and socio-economic factors. *Annals of Human Biology* **29**, 74–88.
- Müller, I., Taime, J., Ibam, E., *et al.* (2002b). Complex patterns of malaria epidemiology in the highlands region of Papua New Guinea. *Papua New Guinea Medical Journal* **45**, 200–5.
- Müller, I., Bockarie, M., Alpers, M. P. and Smith, T. (2003). The epidemiology of malaria in Papua New Guinea. *Trends in Parasitology* **19**, 253–9.
- Müller, I., Tulloch, J., Marfurt, J., Hide, R. and Reeder, J. C. (2005). Malaria control in Papua New Guinea results in complex epidemiological changes. *Papua New Guinea Medical Journal*, **48**, 151–7.
- Naraqi, S., Feling, B. and Leeder, S. R. (2003). Disease and death in Papua New Guinea. *Medical Journal of Australia* **178**, 7–8.
- National AIDS Council Secretariat and Department of Health (2002). *HIV/AIDS Quarterly Report: March 2002*. Port Moresby: National AIDS Council Secretariat.
- National Statistical Office (2002). *Papua New Guinea 2000 Census: Final Figures*. Port Moresby: National Statistical Office of Papua New Guinea.
- (2003a). *Papua New Guinea 2000 Census: National Report*. Port Moresby: National Statistical Office of Papua New Guinea.
- (2003b). *Recent Fertility and Mortality Indices and Trends in Papua New Guinea: A Report Based on the Analysis of 2000 Census Data*. Port Moresby: National Statistical Office of Papua New Guinea.
- Natsuhara, K., Inaoka, T., Umezaki, M., *et al.* (2000). Cardiovascular risk factors of migrants in Port Moresby from the highlands and island villages, Papua New Guinea. *American Journal of Human Biology* **12**, 655–64.
- Norgan, N. G. (1997). Human adaptability studies in Papua New Guinea: original aims, successes and failures. In *Human Adaptability: Past, Present and Future*, ed. S. J. Ulijaszek and R. Huss-Ashmore. Oxford: Oxford University Press, pp. 102–25.
- Ogle, G. D. (2001). Type 2 diabetes mellitus in Papua New Guinea – an historical perspective. *Papua New Guinea Medical Journal* **44**, 81–7.
- Ohtsuka, R. (1986). Low rate of population increase of the Gidra Papuans in the past: a genealogical-demographic analysis. *American Journal of Physical Anthropology* **71**, 13–23.
- Owen, I. L. (2005). Parasitic zoonoses in Papua New Guinea. *Journal of Helminthology* **79**, 1–14.
- Pawley, A. K., Attenborough, R. D., Golson, J. and Hide, R. L. (eds.) (2005). *Papuan Pasts: Cultural, Linguistic and Biological Histories of Papuan-Speaking Peoples*. Canberra: Pacific Linguistics, The Australian National University.
- Peters, W. (1960). Studies on the epidemiology of malaria in New Guinea. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **54**, 242–60.
- Poka, H. and Duke, T. (2003). In search of pigbel: gone or just forgotten in the highlands of Papua New Guinea? *Papua New Guinea Medical Journal* **46**, 135–42.
- Pollard, T. M. (1997). Environmental change and cardiovascular disease: a new complexity. *Yearbook of Physical Anthropology* **40**, 1–24.
- Radford, A. J., van Leeuwen, H. and Christian, S. H. (1976). Social aspects in the changing epidemiology of malaria in the highlands of New Guinea. *Annals of Tropical Medicine and Parasitology* **70**, 11–23.

- Riley, I. D. (1983). Population change and distribution in Papua New Guinea: an epidemiological approach. *Journal of Human Evolution* **12**, 125–32.
- (2002). Pneumonia vaccine trials at Tari. *Papua New Guinea Medical Journal* **45**, 44–50.
- Riley, I. D. and Lehmann, D. (1992). The demography of Papua New Guinea: migration, fertility and mortality patterns. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 67–92.
- Riley, I. D., Lehmann, D. and Alpers, M. P. (1992). Acute respiratory infections. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 281–8.
- Riley, J. C. (2001). *Rising Life Expectancy: A Global History*. Cambridge: Cambridge University Press.
- Secretariat of the Pacific Community (2004). *Pacific Island Populations 2004*. Secretariat of the Pacific Community (Demography/Population Programme). <http://www.spc.int/demog/> (accessed 12 September 2005).
- Serjeantson, S. W., Board, P. G. and Bhatia, K. K. (1992). Population genetics in Papua New Guinea: a perspective on human evolution. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 198–233.
- Sharp, P. T. (1982). Highlands malaria: malaria in Enga Province of Papua New Guinea. *Papua New Guinea Medical Journal* **25**, 253–60.
- Sinnett, P. F. (1977). *The People of Murapin*. Faringdon: E. W. Classey.
- Sinnett, P. F., Kevau, I. H. and Tyson, D. (1992). Social change and the emergence of degenerative cardiovascular disease in Papua New Guinea. In *Human Biology in Papua New Guinea: The Small Cosmos*, ed. R. D. Attenborough and M. P. Alpers. Oxford: Clarendon Press, pp. 373–86.
- Smith, T., Genton, B., Baea, K., et al. (2001). Prospective risk of morbidity in relation to malaria infection in an area of high endemicity of multiple species of *Plasmodium*. *American Journal of Tropical Medicine and Hygiene* **64**, 262–7.
- Specht, J. (2005). Revisiting the Bismarcks: some alternative views. In *Papuan Past: Cultural, Linguistic and Biological Histories of Papuan-Speaking Peoples*, ed. A. K. Pawley, R. D. Attenborough, J. Golson and R. L. Hide. Canberra: Pacific Linguistics, The Australian National University, pp. 235–88.
- Spencer, M. (1999). *Public Health in Papua New Guinea 1870–1939*. Brisbane: Australian Centre for International and Tropical Health and Nutrition.
- Spencer, T. E. T., Spencer, M. and Venters, D. (1974). Malaria vectors in Papua New Guinea. *Papua New Guinea Medical Journal* **17**, 22–30.
- Spriggs, M. (1997). *The Island Melanesians*. Oxford: Blackwell.
- Swadling, P. and Hide, R. L. (2005). Changing landscape and social interaction: looking at agricultural history from a Sepik-Ramu perspective. In *Papuan Past: Cultural, Linguistic and Biological Histories of Papuan-Speaking Peoples*, ed. A. K. Pawley, R. D. Attenborough, J. Golson and R. L. Hide. Canberra: Pacific Linguistics, The Australian National University, pp. 289–327.
- Swadling, P. and Hope, G. (1992). Environmental change in New Guinea since human settlement. In *The Naive Lands: Prehistory and Environmental Change in Australia and the Southwest Pacific*, ed. J. Dodson. Melbourne: Longman Cheshire, pp. 13–42.
- Tefuarani, N., Sleigh, A. and Hawker, R. (2002). Congenital heart diseases: a future burden for Papua New Guinea. *Papua New Guinea Medical Journal* **45**, 175–7.
- Teitelbaum, M. S. (1975). Relevance of demographic transition theory for developing countries. *Science* **188**, 420–5.
- Temu, P. I. (1991). Adult medicine and the 'new killer diseases' in Papua New Guinea: an urgent need for prevention. *Papua New Guinea Medical Journal* **34**, 1–5.
- Townsend, P. K. (1985). Infant mortality in the Saniyo-Hiowe population, Ambunti District, East Sepik Province. *Papua New Guinea Medical Journal* **28**, 177–82.
- Tracer, D. P., Sturt, R. J., Sturt, A. and Braithwaite, L. M. (1998). Two decade trends in birth weight and early childhood growth in Papua New Guinea. *American Journal of Human Biology* **10**, 483–93.
- Trowell, H. C. (1981). Hypertension, obesity, diabetes mellitus and coronary heart disease. In *Western Diseases: Their Emergence and Prevention*, ed. H. C. Trowell and D. P. Burkitt. London: Edward Arnold, pp. 3–32.
- Trowell, H. C. and Burkitt, D. P. (eds.) (1981). *Western Diseases: Their Emergence and Prevention*. London: Edward Arnold.
- Tulu, A. N. (1996). *Determinants of malaria transmission in the highlands of Ethiopia: the impacts of global warming on morbidity and mortality ascribed to malaria*. Ph.D. thesis, University of London.
- Tyson, D. C. (1987). *An ecological analysis of child malnutrition in an Abelam community, Papua New Guinea*. Ph.D. thesis, Australian National University, Canberra.
- Ulijaszek, S. J. (1995). Development, modernisation and health intervention. In *Health Intervention in Less Developed Nations*, ed. S. J. Ulijaszek. Oxford: Oxford University Press, pp. 82–136.
- (1997). Modernization, nutritional status and hypertension in a rural Papua New Guinea population (abstract). *Annals of Human Biology* **24**, 281.
- (2001). Secular trend in birthweight among the Purari delta population, Papua New Guinea. *Annals of Human Biology* **28**, 246–55.
- Ulijaszek, S. J., Koziel, S. and Hermanussen, M. (2005). Village distance from urban centre as the prime modernization variable in differences in blood pressure and body mass index of adults of the Purari delta of the Gulf province, Papua New Guinea. *Annals of Human Biology* **32**, 326–38.
- United Nations Children's Fund (2004a). *The State of the World's Children 2005: Childhood Under Threat*. New York: UNICEF.
- (2004b). *Progress for Children*. New York: UNICEF.
- van de Kaa, D. J. (1971). *The demography of Papua and New Guinea's indigenous population*. PhD thesis, Australian National University, Canberra.
- Vines, A. P. (1970). *An Epidemiological Sample Survey of the Highlands, Mainland and Islands Regions of the Territory of Papua and New Guinea*. Port Moresby: Department of Public Health, Territory of Papua and New Guinea.

- Ward, R. G. and Lea, D. A. M. (eds.) (1970). *An Atlas of Papua and New Guinea*. Port Moresby and Glasgow: University of Papua and New Guinea and Collins – Longman.
- Webb, S. (1995). *Palaeopathology of Aboriginal Australians: Health and Disease across a Hunter-Gatherer Continent*. Cambridge: Cambridge University Press.
- Wiesenfeld, S. L. (1967). Sickle-cell trait in human biological and cultural evolution. *Science* **157**, 1134–40.
- Wigley, S. C. (1977). The first hundred years of tuberculosis in New Guinea. In *The Melanesian Environment*, ed. J. H. Winslow. Canberra: Australian National University Press, pp. 471–84.
- (1991). Tuberculosis and Papua New Guinea: the Australian connection. In *History of Tuberculosis in Australia, New Zealand and Papua New Guinea*, ed. A. J. Proust. Canberra: Brolga Press, pp. 103–17.
- World Health Organization (2000). *The World Health Report 2000 – Health Systems: Improving Performance*. Geneva: World Health Organization.
- World Health Organization Regional Office for the Western Pacific (2005). *Countries and Areas: Papua New Guinea*. World Health Organization. <http://www.wpro.who.int/countries/png/> (accessed 12 September 2005).
- Yen, D. E. (2005). Reflection, refraction and recombination. In *The Sweet Potato in Oceania: A Reappraisal*, ed. C. Ballard, P. Brown, R. M. Bourke and T. Harwood. Pittsburgh and Sydney: Ethnology Monographs and Oceania Monographs, pp. 181–7.
- Zimmerman, P. A., Woolley, I., Masinde, G. L. *et al.* (1999). Emergence of FY*A^{null} in a *Plasmodium vivax* – endemic region of Papua New Guinea. *Proceedings of the National Academy of Sciences of the USA* **96**, 13973–7.
- Zimmerman, P. A., Patel, S. S., Maier, A. G., Bockarie, M. J. and Kazura, J. W. (2003). Erythrocyte polymorphisms and malaria parasite invasion in Papua New Guinea. *Trends in Parasitology* **19**, 250–2.

Index

- ABO blood group 52
- Aboriginal Australians 44–5, 48, 57
- Aboriginal people 247
- accidental trauma 264
- acute infections
- lower respiratory 29
 - viral 265
- adaptation 1, 6, 13, 263, 289
- equilibrium in 263, 289
 - strategies in 51
- adequate prenatal care 213
- adipose tissue 95
- Admiralty Islands 31
- Africa 33, 262
- age-adjusted death rate 195
- age at marriage 239
- agriculture 37, 149, 257
- productivity of 49
 - work in 114
- aid posts 279
- AIDS 287
- alcohol 207
- use of 203
- alpha globin gene 7, 28, 33
- deletions 21, 31, 34
 - haplotypes 33, 34
 - RFLP haplotype 34
 - loci 31
- alpha thalassaemia 7, 30–1, 39, 266
- mutations 38
- ALRI 279
- Alzheimer's disease 197
- American
- Indians 208, 210
 - Samoa 4, 7, 37, 149
- Americas 264
- amniocentesis 212
- AN *see* Austronesian (AN)
- anaemia 7, 72, 75, 210
- anaerobic exercise 96
- anatomically modern
- humans 22
- animal foods 45
- Anopheles* mosquito 284
- anthropometry 103, 134, 226
- characteristics 109
 - measurements 103
 - measures of 88
- Apia 151
- appropriate body size 136
- Arafura
- Plain 46
 - Sea 46
- arboriculture 6, 21
- arboviruses 287
- Arnhem Land 45
- arthritis 264
- Asia-Pacific region 1, 2, 4, 6–8, 12–17, 141, 236, 255
- Asian
- American 12
 - B haplogroup 34–6
 - economic crisis 68
 - financial crisis 68
 - haplotypes 33
 - intrusions 36, 147
- Asian and Pacific Islander (API) 192
- Association of South East Asian Nations (ASEAN) 64
- asthma 14, 261, 278, 289, 290
- AusAID 288
- Australasia 5
- Australia 4, 5, 11–12, 22–3, 95, 127, 246, 261
- aborigines of 13
 - Central 33
 - European settlers in 57, 234
 - Melbourne 219
 - region 5
 - zoogeographic region 47
- Austro-Asiatic family 6
- Austronesian (AN) 25, 31, 44, 101
- language family 31
 - languages 6, 27, 262
 - migrants 47
 - non speakers 6, 38, 44
 - settlers 47
 - speakers 22, 31, 44, 101, 264, 277
 - speaking 50, 51
- B haplogroup 35
- bacterial infection 289
- Balopa 52, 55–6