Research paper

Running title: Cost of *Salmonella* in Australia

The cost of *Salmonella* infections in Australia, 2015

Laura Ford¹, Philip Haywood², Martyn D. Kirk¹, Emily Lancsar³, Deborah A. Williamson⁴,
Kathryn Glass^{1*}

¹National Centre for Epidemiology and Population Health, Research School of Population Health, The Australian National University, Canberra, ACT 2601, Australia

²Centre for Health Economics Research and Evaluation, University of Technology Sydney, PO Box 123, Broadway, NSW 2007, Australia

³Department of Health Services Research and Policy, Research School of Population Health, The Australian National University, Canberra, ACT 2601, Australia

⁴Microbiological Diagnostic Unit Public Health Laboratory, The University of Melbourne at The Peter Doherty Institute for Infection and Immunity, Parkville, VIC 3010, Australia

Keywords: Cost of illness, Incidence, Salmonella Infections, Health Care Costs, Monte Carlo Methods

*Author for correspondence. Tel: +61 2 6125 2468; fax +61 2 6125 0270; E-mail:

Kathryn.glass@anu.edu.au

Abstract

Gastroenteritis caused from infections with *Salmonella enterica* (salmonellosis) causes significant morbidity in Australia. In addition to acute gastroenteritis, approximately 8.8% of people develop irritable bowel syndrome (IBS) and 8.5% of people develop reactive arthritis (ReA). We estimated the economic cost of salmonellosis and associated sequel illnesses in Australia in a typical year circa 2015. We estimated incidence, hospitalizations, other health care usage, absenteeism, and premature mortality for four age groups using a variety of complementary datasets. We calculated direct costs (health care) and indirect costs (lost productivity and premature mortality) using Monte Carlo simulation to estimate 90% Credible Intervals (CrI) around our point estimates. We estimated that 90,833 cases, 4,312 hospitalizations, and 19 deaths occurred from salmonellosis in Australia circa 2015 at a direct cost of AUD 23.8 million (90% CrI 19.3-28.9) and a total cost of AUD 124.4 million (90% CrI 107.4-143.1 million). When IBS and ReA were included, the estimated direct cost was 35.7 million (90% CrI 29.9-42.7) and the total cost was AUD 146.8 million (90% CrI 127.8-167.9 million). Foodborne infections were responsible for AUD 88.9 million (90% CrI 63.9-112.4 million) from acute salmonellosis and AUD 104.8 million (90% CrI 75.5-132.3 million) when IBS and ReA were included. Targeted interventions to prevent illness could considerably reduce costs and societal impact from *Salmonella* infections and sequel illnesses in Australia.

Highlights

- The rate of salmonellosis in Australia is high, but the costs have not previously been assessed.
- Salmonellosis illness and sequelae cost Australia AUD 146.8 million circa 2015.
- Foodborne *Salmonella* infections and sequelae cost AUD 104.8 million.
- Quantifying costs assists with prioritization of public health interventions across the food chain to reduce costs and societal impact.

Acute gastroenteritis from non-typhoidal *Salmonella enterica* (NTS) infection (salmonellosis) causes significant morbidity. Globally, the World Health Organization (WHO) has estimated that NTS is one of the highest burden foodborne pathogens, responsible for approximately 78.7 million illnesses, 59 thousand deaths, and 4 million disability adjusted life years (DALYs) in 2010 (44). In Australia, there were an estimated 262 cases per 100,000 population circa 2010, with 72% of cases transmitted from contaminated food (27, 43). The reported incidence of salmonellosis in Australia has been increasing and is higher than in the US, Canada, the UK, and New Zealand (7, 11, 20, 34, 40).

NTS illness is one of the leading causes of foodborne gastroenteritis-associated hospitalizations and deaths in Australia. In addition to salmonellosis, it has been estimated that approximately 8.8% of people infected with NTS subsequently develop irritable bowel syndrome (IBS) and 8.5% develop reactive arthritis (ReA) (18). These chronic sequelae can be severe and require additional medical tests, treatments, visits to primary care, specialists, or hospitals and time off work.

As the burden of salmonellosis and its sequelae are high, estimated costs of illness are needed to better inform food safety policy and provide inputs for calculating the cost effectiveness of new policies and interventions. Internationally there have been several studies estimating the cost of, or loss of health-related quality of life from, NTS illness, demonstrating that NTS has a substantial economic burden (25, 28, 37, 39). While the overall cost of foodborne disease in Australia circa 2000 was estimated at \$1.2 billion AUD (1), the costs were not assessed at a pathogen level. Given the burden of salmonellosis in Australia, our aim was to provide contemporary estimates of the cost of illness associated with NTS and resulting sequelae (IBS and ReA) in Australia circa 2015.

Materials and Methods

We estimated costs of NTS illness and resulting sequelae illnesses in @Risk (<http://www.palisade.com>) using direct health care costs, productivity loss, and the value of premature mortality for acute and ongoing illness. The estimated direct and indirect costs were based on the incidence of NTS, IBS, and ReA illness and ongoing illness from IBS and ReA in a typical year circa 2015. We estimated incidence and costs by four age groups: 0-4 years, 5-19 years, 20-64 years, and 65+ years. Data sources and model structure are described below with additional information in supplementary material 1 and 2.

Data sources for incidence burden estimates. We used data from the National Notifiable Disease Surveillance System (NNDSS) by sex and 5-year age groups from 2013-2015 (11). Notifications where age and/or sex were unknown were excluded (0.16%). Denominator data were based on the Australian population provided by the Australian Bureau of Statistics for that year (4). As IBS and ReA are rare in children <5 years old (22, 38), we used the ratio of incidence to hospitalizations for IBS and ReA in those aged 5-64 to estimate incidence in children aged <5 years.

Data sources for health care usage burden estimates. For NTS, we used data from the 2008-2009 National Gastroenteritis Survey II (NGSII), a retrospective cross-sectional survey of the population burden of infectious gastroenteritis in that year (8), to estimate the proportion of incident cases that consulted a general practitioner (GP), visited the emergency department (ED), or took medication (antibiotics, diarrhea relief, pain relief, nausea relief, cramp relief). We weighted each of these proportions by the severity of diarrhea, with 1-2 days of diarrhea classified as mild disease, 3-4 days as moderate, and 5 or more days as severe (supplementary material 1 & 2), and assumed that 3% of NTS cases were mild, 14% were moderate and 83% were severe as in Hall et al (21). The number of stool microscopy,

culture and sensitivity (MCS), and/or stool polymerase chain reaction (PCR) tests were estimated from the median number of notifications by age group in the National Disease Surveillance System (NNDSS) (11) from 2013-2015, all of which were confirmed by laboratory testing (supplementary material 1). For IBS, and ReA, we estimated the proportion of cases that see a GP, visit a specialist, receive treatment, or undergo testing from the literature or from expert clinical opinion (see supplementary material 1 & 2 for more information). We made an a priori assumption there would be no ED visits for IBS or ReA following *Salmonella* infection.

We used hospital separation statistics and average length of stay data by principal diagnoses for the financial years 2011-12, 2012-13, and 2013-14 from the Australian Institute of Health and Welfare (AIHW) to estimate NTS, IBS and ReA hospitalizations (5, 6). Diagnostic codes were based on the Australian modification of the 10th International Classification of Diseases and are detailed in Technical Appendix 3 of Kirk et al (27) and Technical Appendix 4 of Ford et al (18). As principal diagnosis data only is available online from AIHW, we imputed the number of additional diagnoses for NTS, IBS, and ReA based on Kirk et al and Ford et al (18, 27).

Data sources for lost productivity burden estimates. Days of lost paid work for illness and caring were estimated from the NGSII for NTS. Using the same approach taken for GP and ED visits, we adjusted data on the number of reported days of paid work missed due to gastroenteritis illness or caring for someone with a gastroenteritis illness for the severity of diarrhea and by age group. We followed Abelson et al (1) in estimating days of lost paid work as days in hospital plus 0.5 days for visiting the GP for IBS and data from Townes et al (42) to estimate a distribution of 3.37 days (90% CrI 2.59-4.3) for ReA. Confidence intervals for days of lost paid work due to NTS, IBS, and ReA illness were generated using cii means with a Poisson distribution in Stata SE 14.2.

Data sources for premature mortality burden estimates. We used data from the Australian Bureau of Statistics on underlying or contributing causes of death in males and females aged 0-14, 15-64, and 65+ from 2001-2010. The diagnostic code used was A02, based on the 10th International Classification of Diseases as detailed in Technical Appendix 3 of Kirk et al (27). We estimated the number of deaths due to NTS infection for our age groups using a statistical model with multipliers for underreporting, domestic acquisition, and foodborne proportion, as in Kirk et al (27). We assumed no premature mortality due IBS or ReA.

Data sources for ongoing illness burden estimates. We used the literature and built on previous work (1), which was reviewed and updated by an expert clinician, to estimate the proportion of cases that have ongoing illness from incident IBS or ReA following *Salmonella* infection in a typical year circa 2015. For IBS, we estimated that 42.9% (95% CI 21.8%-66%) of cases would have continuing symptoms at 12 months (30). For ReA, we estimated that 50% (95% CI 23%-77%) would have continuing symptoms at 3 months (22, 29).

Data sources for health care usage cost estimates. We assumed costs for GP and specialist visits based on the Medicare Benefits Schedule (MBS) (9). We used an estimate from 2009/10 in New South Wales for ED Triage costs (35) and hospital cost data were extracted as Australian Refined Diagnosis Related Group (AR-DRG) costs published by the Independent Hospital Pricing Authority for 2013/14 (26). As pathology costs are generally paid for through national health insurance, we calculated the costs by divided total benefits (monetary value spent) by services (number billed) from Medicare item reports (13) for MBS (9) for relevant item numbers. Costs were extracted from the Pharmaceutical Benefits Schedule (PBS) by dividing total benefits by services (pharmaceuticals) from PBS item reports (14) for PBS item numbers (10) for the calendar year 2015. If the medication or treatment was not listed on the PBS or may commonly be bought over the counter, we

extracted costs from an Australian pharmaceutical store

(<https://www.chemistwarehouse.com.au>). See supplementary material 2 for more details.

Data sources for lost productivity cost estimates. We used a human capital approach to calculate the cost of lost productivity (33) and tested lost productivity in sensitivity analysis. We used the participation rate in the workforce by age group for June 2015 extracted from the Australian Bureau of Statistics (79.2% for 20-64 years and 12.2% for 65 years and older) (2) and multiplied this by the estimated days of lost paid work and by the daily wage rate. As the ABS provided a participation rate for 15-19 year olds (52.6%), we weighted this across our 5-19 years age group to produce a participation rate of 17.5%. We used the participation rate for 20-64 year olds (79.2%) for days of paid work missed by carers reported in the NGSII for the age groups 0-4 years, 5-19 years, and 20-64 years and the participation rate for 65 years and older (12.2%) for the age group 65+ years. To calculate the daily wage rate, we divided the average weekly earnings for May 2015 extracted from the Australian Bureau of Statistics (3) by five to estimate a daily productivity cost of \$227.38.

Data sources for premature mortality cost estimates. To calculate mortality costs we used a value of statistical life (VSL), “the financial value society places on reducing the average number of deaths by one” (24, 32). We used a 2014 VSL from the Australian Office of Best Practice Regulation (32), adjusted to 2015 dollars using the Reserve Bank of Australia inflation calculator (36), of \$4,263,351.50.

Data sources for new ongoing illness cost estimates. To calculate the health care costs of new ongoing illness over 1 year, we used medication costs and specialist visit costs as above. To calculate lost productivity costs from ongoing illness over 1 year, we used the lost productivity cost described above.

Burden estimates. We used the same estimation approach, distributions and multipliers to estimate foodborne NTS, IBS, and ReA incidence, hospitalizations and deaths

as in the foodborne disease estimation study circa 2010 (18, 27). We used simulation techniques in @Risk for each estimate using multiple inputs, each with a level of uncertainty for our four age groups. The final output was a distribution, from which we extracted a median estimate of incidence, hospitalizations and deaths together with 90% credible intervals (CrI). As mortality data were only available for 2000-2010, we added an additional step to these models, which multiplied the estimated median rate of foodborne illness for NTS by the population for 2013-2015. This adjusted the estimate for changes in the population since 2000-2010, although it does not allow us to detect any change in the death rate since 2010. More information about distributions and outputs can be found in supplementary material 1.

Cost estimates. In @Risk, we multiplied burden estimates by costs for health care, lost productivity, and valuation of premature mortality for each age group and for all cases to calculate costs in Australian dollars (AUD) for health care usage, the loss to productivity, and the implied value of life. From the resulting distribution, we extracted a median, mean, and 90% CrIs. We used an expert elicitation of the proportion of foodborne NTS infections (43) to estimate the cost of NTS, IBS, and ReA due to contaminated food.

Sensitivity analyses. We used a probabilistic simulation approach by calculating 90% credible intervals for key parameters known to vary in our burden estimates. We also conducted a one-way sensitivity analysis by halving and doubling individually first health care usage costs, then productivity costs, and finally the VSL values (15). In addition, as estimating the net present value of production changes is difficult (12), we tested the impact of removing all productivity losses from acute illness, and used an alternative participation rate to model the probability that both parents of 0-4 year olds and 5-19 year olds are working (0.627) in calculating lost productivity due to caring. As around 75% of 15-19 year olds are in the labour force part-time (2) and a day of missed paid work may not be the equivalent of a

full-time day, we also performed a sensitivity analysis of halving the daily wage rate for this group.

Results

Estimated burden of NTS in Australia. We estimated 90,833 (90% CrI 51,583-158,265) cases, 4,312 (90% CrI 3,335-11,091) hospitalizations, and 19 (90% CrI 15-22) deaths from salmonellosis in Australia circa 2015 (Table 1). Of these, 64,000 (90% CrI 34,000-117,000) cases, 3,100 (90% CrI 1,829-4,786) hospitalizations, and 13 (90% CrI 10-17) deaths were estimated to be due to contaminated food.

Estimated costs of NTS disease. We estimate that NTS cost a median of AUD 124.4 million and a mean of 124.7 million (90% CrI 107.4-143.1 million) circa 2015, with a median of 23.7 million and a mean of 23.8 million (90% CrI 19.3-28.9 million) from health care usage, a median of 21.3 million and a mean of 22.0 million (90% CrI 13.7-32.6 million) from lost productivity, and a median and mean of 79.0 million (90% CrI 66.0-92.1 million) from premature mortality. When acute and ongoing illness from IBS and ReA following *Salmonella* infection were included, the estimated cost was a median of AUD 146.8 million and a mean of 147.2 million (90% CrI 127.8-167.9 million) (Table 2 and supplementary material 3). NTS illness due to contaminated food were responsible for a median of AUD 88.9 million and a mean of 88.7 million (90% CrI 63.9-112.4 million) of the total cost of NTS illness and a median of 104.8 million and a mean of 104.6 (90% CrI 75.5-132.3 million) when IBS and ReA were included. Premature mortality from NTS as the underlying or contributing cause of death was the highest contributor to the total cost, accounting for 63% of the NTS cost and 54% of the total cost. Lost productivity due to acute NTS infection accounted for 17% of the NTS cost, lost productivity due to acute and ongoing IBS accounted

for 39% of the IBS cost, and lost productivity from acute and ongoing ReA accounted for 59% of the ReA cost.

The total cost of illness and the total cost per case was highest in the 65+ year age group (Table 3, Table 4, and supplementary material 3 for means). Overall, we estimated that the cost of acute salmonellosis was a median of AUD 1,322 per case and AUD 1,559 per case when IBS and ReA were considered (Table 4 and supplementary material 3 for means).

Sensitivity analysis. In probabilistic simulation of the burden estimate parameters, uncertainty was greatest in premature mortality, followed by estimates of lost productivity (Figure 1).

In one-way sensitivity analyses of costs, varying VSL for premature mortality costs by halving and doubling costs had the largest impact on cost values (Figure 2). Additional sensitivity analysis showed that removing costs associated with short term productivity loss from acute NTS, IBS and ReA illness resulted in a AUD 21.1 million drop in the NTS cost to 102.7 million (90% CrI 89.0-116.8 million) and a 26.5 million drop in the total cost estimate to 119.8 million (90% CrI 104.7-135.7 million). Adjusting the participation rate of carers for 0-4 year olds and 5-19 year olds, and halving the daily wage for 5-19 year olds had a minimal impact on the final cost estimates, reducing the estimate by less than 2% and less than 1% respectively.

Discussion

We estimated that the burden of salmonellosis and its sequelae in Australia are substantial, costing a median of 146.8 million Australian dollars in a typical year circa 2015. Of the total cost, 35.7 million (24%) was due to health care costs, 31.7 million (22%) due to costs associated with lost productivity, and 79.0 million (54%) due to premature mortality.

Quantifying these costs allows the burden of NTS to be compared against other illnesses, assisting with prioritization of public health interventions for policy making.

As NTS is mostly transmitted through contaminated food, targeted interventions across the food chain could help to prevent acute and ongoing illness. Australia's Foodborne Illness Reduction Strategy 2018-2021+ aims to reduce the number of illnesses from foodborne salmonellosis (17). While the national strategy does not set a specific reduction target, a 10% reduction in acute foodborne NTS illness incidence would equate to over AUD 8 million immediate reduction in costs, with additional reductions from sequel illnesses expected as well.

Due to the higher proportion of people aged 65+ with salmonellosis taking medications, admitted to hospital, and dying compared with other age groups, the cost per case was highest in the 65+ age group. Therefore interventions targeted at preventing illness in this age group would be most effective in reducing all costs. Our total cost per case for NTS of AUD 1,322 (~USD 995) is generally lower than in studies from the USA from 2013 & 2015 (USD 3,568 (16), USD 5,335 (31)), although higher than an estimate of €691 (~USD 765) by Sundstrom (39) for Sweden in 2016 dollars and the estimate of NZD 220 (~ USD 151) by Lake et al (28) for New Zealand in 2006/07. The estimates from these studies are not directly comparable due to differing years, variables estimated, and estimation methods.

As premature mortality from NTS as the underlying or contributing cause of death was the largest contributor to our cost estimates, accounting for 63% of total NTS cost and 54% of NTS and sequelae cost, the method used to value a lost life is very influential. In one-way sensitivity analysis, premature mortality costs had the largest range of uncertainty, compared with health care usage and lost productivity costs. Methods and values for lost life are also influential when comparing estimates between cost-of-illness studies. For example, while the total cost per NTS case estimated by the Economic Research Service for the USA is

higher than our estimate at USD 3,568, when only costs from healthcare usage and lost productivity are taken into account, the cost estimates are similar at \$383.55 USD for the USA compared to our estimate of AUD 480 (~USD 361) (16).

Costs associated with IBS and ReA following *Salmonella* were also a large contributor to our total cost estimate of \$146.8 million AUD, accounting for 14% of the total cost (10% from IBS and 4% from ReA). While we did account for these sequelae costs in our estimates, we have not included costs of NTS to industry and public health agencies for surveillance and regulation. As NTS can be expensive for the food industry, including costs associated with everyday prevention measures, product recalls, and liability, we have likely underestimated the costs to society.

While we estimated lost productivity costs in line with previously published methods (1), these methods may overestimate costs as we have not considered that either production will be made upon return to work or that employers will have excess capacity in the labor force to cover absenteeism for short term absences (12). In addition, a weakness of the human capital approach is that it overestimates productivity loss compared to the friction cost approach (33). We restricted lost productivity estimates to paid work only, not including the value of lost time for those not employed, such as unpaid carers or housework, or the loss of leisure time. One-way sensitivity analysis of removing productivity loss from acute illness resulted in a \$21.1 million reduction in the estimate for NTS and a \$26.5 million reduction for NTS, IBS, and ReA.

A limitation of this study is that due to a lack of data for IBS and ReA, we relied on expert opinion from a physician to review and update the proportion of cases receiving certain medications, pathology tests, and IBS cases seeing specialists from previous work and the literature. We used uncertainty intervals around these estimates to account for the uncertainty in the data. There is also limited data on duration of ongoing ReA symptoms,

with some studies finding that symptoms had ceased in most patients in less than 12 months (22, 23, 29), while others found cases with long lasting symptoms (19, 41). We used a conservative estimate of the proportion of ReA cases with ongoing symptoms in one year based on Hannu et al (22) and Leirisalo-Repo et al (29). While we were not able to include the decrease in quality of life to calculate quality adjusted life years, we have captured the decrease in quantity of life through the value of statistical life. While our cost of illness estimates provide a baseline, a full economic evaluation that compares costs and benefits of alternative policy interventions is needed to identify how effective new programs and policies would be at reducing burden. Identifying and implementing targeted prevention measures in the food supply chain could considerably reduce overall NTS costs.

In summary, the clinical and economic burden of salmonellosis and its sequelae are high. The cost of illness estimates in this paper will inform food safety policy and can be used in subsequent analysis of cost effectiveness of new policies and interventions aimed at the prevention and control of NTS infection in Australia.

Acknowledgements

Laura Ford is supported by an Australian Government Research Training Program (RTP) Scholarship. Martyn D. Kirk is supported by a National Health & Medical Research Council fellowship (APP1145997). Deborah A. Williamson is supported by a National Health & Medical Research Council fellowship (APP1123854).

Supplementary Material

Supplementary material associated with this article can be found online at: [URL to be completed by the publisher].

References

1. Abelson, P., M. P. Forbes, and G. Hall. 2006. The annual cost of foodborne illness in Australia. Australian Government Department of Health and Ageing, Canberra.
2. Australian Bureau of Statistics. 2018. 6202.0 - Labour Force, Australia, May 2018. GM1 - Labour force status and gross change (flows) by Age, Sex, State and Territory, February 1991 onwards. Available at:
<http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/6202.0May%202018?OpenDocument>. Accessed 3 July 2018.
3. Australian Bureau of Statistics. 2018. 6302.0 - Average Weekly Earnings, Australia, Nov 2017. Available at:
<http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/6302.0Nov%202017?OpenDocument>. Accessed 7 June 2018.
4. Australian Bureau of Statistics. 2016. Australian demographic statistics, Dec 2015. Table 59. Estimated Resident Population by Single Year of Age, Australia. Available at:<http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/3101.0Dec%202015?OpenDocument>. Accessed 20 June 2017.
5. Australian Institute of Health and Welfare. Separation statistics by principal diagnosis (ICD-10-AM 7th edition), Australia, 2011-12 to 2012-13. Available at:
<http://www.aihw.gov.au/hospitals-data/principal-diagnosis-data-cubes/>. Accessed 11 June 2016.
6. Australian Institute of Health and Welfare. Separation statistics by principal diagnosis (ICD-10-AM 7th edition), Australia, 2013-14 to 2014-15. Available at:
<http://www.aihw.gov.au/hospitals-data/principal-diagnosis-data-cubes/>. Accessed 11 June 2016.

7. Centres for Disease Control and Prevention. 2016. National *Salmonella* surveillance annual report, 2013. US Department of Health and Human Services, Atlanta GA.
8. Chen, Y., L. Ford, G. Hall, T. Dobbins, and M. Kirk. 2016. Healthcare utilization and lost productivity due to infectious gastroenteritis, results from a national cross-sectional survey Australia 2008-2009. *Epidemiol Infect.* 144: 241-246. doi: 10.1017/s0950268815001375.
9. Department of Health. MBS Online. Available at: <http://www.mbsonline.gov.au/internet/mbsonline/publishing.nsf/Content/Home>. . Accessed 7 June 2018.
10. Department of Health. The Pharmaceutical Benefits Scheme. Accessed Available at: <http://www.pbs.gov.au/>. Accessed 8 June 2018.
11. Department of Health. National Notifiable Disease Surveillance System. Notifications of a selected disease by age group, sex and year. Available at: http://www9.health.gov.au/cda/source/rpt_5_sel.cfm. Accessed 11 June 2016.
12. Department of Health. 2016. Guidelines for preparing a submission to the Pharmaceutical Benefits Advisory Committee. Australian Government, Canberra.
13. Department of Human Services. Medicare Item Reports. Available at: http://medicarestatistics.humanservices.gov.au/statistics/mbs_item.jsp. Accessed 7 June 2018.
14. Department of Human Services. Pharmaceutical Benefits Schedule Item Reports. Available at: http://medicarestatistics.humanservices.gov.au/statistics/pbs_item.jsp. Accessed 8 June 2018.
15. Drummond, M., M. J. Sculpher, K. Claxton, G. L. Stoddart, and G. W. Torrance. 2015. Methods for the Economic Evaluation of Health Care Programs, 4th edition. Oxford University Press, Oxford.

16. Economic Research Service. Cost estimates of foodborne illness. Available at:
<https://www.ers.usda.gov/data-products/cost-estimates-of-foodborne-illnesses/>.
Accessed 18 May 2017.
17. Food Standards Australia New Zealand. 2018. Australia's Foodborne Illness Reduction Strategy 2018-2021+. Australian Government, Canberra.
18. Ford, L., M. Kirk, K. Glass, and G. Hall. 2014. Sequelae of foodborne illness caused by 5 pathogens, Australia, circa 2010. *Emerg Infect Dis.* 20:1865-1871. doi: 10.3201/eid2011.131316.
19. Fox, R., A. Calin, R.C. Gerber, and D. Gibson. 1979. The chronicity of symptoms and disability in Reiter's syndrome. An analysis of 131 consecutive patients. *Ann Intern Med.* 91:190-193.
20. Government of Canada. Reported cases from 1924-2016 in Canada - Notifiable diseases on-line. Available at: <http://diseases.canada.ca/ndis/charts.php?c=plt>.
Accessed 1 Jan 2019.
21. Hall, G., K. Yohannes, J. Raupach, N. Becker, and M. Kirk. 2008. Estimating community incidence of *Salmonella*, *Campylobacter*, and Shiga toxin-producing *Escherichia coli* infections, Australia. *Emerg Infect Dis.* 14:1601-1609. doi: 10.3201/eid1410.071042.
22. Hannu, T., L. Mattila, A. Siitonen, and M. Leirisalo-Repo. 2005. Reactive arthritis attributable to *Shigella* infection: a clinical and epidemiological nationwide study. *Ann Rheum Dis.* 64:594-598. doi: 10.1136/ard.2004.027524.
23. Hannu, T., L. Mattila, H. Rautelin, P. Pelkonen, A. Siitonen, and M. Leirisalo-Repo. 2002. *Campylobacter*-triggered reactive arthritis: a population-based study. *Rheumatology (Oxford)*. 41:312-318.

24. HM Treasury. 2018. The Green Book. Central government guidance on appraisal and evaluation. UK Government, London.
25. Hoffmann, S., M. B. Batz, and J. G. Morris, Jr. 2012. Annual cost of illness and quality-adjusted life year losses in the United States due to 14 foodborne pathogens. *J Food Prot.* 75:1292-1302. doi: 10.4315/0362-028x.jfp-11-417.
26. Independent Hospital Pricing Authority. 2016. National Hospital Cost Data Collection, Australian Public Hospitals Cost Report, Round 18 (Financial year 2013-14).
27. Kirk, M., L. Ford, K. Glass, and G. Hall. 2014. Foodborne illness, Australia, circa 2000 and circa 2010. *Emerg Infect Dis.* 20:1857-1864. doi: 10.3201/eid2011.131315.
28. Lake, R. J., P. J. Cressey, D. M. Campbell, E. Oakley. 2010. Risk ranking for foodborne microbial hazards in New Zealand: burden of disease estimates. *Risk Anal.* 30:743-752. doi: 10.1111/j.1539-6924.2009.01269.x.
29. Leirisalo-Repo, M., P. Helenius, T. Hannu, A. Lehtinen, J. Kreula, M. Taavitsainen, and S. Koskimies. 1997. Long-term prognosis of reactive salmonella arthritis. *Ann Rheum Dis.* 56:516-520.
30. Marshall, J. K., M. Thabane, M. R. Borgaonkar, and C. James. 2007. Postinfectious irritable bowel syndrome after a food-borne outbreak of acute gastroenteritis attributed to a viral pathogen. *Clin Gastroenterol Hepatol.* 5:457-460. doi: 10.1016/j.cgh.2006.11.025.
31. Minor, T., A. Lasher, K. Klontz, B. Brown, C. Nardinelli, and D. Zorn. 2015. The per case and total annual costs of foodborne illness in the United States. *Risk Anal.* 35:1125-1139. doi: 10.1111/risa.12316.

32. Office of Best Practice Regulation. 2014. Best Practice Regulation Guidance Note: Value of statistical life. Department of the Prime Minister and Cabinet, Australian Government, Canberra.
33. Pike, J., and S. D. Grosse. 2018. Friction Cost Estimates of Productivity Costs in Cost-of-Illness Studies in Comparison with Human Capital Estimates: A Review. *Appl Health Econ Health Policy*. doi: 10.1007/s40258-018-0416-4.
34. Public Health England. 2017. *Salmonella* data 2006 to 2015. November 2016. National laboratory data for residents of England and Wales. Public Health England, London.
35. Reeve, R., and M. Haas. 2013. Estimating the Cost of Emergency Department Presentations in NSW. Ultimo NSW: Centre for Health Economics Research and Evaluation. University of Technology, Sydney.
36. Reserve Bank of Australia. Inflation Calculator. Available at: <https://www.rba.gov.au/calculator/>. Accessed 17 July 2018.
37. Santos, A. C., J. A. Roberts, A. J. Cook, R. Simons, R. Sheehan, C. Lane, G. K. Adak, F. A. Clifton-Hadley, and L. C. Rodrigues. 2011. *Salmonella* Typhimurium and *Salmonella* Enteritidis in England: costs to patients, their families, and primary and community health services of the NHS. *Epidemiol Infect.* 139:742-753. doi: 10.1017/s0950268810001615.
38. Schonberg-Norio, D., L. Mattila, A. Lauhio, M. L. Katila, S. S. Kaukoranta, M. Koskela, S. Pajarre, J. Uksila, E. Eerola, S. Sarna, and H. Rautelin. 2010. Patient-reported complications associated with *Campylobacter jejuni* infection. *Epidemiol Infect.* 138:1004-1011. doi: 10.1017/s0950268809991099.

39. Sundstrom, K. 2018. Cost of illness for five major foodborne illnesses and sequelae in Sweden. *Appl Health Econ Health Policy*. 16:243-257. doi: 10.1007/s40258-017-0369-z.
40. The Institute of Environmental Science and Research Ltd. 2017. Notifiable diseases in New Zealand: Annual report 2016. ESR, Porirua.
41. Thompson, G.T., D.A. DeRubeis, M.A. Hodge, C. Rajanayagam, and R.D. Inman. 1995. Post-Salmonella reactive arthritis: late clinical sequelae in a point source cohort. *Am J Med*. 98:13-21.
42. Townes, J. M., A. A. Deodhar, E. S. Laine, K. Smith, H. E. Krug, A. Barkhuizen, M. E. Thompson, P. R. Cieslak, and J. Sobel. 2008. Reactive arthritis following culture-confirmed infections with bacterial enteric pathogens in Minnesota and Oregon: a population-based study. *Ann Rheum Dis*. 67:1689-1696. doi: 10.1136/ard.2007.083451.
43. Valley, H., K. Glass, L. Ford, G. Hall, M. D. Kirk, C. Shadbolt, M. Veitch, K. E. Fullerton, J. Musto, and N. Becker. 2014. Proportion of illness acquired by foodborne transmission for nine enteric pathogens in Australia: an expert elicitation. *Foodborne Pathog Dis*. 11:727-733. doi: 10.1089/fpd.2014.1746.
44. World Health Organization. 2015. WHO estimates of the global burden of foodborne diseases. Foodborne Diseases Burden Epidemiology Reference Group 2007-2015. Available at: http://www.who.int/foodsafety/areas_work/foodborne-diseases/ferg/en/. Accessed 18 May 2017.

Figure legends

FIGURE 1: Median cost of acute and ongoing illness from NTS, IBS, and ReA from health care usage, lost productivity, and premature mortality, circa 2015, Australia. Ranges reflect 90% credible intervals for burden estimates.

FIGURE 2: One-way sensitivity analysis of health care usage, lost productivity, and premature mortality for NTS, IBS, and ReA cost of illness circa 2015, Australia. Low values represent when costs were halved and high values when costs were doubled around our cost estimate

Tables

TABLE 1: Estimated median number of cases, hospitalizations, and deaths from NTS and resulting IBS and ReA circa 2015, Australia

	Median number (90% CrI)		
	NTS ^a	IBS ^a	ReA ^a
<i>Cases</i>			
0-4 years	20,924 (12,059-36,250)	1.46 (0.82-2.57)	341 (56-898)
5-19 years	14,617 (8,408-25,384)	1,284 (714-2,308)	1,206 (290-3,125)
20-64 years	44,525 (25,012-77,489)	3,906 (2,104-7,077)	3,652 (865-9,626)
65+ years	10,767 (6,104-19,042)	942 (516-1,703)	881 (209-2,310)
Total	90,833 (51,583-158,165)	6,133 (3,335-11,091)	6,080 (1,420-15,959)
<i>Hospitalizations</i>			
0-4 years	984 (661-1,346)	0.29 (0-0.23)	1.84 (1.07-3.52)
5-19 years	607 (391-912)	16 (9-25)	4 (3-6)
20-64 years	1,814 (1,162-2,750)	326 (186-517)	20 (13-26)
65+ years	907 (570-1,382)	67 (39-104)	2.5 (2-4)
Total	4,312 (2,784-6,390)	409 (234-647)	28 (19-40)
<i>Deaths</i>			
0-4 years	0.65 (0.5-0.82)	-	-
5-19 years	1.7 (1.34-2.07)	-	-
20-64 years	4 (2.71-4.38)	-	-
65+ years	13 (10-15)	-	-
Total	19 (15-22)	-	-

^aNTS: non-typhoidal *Salmonella*; IBS: irritable bowel syndrome; ReA: reactive arthritis

TABLE 2: Estimated annual cost of illness (millions AUD) of health care usage, lost productivity, premature mortality, for acute and ongoing illness, circa 2015, Australia

	Median cost in millions AUD (90% CrI)			
	NTS ^a	IBS ^a	ReA ^a	Total ^b
<i>Health care usage</i>				
Acute illness	23.7 (19.3-28.9)	5.6 (4.0-8.0)	1.9 (1.0-3.6)	31.4 (26.5-37.2)
Ongoing illness	-	3.2 (1.8-6.9)	0.77 (0.34-1.7)	4.2 (2.7-6.5)
<i>Lost productivity</i>				
Acute illness	21.3 (13.7-32.6)	2.0 (1.3-3.3)	2.7 (1.0-6.5)	26.5 (18.5-38.4)
Ongoing illness	-	3.6 (1.8-6.9)	1.0 (0.37-2.8)	4.8 (2.7-8.4)
<i>Premature mortality</i>				
Acute illness	79.0 (66.0-92.1)	-	-	79.0 (66.0-92.1)
<i>Total costs</i>				
Acute illness	124.4 (107.4-143.1)	7.6 (5.3-11.3)	4.6 (2.0-10.0)	137.7 (119.7-157.0)
Ongoing illness	-	6.8 (3.9-12.3)	1.8 (0.73-4.4)	9.0 (5.5-14.8)
Total	124.7 (107.4-143.1)	14.5 (9.5-23.0)	6.5 (2.8-14.2)	146.8 (127.8-167.9)

^aNTS: non-typhoidal *Salmonella*; IBS: irritable bowel syndrome; ReA: reactive arthritis

^bNumbers may not sum due to simulation and rounding

TABLE 3: Estimated annual cost of acute and ongoing illness in 1 year (millions AUD) by age group, circa 2015, Australia

Median cost in millions AUD (90% CrI)				
	NTS ^a	IBS ^a	ReA ^a	Total ^b
<i>Health care usage</i>				
0-4 years	4.2 (3.0-5.9)	0.004 (0.002-0.006)	0.14 (0.04-0.37)	4.4 (3.2-6.1)
5-19 years	2.8 (2.0-4.0)	1.6 (0.90-3.0)	0.48 (0.13-1.3)	5.1 (3.7-6.9)
20-64 years	8.6 (5.9-12.4)	5.7 (3.3-9.8)	1.5 (0.46-4.0)	16.3 (11.9-22.0)
65+ years	7.6 (5.1-11.0)	1.4 (0.80-2.4)	0.34 (0.10-0.93)	9.4 (6.8-13.0)
<i>Lost Productivity</i>				
0-4 years	4.9 (2.3-10.0)	0.002 (0.001-0.004)	0.29 (0.07-0.78)	5.2 (2.6-10.3)
5-19 years	1.5 (0.46-4.4)	0.34 (0.17-0.65)	0.23 (0.05-0.61)	2.2 (1.0-5.0)
20-64 years	12.6 (6.7-22.8)	5.0 (2.7-9.4)	3.1 (0.73-8.4)	21.7 (13.9-33.6)
65+ years	1.2 (0.33-3.0)	0.18 (0.10-0.35)	0.11 (0.03-0.31)	1.5 (0.64-3.3)
<i>Premature mortality</i>				
0-4 years	2.8 (2.1-3.5)	-	-	2.8 (2.1-3.5)
5-19 years	7.2 (5.7-8.8)	-	-	7.2 (5.7-8.8)
20-64 years	15.0 (11.6-18.7)	-	-	15.0 (11.6-18.7)
65+ years	53.8 (41.6-66.3)	-	-	53.8 (41.6-66.3)
<i>Total costs</i>				
0-4 years	11.9 (8.5-18.1)	0.006 (0.004-0.009)	0.43 (0.11-1.1)	12.4 (8.9-18.6)
5-19 years	11.8 (9.4-15.3)	2.0 (1.1-3.6)	0.70 (0.09-1.9)	14.7 (11.9-18.6)
20-64 years	36.5 (27.7-49.8)	10.7 (6.0-19.1)	4.6 (1.2-12.2)	53.3 (41.2-70.0)
65+ years	62.9 (50.3-75.9)	1.5 (0.9-2.7)	0.46 (0.12-1.2)	65.2 (52.5-78.2)

^aNTS: non-typhoidal *Salmonella*; IBS: irritable bowel syndrome; ReA: reactive arthritis

^bNumbers may not sum due to simulation and rounding

TABLE 4: Estimated median annual cost of acute and ongoing illness (\$AUD) per case by age group, circa 2015, Australia

Median cost AUD				
	NTS ^a	IBS ^a	ReA ^a	Total
<i>Health care usage only</i>				
0-4 years	200.78	854.05	302.90	208.65
5-19 years	190.85	434.31	287.48	344.33
20-64 years	192.17	491.17	301.1	363.49
65+ years	691.93	491.80	284.09	866.14
Total	250.02	476.64	293.32	377.97
<i>Health care usage with lost productivity</i>				
0-4 years	443.64	1,398.50	920.24	467.78
5-19 years	307.88	525.02	423.95	507.56
20-64 years	480.79	930.31	916.21	845.78
65+ years	814.92	558.66	378.88	1,020.85
Total	479.99	783.99	718.10	713.85
<i>Health care usage, lost productivity, and premature mortality</i>				
0-4 years	579.88	1,398.50	920.24	603.97
5-19 years	807.77	525.02	423.95	1,003.88
20-64 years	818.47	930.31	916.21	1,190.35
65+ years	5,798.29	558.66	378.88	6,003.72
Total*	1,322.14	783.36	718.96	1,558.53

^aNTS: non-typhoidal *Salmonella*; IBS: irritable bowel syndrome; ReA: reactive arthritis