



Original article

Scand J Work Environ Health 2020;46(3):311-320

doi:10.5271/sjweh.3869

Is the association between poor job control and common mental disorder explained by general perceptions of control? Findings from an Australian longitudinal cohort

by [Too LS](#), [Leach L](#), [Butterworth P](#)

After adjusting for general perceptions of control and a range of other covariates, low job control remained significantly associated with an increased risk of common mental disorder. This strengthens evidence that job control is an important contributor to poor mental health, independent of individual differences. Promoting employee job control is an evidence-supported approach to enhance mental health.

Affiliation: Centre for Mental Health, Melbourne School of Population and Global Health, The University of Melbourne, Parkville, Victoria 3052, Australia. tiffany.too@unimelb.edu.au

Refers to the following texts of the Journal: [2002;28\(2\):94-108](#)
[2006;32\(6\):443-462](#)

Key terms: [association](#); [Australia](#); [common mental disorder](#); [control](#); [depression](#); [job control](#); [job stressor](#); [longitudinal cohort](#); [mental disorder](#); [mental health](#); [psychosocial](#); [stress](#)

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/31805200



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Is the association between poor job control and common mental disorder explained by general perceptions of control? Findings from an Australian longitudinal cohort

by Lay San Too, PhD,^{1,2} Liana Leach, PhD,^{3*} Peter Butterworth, PhD²

Too LS, Leach L, Butterworth P. Is the association between poor job control and common mental disorder explained by general perceptions of control? Findings from an Australian longitudinal cohort. *Scand J Work Environ Health*. 2020;46(3):311–320. doi:10.5271/sjweh.3869

Objectives This study sought to examine the influence of general perceptions of control on the association between job control and mental health.

Methods We used four waves of data from a cohort of mid-aged adults from the Personality and Total Health (PATH) Through Life Study (baseline N=2106). Key measures included job control and likelihood of experiencing a common mental disorder (anxiety and/or depression). The data were analyzed using longitudinal random-intercept regression models, controlling for a range of potential confounders including general perceptions of control (ie, not isolated to the work context) via a measure of mastery. The analyses isolated the effect of within-person changes in job control on mental health (apart from between-person differences).

Results The results show that the effect of job control remained significant after adjusting for general perceptions of control and other confounders. The within-person effect in the model demonstrated that, when workers had low job control, they were twice as likely to experience a common mental disorder [odds ratio (OR) 2.04, 95% confidence interval (CI) 1.53–2.73].

Conclusions Individuals' general perceptions of control in life does not account for the association between low job control and poor mental health. The findings add a new layer of evidence to the literature demonstrating that lack of autonomy at work is an independent predictor of employees' mental health. Increasing employee control should be integrated into workplace strategies to promote mental health.

Key terms Australia; depression; job stressor; mental health; psychosocial; stress.

Globally, mental illness is a key contributor to disability burden (1). In an effort to reduce the prevalence of mental illness, risk factor identification (and prevention) in the workplace has become a major focus of research as well as both government and employment policy (2). Systematic reviews and meta-analyses have consistently found that psychologically stressful employment conditions (including jobs with high demands, low control/autonomy, insecurity, and poor social support) are associated with poorer mental health (3–6). This research field is primarily influenced by Karasek's job

demands–control model (7). The model posits that job strain, a state produced by the combination of high job demands and low job control, places employees at an increased risk of health problems. Following on from the job strain model, much of the literature investigating psychosocial job conditions and mental health has focused on combined (rather than single) indicators of psychosocial stress at work. For example, multiple longitudinal studies have shown that high job strain (high demands and low control) precedes declines in mental health (8–12) as do increases in the number of adverse

¹ Centre for Mental Health, Melbourne School of Population and Global Health, The University of Melbourne, Parkville, Victoria, Australia.

² Centre for Research on Ageing, Health & Wellbeing, Research School of Population Health, The Australia National University, Canberra ACT, Australia.

³ National Centre for Epidemiology and Population Health, Research School of Population Health, The Australia National University, Canberra ACT, Australia.

⁴ Melbourne Institute of Applied Economic and Social Research, The University of Melbourne, Parkville, Victoria, Australia.

* These authors contributed equally to this work.

Correspondence to: Dr Lay San Too, Centre for Mental Health, Melbourne School of Population and Global Health, The University of Melbourne, Parkville, Victoria 3052, Australia. [E-mail: tiffany.too@unimelb.edu.au]

psychosocial job conditions experienced (eg, combined experiences of high job demands, low job control, job insecurity) (13, 14).

Research has also examined the individual indicators of poor psychosocial job quality and, in the case of most indicators including low job control, has reliably shown an association with poorer mental health. For example, a UK study by Stansfeld et al (15) showed an increased risk of psychiatric disorders among those with low decision authority. Bentley et al (16) and Butterworth et al (17) analyzed seven or more waves of Australian panel data and showed that mental health improved with increases in job control over time. A recent UK longitudinal study by Harvey et al (12) found that low job control predicted higher odds of subsequent common mental disorder. While there is a substantial body of research linking low job control with poorer mental health, the possibility of residual confounding remains a major barrier to causal interpretations in longitudinal research (12). There are many non-work related mental health risk factors that may be associated with self-reported job control, including psychological resources and personality characteristics. While research by Stansfeld et al (15) and Harvey et al (12) did control for some aspects of personality (eg, adult hostility and adolescent temperament respectively), very little research has considered the role that perceptions of control more broadly (ie, *not isolated to the work context*) might play. Measures of control in the workplace may simply be a proxy for perceptions of control more generally, or alternatively, it may be that control in the workplace is associated with mental health independent of non-work influences. One relevant study by Clark et al (18) used cross-sectional data from the 2007 UK Adult Psychiatric Morbidity Survey to investigate the contribution of both work and non-work stressors to common mental disorders. The findings suggest that the effects of work stressors are not explained by broader non-work stressors such as adverse life events (eg, financial strain, separation/divorce, domestic violence), caring responsibilities, and low social support. However, this research only used data from a single point in time and did not specifically investigate the contribution of control within and outside the workplace.

The current study aims to fill this knowledge gap by using four waves of Australian community-based data to examine the association between job control (one of the main components in Karasek's job demands-control model) and common mental disorder, independent of a broader sense of control in life. It is important to account for the influence of control more broadly, as without doing so, the effect of job control might simply reflect perceptions of low control over life circumstances more generally rather than being specific to work. This study separated within- and between-person differences in

both job control and broader sense of control to show how time-varying change in job control is associated with change in mental health, adjusted for other residual confounders (eg, between-person differences in job control as well as within- and between-person broader sense of control). The findings are reported according to the STROBE statement for cohort studies (19).

Methods

Participants

Participants were from the Personality and Total Health (PATH) Through Life Project, a large longitudinal community survey on individual health and well-being trajectories across the life course. The survey was undertaken by the Centre for Research on Ageing, Health and Wellbeing at the Australian National University (ANU). The sample comprises cohorts of young, midlife, and older adults randomly selected from the Australian Electoral Rolls of Australian Capital Territory and the neighboring town of Queanbeyan (20). The current study was restricted to the midlife cohort who were aged 40–44 years at baseline, and subsequently assessed every (approximately) four years from 2000/2001 (wave 1) to 2012/2013 (wave 4). The participation rate of this cohort at baseline was 65% (2530 participants), where 93% of baseline participants completed the survey at wave 2, 86% at wave 3, and 71% at wave 4 (figure 1). For the first three waves, participants were typically assessed in their own home or at the ANU. They were invited to complete a questionnaire using a laptop computer under the supervision of a trained interviewer. For the fourth wave, participants were invited to complete an online version of the questionnaire and those in the local area were invited to undertake additional face-to-face assessment. All participants provided informed consent to participate at each wave of the study, and each wave of data collection was approved by the Human Research Ethics Committee of the Australian National University.

The current study excluded observations from participants when they (i) were not employed (ie, unemployed or not active in the labor force); (ii) were employed but on long-term leave; or (iii) did not provide information on their status of employment (figure 1). Given the longitudinal focus, participants with fewer than two waves of data were also excluded.

Measures

Common mental disorder. Anxiety and depression were assessed using the Goldberg Anxiety and Depression scales (21) at all four waves. Each scale comprises nine




Figure 1. Study profile

binary items (yes or no) about anxiety or depressive symptoms experienced in the past month. The total score for each measure was computed by summing the number of "yes" responses and dichotomized using the established cut-off points indicative of symptoms of anxiety or depressive disorders (22). A binary measure of common mental disorder was then generated based on the presence of depressive and/or anxiety symptoms. A continuous measure of common mental disorder severity was also calculated based on number of symptoms of anxiety and depression experienced (ranging from 0–18) – this measure was adopted in sensitivity analyses.

Control at work. At all four waves, 15 items from the Whitehall II study (23) assessed aspects of job control (eg, skill discretion: "Do you have a choice in deciding how you do your job?"; decision authority: "Does your job provide you with a variety of interesting things?"). These items offered four response categories: 1=often, 2=sometimes, 3=rarely, and 4=never. Aligned with the

methodology used in previous studies (17, 24), a score for job control was calculated by combining the average scores for skill discretion and decision authority and then divided into tertiles identifying three groups representing high, medium, and low job control. This approach was similarly used in the Whitehall II study (15). In addition to the time-varying measure of job control, a between-person (ie, time-invariant) measure was computed to identify respondents who experienced low job control at some point over the four waves. This measure was used as indicator of individual susceptibility or vulnerability to experience adverse job conditions.

General perceptions of control

General perceptions of control assessed by the Pearlin Mastery Scale (25), which was included in all waves. Perceived mastery is a psychological resource that has been defined as "the extent to which one regards one's life-chances as being under one's own control in contrast

to being fatalistically ruled" (25). The items (eg, "There is really no way I can solve some of the problems I have" and "I have little control over the things that happen to me.") offered four response categories: 1=strongly agree, 2=agree, 3=disagree, and 4=strongly disagree. The seven items were summed and the total score categorized into tertiles, representing high, medium, and low general control. The analyses also included a between-person (ie, time-invariant) measure of general control, derived by calculating respondents average mastery scale score over the survey waves.

Other covariates

Work-related. Four items from the Whitehall II study (23) assessing aspects of job demands (eg, "Do you have to work very intensively") with four response categories (1=often, 2=sometimes, 3=rarely, and 4=never) were included from all waves. As with job control, we computed the average score for this measure and then used tertiles to identify three groups representing high, medium, and low job demands. One item: "How secure do you feel about your job or career future in your current workplace?" with four response categories ("not at all secure", "moderately secure", "secure", "extremely secure") was used to assess job insecurity. The former two categories were coded as high insecurity and the latter two as low insecurity.

Sociodemographic and health covariates. We included other variables that may potentially confound the association between job control and common mental disorders (26–28): time-invariant measures from baseline (ie, gender, educational attainment) and time-varying measures from each wave (ie, partner status, occupational skill level, parental responsibilities, childhood adversity, financial hardship, chronic physical health conditions, smoking status, alcohol consumption, and physical exercise).

Seventeen items on childhood adversity were adopted and adapted from the Parental Bonding Instrument (29), the British National Survey of Health and Development (30), the US National Comorbidity Survey (31), and from open-ended responses in a previous cross-sectional study (32). These items comprised six covering parental affection, parental nervous/emotional trouble, and parental drinking/drug problems; two covering conflict in the household and parental divorce/separation; eight covering neglect, authoritarian upbringing, witnessing physical/sexual abuse, as well as verbal abuse, psychological abuse, physical abuse, physical punishment, and sexual abuse by a parent; and one item covering childhood poverty/financial hardship. Participants were categorized as having childhood adversity if they responded yes to any of these items. Current financial hardship was generated based on one item: "Have you or your family had to go

without things you really needed in the last year because you were short of money?" Respondents were classified as having financial hardship if they responded yes, often or yes, sometimes to this item.

Chronic physical health conditions including heart problems, hypertension, cancer, arthritis, thyroid problems, epilepsy, asthma, diabetes, and stroke were considered and coded as a summary variable representing the experience of 0, 1, or ≥ 2 of these conditions. Hazardous/harmful alcohol consumption (33) was derived from the Alcohol Use Disorders Identification Test (34) and classified as yes and no. The hours respondents engaged in moderate or vigorous physical exercise per week was assessed by items in the Whitehall II study (35) and categorized into five groups: 0, <1.5, 1.5–3, 3.1–5.5, and >5.5 hours. The categories of covariates not described in detail above are presented in table 1.

Statistical analyses

Sample characteristics at baseline and descriptive information for job control and common mental disorder at each wave were reported. Chi-square test was used to assess the relationship between job control and general control. To appropriately model the lack of independence in the data, given repeated measures from the same individuals, longitudinal random-intercept logistic regression models with two levels (occasion clustered within individuals) were used to assess the association between job control and common mental disorder over time (36). These models also facilitate the use all available data (rather than restriction to complete case analysis), and permit the decomposition of time-varying covariates into elements representing within-person change over time and consistent between-person differences (36). The model fitted a fixed (average) regression slope for common mental disorder over time while permitting the intercept to vary between respondents to reflect differences between individuals in their likelihood of common mental disorder. The longitudinal models first included time-varying job control (model 1) and general control (model 2), followed by between-person terms for job control and general control (model 3), and finally other work-related covariates, sociodemographic, and health variables (model 4). Survey wave (a category term) was included in all models. We also assessed the association of each covariate with common mental disorder, adjusted for survey wave only.

In addition, we compared the model comprising only the main effects of job control and general control with a model that also incorporated the interaction term using the likelihood ratio test. We did the same test for job control and gender, and for general control and gender.

The proportion of observations with missing data on all variables was low, ranging from 0–1.5% (financial

Table 1. Sample characteristics at baseline (N=2106).

	Overall		Low job control		Medium job control		High job control	
	N	%	N	%	N	%	N	%
Sex								
Male	1052	49.9	266	38.6	387	52.4	399	58.8
Female	1054	50.1	423	61.4	351	47.6	280	41.2
Partner status								
No partner (separated/divorced/widowed/never married)	414	19.7	154	22.4	153	20.7	107	15.8
Having a partner (married/in a de facto relationship)	1692	80.3	535	77.7	585	79.3	572	84.2
Education completion								
Incomplete high school	561	26.6	248	36.0	196	26.6	117	17.2
Completion of high school	695	33.0	257	37.3	243	32.9	195	28.7
Completion of tertiary study	850	40.4	184	26.7	299	40.5	367	54.1
Occupational skill level ^a								
Low	420	19.9	276	40.1	102	13.8	42	6.2
Medium	564	26.8	192	27.9	220	29.8	152	22.4
High	1122	53.3	221	32.1	416	56.4	485	71.4
Parental responsibilities								
No (no youngest child aged <15 years)	720	34.2	251	36.4	260	35.2	209	30.8
Yes (youngest child aged <15 years)	1386	65.8	438	63.6	478	64.8	470	69.2
Childhood adversities								
No	810	38.5	233	33.8	290	39.3	287	42.3
Yes	1283	60.9	450	65.3	444	60.2	389	57.3
Unknown	13	0.6	6	0.9	4	0.5	3	0.4
Financial hardship								
No	1695	80.5	513	74.5	608	82.4	574	84.5
Yes	380	18.0	167	24.2	121	16.4	92	13.6
Unknown	31	1.5	9	1.3	9	1.2	13	1.9
Number of chronic physical health conditions								
0	1195	56.7	393	57.0	401	54.3	401	59.1
1	704	33.4	225	32.7	268	36.3	211	31.1
≥2	207	9.8	71	10.3	69	9.4	67	9.9
Smoking status								
Never/past smoker	1729	82.1	544	79.0	607	82.3	578	85.1
Current smoker	377	17.9	145	21.0	131	17.8	101	14.9
Hazardous/harmful alcohol consumption								
No	1978	93.9	656	95.2	693	93.9	629	92.6
Yes	128	6.1	33	4.8	45	6.1	50	7.4
Moderate/vigorous physical exercise (hours in the last week) ^b								
0	406	19.3	165	24.0	138	18.7	103	15.2
<1.5	444	21.1	163	23.7	150	20.3	131	19.3
1.5–3.0	357	17.0	115	16.7	117	15.9	125	18.4
3.1–5.5	514	24.4	139	20.2	197	26.7	178	26.2
>5.5	385	18.3	107	15.5	136	18.4	142	20.9
General control (ie, mastery)								
Low	420	19.9	214	31.1	136	18.4	70	10.3
Medium	946	44.9	343	49.8	353	47.8	250	36.8
High	726	34.5	126	18.3	245	33.2	355	52.3
Unknown	14	0.7	6	0.9	4	0.5	4	0.6

^a Low occupational skill level included intermediate production and transport workers; elementary clerical, sales and service workers; and laborers and related workers. Medium occupational skill level included associate professionals; tradespersons and related workers; and advanced clerical and service workers. High occupational skill level included managers and administrators; and professionals.

^b The average hours spent in the last week was 3.6 (SD 6.0) overall; 3.2 (SD 5.9) for low job control; 3.7 (SD 5.6) for medium job control; and 3.9 (SD 6.3) for high job control.

hardship was highest). All analyses were conducted using Stata Statistical Software: release 14 (StataCorp, College Station, TX, USA).

Sensitivity analyses

Sensitivity tests included repeating the analyses (using negative binomial regression models) with a continuous outcome representing severity of common mental disorder to assess whether the results from these models differed from those obtained from the models with

binary outcome of common mental disorder. Another sensitivity test repeated the analyses with a fixed-effect model rather than a random-effect model with between-person terms.

Results

The study included 2106 participants aged 40–44 at baseline (table 1, see descriptions of sample character-

istics by level of job control in table 1 and by level of general control in table S4 in the supplementary material, www.sjweh.fi/show_abstract.php?abstract_id=3869). Of these participants, half were male, 20% did not have a partner, 27% did not complete high school, 20% were in low-skilled occupations, 66% had a youngest child aged <15 years, 61% experienced ≥1 adversity during childhood, and 18% were experiencing financial hardship. In terms of their health, 10% of participants had ≥2 chronic physical health conditions, 18% currently smoked, 6% consumed alcohol at hazardous/harmful levels, and 19% spent less than an hour doing moderate/vigorous physical exercise per week.

The chi-square test showed a significant relationship between job control and general control [$X^2(4, N=2092) = 205.38, P<0.001$]. However, 17% of respondents who reported low job control also reported high general control. Similarly, 17% who reported high job control also reported low general control, suggesting these two measures were not collinear.

There was movement between low, medium, and high categories of job control and general control over the 12-year period (table 2). The proportion of respondents reporting low job control at any point in time (55%) was greater than the proportion at each wave (eg, 35% at wave 4). The proportion of respondents reporting low general control at any point in time (40%) was also greater than the proportion at each wave (eg, 23% at wave 4). There was also variability over time in common mental disorder. Around 42% of respondents reported symptom levels reflective of a common mental disorder at some point over the survey waves.

Job control and common mental disorder over time

Table 3 shows a series of models investigating the asso-

ciation between job control and likelihood of common mental disorder. Job control was significantly associated with common mental disorder in simple and adjusted models, although the magnitude of the effect was attenuated after controlling for general control. After adjusting for all covariates, including between-person differences in job control and general control, the results show that when individuals were in jobs with low control (ie, within-person change) they were at twice the risk of common mental disorder compared to when they held a job with high control (OR 2.04, 95% CI 1.53–2.73). Other variables associated with greater risk of common mental disorder in the final model included current experiences of low general control (eg, within-person effect) and overall average perceptions of general control (eg, between-person effect), high job demands and job insecurity, being female, childhood adversity, financial hardship, having one or more chronic physical conditions, and smoking. Those who had a low skilled occupation appeared at lower risk (although this likely arose from over-adjustment in the final model given the opposite pattern was evident in the univariate analysis). In all models, there was a significant effect of time, suggesting that the odds of common mental disorders decreased after baseline. The interactions between job control and general control, between job control and gender, and between general control and gender were not significant.

Results from the sensitivity analyses

In sensitivity analyses we observed a similar pattern of results with the count outcome as when a binary outcome was adopted. A model adjusting for all covariates showed that low job control was significantly associated with greater severity of common mental disorder [incidence rate ratio (IRR) for low versus high job

Table 2. Descriptive statistics on exposures and outcome at each wave.

	All waves (observations=7644)	2000/2001 (persons=2106)	2004/2005 (persons=2095)	2008/2009 (persons=1984)	2012/2013 (persons=1459)	Any wave ^a (persons=2221)
	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)
Job control						
Low	2475 (32.4)	689 (32.7)	657 (31.4)	624 (31.5)	505 (34.6)	1221 (55.0)
Medium	2581 (33.8)	738 (35.0)	696 (33.2)	678 (34.2)	469 (32.2)	1460 (65.7)
High	2563 (33.5)	679 (32.2)	738 (35.2)	679 (34.2)	467 (32.0)	1197 (53.9)
Unknown	25 (0.3)	0	4 (0.2)	3 (0.2)	18 (1.2)	25 (1.1)
General control (ie, mastery)						
Low	1584 (20.7)	420 (19.9)	450 (21.5)	375 (18.9)	339 (23.2)	878 (39.5)
Medium	3203 (41.9)	946 (44.9)	910 (43.4)	734 (37.0)	613 (42.0)	1627 (73.3)
High	2796 (36.6)	726 (34.5)	723 (34.5)	864 (43.6)	483 (33.1)	1329 (59.8)
Unknown	61 (0.8)	14 (0.7)	12 (0.6)	11 (0.6)	24 (1.6)	59 (2.7)
Common mental disorder						
No	5884 (77.0)	1563 (74.2)	1639 (78.2)	1541 (77.7)	1141 (78.2)	2043 (92.0)
Yes	1720 (22.5)	532 (25.3)	447 (21.3)	432 (21.8)	309 (21.2)	934 (42.1)
Unknown	40 (0.5)	11 (0.5)	9 (0.4)	11 (0.6)	9 (0.6)	38 (1.7)

^a% sum for each variable is more than 100% because individuals can be included in multiple categories.

Table 3. Odds ratios (OR) and 95% confidence intervals (CI) from longitudinal random-intercept logistic regression models assessing the relationship between job control and common mental disorder

	Model 1 ^a	Model 2 ^b	Model 3 ^c	Model 4 ^d
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Job control				
Low	4.33 (3.41–5.51) ^e	2.41 (1.90–3.07) ^e	2.06 (1.56–2.72) ^e	2.04 (1.53–2.73) ^e
Medium	2.37 (1.9–2.96) ^e	1.81 (1.45–2.26) ^e	1.66 (1.32–2.09) ^e	1.59 (1.26–2.00) ^e
High (reference)	1.00	1.00	1.00	1.00
General control (ie, mastery)				
Low	19.36 (14.96–25.05) ^e	15.8 (12.14–20.56) ^e	6.06 (4.45–8.27) ^e	5.94 (4.32–8.18) ^e
Medium	3.38 (2.72–4.20) ^e	3.00 (2.40–3.74) ^e	1.82 (1.43–2.30) ^e	1.84 (1.44–2.34) ^e
High (reference)	1.00	1.00	1.00	1.00
Low control in any wave	3.26 (2.53–4.20) ^e		0.94 (0.70–1.25)	0.87 (0.66–1.17)
Mean of mastery across waves	0.64 (0.61–0.66) ^e		0.76 (0.72–0.80) ^e	0.80 (0.76–0.84) ^e
Job demands				
Low (reference)	1.00			1.00
Medium	1.31 (1.03–1.67) ^f			1.39 (1.08–1.79) ^f
High	2.33 (1.83–2.96) ^e			2.38 (1.85–3.07) ^e
Job insecurity				
Low (reference)	1.00			1.00
High	2.19 (1.82–2.63) ^e			1.46 (1.20–1.77) ^e
Sex				
Male (reference)	1.00			1.00
Female	1.66 (1.29–2.13) ^e			1.40 (1.10–1.77) ^g
Partner status				
No partner	1.51 (1.19–1.92) ^g			1.16 (0.91–1.47)
Having a partner (reference)	1.00			1.00
Education completion				
Incomplete high school (reference)	1.00			1.00
Completion of high school	0.85 (0.62–1.16)			0.92 (0.69–1.24)
Completion of tertiary study	0.55 (0.41–0.76) ^e			0.75 (0.54–1.03)
Occupational skill level				
Low	1.32 (1.01–1.71) ^f			0.73 (0.54–0.99) ^f
Medium	1.41 (1.14–1.75) ^g			1.12 (0.89–1.41)
High (reference)	1.00			1.00
Parental responsibilities				
No (reference)	1.00			1.00
Yes	0.86 (0.70–1.05)			0.95 (0.77–1.18)
Childhood adversities				
No (reference)	1.00			1.00
Yes	3.12 (2.39–4.07) ^e			2.01 (1.58–2.55) ^e
Financial hardship				
No (reference)	1.00			1.00
Yes	3.11 (2.47–3.92) ^e			1.89 (1.49–2.38) ^e
Number of chronic physical health conditions				
0 (reference)	1.00			1.00
1	1.40 (1.15–1.71) ^g			1.33 (1.09–1.63) ^g
≥ 2	1.62 (1.25–2.12) ^e			1.43 (1.09–1.86) ^g
Smoking status				
Never/past smoker (reference)	1.00			1.00
Current smoker	1.82 (1.38–2.40) ^e			1.57 (1.20–2.06) ^g
Hazardous/harmful alcohol consumption				
No (reference)	1.00			1.00
Yes	1.13 (0.80–1.58)			1.09 (0.77–1.54)
Moderate/vigorous physical exercise				
0 (reference)	1.00			1.00
<1.5	0.87 (0.67–1.12)			1.00 (0.77–1.31)
1.5–3.0	0.84 (0.64–1.11)			1.05 (0.79–1.39)
3.1–5.5	0.57 (0.44–0.75) ^e			0.82 (0.62–1.07)
>5.5	0.54 (0.40–0.71) ^e			0.78 (0.58–1.05)
Survey wave				
1 (reference)	1.00	1.00	1.00	1.00
2	0.71 (0.59–0.86) ^e	0.67 (0.55–0.81) ^e	0.67 (0.55–0.81) ^e	0.69 (0.56–0.85) ^e
3	0.77 (0.63–0.93) ^g	0.83 (0.68–1.01)	0.80 (0.65–0.97) ^f	0.82 (0.66–1.03)
4	0.71 (0.57–0.88) ^g	0.65 (0.52–0.81) ^e	0.67 (0.54–0.83) ^e	0.64 (0.49–0.84) ^g

^a Adjusted for time (survey wave). Model 1 reports the results for each variable adjusted for time only. The results on time in Model 1 are for job control.

^b Adjusted for time and general control.

^c Adjusted for time, general control, between-person terms.

^d Adjusted for time, general control, between-person terms, job, sociodemographic status and health.

^e P<0.001.

^f P<0.05.

^g P<0.01.

control was 1.19, 95% CI 1.13–1.25]. The same pattern of results was also found using a fixed-effect model (OR for low versus high job control was 2.23, 95% CI 1.59–3.13).

Discussion

The current study analyzed four waves of community cohort data to examine whether the association between control at work and risk of common mental disorder is independent of general perceptions of control. The findings show that after adjusting for the adverse effect of low perceptions of control generally, being in a job with low control was independently associated with a two-fold risk of common mental disorder. This effect remained after adjusting for between-person differences in (or susceptibility to) low control both within the workplace and more generally, job demands, job insecurity, and sociodemographic and health confounders.

The current study supports previous research indicating that low job control is an important independent predictor of poor mental health. This finding extends existing knowledge from a variety of countries, including Australia (where improvement in job control has been associated with better mental health (16), and in the UK (12, 15). More importantly, the current study makes a unique contribution by showing that the mental health impact of low job control is not simply a measurement artefact, or a subset or reflection of perceptions of low control over life circumstances more broadly. In addition, the separation of within- and between-subject effects in the present study furthers efforts to distinguish the impact of low job control from other residual confounders, including those within and outside the workplace. Together with intervention studies that have shown the mental health benefits of increasing employee control (37), our findings indicate that good job control should be a target for workplace policy and practice. The findings negate concerns that work-based change will be ineffective if broader issues within the personal lives of employees are not addressed.

There are some limitations that should be considered in the interpretation of our findings. First, the sample was recruited from Canberra and Queanbeyan, in Australia. As Canberra is a city that includes many professionals and public servants, (baseline sample comprised of 53% professionals), the results may not be replicated in samples drawn from more disadvantaged communities, nor the broader Australian population. Second, as the study only included data from the path midlife cohort (aged 40–44 at baseline), the findings may not be generalizable to other age cohorts. For example, in adolescents and young adults at the start of their working

lives, norms of job control may be lower and have less impact on mental health. Finally, as the measures of job control, mastery, and mental health were self-reported, the data may be either underreported or overreported due to recall bias.

Concluding remarks

The current study adds to the evidence that low job control is a significant, independent risk factor for common mental disorder, independent of general perceptions of control and other individual differences. The findings underscore the importance of psychologically stressful employment conditions for mental health, and the benefits of considering the workplace as a context for protecting and promoting good mental health.

Acknowledgements

The authors thank the study participants, PATH Interviewers, Karen Maxwell and Trish Jacomb, and other PATH Chief Investigators: Kaarin Anstey, Helen Christensen, Anthony Jorm, Bryan Rodgers, Andrew Mackinnon, Simon Easteal and Nicolas Cherbuin.

Conflict of interests

Authors declare no conflicts of interest.

Funding

The PATH Through Life Study was funded by the National Health and Medical Research Council (973302, 179805, 418139), and the Australian Government Agency — Safe Work Australia. It is currently managed by both the ANU and the University of New South Wales. PB was supported by ARC Future Fellowship (FT130101444) and a University of Melbourne Faculty of Medicine, Dentistry and Health Sciences Research Fellowship. The funders had no involvement in the (i) study design, (ii) the collection, analysis and interpretation of the data, (iii) the writing of the report, and (iv) the decision to submit the paper for publication.

Ethics approval

The Australian National University Human Research Ethics Committee approved the PATH Through Life Project: #M9807; #2002/190; #2006/314 and #2010/542.

References

1. Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M et al. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012 Dec;380(9859):2163–96. [https://doi.org/10.1016/S0140-6736\(12\)61729-2](https://doi.org/10.1016/S0140-6736(12)61729-2).
2. Dollard M, Bailey T, McLinton S, Richards P, McTernan W, Taylor A et al. The Australian Workplace Barometer: report on psychosocial safety climate and worker health in Australia. Magill, SA: University of South Australia; 2012.
3. Theorell T, Hammarström A, Aronsson G, Träskman Bendz L, Grape T, Hogstedt C et al. A systematic review including meta-analysis of work environment and depressive symptoms. *BMC Public Health* 2015 Aug;15:738. <https://doi.org/10.1186/s12889-015-1954-4>.
4. Stansfeld S, Candy B. Psychosocial work environment and mental health—a meta-analytic review. *Scand J Work Environ Health* 2006 Dec;32(6):443–62. <https://doi.org/10.5271/sjweh.1050>.
5. Nieuwenhuijsen K, Bruinvels D, Frings-Dresen M. Psychosocial work environment and stress-related disorders, a systematic review. *Occup Med (Lond)* 2010 Jun;60(4):277–86. <https://doi.org/10.1093/occmed/kqq081>.
6. Bonde JP. Psychosocial factors at work and risk of depression: a systematic review of the epidemiological evidence. *Occup Environ Med* 2008 Jul;65(7):438–45. <https://doi.org/10.1136/oem.2007.038430>.
7. Karasek RA. Job Demands, Job Decision Latitude, and Mental Strain: implications for Job Redesign. *Adm Sci Q* 1979;24:285–308. <https://doi.org/10.2307/2392498>.
8. Stansfeld SA, Shipley MJ, Head J, Fuhrer R. Repeated job strain and the risk of depression: longitudinal analyses from the Whitehall II study. *Am J Public Health* 2012 Dec;102(12):2360–6. <https://doi.org/10.2105/AJPH.2011.300589>.
9. Wang J, Schmitz N, Dewa C, Stansfeld S. Changes in perceived job strain and the risk of major depression: results from a population-based longitudinal study. *Am J Epidemiol* 2009 May;169(9):1085–91. <https://doi.org/10.1093/aje/kwp037>.
10. Åhlin JK, Westerlund H, Griep Y, Magnusson Hanson LL. Trajectories of job demands and control: risk for subsequent symptoms of major depression in the nationally representative Swedish Longitudinal Occupational Survey of Health (SLOSH). *Int Arch Occup Environ Health* 2018 Apr;91(3):263–72. <https://doi.org/10.1007/s00420-017-1277-0>.
11. de Lange AH, Taris TW, Kompier MA, Houtman IL, Bongers PM. Effects of stable and changing demand-control histories on worker health. *Scand J Work Environ Health* 2002 Apr;28(2):94–108. <https://doi.org/10.5271/sjweh.653>.
12. Harvey SB, Sellahewa DA, Wang MJ, Milligan-Saville J, Bryan BT, Henderson M et al. The role of job strain in understanding midlife common mental disorder: a national birth cohort study. *Lancet Psychiatry* 2018 Jun;5(6):498–506. [https://doi.org/10.1016/S2215-0366\(18\)30137-8](https://doi.org/10.1016/S2215-0366(18)30137-8).
13. Butterworth P, Leach LS, Strazdins L, Olesen SC, Rodgers B, Broom DH. The psychosocial quality of work determines whether employment has benefits for mental health: results from a longitudinal national household panel survey. *Occup Environ Med* 2011 Nov;68(11):806–12. <https://doi.org/10.1136/oem.2010.059030>.
14. Leach LS, Butterworth P, Strazdins L, Rodgers B, Broom DH, Olesen SC. The limitations of employment as a tool for social inclusion. *BMC Public Health* 2010 Oct;10:621. <https://doi.org/10.1186/1471-2458-10-621>.
15. Stansfeld SA, Fuhrer R, Shipley MJ, Marmot MG. Work characteristics predict psychiatric disorder: prospective results from the Whitehall II Study. *Occup Environ Med* 1999 May;56(5):302–7. <https://doi.org/10.1136/oem.56.5.302>.
16. Bentley RJ, Kavanagh A, Krnjacki L, LaMontagne AD. A Longitudinal Analysis of Changes in Job Control and Mental Health. *Am J Epidemiol* 2015 Aug;182(4):328–34. <https://doi.org/10.1093/aje/kwv046>.
17. Butterworth P, Leach LS, Rodgers B, Broom DH, Olesen SC, Strazdins L. Psychosocial job adversity and health in Australia: analysis of data from the HILDA Survey. *Aust N Z J Public Health* 2011 Dec;35(6):564–71. <https://doi.org/10.1111/j.1753-6405.2011.00771.x>.
18. Clark C, Pike C, McManus S, Harris J, Bebbington P, Brugha T et al. The contribution of work and non-work stressors to common mental disorders in the 2007 Adult Psychiatric Morbidity Survey. *Psychol Med* 2012 Apr;42(4):829–42. <https://doi.org/10.1017/S0033291711001759>.
19. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007 Oct;370(9596):1453–7. [https://doi.org/10.1016/S0140-6736\(07\)61602-X](https://doi.org/10.1016/S0140-6736(07)61602-X).
20. Anstey KJ, Christensen H, Butterworth P, Eastaer S, Mackinnon A, Jacomb T et al. Cohort profile: the PATH through life project. *Int J Epidemiol* 2012 Aug;41(4):951–60. <https://doi.org/10.1093/ije/dyr025>.
21. Goldberg D, Bridges K, Duncan-Jones P, Grayson D. Detecting anxiety and depression in general medical settings. *BMJ* 1988 Oct;297(6653):897–9. <https://doi.org/10.1136/bmj.297.6653.897>.
22. Kiely KM, Butterworth P. Validation of four measures of mental health against depression and generalized anxiety in a community based sample. *Psychiatry Res* 2015 Feb;225(3):291–8. <https://doi.org/10.1016/j.psychres.2014.12.023>.
23. Bosma H, Marmot MG, Hemingway H, Nicholson AC, Brunner E, Stansfeld SA. Low job control and risk of coronary heart disease in Whitehall II (prospective cohort) study. *BMJ* 1997 Feb;314(7080):558–65. <https://doi.org/10.1136/bmj.314.7080.558>.

- [org/10.1136/bmj.314.7080.558](https://doi.org/10.1136/bmj.314.7080.558).
24. Butterworth P, Leach LS, McManus S, Stansfeld SA. Common mental disorders, unemployment and psychosocial job quality: is a poor job better than no job at all? *Psychol Med* 2013 Aug;43(8):1763–72. <https://doi.org/10.1017/S0033291712002577>.
 25. Pearlin LI, Schooler C. The structure of coping. *J Health Soc Behav* 1978 Mar;19(1):2–21. <https://doi.org/10.2307/2136319>.
 26. Butterworth P, Rodgers B, Windsor TD. Financial hardship, socio-economic position and depression: results from the PATH Through Life Survey. *Soc Sci Med* 2009 Jul;69(2):229–37. <https://doi.org/10.1016/j.socscimed.2009.05.008>.
 27. Stansfeld SA, Clark C, Caldwell T, Rodgers B, Power C. Psychosocial work characteristics and anxiety and depressive disorders in midlife: the effects of prior psychological distress. *Occup Environ Med* 2008 Sep;65(9):634–42. <https://doi.org/10.1136/oem.2007.036640>.
 28. Hatch SL, Harvey SB, Maughan B. A developmental-contextual approach to understanding mental health and well-being in early adulthood. *Soc Sci Med* 2010 Jan;70(2):261–8. <https://doi.org/10.1016/j.socscimed.2009.10.005>.
 29. Parker G. Parental characteristics in relation to depressive disorders. *Br J Psychiatry* 1979 Feb;134:138–47. <https://doi.org/10.1192/bjp.134.2.138>.
 30. Rodgers B. Reported parental behaviour and adult affective symptoms. 1. Associations and moderating factors. *Psychol Med* 1996 Jan;26(1):51–61. <https://doi.org/10.1017/S0033291700033717>.
 31. Kessler RC, Magee WJ. Childhood adversities and adult depression: basic patterns of association in a US national survey. *Psychol Med* 1993 Aug;23(3):679–90. <https://doi.org/10.1017/S0033291700025460>.
 32. Henderson AS, Jorm AF, Korten AE, Jacomb P, Christensen H, Rodgers B. Symptoms of depression and anxiety during adult life: evidence for a decline in prevalence with age. *Psychol Med* 1998 Nov;28(6):1321–8. <https://doi.org/10.1017/S0033291798007570>.
 33. National Health and Medical Research Council. Australian Alcohol Guidelines: Health Risks and Benefits. Canberra: Department of Health and Ageing; 2001.
 34. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction* 1993 Jun;88(6):791–804. <https://doi.org/10.1111/j.1360-0443.1993.tb02093.x>.
 35. Marmot MG, Smith GD, Stansfeld S, Patel C, North F, Head J et al. Health inequalities among British civil servants: the Whitehall II study. *Lancet* 1991 Jun;337(8754):1387–93. [https://doi.org/10.1016/0140-6736\(91\)93068-K](https://doi.org/10.1016/0140-6736(91)93068-K).
 36. Fitzmaurice GM, Laird NM, Ware JH. *Applied Longitudinal Analysis*. 2nd Edition. New Jersey, Wiley; 2011.
 37. Egan M, Bamba C, Thomas S, Petticrew M, Whitehead M, Thomson H. The psychosocial and health effects of workplace reorganisation. 1. A systematic review of organisational-level interventions that aim to increase employee control. *J Epidemiol Community Health* 2007 Nov;61(11):945–54. <https://doi.org/10.1136/jech.2006.054965>.

Received for publication: 8 October 2019