Geochemistry and geochronology of the shallow-level La Esperanza magmatic system (Permian-Triassic), Northern Patagonia

Carmen I. Martínez Dopico^{a,*}, Mónica G. López de Luchi^a, Augusto E. Rapalini^b,
C. Mark Fanning^c, Paul Y.J. Antonio^d

^a Instituto de Geocronología y Geología Isotópica (INGEIS), Departamento de Geología, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires- CONICET, Buenos Aires, Argentina

^b Instituto de Ciencias Básicas, Aplicadas y Ambientales de Buenos Aires (IGEBA), Departamento de Geología, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires- CONICET, Buenos Aires, Argentina

^c Research School of Earth Sciences, The Australian National University, Mills Road, Canberra, Australia

^d Instituto de Astronomía, Geofísica e Ciências Atmosféricas (IAG), Universidade de São Paulo (USP), São Paulo, SP, Brazil

ARTICLE INFO

Keywords:

Patagonia
La Esperanza
Permian- Triassic SLIP
Giménez Granite
U-Pb zircon geochronology

ABSTRACT

The La Esperanza plutonic-volcanic complex is the largest Late Paleozoic-Early Triassic composite magmatic system of northern Patagonia. This paper reports new SHRIMP U-Pb zircon ages and K-Ar muscovite dating as well as whole-rock geochemical data for selected units. In addition, we present some new and reprocessed whole-rock Sr isotopic compositions. On the basis of the new and published data, three compositionally and isotopically distinct high-K magnesian calc-alkaline series were distinguished. Two of these are characterized by high Ba-Sr: (i) biotite and muscovite bearing rhyolites and granites (265 ± 2 Ma; 260 ± 2 Ma) and (ii) metaluminous amphibole-biotite bearing granodiorites (273 ± 2 Ma), monzogranites (255 ± 2 Ma), dacites (253 ± 2 Ma), and slightly peraluminous granites (dated herein as 251 ± 2 Ma). There is also a low Ba-Sr series of high-silica metaluminous rocks (granites and acid dike swarms; 250 ± 2 Ma and $\approx 244 \pm 2$ Ma). Geochemistry coupled with geochronology revealed a pulsatory multi-sourced open magmatic system with mafic magma replenishment and reactivation of crystal mushes that occurred before upward migration to upper crustal levels. Mafic magmas alternated with crust-derived magmas incrementally assembled in subvolcanic levels over 30 Ma. Zircon crystallization and mica cooling ages in the granite units allowed detection of two magmatic lulls, between 270 and 265 Ma and between 260 and 255 Ma. Both episodes coincide with a period of exhumation in upper crustal levels. The new temporal and geochemical constraints allow correlation of the La Esperanza plutonic-volcanic complex with the Los Menucos Group (258–248 Ma), encompassing a volume of magmatism comparable to a moderately sized large igneous silicic province. These mid-to-late Permian to Middle Triassic rocks record the transition between subduction-related magmatism (> 273 Ma) and post-orogenic extensional magmatism (< 250 Ma) in the Gondwana margin. Even though this magmatism would be coeval with the proposed collision of the Patagonia terrane, no expected syn-collisional magmatism or associated deformation were found in upper crustal levels. However, the different nature and melting conditions of the inferred sources of the magmas that crystallized before 270 Ma, between 265 and 260 Ma, and from 255 to 245 Ma, suggest that the La Esperanza plutonic-volcanic complex was assembled during a 30 Ma period of major plate reorganization.

1. Introduction

The igneous rocks in the La Esperanza area have received much attention since Ramos (1984) proposed the allochthonous origin of Patagonia. Their isolated location far away from already suspected Paleozoic active margins, the calc-alkaline character of the rocks and alleged Carboniferous ages (subsequently corrected to Permo-Triassic;

Pankhurst et al., 1992) together with the deformation in Late Paleozoic rocks of Sierra de la Ventana, were the three strongest arguments that the rocks in La Esperanza were an active magmatic arc with pre-to syn-collisional stages. Pankhurst et al. (2006) proposed that this magmatism was the result of a thermal anomaly in the upper plate (i.e. post-collisional magmatism) postdating the earlier Carboniferous collision of the Deseado Massif and the North Patagonian Massif (Fig. 1a). Recently,

* Corresponding author.

E-mail address: carmen.martinez.dopico@gmail.com (C.I. Martínez Dopico).

<https://doi.org/10.1016/j.jsames.2019.102347>

Received 4 May 2019; Received in revised form 2 September 2019; Accepted 2 September 2019

Available online 10 September 2019

0895-9811/ © 2019 Elsevier Ltd. All rights reserved.

Fig. 1. a) Location of the La Esperanza area in Northern Patagonia; b) Geological map of the La Esperanza area and surroundings. The yellow stars indicate the localities that were previously dated, as either U-Pb zircon SHRIMP dates or mica K-Ar cooling ages, whereas the black stars show the localities dated in this paper. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

Luppo et al. (2019) found anomalous paleomagnetic pole positions for the 265–252 Ma volcanic rocks around La Esperanza, whereas the paleomagnetic poles of rhyolite dikes dated at 244 ± 2 Ma were consistent with the pole position for Middle-Triassic in most reference paths for South America.

The aim of this paper is to present the magmatic stratigraphy, geochronological and geochemical backgrounds of a very well exposed example of a shallow composite magmatic system, the La Esperanza plutonic-volcanic complex of northern Patagonia (Llambías and Rapela, 1984; Martínez Dopico et al., 2013a,b; 2017a), to provide a framework to test future hypotheses regarding the potential cogenetic evolution of the magmas. We report new SHRIMP U-Pb zircon data and two K-Ar muscovite dating as well as WR geochemical data for selected units. In addition, we present some new and reprocessed whole-rock Sr isotopic compositions and review Hf and O (in zircon) and whole rock Nd-isotope data published for the area.

2. Geological background

The La Esperanza plutonic-volcanic complex crops out in the surroundings of La Esperanza settlement (“Estancia La Esperanza Nueva”, 19G 543350 E 5525800 UTM coordinates), in central Río Negro province, Patagonia, Argentina. The Estancia La Esperanza is located 170 km to the south of Neuquén and 60 km to the north of Los Menucos (Fig. 1a). Permian to Triassic igneous rocks are distributed in an area of over 1000 km² (conservatively), although recent regional correlations

(Luppo et al., 2018, 2019) suggest that the magmatic event could have extended over more than 3000 km² beneath large areas covered by the Cenozoic Somun Curá volcanic plateau.

The geology of the La Esperanza area was described in detail by Llambías and Rapela (1984) who proposed the terms “La Esperanza plutonic complex” and “Dos Lomas volcanic complex” to distinguish the plutonic suites from the volcanic and volcanoclastic rocks, subvolcanic leucogranites and dikes. These complexes were thought to be separated by a regional unconformity. This former stratigraphy, followed by Cucchi et al. (2001), was modified by Martínez Dopico et al. (2013b) with the hypothesis of a single magmatic plumbing system built by the assembly of magma batches and termed it the La Esperanza plutonic-volcanic complex. Rapela and Llambías (1985) and Martínez Dopico et al. (2013a,b, 2014) studied the geochemistry of the pre-Jurassic units, describing a meta- to slightly peraluminous, magnesian, high-K calc-alkaline granite series that evolved through a combination of processes such as mixing and fractional crystallization of magmas from at least two different sources (Martínez Dopico et al., 2013a,b).

The first reliable time constraints for these rocks were established with WR Rb-Sr isochrons yielding late Permian to Middle Triassic ages (Pankhurst et al., 1992), superseding previous mistaken Carboniferous ages. U-Pb zircon crystallization ages were provided by Pankhurst et al. (2006), and after by Martínez Dopico et al. (2017b) and Luppo et al. (2019). Together with thermochronological data (Martínez Dopico et al., 2013a,b; 2017b), mica cooling ages bracketed the evolution of the rocks between mid-Permian (Kungurian-Roadian) and Middle

Triassic (Anisian). Sm-Nd, Lu-Hf and O isotopic data are available from Pankhurst et al. (2006), Fanning et al. (2011) and Castillo et al. (2017).

3. La Esperanza plutonic-volcanic complex

The exposures of igneous rocks around Estancia La Esperanza cover an area larger than 2500 km², where mid-Permian to Middle Triassic intrusions and associated volcanic products were mapped as La Esperanza plutonic-volcanic complex (LEPVC) (Fig. 1b). The intrusive rocks of the LEPVC crop out as small scattered and flat bodies, whereas the volcanic and subvolcanic counterparts constitute a dome (known as a *rhyolite dome* in Llambías and Rapela, 1984), and a large mostly eroded plain (Fig. 1b). These rocks are intruded by several generations of large super-acidic dike swarms and leucogranite plugs that stand out above the regional topography as ridges. Volcanic rocks are distributed along a N-S elevated axis that extends 45 km to the south, from the puesto Llanquil to the Piche Graben (outside our area of interest). Spatial distribution is affected by E-W transform faults (Giacosa et al., 2007). The regional host rocks of the LEPVC are the Early Paleozoic phyllites and quartzitic schists of the Colo Niyeu Formation (Labudía and Bjerg, 1994; Martínez Dopico et al., 2017), which are intruded by the oldest unit of the complex, the Prieto Granodiorite, 30 km to the SW of Estancia La Esperanza.

The geochemical data provided in this paper is a compilation of the whole-rock major and minor elements of Rapela and Llambías (1985), major, minor and trace elements of Martínez Dopico et al. (2013a,b), mineral chemistry of Martínez Dopico et al. (2013a) and new data representative of the main units of the LEPVC.

3.1. Intrusive units

Geochemical and field data allow recognition of four main mappable units among the subvolcanic intrusions: 1) Prieto Granodiorite; 2) Donosa Granite and muscovite-bearing aplite dikes; 3) La Esperanza Monzogranite; and 4) Giménez Granite (Fig. 1b).

3.1.1. Prieto Granodiorite

This unit crops out to the north and south of the Estancia La Esperanza and its surrounding areas (Fig. 1b) and partially corresponds to Prieto Granodiorite of Llambías and Rapela (1984). The granodiorite dated at 273 ± 2 Ma by Pankhurst et al. (2006) belongs to this unit. The granodiorite bodies were emplaced as large horizontal layers in the border of the complex or as N-S aligned vertical feeders (magma conduits, see Martínez Dopico et al., 2017a). The Prieto Granodiorite is in fault contact with the Donosa Granite and is covered by younger volcanic and subvolcanic rocks to the east of Estancia La Esperanza. Biotite-amphibole granites, granodiorites and minor diorites (modal classification) displaying a wide range of textures and compositions make up the Prieto Granodiorite (63–70% SiO₂; $3 < K_2O < 4\%$; MgO $> 2\%$; $0.84 < ASI < 0.93$; La/Yb_N ≈ 13). The rocks are dominantly dark grey, ferromagnetic (10^{-2} SI) and exhibit medium-grained equigranular textures with variable modal contents of mafic minerals (Fig. 2a). They are characterized by euhedral zoned plagioclase (cores An_{45–55} to rims An_{20–25}; see details in Table 6 Supplementary data) (30–35% modal) together with magnesian hornblende (8–15%) and biotite (10%) clots, each of similar size, embedded in K-feldspar (20–30%) and anhedral quartz (20–25%) pools. Pyroxene was occasionally observed as cores of amphibole, particularly in coarse-grained microgranular mafic enclaves. Apatite, zircon, magnetite and titanite are accessory minerals.

3.1.2. La Esperanza monzogranite

This is a composite pluton that covers half of the studied area. Based on its texture, type of magmatic enclaves, magnetic susceptibility and age, the monzogranites and granites are distinguished from the original “Prieto Granodiorite” of Llambías and Rapela (1984). Two granitoid

facies with transitional contacts were recognised; the most extended facies is inequigranular, whereas the other encompasses porphyritic granites and crops out exclusively around puesto Calvo. The inequigranular facies crops out in the easternmost sector of La Esperanza as a subhorizontal layer or sill, whereas in the west, the spatial distribution of its outcrops reflects the NNW-SSE and NW-SE anisotropies of the basement (Martínez Dopico et al., 2017a). This facies comprises grey, variably ferromagnetic (10^{-3} – 10^{-2} SI), inequigranular granodiorites and monzogranites (63–68% SiO₂; $3 < K_2O < 4\%$; MgO $< 2\%$; $0.90 < ASI < 1.13$; $20 < La/Yb_N < 30$) in which euhedral plagioclase (or K-feldspar) is embedded in a matrix of granitic composition. It is characterized by euhedral zoned plagioclase (cores An_{35–45} to rims An_{20–25}; Table 6 Supplementary data, 30–45% modal), euhedral biotite and Mg-hornblende clots (color index $< 18\%$), poikilitic filiform perthitic K-feldspar (20–30%) and anhedral quartz (20–30%). Allanite, zircon and titanite are accessory minerals. In the Arroyo del Corral (Fig. 1b), the inequigranular monzogranites of La Esperanza are observed intruding the Donosa Granite. Close to puesto Calvo, K-feldspar megacrysts are developed, amphibole disappears and the color index drops to $< 5\%$ (porphyritic granite; 68–71% SiO₂; $3 < K_2O < 4\%$; MgO $< 1.2\%$; $0.9 < ASI < 1.0$; $K \approx 10^{-3}$ SI), and porphyritic microgranular mafic enclaves are also present (Fig. 2b). In the field, these granitoids are isotropic. However, in several localities north of puesto Donosa and to the east of puesto Linconao very fine-grained rocks exhibit macroscopic magmatic foliation due to a weak shape-preferred orientation of ferrosilicates or plagioclase crystals: in such rocks, the color index rises to 25%. Rounded mafic microgranular enclaves are very abundant and display two types: (i) up to 15 cm long rounded and very dark, porphyritic (plagioclase) enclaves (Fig. 2b) and (ii) small fine-grained equigranular enclaves. The mineralogical composition of the enclaves is identical to that of the host. The inequigranular monzogranite facies was dated as 255 ± 2 Ma using U-Pb SHRIMP methodology on magmatic zircons (Martínez Dopico et al., 2017a). This age agrees with K-Ar mica cooling ages and stratigraphic relations (Fig. 1b); see Martínez Dopico et al. (2017b) for further discussion.

3.1.3. Donosa Granite

This unit is the core of the complex. It crystallized at 260 ± 2 Ma (Martínez Dopico et al., 2017a) and comprises a pink euhedral K-feldspar megacrystic granite. Its discontinuous outcrops describe a large N-S elongated body with NW-SE and SW-NE, fault-related, rectilinear borders. Several E-W and SW-NE dextral faults dismember the pluton and there are isolated outcrops along the E-W and NE-SW valleys to the west of Estancia La Esperanza. The Donosa Granite is in fault contact with the Prieto Granodiorite in its northern and western margins and it is covered by the Collinao Dacite to the east. Donosa granites exhibit oversized pink euhedral K-feldspar megacrysts up to 15×4 cm. Field measurements indicate a dominant NNW-SSE orientation of megacrysts with a slight NW plunge. Excluding this magmatic lineation, the main N-S body is mostly isotropic. However, along the Arroyo del Corral valley, locally ductile magmatic shear bands and porphyritic synmagmatic dikes are common. These bands are defined by the alignment of biotite and pseudotachylite material that crosscut the rock and create areas of orthogneiss. Some K-feldspar megacrysts display a domino alignment when associated with shear band sense of movement. These deformational features were usually detected close to the contact with the more dioritic components of the Prieto Granodiorite. Other evidence of brittle and ductile deformation such as autobrecciation and development of decimetric zones of orthogneiss were found. Fine-grained pink muscovite-bearing leucogranite dikes are associated with this unit (e.g. sample DZ35a; Table 3).

The most outstanding feature of the Donosa granites ($K \approx 10^{-3}$ – 10^{-4} SI; 71–76% SiO₂; $3.4 < K_2O < 4.3\%$; MgO $< 1\%$; $0.95 < ASI < 1.10$) is the presence of oversized K-feldspar megacrysts. They occur as euhedral perthitic microcline locally showing macroscopic Carlsbad twinning, sodium-rich rims and poikilitic inclusions of

Fig. 2. Plutonic units of the LEPVC a) Prieto Granodiorite at locality 19G 539094 m E, 5524100 m S, note the absence of enclaves; b) Plagioclase porphyritic magmatic microgranular enclave in La Esperanza granodiorites at 539934 m E, 5521493 m S; c) Giménez Granite (equigranular texture) in the locality type, nearby puesto Giménez (536362 m E, 5519878 m S); d) Megacrystic facies of Giménez granites in the locality dated herein (534601 m E, 5525913 m S). Donosa Granite and Calvo Granite are not shown here; the reader is referred to [Martínez Dopico et al. \(2017a\)](#).

quartz. The matrix is monzogranitic and displays equigranular coarse-to-medium grained texture that consists of quartz (20–30% modal), plagioclase (30–40%), perthitic microcline (25–35%) and biotite (< 10% modal). Apatite, zircon, monazite, magnetite and muscovite (along borders and cleavage traces of biotite) are accessory minerals. Plagioclase crystals are mostly subhedral, complexly twinned, and show zoning from core (An_{30-20}) to rim (An_{20-15}) ([Table 6 Supplementary material](#)). Quartz crystals are subhedral to anhedral, normally forming interstitial clusters and sometimes develop parallel subgrains and chessboard patterns. Biotite is subhedral and greenish-brown, locally intergrown with magnetite. Magnetite is mostly euhedral and free of alteration, although in some sampling sites hematite replacement is widespread. Tourmaline traces were found where decimetric magmatic shear zones develop. Except for local deformation along the E-W Arroyo del Corral, the overall magmatic texture remains unmodified. The transition to the dominant undeformed rock is sharp. In outcrop scale subrounded granite biotite-rich clots occasionally appear. Well-rounded microgranular granodiorite enclaves are rare.

3.1.4. Giménez Granite

This unit crops out to the north of the puesto Giménez area and shows transitional contacts with the inequigranular monzogranites of the La Esperanza Monzogranite. It is typically a fine-grained equigranular pink to pinkish grey granite (70–76% SiO_2 ; 3 < K_2O < 5%; MgO < 1%; $0.90 < ASI < 1.15$) ([Fig. 2c](#)). A geochemical transition between these two units was proposed by [Rapela and Llambías \(1985\)](#). Rocks are slightly ferromagnetic ($K \approx 10^{-3} - 10^{-4}$ SI). Close to the puesto Giménez, the granite contains K-feldspar (30–40%), plagioclase (25–35%) and quartz (25–30%) with variable amounts of biotite (5–15%). Magnetite, apatite, zircon and scarce monazite are accessory minerals. K-feldspar is euhedral, characteristically zoned and perthitic.

Plagioclase crystals are subhedral and zoned, with core composition ranging from An_{42} to An_{38} and strongly sodic rims (< An_{10}), sometimes myrmekitic. Quartz is anhedral and exhibits low-to-medium temperature deformation features such as chessboard extinction, subgrain development and bulging structures. Biotite is subhedral, brown-yellow colored. In the field, these granites are mostly isotropic, although NW-SE (130–150°) trending magmatic foliation with variable dip was observed in a few localities.

Further north of the puesto Giménez, around Ruta Nacional 67, the granite develops a porphyritic texture with K-feldspar megacrysts up to 12 cm-long embedded in a granite matrix (megacrystic granite facies; [Fig. 2d](#)). The matrix is composed of K-feldspar, plagioclase, quartz, biotite, minor quantities of muscovite and accessories such as zircons, magnetite and apatite. The K-feldspar megacrysts are euhedral and locally exhibit shadow tails. They are inhomogeneously distributed but locally concentrated with their major axes defining steep lineations. K-feldspar megacrysts are thought to concentrate through mechanical accumulation during constricted magma flow in pipes ([Paterson et al., 2005](#)). Flattened equigranular biotite rich-schlieren and plagioclase aggregates up to 10 cm are abundant. The schlieren consist mostly of biotite and andalusite (plus cordierite) surrounded (replaced) by muscovite (chlorite). Andalusite might be considered as a restitic or peritectic phase whereas individual crystals of euhedral biotite apparently crystallized from the melt. Foliated metamorphic enclaves are also observed. Further northeast, these rocks are transitionally interfingering with a grey banded porphyritic muscovite-biotite granodiorite in syn-magmatic folds. The attitude of these banded rocks is NNW-SSE, dipping to the west. The megacrystic granites are sharply interlayered and intruded by a grey biotite-bearing tonalite (herein named Álvarez Tonalite; 66–70% SiO_2 ; 2 < K_2O < 3%; MgO < 2%; $1.03 < ASI < 1.12$) in which magmatic foliation is depicted by the alignment of plagioclase

Fig. 3. a) Banded rhyolite from the top of the of the rhyolite dome; b) Syn-sedimentary folding in the Llanquil Formation; c) Collinao porphyritic dacite; d) Coarse-grained leucocratic Calvo granites in the surroundings of puesto Calvo (see Fig. 1b).

crystals.

3.2. Extrusive units

Among (semi)extrusive rocks formerly known as the Dos Lomas plutonic-volcanic complex (Llambías and Rapela, 1984), seven units have been distinguished: 1) Pailemán Rhyolite composed of rhyolite, ignimbritic rhyolite and vitrophyre, 2) Pailemán breccias, and 3) Llanquil Fm. making up the rhyolite dome; 4) andesite dikes; 5) Collinao Dacite made up of (a) ignimbritic dacite and rhyodacite, and (b) dacite; 6) Calvo Granite; and 7) Acidic dikes of (a) aplite and leucogranite and (b) porphyritic rhyolite (Fig. 1b).

3.2.1. Pailemán Rhyolite and related lithologies

The Pailemán Rhyolite crops out in the central to northern parts of a dome-shaped body (the “rhyolite dome” of Llambías and Rapela, 1984) ca. 8 km northeast of Estancia La Esperanza. The unit corresponds to the mid and upper sections of high-viscosity rhyolitic lava plugs and black and purple vitrophyres (Fig. 3a). The Pailemán rhyolites are porphyritic with a high proportion of phenocrysts to matrix. They consist of subhedral to anhedral quartz, subhedral and partially broken crystals of zoned plagioclase, variable amounts of K-feldspar, muscovite, ferromagnesian minerals (biotite) and opaque minerals set in a devitrified groundmass. Feldspar phenocrysts are perthitic and partially replaced by sericite and clay minerals whereas ferromagnesian minerals are replaced by chlorite aggregates. A representative rock of the rhyolite dome, a muscovite-bearing rhyolite was dated with U-Pb SHRIMP zircon data at 264 ± 2 Ma (Pankhurst et al., 2006).

The Llanquil Formation is an epiclastic volcanic microbreccia (Fig. 3b) associated with the volcanic dome that comprises a series of very small outcrops in the western slope of the southernmost part of a SSE-NNW valley that dissects the Pailemán dome. It is crosscut by felsic microgranite dikes.

In the eastern part of the rhyolite dome, an autoclastic rhyodacitic breccia (herein called Pailemán Breccia) overlies the Prieto Granodiorite. It is a porphyroclastic rock with a high proportion of phenocrasts to matrix (> 60%) which shows several episodes of

brecciation. Phenocrasts show great variation in size, reaching up to a centimetre; the phenoclastic fraction is composed of mainly subhedral rounded and partially broken quartz and subhedral non-zoned plagioclase, biotite (their proportions are widely variable) and very minor amount of opaque minerals. Alteration is pervasive and mainly affects plagioclase and biotite. Secondary muscovite was seen. The groundmass is inhomogeneous showing different degrees of crystallinity with perlitic, felsitic and microgranular textures. There are amphibole bearing-xenoliths with pilotaxitic groundmass textures.

3.2.2. Collinao Dacite

This comprises a mainly greenish-grey porphyritic dacite with occasional eutaxitic textures ($62\text{--}69\%$ SiO_2 ; $\text{K}_2\text{O} < 4\%$; $5 < \text{MgO} + \text{Fe}_2\text{O}_{3t} < 8\%$; $0.8 < \text{ASI} < 1.1$) (Fig. 3c). The original name of the unit given by Llambías and Rapela (1984) was Collinao dacitic ignimbrite. However, examination in more than 15 localities only revealed one with ignimbritic texture, so we consider it more accurately referred to as a lava rather than an ignimbrite. The phenocryst mineral assemblage is dominated by plagioclase (up 50 vol % in the coarser grained types), amphibole, biotite, Fe–Ti oxides, quartz and apatite (trace). The groundmass varies from fine-grained micropoikilitic crystalline with the development of snowflake textures and interstitial quartz to felsitic–microgranular in the subvolcanic types. Groundmass minerals are feldspar and quartz with small grains of amphibole, biotite and opaque minerals, titanite and apatite microlites. Amphibole is the dominant ferromagnesian mineral in the quartz-poor dacites. Close to the puesto Collinao (Fig. 1b), the groundmass comprises plagioclase microlites set in the poikilitic groundmass and the only ferromagnesian phase is biotite. Apatite is abundant and appears either as stubby prisms when associated with amphibole or as acicular crystals in the groundmass. The unit is dated at 253 ± 2 Ma (U-Pb SHRIMP in zircon; Luppo et al., 2019). In the southernmost outcrops of the Collinao Dacite, a subvolcanic facies is exposed comprising a light grey equigranular felsic rock ($74\text{--}77\%$ SiO_2 ; $\text{K}_2\text{O} < 5\%$; $5 < \text{MgO} + \text{Fe}_2\text{O}_{3t} < 2\%$), in which tourmaline aggregates are present. Biotite microgranite dikes with localized coarse-grained pods with quartz–tourmaline miaroles are observed intruding the Collinao Dacite.

3.2.3. Andesitic dikes

All the pre-253 Ma magmatic units are crosscut by < 1 m thick, porphyritic to aphyric, dark grey mafic dikes (49–55% SiO₂; K₂O < 2%; Fe₂O_{3t} + MgO < 15%) which, in turn, are crosscut by acid microgranite dikes (Luppo et al., 2019). Mafic dikes with a preferred NE-SW trend (Fig. 1b) intrude the Prieto Granodiorite, Donosa Granite and the base of the rhyolite dome. Most are aphyric and composed of euhedral plagioclase in a widely altered fine-grained hornblende and biotite bearing matrix. Other mafic dikes show E-W trends, such as those that intrude the La Esperanza monzogranites and exhibit up to 1 cm euhedral hornblende crystals embedded in pilotaxitic to intersertal matrix (lamprophyres).

3.2.4. Calvo Granite and acidic dike swarms

The Calvo Granite is a silica-rich granite (U-Pb SHRIMP zircon age of 250 ± 2 Ma, Pankhurst et al., 2006) associated with aplite and concentric and radial leucogranitic dikes that constitute the last plutonic activity exposed in the area (246 ± 2 Ma; 244 ± 2 Ma U-Pb SHRIMP zircon ages in Pankhurst et al., 2006; Luppo et al., 2019). The granite crops out as rounded plugs in the surroundings of La Esperanza. It is a medium to coarse grained equigranular pink leucogranite (75–80% SiO₂; 4 < K₂O < 5%; MgO + Fe₂O_{3t} < 3%; ASI < 1.1) (Fig. 3d). In the southern N-S trending outcrop, the granite contains occasional K-feldspar megacrysts and/or the amount of biotite increases, sometimes resembling the Donosa granites. It contains quartz, K-feldspar and plagioclase, minor amounts of biotite and magnetite (< 5%). Garnet has been described to the northeast of Estancia La Esperanza but was not seen by us.

Fine- to very fine-grained granite and banded and/or porphyritic rhyolite dikes intrude all the previous units of this complex (either plutonic or volcanic). In the easternmost sector they are mostly E-W, but in the surrounding of puesto Calvo they radiate following a NE-SW and E-W trend (Fig. 1b). In most cases their length is more than 5 km long and their width between 1 and 2 m, but porphyritic rhyolite dikes are thicker than the rest and normally form fold hinges in the field (Fig. 1b). The crosscutting relations between dikes of different orientations are not clear since they change trend imitating a ring structure around the subvolcanic Calvo Granite and thus seem to be related to its emplacement.

4. Geochronology

We selected three samples to complete the geochronological framework already published in Pankhurst et al. (2006), Martínez Dopico et al. (2017b) and Luppo et al. (2019) (Table 1).

4.1. U-Pb SHRIMP zircon dating

A sample of porphyritic biotite granite 10 km west of Estancia La Esperanza (megacrystic facies of the Giménez Granite) (Z9; 19G 534600 m. E 5525900 m. S) was collected for U-Pb dating on zircons (Fig. 1). Once extracted, zircons were mounted in epoxy together with chips of the Temora standard zircon (Black et al., 2003). Cathodoluminescence (CL) images were used to reveal the internal structures of the polished grains (Fig. 4a). Most zircons were euhedral to subhedral, predominantly translucent and exhibited magmatic zonation (Fig. 4a). Analyses were performed in the SHRIMP RG (Sensitive High-Resolution Ion Microprobe) at the Research School of Earth Sciences of the Australian National University in Canberra according to the protocol outlined in Williams (1998). Data were reduced, statistically analysed and plotted with Isoplot 4.1 (Ludwig, 2005). Common Pb corrections were made using ²⁰⁷Pb (Williams, 1998). Most zircons were largely euhedral to subhedral, predominantly translucent and exhibit magmatic zonation (Fig. 4a). Twenty-two zircons (27 pits in total) were studied in detail, concentrating on rims but testing potential inheritance in some core-rim pairs (grains #2, 14, 17, 18 and 20; Table 2). We only

found Middle Devonian inheritance in only one core of zircon grains. The rest of the cores have shown to have ages between 250 and ca. 260 Ma, produced during the same magmatic event. One result (for #14.1) was discarded due to high ²⁰⁶Pb (> 5%). The other zircons frequently exhibited parallel zoning and all had Th/U ratios > 0.1 and were attributed to a magmatic origin as proposed by Maas et al. (1992). In the Tera-Wasserburg diagram, a tight coherent group on Concordia indicates the crystallization age of this facies of the pluton: 21 ²³⁸U/²⁰⁶Pb ages gave a weighted mean of 251 ± 2 Ma (95% confidence level) with a MSWD of 1.4 (Fig. 4b and c). Six zircons ages were ignored because they were either isolated younger (i.e., grain #6 242 Ma) or somewhat older ages (i.e., grains #3, 20 ≈ 260 Ma), or they had high common lead (i.e., grain #14).

4.2. K-Ar white mica dating

Two samples were selected for K-Ar mineral dating from fresh and well exposed areas of acidic and basic dikes from the La Esperanza area. These were Z67b, a coarse-grained variety of the Donosa Granite in Estancia La Esperanza (5443311 m E 55527092 m S) and D35a, a medium-grained muscovite-bearing leucogranite dike intruding Prieto Granodiorite 4 km southeast of Puesto Llanquil (554280 m E 5531743 m S) (Table 3).

The samples were crushed in a steel jaw crusher and sieved to isolate the 300–400 µm size fractions. After magnetic separation, muscovite grains from each sample were handpicked under a binocular microscope to obtain homogeneous microcrystalline separates. The purity of the mineral separates is > 99%. Clean micas were ground in pure alcohol to remove the possible altered rims that might have suffered a loss of potassium or argon. The K-Ar methodology used is described in Solé (2009). For potassium analysis X-ray fluorescence with high-dilution fused pearls was used to minimize the matrix effects (Solé and Enrique, 2001). For argon determination a CO₂ laser system was used for sample fusion, followed by gas purification (Solé, 2009). Measurements were performed in a MM1200B noble gas mass spectrometer (Instituto de Geología, UNAM). Age errors are reported at the 2-σ level (Table 3).

Both samples yielded muscovite-cooling ages of 260 ± 6 Ma, showing rapid cooling of the Donosa Granite, which has a U-Pb zircon crystallization age of 260 ± 2 Ma (Martínez Dopico et al., 2017b).

5. Geochemistry

We present the first complete comparative WR chemical analysis of major, minor and trace elements for the Prieto Granodiorite, La Esperanza Monzogranite, Donosa Granite, Collinao Dacite, Paileman Rhyolite, Calvo Granite, and two acidic dikes from the area of La Esperanza (Table 4). The chemical analyses were conducted at ActLabs, Ontario, Canada under their WRA + trace 4Lithoresearch program. Determinations were performed on samples of up to 5 kg that were screened for alteration in hand specimen and thin sections. Samples were broken using an iron hammer and reduced using an iron-plated jaw crusher and agate mills. X-ray fluorescence spectrometry (XRF) was used for major elements and ICP-MS for trace elements. The precision is better than 1% for major elements and better than 5% for trace elements. The collection was complemented with previous major and some minor elementary data of Rapela and Llambías (1985).

Information was processed using GCDKit 5.0 (Janousek et al., 2006) and the scripts of Janousek et al. (2016). In the following text, compositions are expressed recalculated to 100% anhydrous to minimize the effect of alteration on the samples. However, because samples show scatter in alkali contents, we tested the potential effect of secondary processes using the chemical weathering index of Ohta and Arai (2007) (not shown). The great majority of samples follow the predicted igneous trend, except those of the Las Pampas rhyolite and some of the rhyolite dome of Rapela and Llambías (1985), which follow the linear trends for

Table 1
Summary of zircon crystallization ages (U-Pb SHRIMP) and biotite/white mica cooling ages in the La Esperanza area. References: P92, P06- Pankhurst et al. (1992, 2006); C13, C17, C19- Martínez Dopico et al. (2013a,b, 2017a, Martínez Dopico et al., 2017; L18, L19 -Luppo et al. (2018, 2019).

Sample	Lithology	Method	Material	Reference	Interpretation	MDA	Age (Ma)	Depo age 528 ± 2 Ma	Observation
Cambrian Metamorphic Basement									
Colo Niyen	MB	U-Pb SHRIMP	Zrc	C17c	MDA	MDA	530		
Prieto Granodiorite	La Esperanza Plutono- Volcanic Complex								
	LES-118	U-Pb SHRIMP	Zrc	P06	Crystallization		273		
Paillemán Rhyolite	Equigranular Bt-Amp Granodiorite	K-Ar	Bio	C13	Cooling at 300°C		259		Next to the contact with Donosa Granite
	LES-125	U-Pb SHRIMP	Zrc	P06	Crystallization		265		
Donosa Granite	Rhyolite	U-Pb SHRIMP	Zrc	C17	Crystallization		260		
	Z67	Rb-Sr	WR	P92	Cooling at 500°C		259		Next to the contact with Calvo Granite
	Megacrystic Granite	K-Ar	Ms	C17	Cooling at 450°C		265		
	Z67-b	K-Ar	Bio	C13	Cooling at 300°C		237		
	Z12	U-Pb SHRIMP	Zrc	C17	Crystallization		255		
La Esperanza Monzogranite	Inequigranular Bt-Amp Monzogranite	K-Ar	Bio	C17	Cooling at 300°C		248		
	Z20	K-Ar	Bio	C17	Cooling at 300°C		251		
Collinao Dacite	Inequigranular Bt-Amp Monzogranite	U-Pb SHRIMP	Zrc	L19	Crystallization		253		
	E5	K-Ar	Bio	C17	Cooling at 300°C		251		
Giménez Granite	Rhyodacite	U-Pb SHRIMP	Zrc	this study	Crystallization		251		
	Z9	Rb-Sr	WR	P92	Cooling at 500°C		258		
	Metatexite	K-Ar	Bio	C17	Cooling at 300°C		232		
	Z305b	U-Pb SHRIMP	Zrc	P06	Crystallization		250		
Calvo Granite	Banded Granite	Rb-Sr	WR	P92	Cooling at 500°C		239		
	LES-119	K-Ar	Bio	C17	Cooling at 300°C		232		
	Leucogranite	U-Pb SHRIMP	Zrc	P06	Crystallization		250		
	Leucogranites	Rb-Sr	WR	P92	Cooling at 500°C		239		
Acidic dike	Ms-granite	K-Ar	Ms	this study	Crystallization		260		
	D35a	U-Pb SHRIMP	Zrc	P06	Crystallization		246		
Acidic dike	Volcanic rock with devitrified pumice	U-Pb SHRIMP	Zrc	L19	Crystallization		244		
	DZ3	U-Pb SHRIMP	Zrc	L19	Crystallization		244

Fig. 4. U-Pb zircon dating of the Giménez Granite a) Cathodoluminescence images of the dated zircons. Note the inherited crystal-cores and dissolved areas; b) U-Pb concordia plot; c) Weighted average $^{238}\text{U}-^{206}\text{Pb}$ (SHRIMP) crystallization age for Giménez Granite. A summary of the analytical data is in Table 2.

weathering of rhyolites.

Four unpublished Sr isotopic determinations were taken from the data repository of the Instituto de Geocronología y Geología Isotópica (INGEIS): analytical procedures for these were described in Caminos et al. (1988). Initial $^{87}\text{Sr}/^{86}\text{Sr}$ ratios from the INGEIS data repository, Cingolani et al. (1991) and Pankhurst et al. (1992, 2006) have been recalculated at their respective zircon U-Pb crystallization age and using the ^{87}Rb decay constant of Rotenberg et al. (2012) (Table 5). Whole-rock Nd data from Pankhurst et al. (2006) and zircon-Hf isotopic parameters from Fanning et al. (2011) and Castillo et al. (2017) were recalculated using Sm-Nd CHUR of $^{147}\text{Sm}/^{144}\text{Nd} = 0.1960 \pm 4$ and $^{143}\text{Nd}/^{144}\text{Nd} = 0.512630 \pm 11$, and for Lu-Hf CHUR of $^{176}\text{Lu}/^{177}\text{Hf} = 0.0336 \pm 1$ and $^{176}\text{Hf}/^{177}\text{Hf} = 0.282785 \pm 11$ of Bouvier et al. (2008).

5.1. Geochemical variations

Using the TAS diagram of Middlemost (1994), the Prieto Granodiorite and most of the rocks of the La Esperanza Monzogranite are classified as granodiorites whereas rocks of the base of Collinao Dacite plot as dacites, and andesites. The remaining intrusions and volcanic rocks are compositionally granites or rhyolites.

In the $\text{SiO}_2\text{-K}_2\text{O}$ diagram of Peccerillo et al. (1976), most units follow a wide high K- calc-alkaline trend (Fig. 5a) and overlap with that of the partial melts derived from tonalites (Roberts and Clemens, 1993). Basic dikes show a transitional character to the medium-K calc-alkaline series (Fig. 5a). In the granite classification of Frost et al. (2001) all the samples belong to the magnesian series. In the modified alkali-lime index diagram (Fig. 5b) the majority of the rocks (Prieto, La Esperanza, Giménez, Donosa and Collinao units) span the calc-alkalic spectrum with most of them between calc-alkalic and calcic fields. On the other hand, the granites of Calvo intrusion and some acidic dikes are alkali-calcic to calc-alkalic. According to Frost et al. (2001), plutons that are

genetically related should plot within the same series. In Fig. 5b rocks of Calvo Granite seem to be independent from the rest because at the same SiO_2 values, there is a difference in the alkaline-lime ratio. In both bivariate plots is clear that the trends of Prieto Granodiorite, La Esperanza Monzogranite and Collinao Dacite are not subparallel to the proposed lines of chemical differentiation through fractional crystallization within a series (Roberts and Clemens, 1993). Therefore, a significant open-system process such as magma mixing or multiple magma sources may be invoked to explain the variability in the composition of the units. The rocks are mostly metaluminous to slightly peraluminous ($\text{ASI} \approx 1.1$) and show variable degrees of alumina saturation within the same unit (Fig. 5c). Minor elements important for granites are Ba, Sr and Rb since they replace Ca and K in feldspars. Comparing the Ba-Sr-Rb compositions of the units (Fig. 5d), two groups can be traced: a group characterized by $\text{Ba}/\text{Sr} \gg 1$, $\text{Rb}/\text{Ba} \geq 1$ and $\text{Ce}/\text{Sr} > 1$ consisting of the Calvo Granite and the fine-grained acidic dikes ("Low Ba-Sr granites"), and another with $\text{Rb}/\text{Ba} < 1$, $\text{Ba}/\text{Sr} \geq 1$ $\text{Sr}/\text{Rb} > 1$ and $\text{Ce}/\text{Sr} < 1$ or high Ba-Sr granites (Tarney and Jones, 1994). The high Ba-Sr rocks can be further divided by their $\text{CaO}/\text{K}_2\text{O}$ ratio: the Prieto Granodiorite, La Esperanza Monzogranite, Giménez Granite, Collinao Dacite and porphyritic rhyolite dikes have $\text{CaO}/\text{K}_2\text{O} > 0.5$, in contrast with the Donosa Granite and Pailemán Rhyolite.

Harker diagrams show that TiO_2 , Al_2O_3 , CaO , MgO , FeO_t , and P_2O_5 describe inverse relationships relative to SiO_2 for the plutonic-volcanic suite, whereas K_2O and Na_2O increase or show no correlation with SiO_2 (Fig. 6). Al_2O_3 shows a decoupled behaviour of the high and low Sr-Ba groups with a sudden change in the negative slope, a change replicated in the rest of major element trends, particularly in Na_2O and K_2O the trending slope is inverted between the high and low Sr-Ba groups. The different trends seen with alkali-lime index (Fig. 5b) and the sudden change in the slope of the SiO_2 -major elements trends (Fig. 6) suggests that the low and high Ba-Sr groups are not related to each other by fractional crystallization and might have been sourced from different

Table 2
Summary of the U-Pb SHRIMP results of sample Z9 of Giménez Granite.

Grain.spot	U (ppm)	Th (ppm)	Th/U	²⁰⁶ Pb/ ^a (ppm)	²⁰⁴ Pb/ ²⁰⁶ Pb	f ₂₀₆ %	Total	²³⁸ U/ ²⁰⁶ Pb	1-s ±	²⁰⁷ Pb/ ²⁰⁶ Pb	1-s ±	²⁰⁶ Pb/ ²³⁸ U	Radiogenic	1-s ±	²⁰⁶ Pb/ ²³⁸ U	Age (Ma)	±
1.1	456	170	0.37	15.7	-	0.01	24.90	0.27	0.0514	0.0008	0.0004	0.0402	0.0004	0.0004	253.8	2.8 ^a	
2.1	513	230	0.45	17.3	0.000054	0.03	25.44	0.28	0.0514	0.0006	0.0004	0.0393	0.0004	0.0004	248.5	2.7 ^a	
2.2	147	90	0.61	5.1	0.000436	0.94	24.59	0.32	0.0588	0.0022	0.0005	0.0403	0.0005	0.0005	254.6	3.3 ^a	
3.1	694	348	0.50	24.3	0.000058	< 0.01	24.48	0.27	0.0513	0.0005	0.0004	0.0408	0.0004	0.0004	258.1	2.8	
4.1	639	264	0.41	21.8	0.000059	0.08	25.18	0.27	0.0519	0.0005	0.0004	0.0397	0.0004	0.0004	250.9	2.7 ^a	
5.1	423	419	0.99	14.2	-	< 0.01	25.63	0.28	0.0510	0.0006	0.0004	0.0390	0.0004	0.0004	246.7	2.7 ^a	
6.1	433	202	0.47	14.3	0.000101	< 0.01	26.09	0.29	0.0510	0.0006	0.0004	0.0383	0.0004	0.0004	242.5	2.7	
7.1	443	284	0.64	15.0	-	< 0.01	25.36	0.28	0.0512	0.0006	0.0004	0.0394	0.0004	0.0004	249.3	2.7 ^a	
8.1	253	106	0.42	8.6	0.000277	< 0.01	25.35	0.30	0.0505	0.0008	0.0005	0.0395	0.0005	0.0005	249.6	2.9 ^a	
9.1	452	196	0.43	15.1	-	0.05	25.66	0.29	0.0515	0.0006	0.0004	0.0390	0.0004	0.0004	246.3	2.8 ^a	
10.1	684	341	0.50	22.9	0.000008	0.10	25.66	0.28	0.0519	0.0005	0.0004	0.0389	0.0004	0.0004	246.2	2.7 ^a	
11.1	718	511	0.71	24.2	-	0.04	25.44	0.27	0.0515	0.0005	0.0004	0.0393	0.0004	0.0004	248.4	2.6 ^a	
12.1	585	251	0.43	20.4	0.000065	0.01	24.67	0.27	0.0514	0.0005	0.0004	0.0405	0.0004	0.0004	256.1	2.7 ^a	
13.1	653	334	0.51	22.4	-	< 0.01	25.03	0.27	0.0512	0.0005	0.0004	0.0400	0.0004	0.0004	252.6	2.7 ^a	
14.1	2412	95	0.04	101.7	0.005337	9.93	20.38	0.21	0.1315	0.0011	0.0005	0.0442	0.0005	0.0005	278.8	2.9	
14.2	28	24	0.84	1.5	-	0.04	16.33	0.31	0.0546	0.0020	0.0012	0.0612	0.0012	0.0012	383	7 Inheritance	
15.1	518	242	0.47	17.3	-	0.10	25.65	0.28	0.0519	0.0007	0.0004	0.0390	0.0004	0.0004	246.3	2.7 ^a	
16.1	253	88	0.35	8.7	0.000008	< 0.01	25.08	0.30	0.0506	0.0008	0.0005	0.0399	0.0005	0.0005	252.2	3.0 ^a	
17.1	441	189	0.43	15.1	-	< 0.01	25.03	0.28	0.0512	0.0006	0.0004	0.0400	0.0004	0.0004	252.6	2.8 ^a	
17.2	288	150	0.52	9.8	-	0.02	25.15	0.29	0.0514	0.0007	0.0005	0.0397	0.0005	0.0005	251.3	2.9 ^a	
18.1	438	163	0.37	15.0	-	< 0.01	25.10	0.28	0.0509	0.0006	0.0004	0.0399	0.0004	0.0004	252.0	2.8 ^a	
18.2	164	93	0.57	5.6	0.000113	0.28	25.08	0.32	0.0535	0.0010	0.0005	0.0398	0.0005	0.0005	251.3	3.1 ^a	
19.1	802	669	0.83	27.6	-	< 0.01	24.93	0.26	0.0510	0.0004	0.0004	0.0401	0.0004	0.0004	253.7	2.7 ^a	
20.1	601	254	0.42	21.5	0.000024	< 0.01	24.05	0.26	0.0512	0.0005	0.0005	0.0416	0.0005	0.0005	262.7	2.8	
20.2	198	149	0.75	7.0	-	< 0.01	24.36	0.30	0.0507	0.0009	0.0005	0.0411	0.0005	0.0005	259.5	3.1	
21.1	353	145	0.41	12.3	0.000088	< 0.01	24.63	0.28	0.0507	0.0008	0.0005	0.0406	0.0005	0.0005	256.8	2.9 ^a	
22.1	736	364	0.50	25.7	0.000126	0.23	24.62	0.27	0.0532	0.0005	0.0004	0.0405	0.0004	0.0004	256.0	2.8 ^a	
										± include std: ie							
										external							
										Age							
										251.1							
										1.5							
										0.72							
										1.8							
										MSWD = 1.4 for 21 of 27 areas							
										analysed							

Notes:

1. Uncertainties given at the one s level.
2. Error in Temora reference zircon calibration was 0.41% for the analytical session (not included in above errors but required when comparing data from different mounts).
3. f₂₀₆% denotes the percentage of common Pb (206 Pb).
4. Correction for common Pb for the U/Pb data has been made using the measured ²³⁸U/²⁰⁶Pb and ²⁰⁷Pb/²⁰⁶Pb ratios following Tera and Wasserburg. (1972) as outlined in Williams (1998).

^a Used for age calculation.

Table 3

Summary of the K-Ar results of muscovite of samples Z67 (Donosa Granite) and D35 (Ms-bearing leucogranite dike).

Sample	Unit	Mineral	K ₂ O [Wt. %]	40Ar * [%]	Age [Ma]	2s-Error [Ma]
Z67b	Donosa Granite	Ms	8.41	98.3	259.6	5.8
D35a	Ms-Acid Dike	Ms	8.18	98.7	259.8	6.2

protoliths. The variation of Sr versus SiO₂ (Fig. 7) shows a clockwise-trend that also illustrates the behaviour of Ba, Zr, La and Cerium. It is clear from these bivariate diagrams that there are two different segments within each trajectory and a large scatter in the values. In the Prieto Granodiorite, the La Esperanza Monzogranite and the Collinao Dacite they have a crude positive correlation with SiO₂, while the granites of Donosa and Calvo and the rhyolite dikes show a vertical scatter. Rb and K/Ba plots vs SiO₂ portray anti-clockwise pattern compared to Sr. The variation of Ba, K, Rb within the low-Ba-Sr series (Calvo Granite and fine-grained acidic dikes) suggests K-feldspar fractionation. It is clear from Fig. 7 that there is large scatter of Sr and Ba contents in Donosa granites (385–680; 400–1500 ppm, respectively) within a very restricted range of SiO₂ values (73.4–76.0% weight). The variance of these elements is also large within Giménez granites (265–580; 840–1850 ppm) but within a larger silica range, and with negative correlation between the variables that is not seen in the Donosa Granite. In granites, alkaline earth elements like Sr and Ba are usually mostly contained in feldspars replacing Ca in plagioclase and K in K-feldspar. In both cases, there are rough positive correlations between Sr and atomic Ca but not between Ba and atomic K, suggesting fractionation of Ca-plagioclase. In both cases the span of values in trace elements is rather large to be explained solely by this process. According to Clemens et al. (2010) and Villaros et al. (2009) some alkaline-earth elements, such as Sr, in S-type granites are decoupled from the related major elements because their concentrations in the melts are controlled by local variations in the trace-element contents of the source rocks. Moreover, Sr content and its isotopic composition in the different batches of melt would also depend whether equilibrium melting occurs (see Bea, 1996; Farina et al., 2014, among others). As pointed out by Clemens et al. (2009), melts that escape rapidly from the source have less chance to attain equilibrium and may result in lower trace element concentrations compared to those that remain in contact with the source for a long time.

Rare earth element (REE) contents decrease from La to Lu decrease with SiO₂, yielding very low values for the Calvo and Donosa granites and the fine-grained rhyolite dikes (< 150 ppm). All units are enriched in light rare earth elements (LREE) and display negatively sloping chondrite-normalized REE patterns (Boynnton, 1984) (Fig. 8). The La Esperanza Monzogranite, Collinao Dacite and porphyritic rhyolite dikes (Z19) show very similar patterns with La/Yb_N = 12.2–17.2 and weak negative Eu anomalies (Eu/Eu* = 0.70–0.76), whereas the older Donosa Granite and the rhyolite dome have more fractionated patterns with La/Yb_N = 20.7–27.7 and weak Eu depletion (Eu/Eu* = 0.69–0.76). In contrast, the low Ba-Sr rocks (the Calvo Granite and acidic dikes) show ‘wing-shaped’ chondrite-normalized REE patterns with large Eu negative anomalies (Eu/Eu* = 0.36–0.39).

On multi-element diagrams normalized to primitive-mantle-concentrations, the main rocks of the suite exhibit high contents of LILE such as Cs, Rb, Ba, Pb, Th, U and LREE but negative Nb, Ta and Ti, P anomalies. Coupled with the overall high-K character, this suggests a strong crustal characteristic of the magmas, with some mafic to intermediate compositions that should have been derived from lower crustal sources (see below isotopic compositions). The low Ba-Sr acidic granites and rhyolite dikes (75–80 SiO₂ wt%) have a markedly different pattern with larger negative anomalies in Ba, Sr, P, and positive anomalies in HREE (Dy, Y, Yb, Lu). Compared to the upper Continental Crust (Taylor

and McLennan, 1995), the low Sr-Ba granites and dikes exhibit dramatically lower contents of Ba, Sr, P and Ti in comparison with the high Ba-Sr rocks. Another thing to note in Fig. 8 is that Donosa granites and the rhyolites of the dome have low values of HREE in comparison to the low Ba-Sr rocks at similar SiO₂ contents (i.e. 75% wt). This also suggests that the high and low Sr-Rb rocks cannot be derived from a single parental magma.

5.2. Isotope compositions

The ‘High Ba-Sr’ suite shows a narrow range of mean initial ⁸⁷Sr/⁸⁶Sr values from 0.7065 (Prieto Granodiorite) and 0.7071 (La Esperanza Monzogranite) to 0.7076 (Donosa Granite). The initial ⁸⁷Sr/⁸⁶Sr ratios of the volcanic rocks of the rhyolite dome and Collinao Dacite are slightly higher than the plutonic rocks, ranging from 0.7070 to 0.7084, possibly indicating a more crustal (or hybridized) component than the Prieto Granodiorite (Table 5). Similarly, εNdt values for Calvo Granite and volcanic rocks are largely more negative (−7.5 < εNdt < −5.8) than those of Prieto Granodiorite (−4.8). Zircon εHft values for the ‘High Ba-Sr’ rocks (−2.9 to −4.6; Prieto Granodiorite) are lower than those of Calvo Granite (−5.6 to −9.0) or rhyolite dikes (−7.4 to −8.7). The initial ⁸⁷Sr/⁸⁶Sr ratios of the Calvo Granite at 250 Ma are variable (0.6921–0.7064) ranging to impossibly low values that could indicate post-crystallization open system behaviour. Only initial ⁸⁷Sr/⁸⁶Sr ratio of one sample, LE132, seems more realistic with values of 0.7064.

Fig. 9 shows the initial ⁸⁷Sr/⁸⁶Sr ratio plotted against SiO₂ for the studied rocks. The main rocks of the high Ba-Sr series, the La Esperanza Monzogranite and Giménez Granite, define a horizontal array, suggesting differentiation from a single parent magma, or at least a common source. The initial ⁸⁷Sr/⁸⁶Sr ratios of the Donosa Granite are also coincident with this array, but the age difference between the granites of Giménez (ca. 250 Ma) and Donosa (260 Ma) precludes a comagmatic relationship, although again a similar source would be possible. Another observation is that the span of initial ⁸⁷Sr/⁸⁶Sr ratios of the rhyolite dome is similar to that of Donosa Granite (at identical wt % SiO₂) but unrelated to that of the low Sr-Ba series (considering initial ⁸⁷Sr/⁸⁶Sr values of ca. 0.7064). Epsilon Ndt values of Donosa Granite are very negative (−9.8), indicating that the crustal component would be more important than that of the Prieto Granodiorite and the rhyolite dome. However, given the wide spread of εHft zircon values of the rhyolite dome (−3.8 to −8.7), more data is necessary to extract a conclusion out of this unit.

6. Discussion

6.1. Temporal and spatial evolution of the magmatic system

The oldest unit of the area, the Prieto Granodiorite, crystallized at 273 ± 2 Ma (mid Permian, Cissuralian/Guadalupian) according to a U-Pb SHRIMP zircon dating (Pankhurst et al., 2006) for a sample located less than 1 km north of the fault contact with the Donosa Granite (Figs. 4 and 10). The crystallization age of the La Esperanza Monzogranite (formerly part of Prieto Granodiorite) is 255 ± 2 Ma (late Permian, Wuchiapingian/Changhsingian), with biotite cooling ages of 248 ± 4 Ma and 251 ± 6 Ma (to the east and west of study area) which reinforce this age (Martínez Dopico et al., 2017b). Initial Sr and Hf isotope ratios and geochemical features suggest a lower crustal source for both units, consistent with their compositional and textural similarities (Martínez Dopico et al., 2017a). However, the degree of hybridization of the La Esperanza Monzogranite is less than that of the Prieto Granodiorite: it exhibits abundant evidence for mingling whereas the latter is much more homogeneous suggesting less viscosity contrast (i.e. similar temperatures) between interacting melts. On the other hand, emplacement depth differences between them were negligible according to Al-in-hornblende pressure estimates of < 2 Kbar (Martínez

Table 4

Typical whole-rock major and trace elements geochemical data of several units of the La Esperanza plutonic-volcanic complex.

Sample	Z19	DZ3	LZ13	D55	D39	Z302	Z67	Z1
Locality (UTM m 19G)	555273 E 5506688 S	552708 E 5523514 S	547571 E 5535526 S	552755 E 5521004 S	552977 E 5532826 S	546643 E 5525866 S	544311 E 5527092 S	549840 E 5515478 S
Lithology	porphyritic rhyolite	aphyric rhyolite	leucogranite	hb- bt granite	rhyolite	dacite	megacrystic granite	bt-hb granodiorite
Unit	acidic dike	acidic dike	Calvo Granite	La Esperanza Monzogranite	Paileman dome	Collinao Dacite	Donosa Granite	Prieto Granodiorite
SiO ₂	69.52	78.60	75.89	69.20	74.20	60.06	74	64.08
Al ₂ O ₃	14.42	12.00	12.28	14.70	14.10	16.88	14.29	15.09
Fe ₂ O _{3t}	3.56	2.15	0.73	3.97	1.97	6.02	1.89	5.41
MnO	0.06	0.02	0.08	0.07	0.03	0.05	0.029	0.075
MgO	1.20	0.02	0.11	1.21	0.37	2.41	0.43	2.22
CaO	2.41	0.26	0.62	2.85	1.31	4.78	1.72	3.73
Na ₂ O	3.74	3.27	3.87	3.52	3.34	3.73	3.99	3.42
K ₂ O	4.03	4.97	5.04	3.83	3.89	2.04	3.77	4.83
TiO ₂	0.42	0.09	0.10	0.42	0.18	0.84	0.23	0.57
P ₂ O ₅	0.13	0.01	0.01	0.13	0.03	0.23	0.09	0.15
LOI	1.10	0.12	0.46	0.58	0.89	2.16	0.7	0.86
Total	100.60	101.51	99.19	100.48	100.31	99.19	101.14	100.44
Sc	7	2	4	7	2	13	3	10
Be	2		6	n.d.	n.d.	3	3	3
V	62	57	9	64	19	137	20	99
Cr	20	20	20	20	10	20	20	30
Co	7	1	1	9	2	10	2	11
Ni	20	7	20	7	5	20	20	20
Cu	60	15	10	14	4	10	10	10
Zn	40	9	30	40	31	40	30	60
Ga	17	13.9	19	17.5	18	23	20	19
Ge	1	5	3	5	5	2	1	1
As	5	1.9	5	0.9	1.9	31	5	5
Rb	140	256	348	171.5	158	90	144	225
Sr	402	49.1	35	388	450	593	476	392
Y	14	17.9	17	18.6	8.7	24	6	18
Zr	146	88	80	136	109	204	112	173
Nb	12	18.5	29	15.8	9.4	14	9	11
Mo	2	2	2	1	1	2	2	2
Ag	0.5	0.5	0.5	0.5	0.5	1.1	0.5	0.5
In	0.2	0.006	0.2	0.014	0.005	0.2	0.2	0.2
Sn	3	3	4	3	2	5	2	4
Sb	0.5	0.2	0.5	0.14	0.71	1.3	0.5	0.5
Cs	2.2	1.71	6.9	8.33	9.12	4.5	3.1	9.1
Ba	1102	102.5	110	808	1110	542	913	800
La	38.4	32	22.4	42	24.6	41.9	24.4	50.3
Ce	73.6	59.2	42.3	77.5	46.1	87	47.4	99.1
Pr	7.47	5.42	3.87	8.17	4.91	9.76	4.95	10.1
Nd	26.1	15.8	11.4	27.7	17.7	36.7	17.2	34.8
Sm	4.5	2.58	1.9	4.66	2.66	6.8	3	6
Eu	0.93	0.29	0.2	0.95	0.49	1.35	0.58	1.13
Gd	3.1	2.05	1.5	3.34	1.79	5	1.8	4.2
Tb	0.5	0.4	0.3	0.54	0.29	0.8	0.3	0.6
Dy	2.7	2.24	1.9	3.02	1.48	4.3	1.3	3.3
Ho	0.5	0.56	0.4	0.61	0.29	0.8	0.2	0.6
Er	1.5	1.69	1.5	1.71	0.76	2.3	0.7	1.8
Tm	0.22	0.28	0.31	0.26	0.12	0.34	0.1	0.28
Yb	1.5	2.13	2.6	1.71	0.8	2.3	0.6	1.9
Lu	0.25	0.32	0.48	0.25	0.12	0.36	0.1	0.3
Hf	4.5	3.6	4	3.9	3.2	5.4	3.3	5.5
Ta	1.3	1.5	2.9	1.6	0.9	1.3	1.1	1.2
W	2	2	1	3	7	1	1	5
Tl	0.9	0.08	1.7	0.3	0.13	0.5	0.6	1.2
Pb	18	16	37	21	14	14	22	29
Bi	0.4	0.17	0.4	0.19	0.06	0.5	0.4	0.6
Th	14.6	25.6	33.3	19.1	9.11	23.6	8	28.5
U	4.6	2.28	5.5	3.28	2.11	5.8	1.9	7.6
Eu/Eu*	0.76	0.39	0.36	0.74	0.69	0.71	0.76	0.69
La/YbN	17.26	10.13	5.81	16.56	20.73	12.28	27.42	17.85
La/SmN	5.37	7.8	7.42	5.67	5.82	3.88	5.12	5.27
Sum REE	161.27	124.96	91.06	172.42	102.11	199.71	102.63	214.41

Dopico et al., 2013a). The Prieto Granodiorite is exposed as the host of the rhyolite dome whose basal unit (i.e. crystalloclastic rhyolite) was dated at 264 ± 2 Ma. However, the rhyolite dome is a complex sub-volcanic/volcanic edifice whose age range has not yet been accurately constrained and could extend to Lower Triassic times. Crystallization

and cooling of the Donosa Granite in the central part of the main body have been dated as 260 ± 2 Ma (late Permian, Wuchiapingian/Capitanian, U-Pb on zircon age) with a very fast cooling from 400 to 350 °C to at 259 ± 6 Ma (K-Ar on muscovite age presented here, Table 3, Fig. 10). Textural evidence of near-surface cooling is seen in the 1–2 cm

Table 5
Initial $^{87}\text{Sr}/^{86}\text{Sr}$ ratios from the IR INGHS data repository, P92 Pankhurst et al. (1992, 2006) and C92 Cingolani et al. (1991) have been recalculated at their respective zircon U-Pb crystallization age and using the ^{87}Rb decay constant of Rottenberg et al. (2012). Hf and oxygen-isotopic data belongs to Castillo et al. (2017).

Sample	Unit	Lithology	Crystallization Age	SiO_2	Sr	Rb	$^{87}\text{Rb}/^{86}\text{Sr}$	$^{87}\text{Sr}/^{86}\text{Sr}$	2s	ϵ_{Sr} (WR)	Initial $^{87}\text{Sr}/^{86}\text{Sr}$	Reference	ϵ_{Nd} (WR)	Initial $^{143}\text{Nd}/^{144}\text{Nd}$	Reference	ϵ_{Hf} (Zrc)	$\delta^{18}\text{O}\text{‰}$	Reference
LES-125	Rhyolite dome	rhyolite	264	73.0	123	427	6.500	0.731292	n.d.	3.8	0.707275	P06	-5.8	0.511992	P06	-3.8 to -8.7	6.6 to 7.0	Ca17
LE 101	Rhyolite dome	rhyolite	264	70.9	167	394	6.8300	0.73450	.050	4.3	0.709264	IR						
LE 102	Rhyolite dome	rhyolite	264	70.8	171	473	6.4600	0.73230	.050	3.9	0.708431	IR						
LE 146	Rhyolite dome	rhyolite	264	73.0	193	171	2.4600	0.71610	.050	1.6	0.707011	IR						
LES-118	Calvo Granite	leucogranite	250	75.9	35	377	29.096	0.807041	n.d.	14.6	0.705246	P06	-7.5	0.511922	P06	-5.6 to -9.0	4.4 to 5.8	Ca17
LE 185	Calvo Granite	leucogranite	250	78.2	13.4	228	50.0800	0.87735	.015	24.5	0.702135	P92*						
LE 126A	Calvo Granite	leucogranite	250	76.8	7.4	232	93.1900	1.01821	.020	44.5	0.692178	P92*						
LE 126B	Calvo Granite	leucogranite	250	76.8	7.4	225	91.0400	1.01775	.020	44.5	0.699238	P92*						
LE 127	Calvo Granite	leucogranite	250	77.3	15.8	241	44.6900	0.85906	.015	21.9	0.702704	P92*						
LE 131	Calvo Granite	leucogranite	250	75.0	12.9	224	51.0300	0.88252	.0150	25.3	0.703990	P92*						
LE 132A	Calvo Granite	leucogranite	250	74.9	90.5	256	8.1990	0.73505	.010	4.3	0.706365	P92						
LE 132B	Calvo Granite	leucogranite	250	74.9	88.8	250	8.1680	0.73503	.010	4.3	0.706454	P92						
LES-120	Donosa Granite	granite	260	73.0	480	139	6.998	0.710168	n.d.	0.8	0.707628	P06	-9.8	0.511794	P06			
LE 120	Donosa Granite	granite	260	73.1	532	133	0.7249	0.71022	.010	0.8	0.707582	P92						
LE 121	Donosa Granite	granite	260	74.3	377	137	1.0554	0.71149	.010	1.0	0.707646	P92						
LE 134	Donosa Granite	granite	260	75.2	321	186	1.6788	0.71374	.010	1.3	0.707627	P92						
LE 137A	Gimenez Granite	granite	251	75.3	239	209	2.5378	0.71645	.010	1.7	0.707535	P92						
LE 137B	Gimenez Granite	granite	251	75.3	233	204	2.5420	0.71646	.010	1.7	0.707526	P92						
LE 139	Gimenez Granite	granite	251	71.4	475	153	0.9351	0.71056	.010	0.9	0.707275	P92						
LES-119	Prieto Granodiorite	granodiorite	273	67.0	440	161	1.065	0.710794	n.d.	0.9	0.706725	P06	-4.8	0.512033	P06	-2.9 to -4.6	6.0 to 7.3	Ca17
CON-88.48	Prieto Granodiorite	granodiorite	273	67.0	445	175	1.1390	0.71084	.010	0.9	0.706488	C91						
LE 165	Prieto Granodiorite	granodiorite	273	66.8	421	183	1.2570	0.71145	.010	1.0	0.706649	P92						
LE 169	Prieto Granodiorite	granodiorite	273	67.3	418	179	1.2384	0.71122	.010	1.0	0.706487	P92						
LE 152	Collinao Dacite	dacite	252	65.5	445	206	1.3000	0.71310	.050	1.2	0.708515	IR						
LE 99 A	La Esperanza	granodiorite	255	68.1	542	179	0.8688	0.71028	.010	0.8	0.707179	P92						
LE 99 B	La Esperanza	granodiorite	255	68.1	426	129	0.8731	0.71024	.010	0.8	0.707122	P92						
LE 130	Monzogranite	granite	255	71.8	323	164	1.4651	0.71224	.010	1.1	0.707006	P92						
LE 136	Monzogranite	monzogranite	255	70.9	429	102	0.6905	0.70969	.010	0.7	0.707226	P92						
465	La Esperanza	granodiorite	255	67.0	394	195	1.4333	0.71193	.010	1.1	0.706814	P92*						
LES-122	Rhyolite dike	rhyolite	245	?												-7.4 to -8.7	5.1 to 6.6	Ca17

* not considered. n.d. not reported.

Fig. 5. Major element geochemistry of the La Esperanza plutonic-volcanic complex (this paper; Rapela and Llambías, 1985). a) Binary diagram of $(K_2O + Na_2O)$ vs SiO_2 (Middlemost, 1994) shows that most units follow a wide high K- calc-alkaline trend. The black and white stars represent average I- and S-type granite compositions reported by Chappell and White (1983) whereas fields a, b and c show the trends of differentiation of three examples of plutonic magma series: transitional, calc-alkaline and high-K rocks, respectively shown as in Roberts and Clemens (1993); b) Binary modified-alkali-lime index (Frost et al., 2001) vs SiO_2 ; note that the Prieto, La Esperanza, Giménez, Donosa and Collinao units extend over the calc-alkalic and calcic fields whereas Calvo Granite and some acidic dikes belong to the alkali-calcic to calc-alkalic domains; c) Alumina saturation indexes $Al_2O_3/[(CaO + Na_2O + K_2O) - (1.67 \cdot CaO)]$ vs $Al_2O_3/(Na_2O + K_2O)$ (mol. %); and d) Sr-Rb-Ba Ternary diagram showing high and low Ba-Sr groups.

euhedral crystals of quartz in localities around Arroyo del Corral (Fig. 1b), which indicate very fast cooling of the unit and its very shallow character. Muscovite-bearing microgranite dikes (Table 3; sample D35b) could be associated either with the Pailemán Rhyolite or with the Donosa Granite (both rocks contain primary muscovite), and they intrude the Prieto Granodiorite.

Slightly after intrusion of the La Esperanza Monzogranite, the Collinao Dacite was extruded at 253 ± 2 Ma (Luppo et al., 2019) along a N-S trending ridge. The Collinao Dacite clearly overlies the main body of Donosa granites, but contact with the La Esperanza Monzogranite occurs where the granite is highly fractured and altered (Fig. 1b). The following event is the intrusion of Calvo Granite, dated at 250 ± 2 Ma by Pankhurst et al. (2006). Although there are no available

geochronological constraints on the mafic NW-SE dyke swarm, these dikes intrude the La Esperanza Monzogranite and the Giménez Granite. Therefore, these dikes are constrained as close in time to the extrusion of Collinao Dacite, after the intrusion of Giménez Granite at 251 ± 2 Ma (Fig. 1b), but before the ≈ 244 Ma acid dikes (Luppo et al., 2019). The ≈ 244 Ma acidic dikes clearly crosscut all the units of the LEPVC, following E-W, NE-SW and NW-SE trends and are geochemically related to the Calvo Granite (250 ± 2 Ma, Pankhurst et al., 2006).

6.2. The unconformity between the La Esperanza plutonic and volcanic rocks

In the first approach to the geology of the La Esperanza magmatic

Fig. 6. Major element (wt%) variation diagrams versus SiO_2 for entire chemical data set of the granitoid and volcanic groups. Yellow and pink lines indicate suggested evolutionary trends of low and high Sr-Ba groups, respectively. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

system, Llambías and Rapela (1984) and Rapela and Llambías (1985) divided the rocks around Estancia La Esperanza in two cycles, the first entirely plutonic and the second volcanic and subvolcanic including the Calvo Granite and acidic dikes. Llambías and Rapela (1984) claim “The second Cycle begins after a short erosive period that uncovers the plutonic rocks of the first Cycle and on which in almost horizontal surfaces the first ignimbritic eruptions of dacitic composition (Collinao) were deposited.” In the light of the new stratigraphic arrangement (Fig. 10), this simple subdivision does not hold anymore. There is an erosion surface between the Donosa Granite and the Collinao Dacite which is easily seen around Estancia La Esperanza. A second erosion surface might occur between the Permian rocks and the Calvo Granite

and associated dikes (< 250 Ma). In any case, in the La Esperanza area none of these erosion surfaces seem to represent more than minor adjustments associated with the evolution of the caldera rather than substantial discordances. Further south, rhyolitic ignimbrites, dacite lavas and rhyolitic tuffs at the base and middle sections of the coeval Los Menucos Group (252–258 Ma) were tilted (30° to the east) and intruded by a felsic rhyolite dike swarm equivalent to the group of acidic dikes in La Esperanza. The acidic dikes of the La Esperanza area (Fig. 1) were dated as old as 244 ± 2 Ma (crystallization age of a fine-grained rhyolite dike; Luppo et al., 2019), suggesting that regional uplift and erosion occurred between the emplacement of Calvo Granite (250 ± 2 Ma crystallization age of the northern plug of the Calvo

Fig. 7. Selected trace elements (Sr, Rb, Ba, Y, Zr, La, Ce, Lu; ppm) versus SiO_2 variation diagrams for the entire chemical data set of the granitoid and volcanic groups.

Granite; Pankhurst et al., 2006) and the diking.

6.3. Petrogenesis and magmatic kinships

The early Permian Prieto Granodiorite (273 ± 2 Ma) shares major and trace element geochemical characteristics with the late Permian monzogranites and granites of the La Esperanza Monzogranite (255 ± 2 Ma) and its extrusive counterpart, the Collinao Dacite (253 ± 2 Ma). All these rocks contain early crystallizing phases such as orthopyroxene, magnesian clinopyroxene and plagioclase, followed by

late hornblende and biotite, titanite and allanite, but lack peraluminous minerals, suggesting that their parental magmas were hydrated and metaluminous. According to the amphibole chemistry of the granodiorites and monzogranites (data in Martínez Dopico et al., 2013a,b) and following the equations of Ridolfi et al. (2010) and Ridolfi and Renzulli (2012), crystallization occurred in similar oxidizing ($f_{\text{O}_2} \approx \text{NNO}$) and hydrous conditions ($4 > \text{H}_2\text{O} > 5\%$ wt). Minimum crystallization temperatures estimated using the zircon saturation thermometer (TZr, Watson and Harrison, 1983) yielded temperatures of ca. 760 °C for Prieto and La Esperanza granodiorites and ca. 800 °C for the

Fig. 8. Chondrite-normalized REE patterns normalized to values of Boynton (1984) (left) and Spider plot of values normalized to the Upper Continental Crust of Taylor and McLennan (1995) (right).

Fig. 9. Initial $^{87}\text{Sr}/^{86}\text{Sr}$ ratio vs SiO_2 of the units of the La Esperanza plutonic-volcanic complex. Data from the INGEIS data repository, Pankhurst et al. (1992, 2006), and Cingolani et al. (1991) were recalculated at the crystallization ages using the ^{87}Rb decay constant of Rotenberg et al. (2012). Data table as supplementary material and symbols as in Fig. 7.

later porphyritic granite facies of La Esperanza Monzogranite and Collinao dacites, indicating that magma temperature increased with time. The lack of inherited zircon ages (see Martínez Dopico et al., 2017a; Luppo et al., 2019 and Table 2) and the absence of correlation between Zr and SiO_2 suggest that these are minimum temperatures for the magmas.

Initial $^{87}\text{Sr}/^{86}\text{Sr}$ ratios, mineralogy as well as crystallization ages suggest that the Giménez Granite is an evolved magma batch ($> 70\%$ SiO_2) derived from the fractionation of La Esperanza Monzogranite magmas. Major element trends show that it was formed after extensive amphibole and Ca-plagioclase fractionation. All rocks with $< 70\%$ SiO_2 show coherent Sm/Yb ratios from 3.0 to 4.5, indicating that pyroxene and mainly amphibole were present in the source. This would suggest that the first batch of melt had left hornblende and pyroxene as major

residual phases at high T and low P melting conditions of a mafic metaigneous source. Whole-rock initial $^{87}\text{Sr}/^{86}\text{Sr}$ (0.7064–0.7067; Table 5), Nd ($\epsilon\text{Nd} = -4.8$) and zircon Hf isotope compositions (ϵHf of -2.9 to -4.6 ; Castillo et al., 2017) for the intermediate rocks of the Prieto Granodiorite are compatible with the interpretation that these magmas were extracted from a mafic or intermediate source with long-term crustal residence. Even though there is a 20 Ma gap between the pulses, the La Esperanza Monzogranite and Giménez Granite seem to share a common ancestry with the Prieto Granodiorite at the same crustal level. However, the higher initial $^{87}\text{Sr}/^{86}\text{Sr}$ of the more evolved members of the La Esperanza Monzogranite (porphyritic facies; 0.7070–0.7072) and Giménez Granite (0.7072–0.7075; Table 5) suggest that the felsic component increased with time. The higher abundance of mafic microgranular enclaves in the monzogranites and granites of La Esperanza in comparison with the older rocks indicates that magma hybridization was inefficient. Low zircon $\delta^{18}\text{O}$ values for the whole series of rocks of La Esperanza (4.4‰–7.3‰; Castillo et al., 2017) confirm an I-type origin (Valley, 2003). Trace element-based granite tectonic discrimination diagrams (Pearce et al., 1984; Pearce, 1996; Harris et al., 1986) also point to I-type volcanic arc sources (syn to post-collisional fields) (not shown; see Martínez Dopico et al., 2014).

Granodiorite/dacite, monzogranite and granite of the La Esperanza area yield a wide range of major and trace element concentrations, suggesting variable degrees of partial melting and probably a rapid escape of the magma that prevented further chemical equilibration with the source. The rough negative correlation between SiO_2 and FeO , MgO , MnO and TiO_2 , and La/Yb_N , Sr , Ba , K , Y and Zr indicates that the fractionation of amphibole and plagioclase might have operated for the crystallization of the Prieto Granodiorite, La Esperanza Monzogranite and Collinao Dacite intermediate magmas, as well as the Giménez Granite. In turn, biotite does not seem to have had an important role since Rb and Ba (both compatible elements in biotite but not in amphibole) do not decrease with SiO_2 or K_2O (Fig. 7). The decrease in the Ba content within Donosa would be related to the grain size of the K-feldspar crystals, and, perhaps, related to K-feldspar fractionation. Minor phases that might have controlled the REE fractionation are allanite or monazite. In turn, the middle-Permian Donosa Granite, Ms-bearing leucogranites and the rhyolites of the dome are high- SiO_2 I-type magmas that contain plagioclase as an early crystallizing phase. The growth of muscovite, acid plagioclase and subhedral quartz as well as a disharmonic growth of K-feldspar megacrysts occurred during low-nucleation high-growth rate episode at high-temperature conditions. Hf

Fig. 10. Chronostratigraphic chart of the intrusive and extrusive rocks of La Esperanza, Los Menucos (25 km to the south of La Esperanza), Yaminué (100 km to the East) and Sierra Grande areas (250 km to the East), showing crystallization (U-Pb zircon) and cooling (K-Ar and Ar-Ar mica) ages (Pankhurst et al., 2006, 2014; Chernicoff et al., 2013; Martínez Dopico et al., 2017a,b; Martínez Dopico et al., 2017; Luppo et al., 2018, 2019; González et al., 2017, this paper), and isotope parameters (Pankhurst et al., 2006; Castillo et al., 2017). Note the similarity of the crystallization-time intervals of the plutonic rocks.

data in zircons of the rhyolite dome aged between 260 and 269 Ma show a large variability ($-4 < \epsilon_{\text{Hf}} < 11$; Castillo et al., 2017, Table 5), suggesting either a mixing of sources (mid crustal vs lower crustal) or isotopic disequilibrium melting. Three arguments would preclude a potential derivation of the felsic magmas of the granites of Donosa and rhyolites of the dome from the same source as the Prieto Granodiorite. First, the large REE decoupling between the felsic Donosa Granite (average 73 wt% SiO_2) and the rocks of the rhyolite dome (74 wt% SiO_2) and the intermediate rocks of the La Esperanza Monzogranite and Giménez Granite (72–75 wt% SiO_2). Second, the different whole-rock trace element compositions (high vs low Ba-Sr) with similar initial Sr ratios (Table 5). And third, the Nd isotope difference between Prieto Granodiorite ($\epsilon_{\text{Nd}} = -4.8$) and Donosa Granite ($\epsilon_{\text{Nd}} = -9.8$). Thus, the Prieto Granodiorite and the La Esperanza Monzogranite could have a mixed crustal - mafic source whereas the Donosa Granite and the Pilemán Rhyolite have a stronger crustal signature. On the other hand, there is no overlap in Sr, Nd and scant overlap in Hf isotopic composition between the < 250 Ma units, the Calvo Granite and rhyolite dikes, and any other group of rocks. The non-disturbed initial $^{87}\text{Sr}/^{86}\text{Sr}$ ratios for the early Triassic rocks (< 250 Ma) are low (0.7052–0.7064 at 75% SiO_2), coupled with mantle-like O^{18} values (4.4–6.6‰) for zircon with $-5.6 < \epsilon_{\text{Hf}} < -9.0$ units giving the rocks a distinctive signature compared to the older rocks.

6.4. Inferences for the tectonic setting and connections with other plutonic or plutonic-volcanic complexes in Patagonia and surroundings

Pankhurst et al. (2006) proposed that the genesis of these calc-alkaline high-K rocks was related to a crustal thickening in the upper plate that followed the ca 320–305 Ma continental collision of the De-sado Massif and the North Patagonian Massif. The high-K character should not be interpreted as indicating a particular tectonic environment, but rather as a product of the mineralogical and elementary content of mafic sources combined with different degrees of hybridization with felsic materials (e.g., mixing and mingling processes). A primary interpretation of the chemical and isotopic features of these rocks would be that the suitable protoliths for I-type granitic magmas could be related to a previous nearby subduction. Although the volume of mafic rocks within the age range of La Esperanza plutonic-volcanic complex related to the high-K rocks is not significant (e.g., mafic microgranular enclaves and basic dikes), mafic underplating could be invoked as a source of the heat responsible for the magma generation. An alternative process allowing heat in the lower crust is the juxtaposition of the asthenospheric mantle against the base of the crust (i.e. detachment of a subducting slab) and does not require a continental collision. An element that we should start looking at when further isotopic data becomes available is the Hf-isotopic compositions of the

inherited zircons aged between 300 and 280 Ma. The ϵ_{Hf} values of two zircon rims (inherited) aged 302 and 290 Ma (grains 3.2 and 11.2 of Fanning et al., 2011) of the Calvo Granite (low crystallization $\text{T}_{\text{Zr}} \approx 725$ Ma) are -3.7 and -3.8 (recalculated after Fanning et al., 2011) whereas those of igneous zircon in the Prieto Granodiorite vary between -2.9 and -4.6 units. This suggests that zircons aged between 300 and 270 Ma are showing magmas with more positive Hf signatures than those of the latest Permian. At this age, c. 300 Ma, massive diorite bodies were dated in the proto-Andean open margin of Gondwana at the same latitude of Estancia La Esperanza, such as the Rahue diorite (WR-mineral Rb-Sr composite isochrons 296 ± 2 Ma, 300 ± 3 Ma; Lucassen et al., 2004) and, further west, the Coastal Batholith (Deckart et al., 2014). Whole-rock initial $^{143}\text{Nd}/^{144}\text{Nd}$ and zircon initial $^{177}\text{Hf}/^{178}\text{Hf}$ ratios also point towards a common origin for the Prieto Granodiorite and other older-than-260 Ma plutonic complexes such as Quintuleu Granodiorite in the Mamil Choique complex (281 ± 2 Ma; $-3.6 < \epsilon_{\text{Hf}} < -4.2$, recalculated after Fanning et al., 2011) in the southwest of the North Patagonian Massif and the granodiorites of the Navarrete plutonic complex (282 ± 2 Ma, Pankhurst et al. (2006); $-3.5 < \epsilon_{\text{Hf}} < 2.9$, recalculated after Fanning et al. (2011); see discussion in Martínez Dopico et al. (2011, 2017a) and Castillo et al. (2017)) (Fig. 10). In turn, for the magmatism after 260 Ma in La Esperanza area, represented by Donosa Granite, La Esperanza Monzogranite and Calvo Granite, isotope data show that the felsic component is higher in the magmatic precursors than those previously molten. The volume of this second stage is larger than the first and corresponds to a major tectonic change with respect to the older conditions. Recent dating and stratigraphic studies (Luppo et al., 2018, 2019) suggest that the plutonic-volcanic event in La Esperanza area is synchronous and geochemically comparable with the dacites, rhyolites and rhyolitic and dacitic ignimbrites represented by the northernmost expression of the Los Menucos Group (258–252 Ma) (Fig. 10). We propose that La Esperanza-Los Menucos magmatic system meets the criteria proposed by Bryan et al. (2002) to consider it as a “Silicic Large Igneous Province” (SLIP): 1) its bulk magmatism would have reached at least 4000 km²; 2) extrusive volumes are larger than 8000 km³; 3) it is volumetrically dominated ($> 75\%$) by dacite to rhyolite igneous rock that have calc-alkaline I-type signatures; 4) it is lithologically dominated by silicic ignimbrites; 5) igneous activity over long periods (> 30 Ma); and 6) a spatial and temporal relationship to continental rifting, plate break-up and potentially, other mafic large igneous provinces. The La Esperanza-Los Menucos magmatic system is not only temporally related to the Choiyoi magmatic province in western Argentina (Strazzere et al., 2006; Kleiman and Japas, 2009; Rocha-Campos et al., 2011; Sato et al., 2015; Rocher et al., 2015), but also spatially associated with other late Permian to Triassic magmatic igneous complexes in Patagonia such as the Mamil Choique complex, Navarrete plutonic complex, Ramos Mejía and Yaminué igneous complexes (see Pankhurst et al., 2006, 2014; Martínez Dopico et al., 2011; 2017a). Further studies of the plutonic connections of the Choiyoi magmatic event and contemporaries should be focused on isotope, whole-rock and mineral geochemistry in order to understand the triggering factors for this regional magmatism.

7. Final remarks

La Esperanza plutonic-volcanic complex (LEPVC) and its temporal-lithological eruptive counterpart, the Los Menucos Group, meet the requirements for a Silicic Large Igneous Province (albeit rather smaller than most global examples) that is bracketed in age between 273 and 244 Ma. Several compositionally and isotopically distinct, high-K, magnesian, calc-alkaline series were distinguished: High Ba-Sr (i) metaluminous amphibole-biotite bearing granodiorites (273 ± 2 Ma); (ii) biotite and muscovite-bearing rhyolites and granites (265 ± 2 Ma; 260 ± 2 Ma) and (iv) and metaluminous mafic microgranular enclave-bearing amphibole-biotite monzogranites (255 ± 2 Ma), dacites

(253 ± 2 Ma), and slightly peraluminous granites; and, finally, Low Ba-Sr high silica metaluminous rocks (granites and acid dike swarms 250 ± 2 Ma and $\approx 244 \pm 2$ Ma represented by the subvolcanic Calvo Granite and the acidic dike swarm). The episode of shallow intermediate to acidic granite magmatism in LEPVC is now dated with the new U-Pb SHRIMP zircon data for the Giménez Granite as old as 251 ± 2 Ma. Geochemistry revealed a multi-sourced open magmatic system evidenced by rocks with very different major and trace element contents and initial Sr and Nd ratios at equivalent SiO_2 intervals. The magmatic system underwent mafic magma replenishment (shown by the La Esperanza granodiorites and their mafic microgranular enclaves and the Collinao dacites) following an episode of crust-derived magmas represented by the Donosa granites and the Pailémán rhyolites. Available crystallization ages suggest the magmatism spanned in time over 30 Ma with two magmatic lulls (Fig. 10) that coincided with exhumation in upper crustal levels.

The new temporal, lithological, isotopic and geochemical features allow correlation of the La Esperanza plutonic-volcanic complex with the Los Menucos Group, encompassing a volume of magmatism comparable to a small-sized Silicic Large Igneous Province (Fig. 10). The mid-late Permian to Middle Triassic rocks in northern Patagonia record a transition between subduction-related magmatism (> 273 Ma) associated with mafic magmatic sources with limited interaction with a felsic component, to post-orogenic extensional, mostly felsic hybridized sources (< 260 Ma) in the Gondwana margin. Even though La Esperanza – Los Menucos magmatism would be coeval with the collision of the Patagonia terrane, syn-collisional magmatism or associated deformation were not found in upper crustal levels, as expected. However, the different nature and melting conditions of the inferred sources of the magmas that crystallized before 270 Ma, between 265 and 260 Ma, and from 255 to 245 Ma, suggest that the La Esperanza plutonic-volcanic complex was assembled during a 30 Ma period of major plate reorganization.

Acknowledgements

This work was supported by the Agencia Nacional de Promoción Científica y Tecnológica (Argentina) under grants PICT2013-1162 and PICT2016-2188, PICT2016-3148. We thank an anonymous reviewer and R. Pankhurst for thorough and constructive reviews and A. Folguera for his editorial handling. C.I. Martínez Dopico would like to thank T. Luppo, G. Giordanengo, and Fundaleu (Fundación de Lucha contra la Leucemia).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jsames.2019.102347>.

References

- Bea, F., 1996. Residence of REE, Y, Th and U in granites and crustal protoliths; implications for the chemistry of crustal melts. *J. Petrol.* 37 (3), 521–552. <https://doi.org/10.1093/petrology/37.3.521>.
- Black, L.P., Kamo, S.L., Allen, C.M., Aleinikoff, J.N., Davis, D.W., Korsch, R.J., Foudoulis, C., 2003. TEMORA 1: a new zircon standard for Phanerozoic U-Pb geochronology. *Chem. Geol.* 200, 155–170.
- Bouvier, A., Vervoort, J.D., Patchett, P.J., 2008. The Lu-Hf and Sm-Nd isotopic composition of CHUR: constraints from unequilibrated chondrites and implications for the bulk composition of terrestrial planets. *Earth Planet. Sci. Lett.* 273, 48–57.
- Boynton, N.V., 1984. Cosmochemistry of the rare earth elements: meteorite studies. In: *Rare Earth Element Geochemistry*. In: Henderson, P. (Ed.), *Development in Geochemistry*. vol. 2. Elsevier, pp. 63–114.
- Bryan, S.E., Riley, T.R., Jerram, D.A., Leat, P.T., Stephens, C.J., 2002. Silicic volcanism: an under-valued component of large igneous provinces and volcanic rifted margins. In: Menzies, M.A., Klempner, S.L., Ebinger, C.J., Baker, J. (Eds.), *Magmatic Rifted Margins*. vol. 362. Geological Society of America Special Paper, pp. 99–120.
- Caminos, R., Llambías, E.J., Rapela, C.W., Párica, C.A., 1988. Late Paleozoic-Early Triassic magmatic activity of Argentina and the significance of new Rb-Sr ages from northern Patagonia. *J. South Am. Earth Sci.* 1, 137–145.

- Castillo, P., Fanning, C.M., Pankhurst, R.J., Hervé, F., Rapela, C.W., 2017. Zircon O- and Hf-isotope constraints on the genesis and tectonic significance of Permian magmatism in Patagonia. *J. Geol. Soc.* 10 1144/jgs2016-152.
- Chappell, B.W., White, A.J.R., 1983. Granitoid types and their distributions in the Lachlan Fold Belt, southeastern Australia. *Geol. Soc. Am. Mem.* 159, 21–34.
- Chernicoff, C.J., Zappettini, E.O., Santos, J.O.S., Belousova, E., McNaughton, N.J., 2013. Combined U-Pb SHRIMP and Hf isotope study of the Late Paleozoic Yaminué Complex, Río Negro province, Argentina. Implications for the origin and evolution of the Patagonia composite terrane. *Geosci. Front.* 4 (6), 37–56.
- Cingolani, C., Dalla Salda, L., Hervé, F., Munizaga, F., Pankhurst, R.J., Parada, M.A., Rapela, C.W., 1991. The magmatic evolution of northern Patagonia; New impressions of pre-andean tectonics. *Geol. Soc. Am. Spec. Pap.* 265, 29–43.
- Clemens, J.D., Darbyshire, D.P.F., 2009. Sources of post-orogenic calcalkaline magmas: The Arrochar and Garabal Hill–Glen Fyne complexes, Scotland. *Lithos* 112, 524–542.
- Clemens, J.D., Helps, P.A., Stevens, G., 2010. Chemical structure in granitic magmas – a signal from the source? *Earth Env. Sci. Trans. R. Soc. Edinburgh* 100 (1–2), 159–172. <https://doi.org/10.1017/s1755691009016053>.
- Cucchi, R., Lema, H., Busteros, A., 2001. Hoja Geológica 4169-II, Los Menucos. Provincia de Río Negro. Instituto de Geología y Recursos Minerales, Servicio Geológico Minero Argentino. *Boletín* 265, 67pp. Buenos Aires.
- Deckart, K., Hervé, F., Fanning, M., Ramírez, V., Calderón, Godoy, M., 2014. U–Pb geochronology and Hf–O isotopes of zircons from the Pennsylvanian coastal Batholith, south-Central Chile. *Andean Geol.* 41, 49–82. <https://doi.org/10.5027/andgeoV41n1-a03>.
- Fanning, C.M., Hervé, F., Pankhurst, R.J., Rapela, C.W., Kleiman, L.E., Yaxley, G.M., Castillo, P., 2011. Lu–Hf isotope evidence for the provenance of Permian detritus in accretionary complexes of western Patagonia and the northern Antarctic Peninsula region. *J. South Am. Earth Sci.* 32, 485–496.
- Farina, F., Stevens, G., Gerdess, A., Frei, D., 2014. Small-scale Hf isotopic variability in the Peninsula pluton (South Africa): the processes that control inheritance of source 176Hf/177Hf diversity in S-type granites. *Contrib. Mineral. Petrol.* 168, 1065. <https://doi.org/10.1007/s00410-014-1065-8>.
- Frost, B.R., Barnes, C.G., Collins, W.J., Arculus, R.J., Ellis, D.J., Frost, C.D., 2001. A geochemical classification for granitic rocks. *J. Petrol.* 42, 2033–2048.
- Giacosa, R.E., Lema, H., Busteros, A., Zubia, M., Cucchi, R., Tommaso, D.I., 2007. Estructura del Triásico de la región norte Macizo Nordpatagónico (40°–41°S, 67°30'–69°45'O) Río Negro. *Rev. Asoc. Geol. Argent.* 62 (3), 355–365.
- González, S.N., Greco, G.A., Sato, A.M., Llambías, E.J., Basei, M.A.S., González, P.D., Díaz, P.E., 2017. Middle Triassic trachytic lava flows associated with coeval dyke swarm in the North Patagonian Massif: a postorogenic magmatism related to extensional collapse of the Gondwanide orogen. *J. South Am. Earth Sci.* 75, 134–143.
- Pearce, J.A., Harris, N.W., Tindle, A.G., 1984. Trace element discrimination diagrams for the tectonic interpretation of granitic rocks. *J. Petrol.* 25, 956–983.
- Harris, N.B.W., Pearce, J.A., Tindle, A.G., 1986. Geochemical characteristics of collision-zone magmatism. In: Coward, M.P., Ries, A.C. (Eds.), *Collision Tectonics*. vol. 19. Geological Society London Special Publication, pp. 67–81.
- Janousek, V., Farrow, C.M., Erban, V., 2006. Interpretation of whole-rock geochemical data in igneous geochemistry: introducing Geochemical Data Toolkit (GCDKit). *J. Petrol.* 47 (6), 1255–1259. <https://doi.org/10.1093/petrology/egl013>.
- Janousek, V., Moyer, J.F., Martin, H., Erban, V., Farrow, C., 2016. Geochemical Modelling of Igneous Processes – Principles and Recipes in R Language. Bringing the Power of R to a Geochemical Community. Springer-Verlag, Berlin, Heidelberg, pp. 1–346.
- Kleiman, L.E., Japas, M.S., 2009. The Choiyoi volcanic province at 34°–36°S (San Rafael, Mendoza, Argentina): implications for the late Paleozoic evolution of the south-western margin of Gondwana. *Tectonophysics* 473, 283–299.
- Labudía, C.H., Bjerg, E.A., 1994. Geología del Sector Oriental de la Hoja Bajo Hondo (39e), Provincia de Río Negro. *Rev. Asoc. Geol. Argent.* 49 (3/4), 284–296.
- Llambías, E.J., Rapela, C.W., 1984. Geología de los complejos eruptivos del Paleozoico superior de La Esperanza, provincia de Río Negro. *Rev. Asoc. Geol. Argent.* 39 (3–4), 220–243.
- Lucassen, F., Trumbull, R., Franz, G., Creixell, C., Vázquez, P., Romer, R.L., Figueroa, O., 2004. Distinguishing crustal recycling and juvenile additions at active continental margins: the Paleozoic to recent composition of the Chilean Pacific margin (36–41°S). *J. South Am. Earth Sci.* 17, 103–119.
- Ludwig, K.R., 2005. ISOPLOT/Ex A Geochronological Toolkit for Microsoft Excel. Berkeley Geochronology Center Special Publication N°1a, Berkeley.
- Luppo, T., López de Luchi, M.G., Rapalini, A.E., Martínez Dopico and, C.I., Fanning, C.M., 2018. Geochronologic evidence of a large magmatic province in northern Patagonia encompassing the Permian-Triassic boundary. *J. South Am. Earth Sci.* 82, 346–355.
- Luppo, T., Martínez Dopico, C.I., Rapalini, A.E., López de Luchi, M.G., Miguez, M., Fanning, C.M., 2019. Paleomagnetism of Permo-Triassic volcanic units in northern Patagonia: are we tracking the final stages of collision of Patagonia? *Int. J. Earth Sci.* 108 (2), 621–647.
- Maas, R., Kinny, P.D., Williams, I.S., Froude, D.O., Compston, W., 1992. The Earth's oldest known crust: A geochronological and geochemical study of 3900–4200 Ma detrital zircons from Mt Narryer and Jack Hills, Western Australia: *Geochimica et Cosmochimica Acta*. *Geochim. Cosmochim. Acta* 56, 1281–1300. [https://doi.org/10.1016/0016-7037\(92\)90062-N](https://doi.org/10.1016/0016-7037(92)90062-N).
- Martínez Dopico, C., López de Luchi, M., Rapalini, A.E., Hervé, F., Fuentes, F., Fanning, M., 2017. U-Pb SHRIMP Dating of Detrital Zircon Grains of the Colo Niyue Formation: Extending the Neoproterozoic to Early Cambrian Basins to Central Western Northern Patagonia. XX Congreso Geológico Argentino, Tucumán, pp. 6.
- Martínez Dopico, C., Lopez de Luchi, M., Rapalini, A.E., Kleinhans, I.C., 2011. Crustal segments in the North Patagonian Massif, Patagonia: an integrated perspective based on Sm/Nd isotope systematics. *J. South Am. Earth Sci.* 31, 324–341.
- Martínez Dopico, C.I., López de Luchi, M.G., Wemmer, K., Rapalini, A.E., 2013a. Composición química de biotita y hornblenda y edades de enfriamiento como indicadores de las condiciones de emplazamiento del Complejo Plutónico La Esperanza (Pérmico Superior), Macizo Norpatagónico. *Rev. Asoc. Geol. Argent.* 70, 3–15.
- Martínez Dopico, C.I., Rapalini, A.E., López de Luchi, M.G., Wemmer, K., 2013b. Assembly of shallow intrusions from multiple magma pulses in La Esperanza plutonic complex, Northpatagonian Massif, Argentina. *Latinmag Lett.* 3 OB06, 5pp.
- Martínez Dopico, C.I., López de Luchi, M.G., Rapalini, A.E., 2014. Geoquímica del Complejo Plutónico La Esperanza (Pérmico), Provincia de Río Negro. XIX Congreso Geológico Argentino.
- Martínez Dopico, C.I., López de Luchi, M.G., Rapalini, A.E., Wemmer, K., Fanning, M., Basei, M., 2017a. Emplacement and temporal constraints of the gondwanian intrusive complexes of Northern Patagonia: La Esperanza plutono-volcanic case. *Tectonophysics* 712–713, 249–269. <https://doi.org/10.1016/j.tecto.2017.05.015>.
- Martínez Dopico, C.I., Tohver, E., López de Luchi, M.G., Wemmer, K., Rapalini, A.E., Cawood, P.A., 2017b. Jurassic cooling ages in Paleozoic to early Mesozoic granitoids of northeastern Patagonia: 40Ar/39Ar, 40K–40Ar mica and U–Pb zircon evidence. *Int. J. Earth Sci.* <https://doi.org/10.1007/s00531-016-1430-0>.
- Middlemost, E.A.K., 1994. Naming materials in the magma/igneous rock system. *Earth Sci. Rev.* 37, 215–224.
- Ohta, T., Arai, H., 2007. Statistical empirical index of chemical weathering in igneous rocks. *Chem. Geol.* 18, 280–297.
- Pankhurst, R.J., Rapela, C.W., Caminos, R., Llambías, E., Parica, C., 1992. A revised age for the granites of the central Somuncura Batholith, North Patagonian Massif. *J. South Am. Earth Sci.* 5 (3/4), 321–325.
- Pankhurst, R.J., Rapela, C.W., Fanning, C.M., Márquez, M., 2006. Gondwanide continental collision and the origin of Patagonia. *Earth Sci. Rev.* 76, 235–257. <https://doi.org/10.1016/j.earscirev.2006.02.001>.
- Pankhurst, R.J., Rapela, C.W., Lopez De Luchi, M.G., Rapalini, A.E., Fanning, C.M., Galindo, C., 2014. The Gondwana connections of northern Patagonia. *J. Geol. Soc.* 171, 313e328. <https://doi.org/10.1144/jgs2013-081>.
- Paterson, S.R., Vernon, R.H., Zak, J., 2005. Mechanical Instabilities and physical accumulation of K-feldspar megacrysts in granitic magma, Tuolumne Batholith, California, USA. *J. Virtual Explor.* 18, 1.
- Pearce, J.A., 1996. Sources and settings of granitic rocks. *Episodes* 19 (4), 120–125.
- Peccerillo, A., Taylor, S.R., 1976. Geochemistry of Eocene calc-alkaline volcanic rocks from the Kastamonu area, northern Turkey. *Contrib. Mineral. Petrol.* 58, 63–81. <https://doi.org/10.1007/BF00384745>.
- Ramos, V.A., 1984. Patagonia: ¿Un continente paleozoico a la deriva? 9° Congreso Geológico Argentino. San Carlos de Bariloche, Actas 2, 311–325.
- Rapela, C.W., Llambías, E.J., 1985. Evolución magmática y relaciones regionales de los Complejos Eruptivos de La Esperanza, provincia de Río Negro. *Rev. Asoc. Geol. Argent.* 40 (1–2), 4–25.
- Ridolfi, F., Renzulli, A., 2012. Calcic amphiboles in calc-alkaline and alkaline magmas: thermobarometric and chemometric empirical equations valid up to 1130 °C and 2.2 GPa. *Contrib. Mineral. Petrol.* 163, 877–895.
- Ridolfi, F., Renzulli, A., Puerini, M., 2010. Stability and chemical equilibrium of amphibole in calc-alkaline magmas: an overview, new thermobarometric formulations and application to subduction-related volcanoes. *Contrib. Mineral. Petrol.* 160, 45–66.
- Roberts, M., Clemens, J.D., 1993. Origin of high-potassium, calc-alkaline, I-type granitoids. *Geology* 21, 825–828.
- Rocha-Campos, A.C., Basei, M.A., Nutman, A.P., Kleiman, L.E., Varela, R., Llambías, E., Canile, F.M., da Rosa, O., de, C.R., 2011. 30 million years of Permian volcanism recorded in the Choiyoi igneous province (W Argentina) and their source for younger ash fall deposits in the Paraná Basin: SHRIMP U-Pb zircon geochronology evidence. *Gondwana Res.* 19, 509–523.
- Rocher, S., Vallecillos, G., Castro de Machuca, B., Alasino, P., 2015. El Grupo Choiyoi (Pérmico temprano-medio) en la Cordillera Frontal de Calingasta, San Juan, Argentina: volcanismo de arco asociado a extensión. *Rev. Mex. Ciencias Geol.* 32 (3) 451–432.
- Rotenberger, E., Davis, D.W., Amelin, Y., Ghosh, S., Bergquist, B.A., 2012. Determination of the decay-constant of ⁸⁷Rb by laboratory accumulation of ⁸⁷Sr. *Geochim. Cosmochim. Acta* 85, 41–57.
- Sato, A.M., Llambías, E.J., Basei, M.A.S., Castro, C.A., 2015. Three stages in the Late Paleozoic to Triassic magmatism of southwestern Gondwana, and the relationships with the volcanogenic events in coeval basins. *J. South Am. Earth Sci.* 63, 48–69.
- Solé, J., 2009. Determination of K-Ar ages in milligram samples using an infrared laser for argon extraction. *Rapid Commun. Mass Spectrom.* 23, 3579–3590. <https://doi.org/10.1002/rcm.4280>.
- Solé, J., Enrique, P., 2001. X-ray fluorescence analysis for the determination of potassium in small quantities of silicate minerals for K-Ar dating. *Anal. Chim. Acta* 440, 199–205. [https://doi.org/10.1016/S0003-2670\(01\)01060-1](https://doi.org/10.1016/S0003-2670(01)01060-1).
- Strazzere, L., Gregori, D.A., Dristas, J.A., 2006. Genetic evolution of Permo-Triassic volcaniclastic sequences at Uspallata, Mendoza Precordillera, Argentina. *Gondwana Res.* 9, 485–499.
- Tarney, J., Jones, C.E., 1994. Trace element geochemistry of orogenic igneous rocks and crustal growth models. *J. Geol. Soc.* 151, 855–868.
- Taylor, S.R., McLennan, S.M., 1995. The geochemical evolution of the continental crust. *Rev. Geophys.* 33, 241–265.
- Tera, F., Wasserburg, G.J., 1972. U–Th–Pb systematics in three Apollo 14 basalts and the problem of initial Pb in lunar rocks. *Earth Planet. Sci. Lett.* 14, 281–304. [https://doi.org/10.1016/0012-821X\(72\)90128-8](https://doi.org/10.1016/0012-821X(72)90128-8).
- Valley, J.W., 2003. Oxygen isotopes in zircon- Reviews in Mineralogy and Geochemistry. 53 (1). <https://doi.org/10.2113/0530343>.
- Villaras, A., Stevens, G., Moyer, J.-F., Buick, I.S., 2009. The trace element compositions of

- S-type granites: evidence for disequilibrium melting and accessory phase entrainment in the source. *Contrib. Mineral. Petrol.* 158 (4), 543–561.
- Watson, E.B., Harrison, T.M., 1983. Zircon saturation revisited: temperature and composition effects in a variety of crustal magma types. *Earth Planet. Sci. Lett.* 64, 295–304.
- Williams, I.S., 1998. U-Th-Pb Geochronology by Ion Microprobe. In: en McKibben, M.A., Shanks III W.C., Ridley, W.I. (Eds.), *Applications of Microanalytical Techniques to Understanding Mineralizing Processes*. Littleton, Society of Economic Geologists. *Reviews in Economic Geology*. vol. 7. pp. 1–35.