Author Affiliations: Departments of Physical Medicine and Rehabilitation (Drs Peterson and Gordon) and Urology & Epidemiology (Dr Sarma), University of Michigan, Ann Arbor.

Correspondence: Dr Peterson, Department of Physical Medicine and Rehabilitation, University of Michigan Hospital and Health Systems, 325 E Eisenhower Pkwy, Ann Arbor, MI 48108 (mdpeterz@med.umich.edu).

Financial Disclosure: None reported.

1. van der Ploeg HP, Chey T, Korda RJ, Banks E, Bauman A. Sitting time and all-cause mortality risk in 222 497 Australian adults. *Arch Intern Med*. 2012; 172(6):494-500.
2. Gardiner PA, Healy GN, Eakin EG, et al. Associations between television viewing time and overall sitting time with the metabolic syndrome in older men and women: the Australian Diabetes, Obesity and Lifestyle study. *J Am Geriatr Soc*. 2011;59(5):788-796.
3. Hamilton MT, Hamilton DG, Zderic TW. Role of low energy expenditure and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease. *Diabetes*. 2007;56(11):2655-2667.
4. Katzmarzyk PT, Church TS, Craig CL, Bouchard C. Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Med Sci Sports Exerc*. 2009;41(5):998-1005.
5. Dunstan DW, Kingwell BA, Larsen R, et al. Breaking up prolonged sitting reduces postprandial glucose and insulin responses. *Diabetes Care*. 2012;35(5):976-983.

It Is Also What You Do When Sitting

The article by van der Ploeg et al¹ in the March 26, 2012, issue of the *Archives* presents further evidence of all-cause mortality risk and time spent sitting. Although the authors considered occupation as a potential confounding factor, only 1 aspect of occupation was considered, namely, time spent sitting in one's occupation. The medical literature on occupational health is replete with evidence of additional risk factors associated with one's occupation, apart from being sedentary, that adversely affect all-cause mortality. Examples of such risk factors—exclusive of workplace injuries and violence—include the following: occupational exposure to airborne particulates and incidence of stroke²; occupational stress and cardiovascular disease³; and occupational cancer-related mortality.^{4,5} Appropriate consideration for the potential confounding effect of occupational hazardous exposures is warranted, albeit challenging, when analyzing data on all-cause mortality.⁶

George J. Piligian, MD, MPH

Author Affiliation: The Mount Sinai-Irving J. Selikoff Center for Occupational & Environmental Medicine, Department of Preventive Medicine, The Mount Sinai School of Medicine, New York, New York.

Correspondence: Dr Piligian, Department of Preventive Medicine, Mount Sinai School of Medicine, One Gustave L. Levy Place, New York, NY 10029-6574 (george.piligian@mssm.edu).

Financial Disclosure: None reported.

Additional Contributions: Debra J. Milek, MD, PhD, MPH, suggested the title.

1. van der Ploeg HP, Chey T, Korda RJ, Banks E, Bauman A. Sitting time and all-cause mortality risk in 222 497 Australian adults. *Arch Intern Med*. 2012; 172(6):494-500.
2. Sjogren B, Lonn M, Feychting M, et al. Occupational exposure to particles and incidence of stroke. *Scand J Work Environ Health*. 2008;34(1):40-47.
3. Landsbergis PA, Schnall PL, Belkic KL, Baker D, Schwartz JE, Pickering TG.

The workplace and cardiovascular disease: relevance and potential role for occupational health psychology. In: Quick J, Campbell T, Lois E, eds. *Handbook of Occupational Health Psychology*. Washington, DC: American Psychological Association; 2003:265-287.

4. Health and Safety Executive. Cancer. <http://www.hse.gov.uk/statistics/causdis/cancer/index.htm>. Accessed April 17, 2012.
5. Robinson CF, Sullivan PA, Li J, Walker JT. Occupational lung cancer in US women, 1984-1998. *Am J Ind Med*. 2011;54(2):102-117.
6. Geyer S, Peter R. Occupational status and all-cause mortality. *Eur J Public Health*. 1999;9(2):114-118.

Is Sitting Harmful to Health? It Is Too Early to Say

Van der Ploeg et al¹ analyzed prospectively collected data from a large population-based cohort and found that, regardless of age, sex, body mass index, self-rated health, level of physical activity, baseline cardiovascular disease and diabetes, and other confounders, longer hours of sitting were dose-dependently associated with greater risk of all-cause mortality across an average of 2.8 years of follow-up. They described mechanisms and cited previous literature to support a hypothesis that excessive sitting, independent of physical activity, is harmful to health.

We suggest that the following possibilities must first be considered to explain the observed relationship between sitting and mortality:

1. *Imprecise measurement of self-rated health as a covariate.* Different persons may use different anchors to define good health, and these anchors may be subject to recalibration, reconceptualization, and reprioritization based on recent events and experiences,² resulting in poor health receiving a better self-appraisal than warranted. Poor insight and self-deception may also result in optimistic self-appraisals. Consequently, assertions of good health notwithstanding, greater sitting may be a result of poor health rather than its cause, and analyzing data from only those who complete a year of follow-up may not necessarily exclude all persons who were at higher risk of death at baseline.

2. *Residual confounding.* People who sit more may snack on unhealthy food and beverages, compensate by down-sizing healthy elements in their diet, work in more stressful desk jobs, or show predisposition to other behaviors and experiences that compromise health and life expectancy. None of these variables were measured and controlled for.

These possibilities were not considered by van der Ploeg et al¹ and the authors of the accompanying commentary.³ Finally, behaviorists know how difficult it is to implement even small changes, so, assuming that sitting is pathoplastic, how easy could it be to start moving for several hours a day in order to enter a less sedentary bracket of sitting behavior, as operationalized in the study?¹

Chittaranjan Andrade, MD
Praveen Fernandes, MD

Author Affiliations: Department of Psychopharmacology, National Institute of Mental Health and Neurosciences, Bangalore, India (Dr Andrade); and Department

of Psychiatry, Creighton University School of Medicine, Omaha, Nebraska (Dr Fernandes).

Correspondence: Dr Andrade, Department of Psychopharmacology, National Institute of Mental Health and Neurosciences, Bangalore 560 029, India (andradec@gmail.com).

Financial Disclosure: Dr Andrade has completed nearly 2 dozen clinical trials (investigator-initiated, industry-sponsored studies funded by Phyto-Pharma, Gufic Ltd, Zandu Pharmaceuticals, Aristo Pharmaceuticals, Cybele Laboratories, Himalaya Drug Company, Natural Remedies, Arya Vaidya Nilayam, Lupin Laboratories, Corcept Inc, and Natreon Inc; phase 2 and phase 3 clinical trials conducted for GlaxoSmithKline); has received lecture fees from Pfizer (payment made directly to a registered charity); publishes an e-newsletter, which is supported by Sun Pharmaceuticals (payments are made directly to registered charities); has provided advice to Astra Zeneca, Wyeth, Sun Pharma, Intas Pharma, and Torrent Pharma (payments either in the form of textbooks or other academic materials or directed to charities); and is principal investigator and medical monitor for a multicenter, investigator-initiated Indian investigation, funded by Natreon Inc, of the efficacy of Sensoril, a branded formulation of Ashwagandha. Dr Andrade authors and receives authorship payments for a Critical Readings in Psychiatry series (current publisher, Zydus Neurosciences).

1. van der Ploeg HP, Chey T, Korda RJ, Banks E, Bauman A. Sitting time and all-cause mortality risk in 222 497 Australian adults. *Arch Intern Med.* 2012; 172(6):494-500.
2. Howard JS, Mattacola CG, Howell DM, Lattermann C. Response shift theory: an application for health-related quality of life in rehabilitation research and practice. *J Allied Health.* 2011;40(1):31-38.
3. Dunstan DW, Owen N. New exercise prescription: don't just sit there: stand up and move more, more often. *Arch Intern Med.* 2012;172(6):500-501.

In reply

The responses to our recent article on sitting time and mortality¹ highlight the interest in, and potential importance of, sedentary behaviors and health. Our original article found that prolonged sitting was associated with increased all-cause mortality risk, independent of physical activity.¹ A study limitation was the relatively short follow-up time (mean, 2.8 years), potentially leading to confounding by occult disease. However, findings persisted after adjustments (for demographic variables, physical activity, body mass index, smoking, self-rated health, and disability), chronic disease stratification, and sensitivity analyses excluding deaths in the first year. Nevertheless, neither reverse causality due to occult disease nor residual confounding could be completely excluded.¹

The correspondents suggest additional potential sources of confounding and reverse causality from occupational factors including "exposure to airborne particulates and incidence of stroke; occupational stress and cardiovascular disease; occupational cancer-related mortality"; and poor diet. Although these factors might hypothetically play a role in the relationship between sitting and mortality, our study considered total sitting time and not "occupational sitting time." More-

over, only 49% of participants in the study were in paid work. Nonoccupational sedentary behaviors (especially television viewing) have been shown to be associated with all-cause mortality,^{1,2} making the proposed residual confounding less likely.

Furthermore, a recent similar study with an 8.5-year follow-up showed that sitting for 9 hours per day or more was associated with an all-cause mortality hazard ratio of 1.19 (95% CI, 1.12-1.27) compared with people sitting less than 3 hours per day, after extensive adjustment for confounding, including diet quality; excluding deaths in the first 3 years of follow-up suggested no evidence of reverse causality.³ Importantly, other research has suggested plausible physiological mechanisms for the detrimental effects of prolonged sitting, and findings from intervention studies indicate metabolic benefits from reductions in time spent sitting.^{1,4}

Finally, the substantial difference between the unadjusted population attributable fraction (20.9%) calculated by Peterson and colleagues in their letter, and our adjusted population attributable fraction (6.9%)¹ illustrates the importance of adjustment for confounders and suggests caution in the interpretation of unadjusted attributable risks.

Our study is one piece in the accumulating evidence that prolonged sitting may be detrimental to health. It is unclear if and when this evidence will become sufficient to warrant widespread changes in policy and behavior. We agree that achieving far-reaching change is likely to be challenging, but not impossible, especially when such important health outcomes are at stake.

Hidde P. van der Ploeg, PhD
Emily Banks, MBBS, PhD
Adrian Bauman, MBBS, PhD

Author Affiliations: Sydney School of Public Health, University of Sydney, Sydney, Australia (Drs van der Ploeg and Bauman); Department of Public and Occupational Health, VU University Medical Center Amsterdam, the Netherlands (Dr van der Ploeg); and National Centre for Epidemiology and Population Health, Australian National University, Acton, Australia and The Sax Institute, Sydney (Dr Banks).

Correspondence: Dr van der Ploeg, Department of Public and Occupational Health, VU University Medical Center Amsterdam, van der Boechorststraat 7, 1081BT Amsterdam, the Netherlands (hp.vanderploeg@vumc.nl).

Financial Disclosure: None reported.

Additional Contributions: Tien Chey, MAS, and Rosemary Korda, PhD, assisted with the preparation of this letter to the editor.

1. van der Ploeg HP, Chey T, Korda RJ, Banks E, Bauman A. Sitting time and all-cause mortality risk in 222 497 Australian adults. *Arch Intern Med.* 2012; 172(6):494-500.
2. Grøntved A, Hu FB. Television viewing and risk of type 2 diabetes, cardiovascular disease, and all-cause mortality: a meta-analysis. *JAMA.* 2011;305(23):2448-2455.
3. Matthews CE, George SM, Moore SC, et al. Amount of time spent in sedentary behaviors and cause-specific mortality in US adults. *Am J Clin Nutr.* 2012; 95(2):437-445.
4. Dunstan DW, Kingwell BA, Larsen R, et al. Breaking up prolonged sitting reduces postprandial glucose and insulin responses. *Diabetes Care.* 2012;35(5):976-983.