SMALL TIME ALMOST SURE BEHAVIOUR OF EXTREMAL PROCESSES

ROSS A. MALLER,^{*} *The Australian National University*

PETER SCHMIDLI,^{**} *The Australian National University*

Abstract

An r^{th} order extremal process $\Delta^{(r)} = (\Delta_t^{(r)})_{t \geq 0}$ is a continuous time analogue of the r^{th} partial maximum sequence of a sequence of i.i.d. random variables. Studying maxima in continuous time gives rise to the notion of limiting properties of $\Delta_t^{(r)}$ as $t \downarrow 0$. Here we describe aspects of the small time behaviour of $\Delta^{(r)}$ by characterising its upper and lower classes relative to a nonstochastic nondecreasing function $b_t > 0$ with $\lim_{t \downarrow 0} b_t = 0$. We are then able to give an integral criterion for the almost sure relative stability of $\Delta_t^{(r)}$ as $t \downarrow 0$, $r = 1, 2, \dots$, or, equivalently, as it turns out, for the almost sure relative stability of $\Delta_t^{(1)}$ as $t \downarrow 0$.

Keywords: Extremal process; upper and lower classes; relative stability; Lévy flight; Lévy dust; fractal properties.

2010 Mathematics Subject Classification: Primary 60G51; 60G55; 60G70

Secondary 60F15

1. Introduction

We study the small time (as $t \downarrow 0$) almost sure (a.s.) behaviour of an r^{th} order extremal process $\Delta^{(r)} = (\Delta_t^{(r)})_{t \geq 0}$, the continuous time analogue of the sequence of

---

^{*} Postal address: Ross Maller, Research School of Finance, Actuarial Studies and Statistics, Australian National University, Canberra, ACT, 0200, Australia.

^{*} Email address: Ross.Maller@anu.edu.au

^{**} Postal address: Peter Schmidli, Research School of Finance, Actuarial Studies and Statistics, Australian National University, Canberra, ACT, 0200, Australia.

^{**} Email address: Peter.Schmidli@anu.edu.au

r^{th} largest observations in a sequence of i.i.d. random variables. Our focus is on deriving criteria for lower and upper classes for $\Delta^{(r)}$ which generalise among others a definitive result of Klass [11, 12] relating to the large time behaviour of the partial maximum of a sequence of i.i.d. random variables. As an application, we give an integral criterion for the almost sure relative stability at 0 of $(\Delta_t^{(r)})$ as $t \downarrow 0$, i.e., having the property that there exists a nonstochastic nondecreasing function $b_t > 0$ such that $\lim_{t \downarrow 0} \Delta_t^{(r)} / b_t = 1$ a.s. The corresponding weak version: $\Delta_t^{(r)} / b_t \xrightarrow{P} 1$ as $t \downarrow 0$ for some nonstochastic nondecreasing function $b_t > 0$, with $\lim_{t \downarrow 0} b_t = 0$, is also characterised.

Our interest in this area was partly motivated by the ideas of Lévy flight and Lévy dust, which occur in various applications areas, in particular in Physics. As one interesting example, these kinds of models are being used to describe the self-similar or scale invariant spatial distribution of galaxies in the universe, and their fractal properties. The corresponding extremal processes would then refer to the distances between galaxies, or their order statistics, and the small time aspect would relate to small scale or local properties. At the other end of the spectrum, similar processes occur in the study of transmission of light or other particles through thin films at the nanoscale. We expect that our results will have application in these areas. Buchmann, Fan and Maller [2] provides some further context to the problems we consider.

The paper is structured as follows. In Section 2 we review some background in Poisson point processes required to define the r^{th} -order extremal process. Section 3 derives a characterisation of the lower class of $\Delta^{(r)}$ at 0. Section 4 gives a version for the upper class, and then combines both lower class and upper class results to give a criterion for the almost sure relative stability of $\Delta^{(r)}$. Some examples are in Section 5, and further motivating background and a brief review of related discrete time results is given in a concluding Section 6.

2. Poisson Point Processes and Extremal Processes

Herein we establish the necessary framework of Poisson point processes and extremal processes which are used throughout the sections that follow. We draw on the methodology of Buchmann et al. [2]. Although the corresponding $(\Delta_t)_{t \geq 0}$ in their paper is the

jump process of a Lévy process, their point process setup is quite general.

Let Π be a Borel measure on $(0, \infty)$, locally finite at infinity, thus having finite-valued tail function $\bar{\Pi} : (0, \infty) \rightarrow (0, \infty)$, defined by $\bar{\Pi}(x) := \Pi(x, \infty)$, $x > 0$. This is a right-continuous and non-increasing function. Let

$$\mathbb{D} = \sum_{s>0} \delta_{(s, \Delta_s)}$$

be a Poisson point process on $(0, \infty) \times (0, \infty)$ with intensity measure $dt \otimes \Pi(dx)$ defined on a probability space $(\Omega, \mathcal{F}, \mathbf{P})$. The measure Π need not be a Lévy measure but as a special case $(\Delta_t)_{t \geq 0}$ could be the jump process $(X_t - X_{t-})_{t \geq 0}$ of a Lévy process $(X_t)_{t \geq 0}$ on $\mathbb{R}$, as in [2].

We assume throughout that $\Pi\{(0, \infty)\} = \bar{\Pi}(0+) = \infty$. The number of points $(\Delta_s)_{0 < s \leq t}$ exceeding a value $\varepsilon > 0$ in time interval $[0, t]$ is distributed as Poisson with parameter $t\bar{\Pi}(\varepsilon)$, which increases without bound as $\varepsilon \downarrow 0$, so in this sense there is an arbitrarily large number of points of Δ in any finite time interval, a.s. Fix $t > 0$ and let $T^{(1)}(\mathbb{D}_t)$ be randomly chosen, independently of $(\Delta_t)_{t \geq 0}$, according to the discrete uniform distribution in the set $\{0 \leq s \leq t : \Delta_s = \sup_{0 \leq u \leq t} \Delta_u\}$, which is almost surely finite. Then define $\Delta_t^{(1)} := \Delta_{T^{(1)}(\mathbb{D}_t)}$ and the *maximum-trimmed point process* on $[0, t] \times (0, \infty)$ by

$$^{(1)}\mathbb{D}_t := \mathbb{D}_t - \delta_{(T^{(1)}(\mathbb{D}_t), \Delta_t^{(1)})}.$$

Letting $r = 2, 3, \dots$, proceed to define iteratively $T^{(r)}(\mathbb{D}_t) := T^{(1)}(^{(r-1)}\mathbb{D}_t)$ and $\Delta_t^{(r)} := \Delta_{T^{(r)}(\mathbb{D}_t)}$. The *r-fold trimmed extremal process* is then given by

$$^{(r)}\mathbb{D}_t := \mathbb{D}_t - \sum_{i=1}^r \delta_{(T^{(i)}(\mathbb{D}_t), \Delta_t^{(i)})}.$$

By this procedure we have specified a sequence of ordered pairs

$$(T_t^{(1)}, \Delta_t^{(1)}), (T_t^{(2)}, \Delta_t^{(2)}), (T_t^{(3)}, \Delta_t^{(3)}), \dots \in (0, t] \times (0, \infty),$$

in which $\{\Delta_t^{(1)} \geq \dots \geq \Delta_t^{(r)}\}$ represent the r largest order statistics of $\mathbb{D}$, sampled on the time interval $[0, t]$, possibly with ties, located at times $T_t^{(i)} = T^{(i)}(\mathbb{D}_t)$, $i \geq 1$. The

---

An alternative approach to defining an extremal process as a Markov process in terms of its finite dimensional distributions is given in Resnick [16]. Further details about the structure and properties of extremal processes are in Resnick and Tomkins [18] and Resnick [15].

process $(\Delta_t^{(r)})_{t \geq 0}$ is the r^{th} -order extremal process, $r = 1, 2, \dots$. As an extension, the notation $\Delta_I^{(r)}$ will be used to denote the r -th largest order statistic of $\mathbb{D}$ sampled on an interval I of the form (a, b) or $(a, b]$, constructed similarly.

Let $(\mathcal{F}_t)_{t \geq 0}$ be the natural filtration of $\Delta = \{\Delta_t : t \geq 0\}$ completed with the null sets of $\mathcal{F}$, so that $\mathcal{F}_0 = \mathcal{F}_{0+}$ (the “usual” hypothesis). We refer to [2] for more background information on the properties of $(\Delta_t^{(r)})_{t \geq 0}$. In particular, using the Poisson property, we can calculate

$$\mathbf{P}(\Delta_t^{(r)} \leq x) = e^{-t\bar{\Pi}(x)} \sum_{n=0}^{r-1} \frac{(t\bar{\Pi}(x))^n}{n!}, \quad t > 0, x > 0, r = 1, 2, \dots \quad (2.1)$$

The $\Delta_t^{(r)}$ are nondecreasing in $t > 0$ for each $r = 1, 2, \dots$ and nonincreasing in r for each $t > 0$, and since $\Pi\{(0, \infty)\} = \infty$, we have $\Delta_t^{(r)} > 0$ a.s. for all $t > 0$ for each $r = 1, 2, \dots$, and $\Delta_t^{(r)} \downarrow 0$ a.s. as $t \downarrow 0$ for each $r = 1, 2, \dots$.

3. Lower classes for $\Delta^{(r)}$ at 0

Throughout, b_t will be a positive non-decreasing nonstochastic function on $(0, \infty)$ with $\lim_{t \downarrow 0} b_t = 0$ and r is a fixed positive integer. We say that b_t is in the *lower class* for $\Delta_t^{(r)}$ at 0 if

$$\mathbf{P}(\Delta_t^{(r)} \leq b_t \text{ i.o. as } t \downarrow 0) = 1. \quad (3.1)$$

If this is not the case then the probability in (3.1) is 0. (This follows from Blumenthal’s zero-one law for Markov processes.) In this section we derive a criterion to distinguish between these situations.

Theorem 3.1. *Assume $\bar{\Pi}(0+) = \infty$ and let r and b_t be as specified.*

(i) *Suppose $\liminf_{t \downarrow 0} t\bar{\Pi}(b_t) < \infty$. Then*

$$\mathbf{P}(\Delta_t^{(r)} \leq b_t \text{ i.o. as } t \downarrow 0) = 1. \quad (3.2)$$

(ii) *Alternatively, suppose $\lim_{t \downarrow 0} t\bar{\Pi}(b_t) = \infty$. Then (3.2) holds iff*

$$J_r := \int_0^1 t^{r-1} \bar{\Pi}^r(b_{t-}) e^{-t\bar{\Pi}(b_{t-})} dt = \infty. \quad (3.3)$$

Our usage of this terminology varies from some in the (random walk) literature. We use “lower class” to denote the behaviour in (3.1). “Upper class” for us will denote the behaviour in (4.1) below.

Remarks. (i) In Part (ii) of Theorem 3.1, $t\bar{\Pi}(b_t)$ is large for small t , so $J_r < \infty$ implies $J_{r-1} < \infty$, $r = 2, 3, \dots$. Thus $\Delta_t^{(r)} > b_t$ all but finitely often, a.s., as $t \downarrow 0$, implies $\Delta_t^{(r-1)} > b_t$ all but finitely often, a.s., as $t \downarrow 0$, as of course it should.

(ii) It is not difficult to show that (3.3) is equivalent under the assumptions of Theorem 3.1 to

$$\int_0^1 t^{r-1} \bar{\Pi}^r(b_t) e^{-t\bar{\Pi}(b_t)} dt = \infty.$$

(iii) The monotonicity requirement on b_t can be relaxed somewhat, analogous to the discrete situation, as discussed by Klass [11]. We omit details of this.

Proof of Theorem 3.1: Throughout, assume $\bar{\Pi}(0+) = \infty$ and let $b_t \downarrow 0$ as $t \downarrow 0$.

(i) Suppose first that $\liminf_{t \downarrow 0} t\bar{\Pi}(b_t) < \infty$. Then there exists $c > 0$ and a sequence $t_k \downarrow 0$ as $k \rightarrow \infty$ such that

$$\lim_{k \rightarrow \infty} \mathbf{P}(\Delta_{t_k}^{(1)} \leq b_{t_k}) = \lim_{k \rightarrow \infty} e^{-t_k \bar{\Pi}(b_{t_k})} = e^{-c} > 0.$$

This implies

$$\begin{aligned} \mathbf{P}(\Delta_{t_k}^{(1)} \leq b_{t_k} \text{ i.o. as } k \rightarrow \infty) &= \lim_{N \rightarrow \infty} \mathbf{P}\left(\bigcup_{k \geq N} \{\Delta_{t_k}^{(1)} \leq b_{t_k}\}\right) \\ &\geq \lim_{N \rightarrow \infty} \mathbf{P}(\Delta_{t_N}^{(1)} \leq b_{t_N}) \\ &= e^{-c} > 0. \end{aligned}$$

Hence by the zero-one law, $\mathbf{P}(\Delta_{t_k}^{(1)} \leq b_{t_k} \text{ i.o. as } k \rightarrow \infty) = 1$. Then since $\Delta_t^{(r)} \leq \Delta_t^{(1)}$, we have $\mathbf{P}(\Delta_{t_k}^{(r)} \leq b_{t_k} \text{ i.o. as } k \rightarrow \infty) = 1$ and (3.2) follows.

(ii) From now on assume $\lim_{t \downarrow 0} t\bar{\Pi}(b_t) = \infty$ and without loss of generality that $t\bar{\Pi}(b_t) \geq a_r$ for all $t \in (0, 1]$, where $a_r > 4r^2$ is a constant to be specified later. We will show that (3.2) and (3.3) are equivalent.

Assume that (3.3) holds. Let $t_0 = 1 = b_{t_0}$ and define the sequence t_k recursively by

$$t_{k+1} = t_k - 1/\bar{\Pi}(b_{t_k}), \quad k = 0, 1, 2, \dots \quad (3.4)$$

Clearly $0 < t_{k+1} < t_k$ and $t_k \downarrow 0$ as $k \rightarrow \infty$ (because if $t_k \downarrow T > 0$ then we would have $T + 1/\bar{\Pi}(b_{T+}) = T$, which is not possible since $b_{T+} > 0$ and $\bar{\Pi}(b_{T+}) > 0$).

Using the fact that $f(x) = x^r e^{-tx}$ is decreasing for $x > r/t$ together with (3.4) we have

$$\begin{aligned}
\int_0^1 t^{r-1} \bar{\Pi}^r(b_t) e^{-t\bar{\Pi}(b_t)} dt &= \sum_{k=0}^{\infty} \int_{t_{k+1}}^{t_k} t^{r-1} \bar{\Pi}^r(b_t) e^{-t\bar{\Pi}(b_t)} dt \\
&\leq \sum_{k=0}^{\infty} \int_{t_{k+1}}^{t_k} t^{r-1} \bar{\Pi}^r(b_{t_k}) e^{-t\bar{\Pi}(b_{t_k})} dt \\
&\leq \sum_{k=0}^{\infty} (t_k - t_{k+1}) \bar{\Pi}(b_{t_k}) (t_k \bar{\Pi}(b_{t_k}))^{r-1} e^{-t_{k+1} \bar{\Pi}(b_{t_k})} \\
&= e \sum_{k=0}^{\infty} (t_k \bar{\Pi}(b_{t_k}))^{r-1} e^{-t_k \bar{\Pi}(b_{t_k})}.
\end{aligned}$$

Since (3.3) holds, we conclude

$$\sum_{k=0}^{\infty} (t_k \bar{\Pi}(b_{t_k}))^{r-1} e^{-t_k \bar{\Pi}(b_{t_k})} = \infty. \quad (3.5)$$

Recall $\bar{\Pi}(b_t) \geq 4r^2/t$ for $t \in (0, 1]$. Construct a subsequence $\{s_k\}$ of $\{t_j\}$ as follows. Let $s_1 = t_1$ and assume s_k has been defined. Note that for any $j \geq 1$ we have $t_{j+1} \bar{\Pi}(b_{t_j}) = t_j \bar{\Pi}(b_{t_j}) - 1 \geq 4r^2 - 1 \geq 1$, so the quantity

$$s_{k+1} := \min\{t_j : t_j \bar{\Pi}(b_{s_k}) \geq 1\}, \quad k = 0, 1, 2, \dots,$$

satisfies $0 < s_{k+1} < s_k$, and we have $s_k \downarrow 0$ as $k \rightarrow \infty$. Further,

$$t_j \bar{\Pi}(b_{s_k}) \leq 1 \text{ for all } t_j < s_{k+1}, \quad (3.6)$$

and

$$s_{k+1} \bar{\Pi}(b_{s_k}) \geq 1. \quad (3.7)$$

Define a sequence of events A_k , $k = 1, 2, \dots$, by

$$A_k := \{\Delta_{t_i}^{(r)} \leq b_{t_i} \text{ for some } t_i \in [s_{k+1}, s_k]\}.$$

Take $k \geq 1$ and $m > k + 2$. Then $s_m < s_{k+2}$ and $b_{s_m} \leq b_{s_{k+2}}$. Also define

$$A_k^*(m) := \left\{ \Delta_{(s_m, t_i]}^{(r)} \leq b_{t_i} \text{ for some } t_i \in [s_{k+1}, s_k] \right\}$$

(recall that $\Delta_{(s_m, t_i]}^{(r)}$ is the r -th largest of the Δ_s for $s \in (s_m, t_i]$). Then $A_k \subseteq A_k^*(m)$ and $A_k^*(m)$ and A_m are independent. Hence

$$\mathbf{P}(A_m \cap A_k) \leq \mathbf{P}(A_m \cap A_k^*(m)) = \mathbf{P}(A_m) \mathbf{P}(A_k^*(m)).$$

If $A_k^*(m)$ occurs and $\Delta_{s_m}^{(1)} \leq b_{s_{k+1}}$ then $\Delta_{t_i}^{(r)} \leq b_{t_i}$ for some $t_i \in [s_{k+1}, s_k)$, and so A_k occurs. Thus we have the inequality

$$\mathbf{P}(A_k^*(m))\mathbf{P}(\Delta_{s_m}^{(1)} \leq b_{s_{k+1}}) \leq \mathbf{P}(A_k),$$

and we get

$$\begin{aligned} \mathbf{P}(A_m \cap A_k) &\leq \mathbf{P}(A_m)\mathbf{P}(A_k^*(m)) \leq \frac{\mathbf{P}(A_m)\mathbf{P}(A_k)}{\mathbf{P}(\Delta_{s_m}^{(1)} \leq b_{s_{k+1}})} \\ &= \mathbf{P}(A_m)\mathbf{P}(A_k)e^{s_m \bar{\Pi}(b_{s_{k+1}})} \\ &\leq e\mathbf{P}(A_m)\mathbf{P}(A_k) \quad (\text{using (3.6), and } m > k + 2). \end{aligned}$$

Hence $\mathbf{P}(A_m \cap A_k) \leq e\mathbf{P}(A_m)\mathbf{P}(A_k)$ for all $m > k + 2$.

Suppose

$$\sum_{k=0}^{\infty} \mathbf{P}(A_k) = \infty. \quad (3.8)$$

Then for $n \geq 4$,

$$\begin{aligned} \sum_{m=2}^n \sum_{k=1}^{m-1} \mathbf{P}(A_m \cap A_k) &\leq 2 \sum_{m=1}^n \mathbf{P}(A_m) + \sum_{m=4}^n \sum_{k=1}^{m-3} \mathbf{P}(A_m \cap A_k) \\ &\leq 2 \sum_{m=1}^n \mathbf{P}(A_m) + e \sum_{m=1}^n \sum_{k=1}^n \mathbf{P}(A_m)\mathbf{P}(A_k) \\ &\sim e \left(\sum_{m=1}^n \mathbf{P}(A_m) \right)^2, \text{ as } n \rightarrow \infty. \end{aligned}$$

With this, Spitzer's converse Borel-Cantelli Lemma (Spitzer [19], p.317) for the events $\{A_k\}_{k \in \mathbb{N}}$ allows us to deduce from (3.8) that $\mathbf{P}(A_k \text{ i.o. as } k \rightarrow \infty) = 1$, which implies (3.2).

To complete this part of the proof, it remains to prove (3.8), knowing (3.5). For $k = 1, 2, \dots$ and $j = 1, 2, \dots, \ell(k)$, where $\ell(k) = \max\{j : js_{k+1} \leq s_k\}$, define a sequence of sub-events of $\{A_k\}$ by

$$A_{k,j} := \{\Delta_{t_i}^{(r)} \leq b_{t_i} \text{ for some } t_i \in [js_{k+1}, ((j+1)s_{k+1}) \wedge s_k)\},$$

(If there is no t_i in $[js_{k+1}, ((j+1)s_{k+1})$ set $A_{k,j} = \emptyset$.) Define another sequence of events by

$$B_{k,j} := \left\{ \Delta_{((j+1)s_{k+1}, (j+2)s_{k+1}]}^{(r)} > b_{s_k} \right\}, \quad j, k, = 1, 2, \dots$$

Clearly $A_{k,j}$ and $B_{k,j}$ are independent for each j and k . In addition, for each k , $B_{k,j}$ and $A_{k,j'}$ are disjoint when $j' \geq j+2$. This is true because, on $B_{k,j}$, we have $\Delta_t^{(r)} > b_{s_k}$ for all $t \geq (j+2)s_{k+1}$. But on $A_{k,j'}$, we have $\Delta_{t_i}^{(r)} \leq b_{s_k}$ for some $t_i \geq j's_{k+1}$. This is impossible when $j' \geq j+2$.

Using (2.1), $\mathbf{P}(B_{k,j})$ can be bounded away from 0, because

$$\begin{aligned}
\mathbf{P}(B_{k,j}) &= 1 - \mathbf{P}(\Delta_{s_{k+1}}^{(r)} \leq b_{s_k}) \\
&= 1 - e^{-s_{k+1}\bar{\Pi}(b_{s_k})} \sum_{n=0}^{r-1} \frac{(s_{k+1}\bar{\Pi}(b_{s_k}))^n}{n!} \\
&\geq 1 - e^{-1} \sum_{n=0}^{r-1} \frac{1}{n!} \quad (\text{by (3.7)}) \\
&=: M_1 > 0.
\end{aligned} \tag{3.9}$$

In the final inequality we used that $f(x) = 1 - e^{-x} \sum_{n=0}^{r-1} \frac{x^n}{n!}$ is an increasing function of x . (3.9) allows us to bound $\mathbf{P}(A_k)$ from below as follows: for each $k = 1, 2, \dots$,

$$\begin{aligned}
2\mathbf{P}(A_k) &= 2\mathbf{P}\left(\bigcup_{j=1}^{\ell(k)} A_{k,j}\right) \\
&\geq \mathbf{P}\left(\bigcup_{j \text{ even}} (A_{k,j} \cap B_{k,j})\right) + \mathbf{P}\left(\bigcup_{j \text{ odd}} (A_{k,j} \cap B_{k,j})\right) \\
&= \sum_{j=1}^{\ell(k)} \mathbf{P}(A_{k,j} \cap B_{k,j}) = \sum_{j=1}^{\ell(k)} \mathbf{P}(A_{k,j})\mathbf{P}(B_{k,j}) \\
&\geq M_1 \sum_{j=1}^{\ell(k)} \mathbf{P}(A_{k,j}).
\end{aligned} \tag{3.10}$$

In the second equality we used the disjointness of $B_{k,j}$ and $A_{k,j'}$ for $j' \geq j+2$ and in the third we used the independence of $A_{k,j}$ from $B_{k,j}$.

Now set

$$C_i = \{\Delta_{t_i}^{(r)} \leq b_{t_i}\}, \quad i = 1, 2, \dots$$

By (2.1) and (3.5) we have

$$\begin{aligned}
\sum_{i=1}^{\infty} \mathbf{P}(C_i) &= \sum_{i=1}^{\infty} e^{-t_i \bar{\Pi}(b_{t_i})} \sum_{n=0}^{r-1} \frac{(t_i \bar{\Pi}(b_{t_i}))^n}{n!} \\
&\geq \sum_{i=1}^{\infty} e^{-t_i \bar{\Pi}(b_{t_i})} \frac{(t_i \bar{\Pi}(b_{t_i}))^{r-1}}{(r-1)!} = \infty.
\end{aligned} \tag{3.11}$$

Let integers $p = p(j, k) < \dots < q(j, k) = q$ index those t_i occurring in the interval $[js_{k+1}, (j+1)s_{k+1} \wedge s_k)$. (Recall we restrict attention to nonempty $A_{k,j}$. If there is only one t_i in the interval $[js_{k+1}, (j+1)s_{k+1} \wedge s_k)$, set $p = q$.) Then $js_{k+1} \leq t_q < t_{q-1} < \dots < t_p < (j+1)s_{k+1} \wedge s_k$, and $t_i \in [t_q, t_p]$ for any $p \leq i \leq q$. Let

$$D_i = \{\text{exactly } r \text{ values of } s \in (t_i, t_{i-1}) \text{ satisfy } \Delta_s > b_{t_p}\}, \quad i = 1, 2, \dots$$

On D_j , for $j \geq i+1$, we have $\Delta_{t_j}^{(r)} > b_{t_p}$, hence $\Delta_{t_i}^{(r)} > b_{t_p}$, but on C_i we have $\Delta_{t_i}^{(r)} \leq b_{t_i} \leq b_{t_p}$. Thus D_j is disjoint from C_i for $j \geq i+1$, $i \geq p$. Additionally, C_i is independent of D_i , $i = 1, 2, \dots$, and

$$\mathbf{P}(D_i) = ((t_{i-1} - t_i)\bar{\Pi}(b_{t_p}))^r e^{-(t_{i-1} - t_i)\bar{\Pi}(b_{t_p})} / r!. \quad (3.12)$$

We can now bound $\mathbf{P}(A_{k,j})$ from below as follows:

$$\begin{aligned} 2\mathbf{P}(A_{k,j}) &\geq \mathbf{P}\left(\bigcup_{i=p+1}^q C_i\right) + \mathbf{P}(C_p) \\ &\geq \mathbf{P}\left(\bigcup_{i=p+1}^q (C_i \cap D_i)\right) + \mathbf{P}(C_p) \\ &= \sum_{i=p+1}^q \mathbf{P}(C_i \cap D_i) + \mathbf{P}(C_p) \\ &= \sum_{i=p+1}^q \mathbf{P}(C_i)\mathbf{P}(D_i) + \mathbf{P}(C_p). \end{aligned} \quad (3.13)$$

Here we used the disjointness of D_j from C_i for $j \geq i+1$, $i \geq p$, together with the independence of C_i and D_i .

If $p = q$ let $m = q$, otherwise set $m = \min\{i : p < i \leq q, 4\bar{\Pi}(b_{t_p}) \leq \bar{\Pi}(b_{t_i})\}$ with $m = q$ if $4\bar{\Pi}(b_{t_p}) > \bar{\Pi}(b_{t_i})$ for all $p < i \leq q$. So $4\bar{\Pi}(b_{t_p}) > \bar{\Pi}(b_{t_i}) \geq \bar{\Pi}(b_{t_{i-1}})$ for $p \leq i \leq m-1$. We can now estimate $\mathbf{P}(D_i)$ using (3.12). As an upper bound

$$(t_{i-1} - t_i)\bar{\Pi}(b_{t_p}) = \frac{\bar{\Pi}(b_{t_p})}{\bar{\Pi}(b_{t_{i-1}})} \leq 1,$$

for $i \geq p+1$. For a lower bound, suppose $q > p$, so $m > p$, and keep $p < i < m$. Then $4\bar{\Pi}(b_{t_p}) \geq \bar{\Pi}(b_{t_{i-1}})$ and

$$(t_{i-1} - t_i)\bar{\Pi}(b_{t_p}) \geq (t_{i-1} - t_i)\bar{\Pi}(b_{t_{i-1}})/4 = 1/4.$$

Since $x^r e^{-x}$ increases on $[0, r]$, we get

$$\mathbf{P}(D_i) = ((t_{i-1} - t_i)\bar{\Pi}(b_{t_p}))^r e^{-(t_{i-1} - t_i)\bar{\Pi}(b_{t_p})} / r! \geq 4^{-r} e^{-1/4} / r!.$$

Then as a lower bound for $\mathbf{P}(A_{k,j})$ we deduce from (3.13)

$$\mathbf{P}(A_{k,j}) \geq M_2 \sum_{i=p}^{m-1} \mathbf{P}(C_i), \quad (3.14)$$

where $M_2 = 4^{-r} e^{-1/4} / 2r!$. When $m = p = q$, $\mathbf{P}(A_{k,j}) \geq \mathbf{P}(C_p)$ by (3.13).

The next task is to estimate the summation term in (3.14). For brevity in this calculation, write $y_i = \bar{\Pi}(b_{t_i})$, $i \geq 1$. Define $\mathcal{E}_0 := \{i : p \leq i \leq q, 4y_p > y_i\}$ and for $j \geq 1$ let $\mathcal{E}_j := \{i : m \leq i \leq q, 4^j y_p \leq y_i \leq 4^{j+1} y_p\}$. Then since $(t_i - t_{i+1})y_i = 1$,

$$\begin{aligned} |\mathcal{E}_j| &\leq 1 + 2 \sum_{\{i \in \mathcal{E}_j, i+1 \in \mathcal{E}_j\}} (t_i - t_{i+1})y_i \\ &\leq 1 + 2 \cdot 4^{j+1} y_p \sum_{\{i \in \mathcal{E}_j, i+1 \in \mathcal{E}_j\}} (t_i - t_{i+1}) \\ &\leq 4^{j+2} t_p y_p. \end{aligned}$$

When $i \in \mathcal{E}_j$, we have $t_p/2 \leq (j+1)s_{k+1}/2 \leq js_{k+1} \leq t_i$ and $y_i \geq 4^j y_p$. Thus $t_i y_i \geq 4^j t_p y_p/2 > r-1$ and since $x^{r-1}e^{-x}$ decreases for $x > r-1$ we can calculate

$$\begin{aligned} \sum_{i \in \mathcal{E}_j} \mathbf{P}(C_i) &\leq r \sum_{i \in \mathcal{E}_j} (t_i y_i)^{r-1} e^{-t_i y_i} \\ &\leq r (4^j t_p y_p/2)^{r-1} e^{-4^j t_p y_p/2} |\mathcal{E}_j| \\ &\leq 32r (4^j t_p y_p/2)^r e^{-4^j t_p y_p/2}. \end{aligned} \quad (3.15)$$

In order to sum this expression over j , we use the following estimate, true for any $r \geq 1$, $c > 0$, $a > 1$, with $ca > r$:

$$\begin{aligned} c^r \sum_{j \geq 2} a^{jr} e^{-ca^j} &\leq c^r \sum_{j \geq 2} \int_{j-1}^j a^{xr} e^{-ca^x} dx \\ &\leq \frac{1}{\log a} \int_{ca}^{\infty} y^{r-1} e^{-y} dy \leq \frac{r(ca)^{r-1} e^{-ca}}{\log a}. \end{aligned}$$

Applying this with $a = 4$ and $c = t_p y_p/2$ we find from (3.15)

$$\begin{aligned} \sum_{j=1}^{\infty} \sum_{i \in \mathcal{E}_j} \mathbf{P}(C_i) &\leq 32r (2t_p y_p)^r e^{-2t_p y_p} + \frac{32r^2 (2t_p y_p)^{r-1} e^{-2t_p y_p}}{\log 4} \\ &\leq 64r^2 2^r (t_p y_p)^r e^{-2t_p y_p} \end{aligned}$$

$$\leq \frac{(t_p y_p)^{r-1} e^{-t_p y_p}}{2(r-1)!} \leq \frac{1}{2} \mathbf{P}(C_p)$$

(provided we choose $t_p y_p = t_p \bar{\Pi}(b_{t_p}) > a_r$ large enough so that $128r^2 2^r (r-1)! t_p y_p \leq e^{2t_p y_p}$). Hence

$$\begin{aligned} \sum_{i=p}^q \mathbf{P}(C_i) &= \sum_{i=p}^{m-1} \mathbf{P}(C_i) + \sum_{j=1}^{\infty} \sum_{i \in \mathcal{E}_j} \mathbf{P}(C_i) \\ &\leq \sum_{i=p}^{m-1} \mathbf{P}(C_i) + \mathbf{P}(C_p)/2, \end{aligned}$$

yielding

$$\sum_{i=p}^{m-1} \mathbf{P}(C_i) \geq \frac{1}{2} \sum_{i=p}^q \mathbf{P}(C_i),$$

and, from (3.14),

$$\mathbf{P}(A_{k,j}) \geq \frac{M_2}{2} \sum_{i=p}^q \mathbf{P}(C_i).$$

Finally (3.8) follows from (3.11) and

$$\begin{aligned} 2 \sum_{k=1}^{\infty} \mathbf{P}(A_k) &\geq M_1 \sum_{k=1}^{\infty} \sum_{j=1}^{\ell(k)} \mathbf{P}(A_{k,j}) \quad (\text{by (3.10)}) \\ &\geq M_3 \sum_{k=1}^{\infty} \sum_{j=1}^{\ell(k)} \sum_{i=p}^q \mathbf{P}(C_i) \quad (\text{with } M_3 := M_1 M_2 / 2) \\ &= M_3 \sum_{i=1}^{\infty} \mathbf{P}(C_i) = \infty. \end{aligned}$$

This proves the forward direction of the theorem. For the converse, assume

$$\int_{(0,1]} (t \bar{\Pi}(b_{t-}))^r e^{-t \bar{\Pi}(b_{t-})} \frac{dt}{t} < \infty \quad (3.16)$$

for some $r = 1, 2, \dots$. We will show that

$$\mathbf{P}(\Delta_t^{(r)} \leq b_t \text{ i.o. as } t \downarrow 0) = 0. \quad (3.17)$$

To do this, define a sequence of random times $\{\tau_n\}_{n=0,1,2,\dots}$ by:

$$\begin{aligned} \tau_0 &= 1, \\ \tau_{n+1} &= \inf \left\{ 0 < t < \tau_n : \Delta_t = \Delta_{\tau_n-}^{(r)} \right\}, \quad n = 0, 1, 2, \dots, \end{aligned}$$

where we use the notation $\Delta_{t-}^{(r)} = \sup_{0 < s < t} \Delta_s^{(r)}$. Since $\Delta_t^{(r)} > 0$ a.s. for all $t > 0$ and $\Delta_t^{(r)} \downarrow 0$ a.s. as $t \downarrow 0$, we have $0 < \tau_n \downarrow 0$ as $n \rightarrow \infty$.

For $0 < t \leq 1$ let $n_0(t)$ be the integer such that $\tau_{n_0(t)} \leq t < \tau_{n_0(t)-1}$. Then

$$\begin{aligned} \mathbf{P}(\Delta_t^{(r)} \leq b_t \text{ i.o. as } t \downarrow 0) &= \lim_{t \downarrow 0} \mathbf{P} \left(\bigcup_{0 < s < t} \{\Delta_s^{(r)} \leq b_s\} \right) \\ &= \lim_{t \downarrow 0} \mathbf{P} \left(\bigcup_{n \geq n_0(t)} \bigcup_{\tau_{n+1} \leq s < \tau_n} \{\Delta_s^{(r)} \leq b_s\} \right) \\ &\leq \lim_{t \downarrow 0} \mathbf{P} \left(\bigcup_{n \geq n_0(t)} \{\Delta_{\tau_n-}^{(r)} \leq b_{\tau_n-}\} \right). \end{aligned} \quad (3.18)$$

Here we used that $\tau_{n+1} \leq s < \tau_n$ implies $\Delta_s^{(r)} = \Delta_{\tau_{n+1}}^{(r)} = \Delta_{\tau_n-}^{(r)}$ and $b_s \leq b_{\tau_n-}$. Hence

$$\begin{aligned} \mathbf{P}(\Delta_t^{(r)} \leq b_t \text{ i.o. as } t \downarrow 0) \\ \leq \lim_{t \downarrow 0} \mathbf{P}(\Delta_{s-}^{(r)} \leq b_{s-} \text{ for some } s \in (0, t] \text{ with } \Delta_s > \Delta_{s-}^{(r)}). \end{aligned}$$

To compute the probability on the right hand side we apply the compensation formula (Kyprianou [13], Thm. 4.4, p.95) as follows. Let

$$E_t = \{\Delta_{s-}^{(r)} \leq b_{s-} \text{ for some } s \in (0, t] \text{ with } \Delta_s > \Delta_{s-}^{(r)}\}, \quad t > 0,$$

and let

$$N_t = \int_{(0, t] \times (0, \infty)} 1_{\{\Delta_{s-}^{(r)} \leq b_{s-}, x > \Delta_{s-}^{(r)}\}} \mathbb{D}(ds \times dx)$$

be the number of points (s, Δ_s) which satisfy $\Delta_{s-}^{(r)} \leq b_{s-}$ and $\Delta_s > \Delta_{s-}^{(r)}$ for $s \in (0, t]$.

Then we have, for $0 < t < 1$,

$$\begin{aligned} \mathbf{P}(E_t) &= \mathbf{P}(N_t \geq 1) \leq \mathbf{E}[N_t] \quad (\text{by Markov's inequality}) \\ &= \mathbf{E} \left[\int_{(0, t] \times (0, \infty)} 1_{\{\Delta_{s-}^{(r)} \leq b_{s-}, x > \Delta_{s-}^{(r)}\}} \mathbb{D}(ds \times dx) \right] \\ &= \int_{(0, t] \times (0, \infty)} \mathbf{E} \left[1_{\{\Delta_{s-}^{(r)} \leq b_{s-}, x > \Delta_{s-}^{(r)}\}} \right] ds \Pi(dx) \\ &= \int_{(0, t] \times (0, \infty)} \mathbf{P}(\Delta_{s-}^{(r)} \leq b_{s-}, x > \Delta_{s-}^{(r)}) ds \Pi(dx) \\ &= \int_0^t \left(\int_{0 < x \leq b_{s-}} \mathbf{P}(\Delta_{s-}^{(r)} < x) \Pi(dx) + \mathbf{P}(\Delta_{s-}^{(r)} \leq b_{s-}) \bar{\Pi}(b_{s-}) \right) ds \\ &=: \int_0^t (A(s) + B(s)) ds. \end{aligned} \quad (3.19)$$

(In the third equality we used the compensation formula.)

We consider A and B separately. Keep $s > 0$ small enough for $s\bar{\Pi}(b_{s-}) \geq r$ and write

$$\begin{aligned} A(s) &= \int_{0 < x \leq b_{s-}} \mathbf{P}(\Delta_{s-}^{(r)} < x) \Pi(dx) \\ &= \int_{0 < x \leq b_{s-}} \sum_{i=0}^{r-1} \frac{(s\bar{\Pi}(x-))^i}{i!} e^{-s\bar{\Pi}(x-)} \Pi(dx) \quad (\text{using (2.1)}) \\ &\leq r \int_{0 < x \leq b_{s-}} (s\bar{\Pi}(x-))^{r-1} e^{-s\bar{\Pi}(x-)} \Pi(dx), \end{aligned}$$

in which we note $s\bar{\Pi}(x-) \geq s\bar{\Pi}(b_{s-}) \geq 1$. Define $x = \bar{\Pi}^{-\rightarrow}(y) := \sup\{z > 0 : \bar{\Pi}(z) > y\}$, $y > 0$, and use the change of variable formula (Brémaud [1], p.301) to write the last integral as

$$\begin{aligned} &\int_{\bar{\Pi}(b_{s-})}^{\infty} (s\bar{\Pi}(\bar{\Pi}^{-\rightarrow}(y)-))^{r-1} e^{-s\bar{\Pi}(\bar{\Pi}^{-\rightarrow}(y)-)} dy \\ &\leq \int_{\bar{\Pi}(b_{s-})}^{\infty} (sy)^{r-1} e^{-sy} dy = s^{-1} \int_{s\bar{\Pi}(b_{s-})}^{\infty} y^{r-1} e^{-y} dy. \end{aligned} \quad (3.20)$$

In the last inequality we used $\bar{\Pi}(\bar{\Pi}^{-\rightarrow}(y)-) \geq y$ and that $y^{r-1}e^{-y}$ is decreasing for $y > r-1$, noting that when $0 < x \leq b_{s-}$, $s\bar{\Pi}(x-) \geq s\bar{\Pi}(b_{s-}) \geq r$ by assumption. Estimating the incomplete Gamma function, from (3.20) we obtain for small $s > 0$

$$A(s) \leq rs^{-1} (s\bar{\Pi}(b_{s-}))^{r-1} e^{-s\bar{\Pi}(b_{s-})}.$$

Easier is

$$B(s) = \bar{\Pi}(b_{s-}) e^{-s\bar{\Pi}(b_{s-})} \sum_{i=0}^{r-1} \frac{(s\bar{\Pi}(b_{s-}))^i}{i!} \leq rs^{-1} (s\bar{\Pi}(b_{s-}))^r e^{-s\bar{\Pi}(b_{s-})}.$$

So we have, for small $s > 0$,

$$A(s) + B(s) \leq 2rs^{-1} (s\bar{\Pi}(b_{s-}))^r e^{-s\bar{\Pi}(b_{s-})}.$$

The last expression is integrable over $0 < s \leq t$ by (3.16), so the RHS of (3.19) tends to 0 as $t \downarrow 0$. Thus $\lim_{t \downarrow 0} P(E_t) = 0$ and this means $\mathbf{P}(\Delta_t^{(r)} \leq b_t \text{ i.o. as } t \downarrow 0) = 0$. This completes the proof of Theorem 3.1. $\square$

4. Upper Classes for $\Delta_t^{(r)}$, and Relative Stability

We say that b_t is in the upper class for $\Delta^{(r)}$ at 0 if

$$\mathbf{P}(\Delta_t^{(r)} > b_t \text{ i.o. as } t \downarrow 0) = 1. \quad (4.1)$$

Otherwise $\mathbf{P}(\Delta_t^{(r)} > b_t \text{ i.o. as } t \downarrow 0) = 0$. Recall we assume throughout that $\bar{\Pi}(0+) = \infty$, the nonstochastic function b_t is positive and non-decreasing on $(0, \infty)$ with $\lim_{t \downarrow 0} b_t = 0$, and r is a fixed positive integer. In this section we characterise the upper class for $\Delta_t^{(r)}$ at 0 relative to b_t . We often write $b(\cdot)$ for b_t from now on.

Proposition 4.1. *The following are equivalent:*

$$\mathbf{P}(\Delta_t^{(r)} > b_t \text{ i.o. as } t \downarrow 0) = 0; \quad (4.2)$$

$$\int_0^1 \bar{\Pi}^r(b_t) dt^r < \infty; \quad (4.3)$$

$$\sum_{n \geq 0} (2^{-n} \bar{\Pi}(b(2^{-n})))^r < \infty. \quad (4.4)$$

If any of these holds then

$$\lim_{t \downarrow 0} t \bar{\Pi}(b_t) = 0. \quad (4.5)$$

Proof of Proposition 4.1: This is in Lemma 3.1 of Maller [14]. □

We will also need the following differential formula:

$$\begin{aligned} \frac{|d\bar{\Pi}^r(x)|}{d\bar{\Pi}(x)} &= r \bar{\Pi}^{r-1}(x) 1_{\{\Delta \bar{\Pi}(x)=0\}} + \frac{\bar{\Pi}^r(x-) - \bar{\Pi}^r(x)}{\Delta \bar{\Pi}(x)} 1_{\{\Delta \bar{\Pi}(x) \neq 0\}} \\ &= \sum_{k=0}^{r-1} \bar{\Pi}^k(x) \bar{\Pi}^{r-1-k}(x-), \quad x > 0, \end{aligned}$$

where $\Delta \bar{\Pi}(x) := \bar{\Pi}(x-) - \bar{\Pi}(x)$. (For an analogous formula see Eq. (3.3), p.437 of [7].)

Consequently we have the inequalities

$$\bar{\Pi}^{r-1}(x-) \leq \frac{|d\bar{\Pi}^r(x)|}{d\bar{\Pi}(x)} \leq r \bar{\Pi}^{r-1}(x-). \quad (4.6)$$

The right-continuous inverse of the nonincreasing function $\bar{\Pi}$ is defined by

$$\bar{\Pi}^{\leftarrow}(x) = \inf\{y > 0 : \bar{\Pi}(y) \leq x\}, \quad x > 0.$$

By convention, the inf of the empty set is taken as ∞ . The following properties of the inverse function will be used: for each $x, y > 0$, $\bar{\Pi}^\leftarrow(x) \leq y$ if and only if $\bar{\Pi}(y) \leq x$; $\bar{\Pi}(\bar{\Pi}^\leftarrow(x)) \leq x \leq \bar{\Pi}(\bar{\Pi}^\leftarrow(x)-)$; and $\bar{\Pi}^\leftarrow(\bar{\Pi}(x)) \leq x$.

Proposition 4.1 combined with Theorem 3.1 gives our next result, Theorem 4.1, regarding the almost sure relative stability of $(\Delta_t^{(r)})$ as $t \downarrow 0$. As background, we mention that the relative stability of the order statistics from a sample $\{\xi_n : n \geq 0\}$ of i.i.d. random variables has been studied by a number of authors through the years. The sequence $(\xi_n^{(r)})$ of r^{th} largest values among $(\xi_i)_{1 \leq i \leq n}$ is said to be *relatively stable* if there exists a non-random sequence $b_n > 0$ such that $\xi_n^{(r)}/b_n \rightarrow 1$ as $n \rightarrow \infty$, either in probability or almost surely. These cases are discussed in Gnedenko [8] and Hall [10], among others. Integral criteria for the almost sure relative stability were given in Barndorff-Nielsen [5], Resnick and Tomkins [18] and Goldie and Maller [7]. See also Tomkins [20]. (Some further background is given in Section 6.) We can analogously study the relative stability of $\Delta_t^{(r)}$ as $t \downarrow 0$, and begin by stating necessary and sufficient conditions for the almost sure version of this in the next theorem.

Theorem 4.1. *Assume $\bar{\Pi}(0+) = \infty$ and fix $r = 1, 2, \dots$. There exists a nonstochastic nondecreasing function $b_t > 0$ such that $\lim_{t \downarrow 0} \Delta_t^{(r)}/b_t = 1$ a.s. if and only if*

$$\int_0^1 \left(\frac{\bar{\Pi}(x-)}{\bar{\Pi}((1-\varepsilon)x-)} \right)^r \frac{\bar{\Pi}(dx)}{\bar{\Pi}(x-)} < \infty, \text{ for all } \varepsilon \in (0, 1). \quad (4.7)$$

If this is the case then b_t is slowly varying at 0 and can be chosen such that $b_t = \bar{\Pi}^\leftarrow(1/t)$, $t > 0$.

Further, if (4.7) holds for some $r > 1$ then it holds for $r = 1, 2, \dots$; thus, $\Delta_t^{(r)}$ is relatively stable a.s. as $t \downarrow 0$ if and only if $\Delta_t^{(1)}$ is relatively stable a.s. as $t \downarrow 0$.

Before proving Theorem 4.1 we need to characterise relative stability in probability.

Proposition 4.2. *There exists a nonstochastic nondecreasing function $b_t > 0$ such that $\Delta_t^{(r)}/b_t \xrightarrow{P} 1$ as $t \downarrow 0$ if and only if*

$$\lim_{x \downarrow 0} \frac{\bar{\Pi}((1+\varepsilon)x)}{\bar{\Pi}(x)} = 0, \text{ for all } \varepsilon > 0. \quad (4.8)$$

If these hold then $b_t \sim \bar{\Pi}^\leftarrow(1/t)$ and b_t is slowly varying as $t \downarrow 0$.

Remarks. Since the condition (4.8) does not depend on r , we see that $\Delta_t^{(r)}/b_t \xrightarrow{P} 1$ as $t \downarrow 0$ for any $r = 1, 2, \dots$ if and only if $\Delta_t^{(1)}/b_t \xrightarrow{P} 1$ as $t \downarrow 0$; thus relative stability in probability obtains regardless of which (fixed) value of r is specified. This is perhaps not so surprising, as the same occurs for the relative stability in probability of the order statistics from an i.i.d. sample (see [10], and, for a similar phenomenon, [20]), but it is more surprising that the same holds for almost sure relative stability. This was established in Goldie and Maller [7] for discrete time. Theorem 4.1 shows that the same obtains in our present continuous time setup. We draw on the methods of [7] in that part of the proof of Theorem 4.1.

Proof of Proposition 4.2: For $b_t > 0$ and $\varepsilon > 0$ use (2.1) to write

$$\mathbf{P}(\Delta_t^{(r)} > (1 + \varepsilon)b_t) = 1 - e^{-t\bar{\Pi}((1+\varepsilon)b_t)} \sum_{i=0}^{r-1} \frac{(t\bar{\Pi}((1+\varepsilon)b_t))^i}{i!}. \quad (4.9)$$

(i) Assume $\Delta_t^{(r)}/b_t \xrightarrow{P} 1$ as $t \downarrow 0$ where $b_t > 0$ and $\lim_{t \downarrow 0} b_t = 0$. Then for $\varepsilon > 0$ we have $\mathbf{P}(\Delta_t^{(r)} > (1 + \varepsilon)b_t) \rightarrow 0$ as $t \downarrow 0$. Take a subsequence through which $t\bar{\Pi}((1+\varepsilon)b_t)$ tends to a limit $a \in [0, \infty]$ as $t \downarrow 0$. By (4.9), $a \in (0, \infty]$ is impossible so we have $a = 0$. This is true for all subsequences so we get $\lim_{t \downarrow 0} t\bar{\Pi}((1+\varepsilon)b_t) = 0$. A similar argument gives $\lim_{t \downarrow 0} t\bar{\Pi}((1-\varepsilon)b_t) = \infty$ for $\varepsilon \in (0, 1)$. Substitute these back in (4.9) with $r = 1$ to deduce that $\Delta_t^{(1)}/b_t \xrightarrow{P} 1$ as $t \downarrow 0$. Then from

$$\frac{\Delta_{2t}^{(1)}}{b_{2t}} \leq \frac{\Delta_{2t}^{(1)}}{b_t} = \frac{\max(\Delta_t^{(1)}, \sup_{t < s \leq 2t} \Delta_s)}{b_t} \xrightarrow{P} 1, \text{ as } t \downarrow 0,$$

we obtain $\Delta_{2t}^{(1)}/b_t \xrightarrow{P} 1$ as $t \downarrow 0$, and hence $b_{2t} \sim b_t$, as $t \downarrow 0$. Since b_t is non-decreasing, this means that b_t is slowly varying as $t \downarrow 0$. Given $\delta \in (0, 1)$, set $\eta = \delta/(2 + \delta)$, so that $(1 + \delta)(1 - \eta) = 1 + \eta$. Then choose $x > 0$ small enough and $t = t(x)$ such that

$$(1 - \eta)b(2t) \leq b(t/2) \leq b(t-) \leq x \leq b(t+) \leq b(2t),$$

implying

$$(1 + \eta)b(2t) = (1 + \delta)(1 - \eta)b(2t) \leq (1 + \delta)b(t/2) \leq (1 + \delta)x.$$

Then

$$\frac{\bar{\Pi}((1 + \delta)x)}{\bar{\Pi}((1 - \delta)x)} \leq \frac{t\bar{\Pi}((1 + \eta)b(2t))}{t\bar{\Pi}((1 - \delta)b(2t))}.$$

The RHS tends to 0 because $\Delta_{2t}^{(r)}/b_{2t} \xrightarrow{P} 1$, and hence (replacing $(1 - \delta)x$ by x and $(1 + \delta)/(1 - \delta)$ by $1 + \varepsilon$) we see that (4.8) holds.

(ii) Conversely, assume (4.8) holds and choose $b_t := \bar{\Pi}^{\leftarrow}(1/t)$. Then b_t is nondecreasing and $\bar{\Pi}(b_t) \leq 1/t \leq \bar{\Pi}(b_{t-})$. By (4.8) we have $t\bar{\Pi}((1 + \varepsilon)b_t) = o(t\bar{\Pi}(b_t)) = o(1)$, so by (4.9), $\mathbf{P}(\Delta_t^{(r)} > (1 + \varepsilon)b_t) \rightarrow 0$ as $t \downarrow 0$; while also

$$t\bar{\Pi}((1 - \varepsilon)b_t) \geq \frac{\bar{\Pi}((1 - \varepsilon)b_t)}{\bar{\Pi}(b_{t-})} \geq \frac{\bar{\Pi}((1 - \varepsilon)b_t)}{\bar{\Pi}((1 - \varepsilon/2)b_t)} \rightarrow \infty,$$

so by (2.1), $\mathbf{P}(\Delta_t^{(r)} < (1 - \varepsilon)b_t) \rightarrow 0$ as $t \downarrow 0$. These give $\Delta_t^{(r)}/b_t \xrightarrow{P} 1$ as $t \downarrow 0$ with $b_t = \bar{\Pi}^{\leftarrow}(1/t)$. $\square$

Proof of Theorem 4.1: (i) We will first show that (4.7) implies

$$P\left(\Delta_t^{(r)} > (1 + \varepsilon)b_t \text{ i.o. as } t \downarrow 0\right) = 0, \text{ for all } \varepsilon > 0, \quad (4.10)$$

with b_t defined as $\bar{\Pi}^{\leftarrow}(1/t)$, $t > 0$. Assume (4.7) and define b_t in this way. Then $t\bar{\Pi}(b_t) \leq 1 \leq t\bar{\Pi}(b_{t-})$. Take $\varepsilon > 0$ and use the RHS of (4.6) to estimate

$$\begin{aligned} & \sum_{n \geq 1} \left(2^{-n}\bar{\Pi}((1 + \varepsilon)b(2^{-n})-)\right)^r \\ & \leq r \sum_{n \geq 1} 2^{-nr} \int_{[(1 + \varepsilon)b(2^{-n}), \infty)} \bar{\Pi}^{r-1}(x-)\Pi(dx) \\ & \leq r \sum_{n \geq 1} 2^{-nr} \sum_{j=1}^n \int_{[(1 + \varepsilon)b(2^{-j}), (1 + \varepsilon)b(2^{-j+1})]} \bar{\Pi}^{r-1}(x-)\Pi(dx) + c_r. \end{aligned}$$

Here $c_r := r2^{-r}\bar{\Pi}^r((1 + \varepsilon)b(1-))/(1 - 2^{-r})$ and we interpret intervals of the form $[a, a)$ or $(a, a]$ as empty. Interchange the order of summations in the last expression and perform the inner summation to get an upper bound of the form

$$\frac{r}{1 - 2^{-r}} \sum_{j \geq 1} 2^{-jr} \int_{[(1 + \varepsilon)b(2^{-j}), (1 + \varepsilon)b(2^{-j+1})]} \bar{\Pi}^{r-1}(x-)\Pi(dx) + c_r. \quad (4.11)$$

When $(1 + \varepsilon)b(2^{-j}) \leq x$ we have

$$b(2^{-j}) \leq \frac{x}{1 + \varepsilon \wedge 1} < \frac{x}{1 + (\varepsilon \wedge 1)/2} = (1 - \delta)x,$$

where $\delta = \delta(\varepsilon) = (\varepsilon \wedge 1)/(2 + \varepsilon \wedge 1) \in (0, 1)$. So (since $t\bar{\Pi}(b_t) \leq 1$)

$$2^{-jr} \leq \frac{1}{\bar{\Pi}^r(b(2^{-j}))} \leq \frac{1}{\bar{\Pi}^r((1 - \delta)x-)}.$$

Substitute this in (4.11) to get the upper bound

$$\frac{r}{1-2^{-r}} \sum_{j \geq 1} \int_{[(1+\varepsilon)b(2^{-j}), (1+\varepsilon)b(2^{-j+1})]} \left(\frac{\bar{\Pi}(x-)}{\bar{\Pi}((1-\delta)x-)} \right)^r \frac{\Pi(dx)}{\bar{\Pi}(x-)} + c_r.$$

This is finite by (4.7). This working shows that (4.4) holds with $b(2^{-n})$ replaced by $(1+\varepsilon)b(2^{-n})$, and (4.4) in that form implies (4.10) by Proposition 4.1.

(ii) Next we show that if (4.7) holds and $b_t = \bar{\Pi}^{\leftarrow}(1/t)$ then

$$\mathbf{P}(\Delta_t^{(r)} \leq (1-\varepsilon)b_t \text{ i.o. as } t \downarrow 0) = 0, \text{ for all } \varepsilon \in (0, 1). \quad (4.12)$$

This together with (4.10) will give $\lim_{t \downarrow 0} \Delta_t^{(r)}/b_t = 1$ a.s.

Assume (4.7) and $b_t = \bar{\Pi}^{\leftarrow}(1/t)$. Then as shown in Part (i) we have

$$\sum_{n \geq 1} (2^{-n} \bar{\Pi}((1+\varepsilon)b(2^{-n})-))< \infty \text{ for all } \varepsilon > 0. \quad (4.13)$$

This in turn implies

$$\lim_{x \downarrow 0} \frac{\bar{\Pi}(x)}{\bar{\Pi}((1-\varepsilon)x)} = 0 \text{ for all } \varepsilon \in (0, 1). \quad (4.14)$$

To see this, note that (4.13) implies

$$\lim_{n \rightarrow \infty} 2^{-n} \bar{\Pi}((1+\varepsilon)b(2^{-n})-) = 0 \text{ for all } \varepsilon > 0.$$

Using

$$2^{n-1} \leq \bar{\Pi}(\bar{\Pi}^{\leftarrow}(2^{n-1})-) = \bar{\Pi}(b(2^{-n+1})-)$$

we get that

$$\lim_{n \rightarrow \infty} \frac{\bar{\Pi}((1+\varepsilon)b(2^{-n})-)}{\bar{\Pi}(b(2^{-n+1})-)} = 0.$$

Then, given $x > 0$, choose $n = n(x)$ so that $b(2^{-n}) \leq x \leq b(2^{-n+1})$ and apply this relation to get

$$\lim_{x \downarrow 0} \frac{\bar{\Pi}((1+\varepsilon)x-)}{\bar{\Pi}(x-)} = 0 \text{ for all } \varepsilon \in (0, 1),$$

which is equivalent to (4.14).

Since $\bar{\Pi}(b_t-) \geq 1/t$, (4.14) implies $\lim_{t \downarrow 0} t \bar{\Pi}((1-\varepsilon)b_t-) = \infty$ for all $\varepsilon \in (0, 1)$, hence $\lim_{t \downarrow 0} t \bar{\Pi}((1-2\varepsilon)b_t) = \infty$ for all $\varepsilon \in (0, 1/2)$. Referring to Theorem 3.1, we can thus prove (4.12) by showing

$$\int_0^1 t^{r-1} \bar{\Pi}^r((1-2\varepsilon)b_{t-}) e^{-t \bar{\Pi}((1-2\varepsilon)b_{t-})} dt < \infty \text{ for all } \varepsilon \in (0, 1/2). \quad (4.15)$$

To this end, note that, for $\varepsilon \in (0, 1)$,

$$\begin{aligned} & \int_0^1 t^{r-1} \bar{\Pi}^r((1-\varepsilon)b_t-) e^{-t\bar{\Pi}((1-\varepsilon)b_t-)} dt \\ &= \int_0^1 \frac{t^{-r-1}}{\bar{\Pi}^r((1-\varepsilon)b_t-)} (t\bar{\Pi}((1-\varepsilon)b_t-))^{2r} e^{-t\bar{\Pi}((1-\varepsilon)b_t-)} dt \\ &\leq C \int_0^1 \frac{t^{-r-1}}{\bar{\Pi}^r((1-\varepsilon)b_t-)} dt, \end{aligned} \quad (4.16)$$

where $C := \sup_{x>0} x^{2r} e^{-x}$. Let $\delta = \delta(\varepsilon) = \varepsilon/(1-\varepsilon)$, so $1+\delta = 1/(1-\varepsilon)$, and let A_n denote the intervals

$$A_n = \left\{ t > 0 : \frac{1}{\bar{\Pi}((1+\delta)b(2^{-n-1}))} \leq t < \frac{1}{\bar{\Pi}((1+\delta)b(2^{-n}))} \right\}, \quad n = 1, 2, \dots$$

For $t \in A_n$ we have $1/t > \bar{\Pi}((1+\delta)b(2^{-n}))$, and so

$$b_t = \bar{\Pi}^{\leftarrow}(1/t) \leq \bar{\Pi}^{\leftarrow}(\bar{\Pi}((1+\delta)b(2^{-n}))) \leq (1+\delta)b(2^{-n}).$$

Hence $(1-\varepsilon)b_t \leq b(2^{-n})$ and $\bar{\Pi}((1-\varepsilon)b_t-) \geq \bar{\Pi}(b(2^{-n})-) \geq 2^n$. Thus the integral on the right of (4.16) does not exceed

$$C \sum_{n \geq 0} 2^{-nr} \int_{A_n} t^{-r-1} dt \leq C \sum_{n \geq 0} 2^{-nr} \bar{\Pi}^r((1+\delta)b(2^{-n-1})),$$

and this is finite by (4.13). This implies (4.15), so (4.12) is proved.

(iii) Conversely, assume $\lim_{t \downarrow 0} \Delta_t^{(r)}/b_t = 1$ a.s. for a nonstochastic nondecreasing function $b_t > 0$. Then the same convergence holds in probability, so by Proposition 4.2 we have $b_t \sim \bar{\Pi}^{\leftarrow}(1/t)$, thus without loss of generality we may take $b_t = \bar{\Pi}^{\leftarrow}(1/t)$. Also take $\eta > 0$. Then (4.10) holds with ε replaced by η so (4.4) holds with b_t replaced by $(1+\eta)b_t$. Use the LHS of (4.6) and a similar interchange of order of summations as in (4.11) to write

$$\begin{aligned} & \sum_{n \geq 1} (2^{-n} \bar{\Pi}((1+\eta)b(2^{-n})))^r \\ & \geq a_r \sum_{j \geq 1} 2^{(-j+1)r} \int_{((1+\eta)b(2^{-j}), (1+\eta)b(2^{-j+1}))} \bar{\Pi}^{r-1}(x-) \Pi(dx), \end{aligned} \quad (4.17)$$

where $a_r := 2^{-r}/(1-2^{-r})$. When $x \leq (1+\eta)b(2^{-j+1})$ and $\varepsilon = \eta/(1+\eta)$ we have

$$\bar{\Pi}((1-\varepsilon)x-) = \bar{\Pi}(x/(1+\eta)-) \geq \bar{\Pi}(b(2^{-j+1})-),$$

so

$$2^{-j+1} \geq \frac{1}{\bar{\Pi}(b(2^{-j+1})-)} \geq \frac{1}{\bar{\Pi}((1-\varepsilon)x-)}.$$

Thus a lower bound for the RHS of (4.17) is, apart from constants,

$$\sum_{j \geq 1} \int_{((1+\eta)b(2^{-j}), (1+\eta)b(2^{-j+1})]} \left(\frac{\bar{\Pi}(x-)}{\bar{\Pi}((1-\varepsilon)x-)} \right)^r \frac{\Pi(dx)}{\bar{\Pi}(x-)}.$$

Thus (4.7) holds.

It remains to show that if (4.7) holds for some $r > 1$ then it holds for $r = 1, 2, \dots$. This part of the proof uses key results proved in Goldie and Maller [7]. So assume (4.7) holds with an $r > 1$ and argue as follows. The integral in (4.7) equals

$$\begin{aligned} & \int_{0 < x < 1} \frac{\bar{\Pi}^{r-1}(x-) \Pi(dx)}{\bar{\Pi}^r((1-\varepsilon)x-)} \\ &= \int_{0 < x < 1} \bar{\Pi}^{r-1}(x-) \int_{0 < z < x} d\bar{\Pi}^{-r}((1-\varepsilon)z) \Pi(dx) \\ &= \int_{0 < z < 1} \left(\int_{z < x < 1} \bar{\Pi}^{r-1}(x-) \Pi(dx) \right) d\bar{\Pi}^{-r}((1-\varepsilon)z) \\ &\asymp \int_{0 < z < 1} \bar{\Pi}^r(z) d\bar{\Pi}^{-r}((1-\varepsilon)z) \\ &= \int_{0 < y < 1-\varepsilon} \bar{\Pi}^r((1+\delta)y) d\bar{\Pi}^{-r}(y), \end{aligned} \tag{4.18}$$

where $1 + \delta = 1/(1 - \varepsilon)$ and the symbol “ $\asymp$ ” connecting two quantities means that they are bounded above and below by constant multiples of each other (we used the inequalities (4.6) to obtain this in (4.18)). Now choose $x_0 > 0$ so that $\bar{\Pi}(e^{-x_0}) > 1$, and $\delta > 0$ so small that $\log(1 + \delta) < x_0$, and define

$$\bar{F}(x) = \frac{1}{\bar{\Pi}(e^{-x-})} 1_{\{x \geq x_0\}} + 1_{\{x < x_0\}}. \tag{4.19}$$

$\bar{F}$ is right continuous, nonincreasing on $[0, \infty)$ with $\bar{F}(0+) = 1$ and $\bar{F}(\infty) = 0$, so we can regard it as the tail of a cdf on $[0, \infty)$. Let $\bar{\Pi}_-(x) := \bar{\Pi}(x-)$, $x > 0$. Decomposing Π into continuous and discrete components, convergence or otherwise of the last integral in (4.18) (for all $\delta > 0$) is seen to be equivalent to that of the same integral with $\bar{\Pi}$ replaced by $\bar{\Pi}_-$ (for all $\delta > 0$). Substituting $\bar{F}$ as in (4.19) in this modified integral shows that the convergence of the last integral in (4.18) is equivalent to that of

$$\int_{0 < y < e^{-x_0}} \frac{d\bar{F}^r(-\log y)}{\bar{F}^r(-\log y - c)} = \int_{x > x_0} \frac{|d\bar{F}^r(x)|}{\bar{F}^r(x - c)} \asymp \int_{x > x_0} \frac{\bar{F}^{r-1}(x-) |d\bar{F}(x)|}{\bar{F}^r(x - c)}.$$

Here we set $c := \log(1 + \delta) \in (0, x_0)$ and have made use of analogous inequalities as in (4.6), for $\bar{F}$. In the denominator of the last integral, $\bar{F}(x - c) \leq \bar{F}((x - c) -) \leq \bar{F}(x - c/2)$, so we can replace $\bar{F}(x - c)$ by $\bar{F}((x - c) -)$, then the modified integral is $I_r(c)$ in the notation of Eq. (2.6) of Goldie and Maller [7]. By Equations (2.19) and (2.20) of that paper we have $I_r(\varepsilon) < \infty$ for every $\varepsilon > 0$ if and only if $I_1(\varepsilon) < \infty$ for every $\varepsilon > 0$. Referring back to (4.18) this means that (4.7) holds for $r = 1$, hence for $r = 1, 2, \dots$, as was to be proved. $\square$

5. Examples

We conclude with some examples to illustrate the possibilities in Theorems 3.1 and 4.1. We use the functions $\ell_2(t) = \log |\log t|$, $\ell_{k+1}(t) = \log \ell_k(t)$, $k = 2, 3, \dots$ when $0 < t \leq \delta$ for a sufficiently small $\delta \in (0, 1)$.

Example 1. Let $b_t \equiv t$ and take $\eta \geq 0$. Define Π_η so that

$$t\bar{\Pi}_\eta(b_t) = t\bar{\Pi}_\eta(t) = \ell_2(t) + 2\ell_3(t) + (1 + \eta)\ell_4(t).$$

Then

$$e^{-t\bar{\Pi}_\eta(b_t)} = \frac{1}{|\log t| (\log |\log t|)^2 (\log \log |\log t|)^{1+\eta}},$$

so the integral in (3.3), taken over $(0, \delta)$, converges or diverges with

$$\int_0^\delta \frac{(\log |\log t|)^{r-1}}{|\log t| \log |\log t| (\log \log |\log t|)^{1+\eta}} \frac{dt}{t}.$$

When $r = 1$ this integral converges for $\eta > 0$ and diverges for $\eta = 0$. But when $r = 2, 3, \dots$ it diverges for all $\eta \geq 0$. Thus for an extremal process corresponding to Π_η with $\eta > 0$, $\Delta_t^{(1)} > t$ all but finitely often, a.s., as $t \downarrow 0$, whereas $\Delta_t^{(2)} \leq t$ infinitely often, a.s., as $t \downarrow 0$.

Example 2. Here is an example where $\Delta_t^{(r)}$ is relatively stable in probability but not almost surely. Consider the function

$$f(x) := \exp(|\log x| \ell_2(x)), \quad 0 < x \leq \delta.$$

A calculation gives, for $0 < x \leq \delta$ and $0 < \lambda < 1$,

$$\log f(x\lambda) = |\log(x\lambda)| \log(|\log(x\lambda)|)$$

$$\begin{aligned}
&= (|\log x| + |\log \lambda|) \left(\log |\log x| + \log \left(1 + \frac{|\log \lambda|}{|\log x|} \right) \right) \\
&= \log f(x) + |\log \lambda| \ell_2(x) + O(1), \text{ as } x \downarrow 0.
\end{aligned}$$

Hence

$$\frac{f(x\lambda)}{f(x)} = e^{|\log \lambda| \ell_2(x) + O(1)} \asymp |\log x|^{|\log \lambda|}, \text{ as } x \downarrow 0.$$

It follows that $\lim_{x \downarrow 0} f(x\lambda)/f(x) = \infty$ for each $0 < \lambda < 1$, so if we define Π such that $\bar{\Pi}(x) = f(x)$, then $\bar{\Pi}$ is rapidly varying at 0 and the corresponding $\Delta_t^{(r)}$ is relatively stable in probability at 0.

On the other hand, $f'(x) < 0$ for $0 < x \leq \delta$, and

$$|f'(x)| \sim \frac{f(x) \log |\log x|}{x}, \text{ as } x \downarrow 0,$$

so, considering the integral in (4.7) with $\bar{\Pi}(x) = f(x)$, $r = 1$ and $1 - \varepsilon = \lambda \in (0, 1)$,

$$\begin{aligned}
\int_{(0, \delta]} \frac{\Pi(dx)}{\bar{\Pi}(x\lambda)} &= \int_{(0, \delta]} \left(\frac{|f'(x)|}{f(x\lambda)} \right) dx \\
&\asymp \int_{(0, \delta]} \left(\frac{f(x) \log |\log x|}{f(x) |\log x|^{|\log \lambda|}} \right) \frac{dx}{x} \\
&= \int_{(0, \delta]} \left(\frac{\log |\log x|}{|\log x|^{|\log \lambda|}} \right) \frac{dx}{x}.
\end{aligned}$$

This integral diverges when $|\log \lambda| \leq 1$, i.e., $\lambda > e^{-1}$, so $\Delta_t^{(1)}$ is not almost surely relatively stable at 0.

6. Further Background and Connections

We conclude with some further historical background concerning the discrete case. Barndorff-Nielsen [4] gave a lower class criterion relative to a given nonstochastic sequence for the partial maximum of a sequence of i.i.d. random variables, but needed to impose an unpleasant side condition. (Resnick and Rubinovitch [17] mention continuous time, small and large time, versions of this result.) Klass [11, 12] removed the side condition, thus giving a clean NASC for the partial maximum of a sequence of i.i.d. random variables to be in the lower class (according to our usage of the term).

Godbole [9] re-proved Klass's result by martingale methods, avoiding the need for the delicate (and clever) choices of subsequences in Klass's papers. These authors restricted themselves to the case $r = 1$. Harry Kesten in some unpublished notes

to the first author in the 1990s outlined an extension of Klass's result to r^{th} -order statistics. Our Theorem 3.1 is a continuous time version of these discrete time results, proved for general $r = 1, 2, \dots$, making specific use of the Poisson property of the jumps in the continuous time setup as well as a modification of Klass's subsequence method. Godbole's martingale methods do not seem to transfer easily to our situation because time "runs in reverse" as we let $t \downarrow 0$.

Upper class criteria are much easier to establish and are quoted for example in Barndorff-Nielsen [4] and the earlier literature.

Acknowledgement. We are grateful to two referees for helpful comments. The research was partially supported by ARC grant DP1092502.

References

- [1] Brémaud, P. (1981) *Point Processes and Queues, Martingale Dynamics*, Springer-Verlag, New York.
- [2] Buchmann, B., Fan, Y. and Maller, R.A. (2016) Distributional representations and dominance of a Lévy process over its maximal jump processes, *Bernoulli*, 22, 2325–2371.
- [3] Bingham, N.H., Goldie, C.M. and Teugels, J.L. (1987) *Regular Variation*, Cambridge University Press, Cambridge.
- [4] Barndorff-Nielsen, O. (1961) On the rate of growth of the partial maxima of a sequence of independent identically distributed random variables, *Math. Scand.*, 9, 383–394.
- [5] Barndorff-Nielsen, O. (1963) On the limit behaviour of extreme order statistics, *Ann. Math. Statist.*, 34, 992–1002.
- [6] de Haan, L. and Ferreira, A. (2006) *Extreme Value Theory, An Introduction*. Springer, New York.
- [7] Goldie, C.M., and Maller, R.A. (1996) A point-process approach to almost-sure behaviour of record values and order statistics, *Adv. Appl. Probab.*, 28, 426–462.

- [8] Gnedenko, B.V. (1943) Sur la distribution limite du terme maximum d'une séries aléatoire, *Ann. of Math.*, 44, 423–453.
- [9] Godbole, A.P. (1987) On Klass' series criterion for the minimal growth rate of partial maxima, *Statistics & Probability Letters*, 5, 235–237.
- [10] Hall, P. (1978) On the relative stability of large order statistics, *Math. Proc. Camb. Phil. Soc.*, 86, 467–485.
- [11] Klass, M.J. (1984) The minimal growth rate of partial maxima, *Ann. Probab.*, 12, 380–389.
- [12] Klass, K.J. (1985) The Robbins-Siegmund series criterion for partial maxima, *Ann. Probab.*, 13, 1369–1370.
- [13] Kyprianou, A. (2006) *Introductory Lectures on Fluctuations of Lévy Processes and Applications*, Springer-Verlag Berlin Heidelberg.
- [14] Maller, R.A. (2015) Strong laws at zero for trimmed Lévy processes, *Electron. J. Probab.* 20, no. 88, 1–24.
- [15] Resnick, Sidney I. (1974) Inverses of extremal processes, *Adv. Appl. Prob.* 6, 392–405.
- [16] Resnick, Sidney I. (2008) *Extreme values, Regular Variation and Point Processes*, Springer, New York.
- [17] Resnick, Sidney I. and Rubinovitch M. (1973) The structure of extremal processes, *Adv. Appl. Prob.* 5, 287–307.
- [18] Resnick, S. and Tomkins, R.J. (1973) Almost sure stability of maxima, *J. Appl. Prob.* 10, 387–401.
- [19] Spitzer, Frank (1975) *Principles of Random Walk*, Springer, New York.
- [20] Tomkins, R.J. (1986) Regular variation and the stability of maxima, *Ann. Probab.*, 14, 984–995.