Published ahead of print on June 28, 2007, doi:10.1164/rccm.200607-1005OC*American Journal of Respiratory and Critical Care Medicine* Vol 176. pp. 556-564, (2007)© 2007 [American Thoracic Society](#)

doi: 10.1164/rccm.200607-1005OC

Original Article**Neonatal Chlamydial Infection Induces Mixed T-Cell Responses That Drive Allergic Airway Disease****Jay C. Horvat^{1,2}, Kenneth W. Beagley^{1,2}, Margaret A. Wade^{1,2}, Julie A. Preston^{1,2}, Nicole G. Hansbro^{1,2}, Danica K. Hickey^{1,2}, Gerard E. Kaiko^{1,2}, Peter G. Gibson^{2,3,4}, Paul S. Foster^{1,2,5} and Philip M. Hansbro^{1,2}**

¹ Priority Research Centre for Asthma and Respiratory Disease, School of Biomedical Sciences, Faculty of Health, University of Newcastle, Newcastle, Australia; ² Vaccines, Immunity, Viruses, and Asthma Group, Hunter Medical Research Institute, Newcastle, Australia; ³ Respiratory and Sleep Medicine, John Hunter Hospital, Newcastle, Australia; ⁴ Respiratory and Sleep Medicine, School of Medical Practice, University of Newcastle, Newcastle, Australia; and ⁵ Division of Biosciences, John Curtin School of Medical Research, Australian National University, Canberra, Australia

Correspondence and requests for reprints should be addressed to Philip M. Hansbro, Ph.D., Discipline of Immunology and Microbiology, Level 3, David Maddison Clinical Sciences Building, Royal Newcastle Hospital, Newcastle, Australia 2300. E-mail: philip.hansbro{at}newcastle.edu.au

Rationale: Chlamydial lung infection has been associated with asthma in children and adults. However, how chlamydial infection influences the development of immune responses that promote asthma remains unknown.

Objectives: To determine the effect of chlamydial infection at various ages on the development of allergic airway disease (AAD).

Methods: Mouse models of chlamydial lung infection and ovalbumin-induced AAD were established in neonatal and adult BALB/c mice. Neonatal or adult mice were given a chlamydial infection and 6 weeks later were sensitized and subsequently challenged with ovalbumin. Features of AAD and inflammation were compared between uninfected or unsensitized controls.

Measurements and Main Results: Mild *Chlamydia*-induced lung disease was observed 10–15 days after infection, as evidenced by increased bacterial numbers and histopathology in the lung and a reduction in weight gain. After 6 weeks, infection and histopathology had resolved and the rate of weight gain had recovered. Neonatal but not adult infection resulted in significant decreases in interleukin-5 production from helper T cells and by the numbers of eosinophils recruited to the lung in response to ovalbumin exposure. Remarkably, the effects of early-life infection were associated with the generation of both type 1 and 2 ovalbumin-specific helper T-cell cytokine and antibody responses. Furthermore, although neonatal infection significantly attenuated eosinophilia, the generation of the mixed

This Article

- ▶ [Full Text](#)
- ▶ [Full Text \(PDF\)](#)
- ▶ [Online Supplement](#)
- ▶ **All Versions of this Article:**
[200607-1005OCv1](#)
176/6/556 *most recent*
- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)

Services

- ▶ [Similar articles in this journal](#)
- ▶ [Similar articles in PubMed](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Download to citation manager](#)
- ▶ [© Get Permissions](#)

Citing Articles

- ▶ [Citing Articles via HighWire](#)
- ▶ [Citing Articles via Google Scholar](#)

Google Scholar

- ▶ [Articles by Horvat, J. C.](#)
- ▶ [Articles by Hansbro, P. M.](#)
- ▶ [Search for Related Content](#)

PubMed

- ▶ [PubMed Citation](#)
- ▶ [Articles by Horvat, J. C.](#)
- ▶ [Articles by Hansbro, P. M.](#)

T-cell response exacerbated other hallmark features of asthma: mucus hypersecretion and airway hyperresponsiveness. Moreover, infection prolonged the expression of AAD and these effects were restricted to early-life infection.

Conclusions: Early-life chlamydial infection induces a mixed type 1 and 2 T-cell response to antigen, which differentially affects the development of key features of AAD in the adult.

Key Words: asthma • infection • immunity • *Chlamydia* • T cells

AT A GLANCE COMMENTARY

Scientific Knowledge on the Subject

Chlamydial infection is clinically associated with the onset and exacerbation of asthma in adults and children. However, it is unknown how this Th1-inducing infection is linked with Th2-driven asthma.

What This Study Adds to the Field

Chlamydia infection in early life can differentially drive key features of allergic airway disease. The effects of infection are associated with the generation of both Th1 and Th2 responses to an unrelated Th2-inducing antigen.

This article has been cited by other articles:

AMERICAN JOURNAL OF **Respiratory and Critical Care Medicine**

► HOME

W. C. Moore

Update in Asthma 2007

Am. J. Respir. Crit. Care Med., May 15, 2008; 177(10): 1068 - 1073.

[\[Full Text\]](#) [\[PDF\]](#)

Home
page

The Journal of Immunology

► HOME

M. Jupelli, M. N. Guentzel, P. A. Meier, G. Zhong, A. K. Murthy, and B. P. Arulanandam

Endogenous IFN- γ Production Is Induced and Required for Protective Immunity against Pulmonary Chlamydial Infection in Neonatal Mice

J. Immunol., March 15, 2008; 180(6): 4148 - 4155.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Home
page

The Journal of Immunology

► HOME

G. E. Kaiko, S. Phipps, D. K. Hickey, C. E. Lam, P. M. Hansbro, P. S. Foster, and K. W. Beagley

Chlamydia muridarum Infection Subverts Dendritic Cell Function to Promote Th2 Immunity and Airways Hyperreactivity

J. Immunol., February 15, 2008; 180(4): 2225 - 2232.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

HOME	HELP	FEEDBACK	SUBSCRIPTIONS	ARCHIVE	SEARCH	TABLE OF CONTENTS
----------------------	----------------------	--------------------------	-------------------------------	-------------------------	------------------------	-----------------------------------

Proc. Am. Thorac. Soc.

Am. J. Respir. Cell Mol. Biol.

[Copyright © 2007 American Thoracic Society](#)