Delineation of precursors in murine spleen that develop in contact with splenic endothelium to give novel dendritic-like cells

Jonathan K. H. Tan,¹ Pravin Periasamy,¹ and Helen C. O'Neill¹

¹Research School of Biology, Australian National University, Canberra, Australia

Hematopoietic cell lineages are best described in terms of distinct progenitors with limited differentiative capacity. To distinguish cell lineages, it is necessary to define progenitors and induce their differentiation in vitro. We previously reported in vitro development of immature dendritic-like cells (DCs) in long-term cultures (LTCs) of murine spleen, and in cocultures of spleen or bone marrow (BM) over splenic endothelial cell lines derived from LTCs. Cells produced are phenotypi-

cally distinct CD11b^{hi}CD11c^{lo}CD8⁻MHC-II⁻ cells, tentatively named L-DCs. Here we delineate L-DC progenitors as different from known DC progenitors in BM and DC precursors in spleen. The progenitor is contained within the lineage-negative (Lin)⁻c-kit⁺ subset in neonatal and adult spleen. This subset has multipotential reconstituting ability in mice. In neonatal spleen, the progenitor is further enriched within the c-kit^{lo} and CD34⁺ subsets of Lin⁻c-kit⁺ cells. These cells seed cocul-

tures of splenic endothelial cells, differentiating to give L-DCs that can activate T cells. L-DC progenitors are distinguishable from described splenic CD11c^{lo} DC precursors and from Fms-like tyrosine kinase 3⁺ DC progenitors in BM. Overall, this study confirms that LTCs are a physiologically relevant culture system for in vitro development of a novel DC type from spleen progenitors. (*Blood*. 2010; 115(18):3678-3685)

Introduction

Dendritic cells (DCs) are specialized antigen-presenting cells, which capture and process peripheral antigens for presentation to naive T cells.¹ They exist in peripheral tissues as immature cells, displaying characteristics optimal for antigen uptake and surveillance of local environments. Under these conditions, they present apoptotic self cells to T cells to maintain peripheral tolerance.^{2,3} On exposure to pathogens, DCs can also become immunogenic, adopting a mature, activated phenotype, a state distinct by additional secretion of proinflammatory cytokines, such as interleukin-12, which promote effector T-cell differentiation.⁴

Multiple DC subsets are present in tissues around the body, and in spleen the main subsets include conventional DCs (cDCs) and plasmacytoid DCs (pDCs). cDCs are represented by 2 main populations of CD8α⁻ and CD8α⁺ cDC. Morphologically, cDCs are small, nongranular cells⁵ that express high levels of CD11c and major histocompatibility complex class II (MHC-II) markers.⁶ pDCs are a distinct subset characterized by a small and plasmacytoid morphology, and a pronounced capacity to secrete high levels of IFN-α on viral stimulation.^{7,8}

Isolating rare DC subsets for experimentation is a difficult task, and the isolation and study of DC precursors represent an even more challenging area. It is, however, necessary to delineate precursors to perform lineage experiments to define their downstream progeny. Indeed, examination of DC precursors from early granulocyte-macrophage colony-stimulating factor/interleukin-4 cytokine culture systems has not been possible because of the enforced maturation of precursors.^{9,10} Recently, improved cultures of bone marrow (BM) cells with Fms-like tyrosine kinase 3 ligand (Flt3-L) have facilitated the study of DC development,^{11,12} with in vitro identification of 2 sequential cDC precursors CD11c^{int}CD115⁺c-kit⁻ pre-DCs and CD11c⁻CD115⁺c-kit^{int} pro-

DCs.¹³ Furthermore, in vivo counterparts of pre-DCs have been identified in spleen as CD11c^{int}CD43^{int}SIRP-α^{int} cells,¹⁴ and counterparts to pro-DC have been identified as Lin⁻c-kit^{lo}Flt3⁺ cells, present in BM but not spleen.¹⁵ Success in these studies highlights the value of in vitro culture systems for investigating DC development from precursors.

Previously, development of dendritic-like cells in spleen long-term cultures (LTCs) was demonstrated, which was unique in relation to other culture systems producing DCs. Cultures are continuous, maintaining 2 cell populations distinct in size as “small” and “large” cells.¹⁶ Large cells were previously characterized as a homogeneous population of immature myeloid dendritic-like CD11c^{lo}CD11b^{hi}CD8⁻MHC-II^{-/lo} cells, named LTC-DCs. These were highly endocytic and weakly able to present antigen to CD4⁺ T cells.^{17,18} The identity of a precursor within the small cell population was resolved by sorting and coculturing small cells above a spleen stromal cell line, which supports LTC-DC development.¹⁶ This resulted in development of large, immature DCs.^{16,19} The stromal cells that support myelopoiesis have now been characterized as splenic endothelial cells.²⁰⁻²³ Cloned stromal cell lines support production of distinct dendritic-like cells from precursors contained within BM and spleen equivalent to those produced in LTCs.^{24,25} These have been tentatively named L-DCs.²⁴

This paper characterizes precursors present in spleen that produce L-DCs in coculture over splenic endothelial stroma. Identification of these cells was based on previous evidence that the small cell population of LTCs contained cells expressing c-kit and Sca1.^{16,17} Here, we delineate the splenic precursor of L-DCs as a multipotential hematopoietic progenitor. This study identifies a potentially new pathway for myelopoiesis leading to DC production in spleen from endogenous progenitors.

Submitted June 22, 2009; accepted January 28, 2010. Prepublished online as *Blood* First Edition paper, March 4, 2010; DOI 10.1182/blood-2009-06-227108.

The publication costs of this article were defrayed in part by page charge

payment. Therefore, and solely to indicate this fact, this article is hereby marked “advertisement” in accordance with 18 USC section 1734.

© 2010 by The American Society of Hematology

Methods

Animals

C57BL/6J, C57BL/6.Tg(Tcrb)1100Mjb (OT-I), and C57BL/6.SJL-PtprcaPep^{3b}/BoyJ (B6.SJL) mice were bred at the John Curtin School of Medical Research (Canberra, Australia) under specific pathogen-free conditions. Mice were housed and handled according to protocols approved by the Animal and Experimental Ethics Committee at the Australian National University (ANU, Canberra, Australia).

Coculture assays for DC development

Cells were cultured in Dulbecco Modified Eagle Medium (DMEM) supplemented as described previously (sDMEM).²⁶ Cocultures involved overlay of dissociated spleen or BM cells, or purified subsets of spleen cells, on the STX3 spleen stromal line. STX3 was derived from an LTC established from B10.A(2R) mice, which ceased production of DCs over time with passage.²⁷ STX3 is passaged at 80% to 90% confluence by scraping a section of stroma, pipetting repetitively to disaggregate cells, and then transferring cells into fresh medium. When allogeneic or syngeneic spleen or BM cells were cocultured over STX3, only myeloid dendritic-like cells were produced.^{24,25} To establish cocultures, STX3 stroma was grown to 80% to 90% confluence before addition of overlay cells at 1 to 5×10^5 cells/mL. For coculture maintenance, half medium was exchanged every 3 to 4 days. Cocultures became established by 7 to 10 days and were monitored in culture for cell production by photomicroscopy under phase or bright-field using a DMIRE2 inverted microscope (Leica) equipped with a 40 $\times$ objective and a DFC digital camera (Leica). Images were processed using IM4.0 software (Leica) and prepared for publication using Photoshop CS3 (Adobe Systems Inc). When nonadherent cells were evident, cells were collected for marker analysis by flow cytometry or for functional assays.

Analysis of cell-surface marker expression

Antibody staining and flow cytometry were performed to distinguish cell subsets based on marker expression. To inhibit nonspecific antibody binding via Fc receptors, cells ($\sim 10^6$) were incubated with anti-CD16/32 (FcR block; eBioscience) for 15 minutes on ice in a final volume of 25 μ L of FACS buffer (DMEM/1% fetal calf serum/0.1% NaN₃). Cells were washed, and specific antibody (or isotype control) was absorbed to cells for 20 minutes on ice. Biotin- or fluorochrome-conjugated antibodies were specific for CD11c (N418; allophycocyanin), c-kit (2B8; phycoerythrin [PE] or fluorescein isothiocyanate [FITC]), CD11b (M1/70; PE-Cy7 or FITC), CD45.1 (A20; FITC), CD34 (RAM34; FITC), Flt3 (A2F10; PE), and VCAM (429; FITC; eBioscience). Antibodies specific for Scd1 (E13-161.7; biotin), CD69 (H1-2F3; biotin), CD8 (53-6.7; PE), and MHC-II (25-9-17; biotin) were purchased from BD Biosciences. Isotype controls included hamster IgG (eBio299Arm), rat IgG_{2a} (eBR2a), rat IgG_{2b} (KLH/G2b-1-2), and mouse IgG_{2a} (eBM2a) from eBioscience, and rat IgG_{2a} (R35-95) and hamster IgG1 (G235-2356) from BD Biosciences. After absorption of antibody, cells were washed twice. For addition of second-stage streptavidin-allophycocyanin-Cy7 (eBioscience), the same process was repeated. For analysis, cells were resuspended at 10^7 cells/mL in FACS buffer, and propidium iodide (PI; 1 μ g/mL; Sigma-Aldrich) added for discrimination of live and dead cells. Flow cytometry was performed immediately on a BD LSRII (BD Biosciences) and 3×10^4 to 1×10^6 events collected. BD FACSDiva Software (BD Biosciences) was used to acquire data. Data analysis involved postacquisition gating using FlowJo software (TreeStar).

To monitor proliferation of purified T cells after activation, cells were stained with 5- (and 6-) carboxyfluorescein diacetate succinimidyl ester (CFSE; Invitrogen). Proliferation was measured flow cytometrically by a decrease in fluorescence with cell division. For CFSE labeling, enriched preparations of T cells were washed and resuspended at less than 10^7 cells/100 μ L in sDMEM. CFSE was added at a final concentration of 5 μ M and vortexed immediately to ensure uniform labeling. Cells were incubated at room temperature for 5 minutes with regular agitation before washing twice with sDMEM.

Enrichment of myeloid precursors in spleen

Red blood cells (RBCs) were removed from whole dissociated splenocytes of C57BL/6J mice using lysis buffer (0.14 NH₄Cl, 0.017M Tris-base, pH 7.5) treatment for 5 minutes at 20°C. Cells were then washed twice, resuspended in FACS buffer (10^8 cells/mL), and filtered through a 70- μ m nylon cell strainer (BD Biosciences) before magnetic bead depletion of T and B cells and residual RBCs. For depletion, cells were labeled on ice for 15 minutes with biotinylated monoclonal antibodies specific for CD19 (clone: eBio1D3), Thy1.2 (30-H12), and TER-119, purchased from eBioscience. Cells were washed twice to remove unbound antibody and resuspended at 10^8 cells/mL in separation buffer (phosphate-buffered saline/0.5% bovine serum albumin/2mM ethylenediaminetetraacetic acid/0.1% NaN₃). Cells were incubated with anti-biotin MACS microbeads (13- μ L beads/ 10^8 cells; Miltenyi Biotec) for 20 minutes on ice, washed, and resuspended in separation buffer (500 μ L/ 10^8 cells). Cell depletion involved LS columns (Miltenyi Biotec) in a SuperMACS II Separation Unit (Miltenyi Biotec), washing 3 times with separation buffer and collecting unbound cells as flow-through.

Cell sorting

Sorting was performed using a BD FACSAria (BD Biosciences). Cells were prepared for flow cytometry as previously described, and all incubation and washing steps were performed strictly at 4°C in NaN₃-free FACS buffer. After a final wash, cells were passed through a 70- μ m nylon cell strainer (BD Biosciences) to ensure removal of cell clumps before sorting. Cells were sorted into sDMEM for later culturing.

In vivo reconstitution assays

The hematopoietic potential of cell subsets was tested after adoptive transfer into genetically distinct irradiated host mice. Donor spleen cell subsets (2×10^4) were prepared by sorting from B6.SJL mice (CD45.1⁺). Cells were intravenously injected via the tail vein of C57BL/6J (CD45.2⁺) recipient mice 4 to 6 weeks of age that had been irradiated 3 hours previously with 4.5 Gy (sublethal) or 9.5 Gy (lethal) whole body irradiation using a cobalt⁶⁰ source. Lethally irradiated mice were given 10^5 syngeneic BM cells to ensure survival. Irradiated mice were maintained on antibiotic water (neomycin sulfate, 1.1 g/L, Sigma-Aldrich; and polymyxin B sulfate, 10^6 U/L, Sigma-Aldrich) for 4 weeks. After 10, 20, or 41 days or 25 weeks, mice were killed and progeny cells identified in spleen using flow cytometry.

Positive selection of DCs from spleen

To obtain freshly isolated DCs, splenocytes were labeled with CD11c⁺ microbeads (15 μ L/ 10^8 cells; Miltenyi Biotec) in a total volume of 10^8 cells/mL of separation buffer for 15 minutes on ice followed by washing. To isolate CD11c⁺ DCs, cells were resuspended in separation buffer (500 μ L/ 10^8 cells), run through a SuperMACS II Separation Unit equipped with an MS column (Miltenyi Biotec), and washed 3 times with buffer (500 μ L). Flow-through cells were discarded before addition of 1 mL of separation buffer to the column. The CD11c⁺ cell fraction was collected under pressure.

Assays for antigen-specific T-cell activation by DCs

The antigen-presenting capacity of DCs was assessed using purified T cells from OT-I T cell receptor-transgenic (TCR-tg) mice (TCR-V α 2V β 5.1) specific for the K^b restricted peptide OVA₂₅₇₋₂₆₄²⁸ or OT-II TCR-tg CD4⁺ T cells specific for IA^b/ovalbumin (OVA). CD8⁺ T cells from OT-I mice and CD4⁺ T cells from OT-II mice were isolated from mesenteric lymph nodes by depletion of MHC-II⁺ antigen-presenting cells, B cells, and macrophages. The antibody cocktail contained anti-IA^{b/k} (clone: TIB120), anti-B220 (RA3-6B2), anti-CD4 (GK1.5), or anti-CD8 (53-6.7) and anti-CD11b (M1/70) prepared as hybridoma culture supernatant. Cells were labeled on ice and washed twice with separation buffer. To remove antibody-labeled leukocytes, cells were incubated with sheep anti-rat Ig Dynabeads (50 μ L beads/ 10^7 cells; Dynal) at 4°C for 25 minutes with

Figure 1. Characterization of DC precursors in neonatal spleen. C57BL/6J mice were analyzed for the presence of dendritic-like cells (DCs) and hematopoietic cell subsets by flow cytometric analysis for expression of c-kit, Sca1, CD34, VCAM, Thy1.2, CD69, Flt3, CD11c, and CD11b. Forward scatter (FSC) and side scatter (SSC) analysis was used to gate out debris, and isotype controls were used to determine background binding (shown as density plots) and to set gates. Specific antibody binding is shown by dot plots, and the percentage of cells is indicated in gates. Results are representative of 3 separate analyses.

rotation. For removal of CD11c⁺ DCs, cells were also incubated with anti-CD11c microbeads (1.5- μ L beads/ 10^7 cells; Miltenyi Biotec). After placing cells in a magnetic particle concentrator (Dyna) for 2 minutes, supernatant containing unbound enriched T cells was transferred to a fresh tube for labeling with CFSE.

Antigen-presenting cells were pulsed with antigen (OVA) in vitro at 10^7 cells/mL in sDMEM for 12 hours using either 10 or 100 μ g/mL OVA or hen egg lysozyme (HEL) as a control. To establish T-cell cocultures, enriched CFSE-labeled OT-I T cells were plated at 10^4 to 10^5 cells/well with diluting concentrations of antigen-presenting cells as stimulators in a total volume of 200 μ L of sDMEM. After 4 days, T-cell activation was assessed flow cytometrically to analyze cell division by reduction in CFSE intensity.

Results

Characterization of L-DC precursors in spleen

LTCs have been most successfully established from the spleens of neonatal 8-day-old mice. To investigate the presence of similar precursors in vivo, 8-day-old spleens were examined for cell subsets based on expression of c-kit, Sca1, CD34, and CD11c (Figure 1). In line with LTC precursors distinguishable by c-kit expression, 8-day spleen was found to segregate into clear c-kit⁺, c-kit^{lo}, and c-kit^{hi} subsets. Furthermore, Sca1 was expressed on c-kit^{hi} cells representing 0.02% of total 8-day-old spleen cells and on c-kit⁺ cells (1.3%) in accordance with populations identified among LTC small cells (all data not shown). Subsets of c-kit^{hi} splenocytes also coexpressed CD34 (0.7%) and Thy1.2 (0.03%). CD34 was also expressed on a subset of c-kit^{hi}Sca1⁺ cells (data not shown). Subsets of spleen c-kit^{lo} cells coexpressed Thy1.2 (0.4%), CD34 (0.7%), and VCAM-1 (0.7%). Flt3⁺ cells were barely detectable among the c-kit⁺ subset (0.01%), consistent with an absence of progenitors for cDCs in spleen.²⁹ All of these subsets are reflective of hematopoietic subsets described in BM.³⁰

The expression of lineage markers CD11c and CD11b and the activation marker CD69 was also analyzed in relation to c-kit on cells. Cells expressing c-kit were CD69⁺, consistent with their progenitor rather than activated or effector status. A clear

c-kit^{lo}CD11c^{hi} DC population accounted for 0.2% of total spleen and was consistent with the presence of a small population of cDCs.¹¹ A subset of CD11b⁺ cells also expressed low levels of c-kit, reflecting myeloid precursors. Thus, 8-day-old neonatal spleen contains c-kit^{lo} and c-kit^{hi} populations, which can be investigated for their equivalence with precursors maintained in LTC-DCs.

Neonatal splenocytes were then sorted on the basis of c-kit expression and subsets cocultured over STX3 stroma to determine DC differentiative potential. Overlaid whole BM, known to produce DCs in coculture over STX3 stroma,^{21,24,25} was used as a positive control. Initially, spleen cells were depleted of T and B cells and then sorted on the basis of c-kit and CD11c expression, to yield c-kit⁺CD11c⁺ and c-kit⁺CD11c⁺ subsets for comparison (Figure 2A). After 14 days of coculture over STX3 stroma, progeny cells were collected and stained in a 4-color assay and analyzed by multichannel flow cytometry for markers that distinguish the main DCs and myeloid subtypes, that is, CD11c, CD11b, CD8 α , and MHC-II. Both spleen c-kit⁺CD11c⁺ cells and whole BM yielded productive cocultures as assessed by phase microscopy (Figure 2B). Total cell production was monitored by forward scatter (FSC) analysis of progeny cells. Only control BM and sorted spleen c-kit⁺CD11c⁺ overlay cells produced progeny identifiable as events on FSC versus side scatter (SSC) plots (Figure 2C). Cells developing from the c-kit⁺CD11c⁺ spleen precursor population were heterogeneous and large in size. These were divisible into 2 populations of live (PI⁻) CD11c⁺CD11b⁺CD8 α ⁺MHC-II⁺ LTC-DC-like cells (hereafter referred to as L-DCs) and CD11c⁺CD11b⁺CD8 α ⁺MHC-II⁺ cells (MHC-II⁺ DCs) more reflective of CD8 α ⁺ cDCs or monocyte-derived DCs. These populations made up 60% and 7.2% of live FSC^{hi} cells, respectively. Exactly the same populations developed from cocultures established from total BM, confirming that the c-kit⁺CD11c⁺ spleen cell subset contains DC precursors.

To demonstrate that in vitro derived DCs were functionally competent antigen-presenting cells, cells were collected from cocultures, sorted to give a pure population, and then tested for ability to activate OT-I TCR-transgenic anti-OVA CD8⁺ T cells and OT-II TCR-transgenic anti-OVA CD4⁺ T cells. Cells were collected at 21 days, stained with CD11c, CD11b, and MHC-II and the L-DC population sorted as a CD11c⁺CD11b⁺MHC-II⁺ subset. The population of CD11c⁺CD11b⁺MHC-II⁺ cells was too few for this assay. Freshly isolated splenic CD11c⁺ DCs (f-DCs) were prepared as control antigen-presenting cells. This subset is highly enriched for CD11c⁺CD11b⁺MHC-II⁺ cDCs (data not shown). Coculture-derived cells were sorted to give L-DCs and pulsed with ovalbumin (OVA) or HEL as control antigen, with and without addition of lipopolysaccharide (LPS) as an activator. Antigen-pulsed DCs were then tested for ability to activate purified CFSE-labeled CD8⁺ OT-I T cells or CD4⁺ OT-II T cells in a 4-day assay. Cell division measured by dilution of CFSE intensity showed that sorted coculture-derived L-DCs were capable of inducing OVA-specific CD8⁺ T-cell proliferation, which was not improved by LPS. A similar result was seen for control CD11c⁺ spleen DCs. Antigen-pulsed L-DCs were unable to activate OT-II CD4⁺ T cells, consistent with an MHC-II⁺ phenotype. In contrast, fresh spleen CD11c⁺ cells gave a strong antigen-specific response in both the presence and absence of LPS. Cells developing in stromal cocultures from spleen precursors are functional antigen-presenting cells, although with restricted cross-priming capacity for only CD8⁺ T cells. These are established assays for antigen-

Figure 2. Identification of a DC precursor subset among neonatal spleen cells. (A) Eight-day-old spleen cells were sorted to give 2 populations of $c\text{-kit}^+CD11c^-$ and $c\text{-kit}^-CD11c^-$ cells. (B) Equal numbers of cells were cocultured over STX3 to assess DC development, along with whole red blood cell (RBC)-lysed bone marrow (BM) as positive control, and STX3 stromal cells as negative control. Cocultures were observed under phase microscopy after 14 days. Bar represents 50 μm . (C) Total nonadherent progeny cells were collected from cocultures at 14 days and analyzed flow cytometrically. FSC and SSC parameters were used to distinguish large cells for analysis of relative cell yield and marker analysis. Cells were simultaneously stained for CD11c, CD11b, CD8 α , and MHC-II to delineate DC subsets. Positive staining cells (dot plot) were gated using relevant isotype controls (density plot). Percentage of positive cells relative to all cells collected from cocultures is indicated in quadrants. (D) L-DCs produced in cocultures were tested for the capacity to activate purified CD8 $^+$ OT-I TCR-tg T cells or CD4 $^+$ OT-II TCR-tg T cells. L-DCs were collected from cocultures of $CD11c^-c\text{-kit}^+$ spleen after 21 days. Control f-DCs were isolated from spleen using $CD11c^+$ MACS microbeads. Antigen-presenting cell subsets were pulsed with antigen (10 $\mu\text{g}/\text{mL}$ OVA for OT-I, or 100 $\mu\text{g}/\text{mL}$ OVA for OT-II) in vitro for 12 hours before addition of CFSE-labeled T cells. Cocultures were established at a 1:5 DC:T-cell ratio, in the presence and absence of LPS (10 $\mu\text{g}/\text{mL}$), which was washed off after 12 hours. HEL was control antigen. T-cell activation was measured flow cytometrically after gating live (PI $^-$) $CD11c^-CD8^+$ OT-I T cells, or PI $^-$ $CD11c^-CD4^+$ OT-II T cells). Data are presented as percentage of T cells proliferating on the basis of reduction in CFSE intensity. Data are representative of 2 similar experiments.

specific T-cell activation, and their use here was only limited by the small number of L-DCs produced in cocultures.

Neonatal splenocytes were next sorted into $c\text{-kit}^-$, $c\text{-kit}^{\text{lo}}$, and $c\text{-kit}^{\text{hi}}$ subsets (Figure 3A) and assayed for DC development in cocultures by multicolor flow cytometry. Cell production was analyzed between 13 and 30 days, and common cell populations across multiple plots indicated by either square or round gates. Cell

production was only observed from spleen $c\text{-kit}^{\text{lo}}$ and $c\text{-kit}^{\text{hi}}$ precursors (Figure 3B-C), consistent with previous experiments. In productive cocultures, 2 cell populations were clearly identifiable as $CD11c^{\text{lo}}CD11b^{\text{hi}}MHC\text{-II}^-$ L-DCs and $CD11c^+CD11b^{\text{lo}}MHC\text{-II}^+$ cDCs (Figure 3C). These 2 populations were clearly distinguishable in cocultures established from both $c\text{-kit}^{\text{lo}}$ and $c\text{-kit}^{\text{hi}}$ spleen cells. $CD11c^{\text{lo}}CD11b^{\text{hi}}$ cells were negative for MHC-II (round

Figure 3. Further characterization of precursors in neonatal spleen. (A) Eight-day-old splenocytes were sorted into 3 populations of $c\text{-kit}^-$, $c\text{-kit}^{\text{lo}}$, $c\text{-kit}^{\text{hi}}$ cells (Expt A) and cocultured in equal number over STX3. (B) Cell production was monitored by photography under phase microscopy after 13 days. Bar represents 50 μm . (C) Nonadherent cells were collected from individual cocultures at 13, 23, and 30 days. Cells were simultaneously stained for CD11c, CD11b, CD8 α , and MHC-II. FSC and SSC analysis was used to gate large cells for marker analysis to assess DC production. Isotype controls were used to determine background binding (density plots) and set gates. Square and round gates were used to identify common populations across different plots. Percentage of cells showing specific staining is indicated within gates. Staining for CD8 was negative (data not shown). Results are reflective of 2 similar experiments.

Figure 6. Adult spleen contains progenitors of LTC-DCs. (A) Adult splenocytes depleted of T and B cells were sorted into populations of $c\text{-kit}^-CD11c^-$ and $c\text{-kit}^+CD11c^-$ cells and cocultured over STX3 to evaluate potential for DC development. (B) Nonadherent progeny cells collected from cocultures at 13, 23, and 30 days were assessed for DC production flow cytometrically. Cells were stained simultaneously for CD11c, CD11b, CD8 α , and MHC-II. FSC and SSC parameters were used to distinguish large cells for multichannel marker analysis. Isotype controls were used to define background antibody binding (zebra plots) and set gates. The percentage of cells showing specific binding is indicated within gates. Staining for CD8 was negative (data not shown). (C) DC production from 13-, 23-, and 30-day cocultures established with adult $c\text{-kit}^-CD11c^-$ and $c\text{-kit}^+CD11c^-$ spleen cell subsets is summarized. Cells produced were identified as either $CD11c^+CD11b^+MHC-II^-$ (MHC-II $^-$ LTC-DC) cells or $CD11c^+CD11b^+MHC-II^{lo/+}$ (MHC-II $^+$ DC) cells. Cell production is shown in terms of absolute number and yield of DCs, or percentage DCs among total cells produced in cocultures.

days after birth.³¹ To confirm that the presence of L-DC precursors in 8-day-old spleen was not a temporal phenomenon related to fluctuating progenitor levels, adult spleen was also investigated for the presence of L-DC precursors. Adult splenocytes depleted of T and B cells ($CD19^-Thy1.2^-$) were stained and sorted to isolate the $CD11c^-c\text{-kit}^-$ and $CD11c^-c\text{-kit}^+$ cell populations (Figure 6A). These were cocultured over STX3 stroma for 30 days. Adult spleen $CD11c^-c\text{-kit}^+$ precursors gave rise to large-sized cells (FSC hi ; >93%) over 30-day cultures (Figure 6B). In contrast, the majority of cells arising from $CD11c^-c\text{-kit}^-$ precursors were small cells (FSC lo ; 42%–69%), with very low yield of $CD11c^+$ DCs. Cells recovered, however, had a $CD11c^+CD11b^+MHC-II^-$ phenotype reflecting L-DCs and could have arisen from contaminating $c\text{-kit}^{lo}$ expressing cells not distinguishable by sorting.

The total yield of $CD11c^+$ DCs from cocultures of adult spleen $CD11c^-c\text{-kit}^+$ precursors was 2000-fold higher than from $CD11c^-c\text{-kit}^-$ precursors (Figure 6C). Whereas large-sized cells produced in cocultures of spleen $CD11c^-c\text{-kit}^+$ precursors uniformly expressed $CD11c$ and $CD11b$, progeny were divisible into MHC-II $^-$, MHC-II lo , and MHC-II hi subpopulations. However, MHC-II lo and MHC-II hi populations were only observed early in cultures at 13 and 23 days (Figure 6B), leaving predominantly MHC-II $^-$ LTC-DC-like cells after 30 days. MHC-II $^+$ DCs reduced in number by 4-fold between 23 and 30 days in culture. In contrast, the percentage of MHC-II $^-$ L-DCs increased

2.5-fold over this time. Thus, DC precursor activity was only evident among the $Lin^-CD11c^-c\text{-kit}^+$ cell fraction of adult spleen, and these precursors gave rise to an increasing, stable population of $CD11c^+CD11b^+MHC-II^-$ L-DCs, but a transient population of $CD11c^+CD11b^+MHC-II^+$ cDCs. As with neonatal spleen, the $CD11c^-c\text{-kit}^+$ spleen fraction appears to contain a small number of immediate precursors of MHC-II $^+$ DCs along with a self-renewing progenitor of L-DCs.

The capacity of cocultures to support hematopoiesis could mean either that stromal cell lines are restricted in their hematopoietic support capacity or that $Lin^-c\text{-kit}^+$ precursors are restricted in their differentiative capacity to give only L-DCs. To distinguish these possibilities, the in vivo hematopoietic reconstitution capacity of progenitors was assessed. Adult splenocytes from B6.SJL (CD45.1) mice were depleted of T and B cells ($CD19^-Thy1.2^-$), sorted into $CD11c^-CD11b^-$ (Lin^-) populations of $c\text{-kit}^-$ and $c\text{-kit}^+$ cells. In a first experiment, cells were adoptively transferred into sublethally irradiated C57BL/6J (CD45.2) mice and analyzed at 10, 20, and 41 days for donor reconstitution (Figure 7A). C57BL/6J mice injected with Hanks balanced salt solution only were used as control donors (nil). This study tests the null hypothesis that complete absence of donor cell reconstitution means there are no progenitors among donor cells. A finding of just 1 animal containing donor progeny cells refutes that hypothesis. At 10 days, host mice showed no reconstitution (data not shown); whereas at 20 days, 1 of 4 sublethally irradiated mice given $Lin^-c\text{-kit}^-$ cells showed donor-derived $CD45.1^+$ cells. This result indicates a progenitor frequency of 0.0014% with a lower 95% confidence limit of 0.000064% using the Clopper-Pearson procedure.³² Three of 6 mice given $Lin^-c\text{-kit}^+$ cells displayed donor-derived $CD45.1^+$ cells, evident at 20 days in 1 mouse and at 41 days in 2 others. This represents a progenitor frequency of 0.0035% with a lower 95% confidence limit of 0.00083%. Reconstitution was assessed by staining for lymphoid ($CD8^+CD11c^-$), myeloid ($CD11b^+CD11c^-$), and DC ($CD11c^+MHC-II^+$) subsets.

In a second experiment, donor mice were lethally irradiated and hosts analyzed after 25 weeks. Each of 4 mice given $Lin^-c\text{-kit}^+$ spleen cells showed robust donor cell reconstitution, whereas 1 mouse given donor $Lin^-c\text{-kit}^-$ did not. In this experiment, donor progeny were detected as DCs ($CD11c^+$) and myeloid cells ($CD11b^+CD11c^-$). Donor type B cells ($CD19^+$) and T cells ($CD8^+$) were also identified. Therefore, spleen $Lin^-c\text{-kit}^+$ progenitors demonstrated in vivo differentiative potential for multiple lymphoid, myeloid, and DC subsets.

Overall, the L-DC precursor is contained within the $Lin^-c\text{-kit}^+$ population of spleen, which also contains multipotential progenitors. This study also identifies splenic stromal cocultures as capable of restricted long-term support for differentiation of only L-DCs, and no other DCs or hematopoietic cell type.

Discussion

In this study, we identify the presence of an L-DC precursor among a subset of multipotential progenitors present in spleen. This subset can be demonstrated in both neonatal (8-day) and adult spleen, although neonatal spleen yields higher numbers of these cells. The subset that produces L-DCs in STX3 stromal cocultures appears to be a $Lin^-CD11c^-c\text{-kit}^+CD34^{+/-}Flt3^-$ population, present in both neonatal and adult spleen. In neonatal spleen, L-DC progenitors were present in both the $Lin^-c\text{-kit}^{hi}$ and the $Lin^-c\text{-kit}^{lo}$ subpopulations, although enriched in the latter. They were also present among the $CD34^-$ and $CD34^+$ subpopulations of $Lin^-c\text{-kit}^+$ cells, although they were greatly enriched among the $CD34^+$ subsets. However, L-DC progenitors are present among the $Lin^-c\text{-kit}^+$

Figure 7. In vivo differentiative potential of spleen precursors. Splenocytes of adult B6.SJL mice were depleted of T and B cells and sorted to give CD11c⁻CD11b⁻ (Lin⁻) populations of c-kit⁻ and c-kit⁺ cells. Sorted cells (2×10^4) were transferred intravenously into sublethally irradiated (4.5 Gy) C57BL/6J (CD45.2⁺) recipient mice. Negative "nil" control mice were injected with Hanks balanced salt solution only. For analysis of short-term cell development, spleen was collected at 20 or 41 days and stained with antibodies specific for CD45.1, CD11c, CD11b, CD8 α , and MHC-II. Before flow cytometry, cells were incubated with PI (1 μ g/mL) for gating of live (PI⁻) cells. Splens showing CD45.1⁺ progeny (> 0.05%) were examined for donor myeloid cells (CD11b⁺CD8⁻), cDCs (CD11c⁺MHC-II⁺), and lymphoid cells (CD11b⁻CD8⁺). Numbers in gates indicate percentage positive cells. In a second similar experiment, Lin⁻c-kit⁻ and Lin⁻c-kit⁺ cells were sorted and transferred into lethally irradiated (9.5 Gy) host mice and analyzed for donor progeny (CD45.1⁺) after 25 weeks. T/B-cell depleted spleen was analyzed for DC (CD11c⁺) and myeloid cell (CD11b⁺CD11c⁻) progeny. T/B-cell enriched spleen was analyzed for lymphoid T (CD8⁺) and B (CD19⁺) cells.

subset of adult spleen, which contains hematopoietic stem cells. Because no CD34 subset was found among adult spleen, c-kit, rather than CD34, is a defining marker of L-DC progenitors. The possibility that the L-DC progenitor is endogenous to spleen is supported by evidence that blood does not contain precursors, which seed cocultures for L-DC production.^{24,25}

Although adult Lin⁻c-kit⁺ spleen progenitors produce specifically L-DCs in cocultures, this restricted development was not replicated in vivo. This indicates a limitation of stroma for supporting only the development of L-DCs from overlaid spleen or BM cells.²⁴ Immediate DC precursors have been defined as cells already expressing lineage markers, such as CD11c^{int} pre-cDCs¹⁴ or CD11c⁻CD8 α ⁺ precursors of CD8 α ⁺ cDCs.³³ In addition, in blood, precursors for cDCs and pDCs have been identified as cells

expressing low levels of CD11c.³⁴ The Lin⁻c-kit⁺ subset of spleen does not contain immediate DC precursors because they do not give DC development after 10 days after transfer into mice given sublethal irradiation (4.5 Gy; data not shown). This type of experimentation has been used successfully by others to demonstrate the presence of pre-cDCs and pre-CD8⁺ cDCs in spleen.^{14,33} The earliest evidence of progeny from adoptively transferred cells was at approximately 20 days in sublethally irradiated mice. Dendritic and myeloid cell development was detected, perhaps arising from a myeloid progenitor because there was no evidence of lymphopoiesis. Evidence for multipotentiality was obtained at 40 days, but only on adoptive transfer of Lin⁻c-kit⁺ cells. More complete multilineage reconstitution was confirmed in lethally irradiated hosts held for more than 25 weeks. These experiments verified the potential of Lin⁻c-kit⁺ cells to give rise to lymphoid, myeloid, and cDC subsets, indicative of multipotentiality. Only small numbers of donor cells and the difficulty of identifying donor-derived progeny limited these experiments.

The splenic L-DC progenitor is distinct from the common dendritic progenitor and macrophage dendritic progenitor described in BM. The absence of Flt3 expression on c-kit⁺ L-DC progenitors in spleen is consistent with recent evidence that the common dendritic progenitor, a c-kit⁺Flt3⁺ progenitor of cDCs and pDCs, is not present in spleen.^{15,29} The L-DC progenitor is also distinct from the macrophage dendritic progenitor in BM, a c-kit⁺Flt3⁺ progenitor of macrophages and DCs.^{29,35} Production of L-DCs in cocultures or LTCs occurs independently of Flt3-L because splenic stromal cell lines that support L-DCs do not produce Flt3-L.³⁶ L-DCs appear therefore to develop independently of cDCs and pDCs, which develop both in vitro and in vivo in response to Flt3-L.^{12,37}

Culture of neonatal and adult c-kit⁺ spleen precursors over STX3 stroma resulted in generation of only myeloid CD11b⁺ DCs, consistent with earlier studies demonstrating spleen stroma-directed in vitro development of CD11b⁺ myeloid DCs from BM c-kit⁺ progenitors.^{38,39} However, using defined precursor subsets from spleen, we here distinguish in vitro development of 2 distinct myeloid DC populations: a CD11c⁺CD11b^{lo}MHC-II⁺ CD8 α ⁻ cDCs or monocyte-derived DCs, and a CD11c^{lo}CD11b^{hi}MHC-II⁺ L-DC population. Although populations of MHC-II⁺ DCs were evident in productive cultures, their numbers declined over 30 days, and they became the minority population compared with L-DCs (Figure 6). We have now shown production of L-DCs, but not MHC-II⁺ DCs, in cocultures maintained for 6 to 12 months (data not shown). All data are consistent with the presence of a committed progenitor for L-DCs present in spleen, which we characterize as a Lin⁻c-kit⁺CD34⁻/Flt3⁻ cell subset with lymphomyeloid differentiative capacity. These results identifying DC progenitors within spleen conflict with the common model for DC hematopoiesis where progenitors reside and differentiate in BM, so releasing committed DC precursors into blood that enter spleen for further differentiation. This finding is, however, consistent with a model for DC hematopoiesis endogenous to spleen and dependent on maintenance of early Lin⁻c-kit⁺Flt3⁻ progenitors that are also clearly distinct from the CD11c^{lo} immediate precursors of cDCs and pDCs described previously.¹⁴

Spleen LTCs have been shown to maintain the production of immature, myeloid dendritic-like cells for years.³⁴ This property of cell production is consistent with the maintenance of self-renewing progenitors within the stromal microenvironment. In a previous study, progenitors in LTCs were identified among the heterogeneous small cell population previously essential for continued

production of LTC-DCs.^{16,19,40} Here, similar precursors of L-DCs have been identified as Lin⁻c-kit⁺ cells in both neonatal and adult spleen, consistent with the spleen origin of LTCs. The Lin⁻c-kit^{hi} spleen subset resembles hematopoietic stem cells in BM as Lin⁻Sca1⁺c-kit⁺Flt3⁻⁴¹ or Lin⁻Sca1⁺c-kit⁺CD150⁺ cells.⁴² Long-term spleen cultures, or the spleen stromal cocultures described here, therefore represent a potentially important in vitro model system in which to study both the maintenance of hematopoietic progenitors and the microenvironmental signals supporting hematopoiesis of novel DCs.

Acknowledgments

T. J. O'Neill (Professor of Applied Statistics, ANU) provided statistical support and advice for this project.

References

1. Banchereau J, Steinman RM. Dendritic cells and the control of immunity. *Nature*. 1998;392(6673):245-252.
2. Liu K, Iyoda T, Saternus M, et al. Immune tolerance after delivery of dying cells to dendritic cells *in situ*. *J Exp Med*. 2002;196(8):1091-1097.
3. Vermaelen KY, Carro-Muino I, Lambrecht BN, et al. Specific migratory dendritic cells rapidly transport antigen from the airways to the thoracic lymph nodes. *J Exp Med*. 2001;193(1):51-60.
4. Cella M, Salio M, Sakakibara Y, et al. Maturation, activation, and protection of dendritic cells induced by double-stranded RNA. *J Exp Med*. 1999;189(5):821-829.
5. Xu Y, Zhan Y, Lew AM, et al. Differential development of murine dendritic cells by GM-CSF versus Flt3 ligand has implications for inflammation and trafficking. *J Immunol*. 2007;179(11):7577-7584.
6. Vremec D, Pooley J, Hochrein H, et al. CD4 and CD8 expression by dendritic cell subtypes in mouse thymus and spleen. *J Immunol*. 2000;164(6):2978-2986.
7. Asselin-Paturel C, Boonstra A, Dalod M, et al. Mouse type I IFN-producing cells are immature APCs with plasmacytoid morphology. *Nat Immunol*. 2001;2(12):1144-1150.
8. Nakano H, Yanagita M, Gunn MD. CD11c(+)B220(+)Gr-1(+) cells in mouse lymph nodes and spleen display characteristics of plasmacytoid dendritic cells. *J Exp Med*. 2001;194(8):1171-1178.
9. Inaba K, Inaba M, Romani N, et al. Generation of large numbers of dendritic cells from mouse bone marrow cultures supplemented with granulocyte/macrophage colony-stimulating factor. *J Exp Med*. 1992;176(6):1693-1702.
10. Sallusto F, Lanzavecchia A. Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor plus interleukin 4 and downregulated by tumor necrosis factor alpha. *J Exp Med*. 1994;179(4):1109-1118.
11. Brasel K, De Smedt T, Smith JL, et al. Generation of murine dendritic cells from flt3-ligand-supplemented bone marrow cultures. *Blood*. 2000;96(9):3029-3039.
12. Naik SH, Proietto AI, Wilson NS, et al. Cutting edge: generation of splenic CD8⁺ and CD8⁻ dendritic cell equivalents in Fms-like tyrosine kinase 3 ligand bone marrow cultures. *J Immunol*. 2005;174(11):6592-6597.
13. Naik SH, Sathe P, Park HY, et al. Development of plasmacytoid and conventional dendritic cell subtypes from single precursor cells derived in vitro and in vivo. *Nat Immunol*. 2007;8(11):1217-1226.
14. Naik SH, Metcalf D, van Nieuwenhuijze A, et al. Intraspinal steady-state dendritic cell precursors that are distinct from monocytes. *Nat Immunol*. 2006;7(6):663-671.
15. Onai N, Obata-Onai A, Schmid MA, et al. Identification of clonogenic common Flt3+M-CSFR+ plasmacytoid and conventional dendritic cell progenitors in mouse bone marrow. *Nat Immunol*. 2007;8(11):1207-1216.
16. Wilson HL, Ni K, O'Neill HC. Identification of progenitor cells in long-term spleen stromal cultures that produce immature dendritic cells. *Proc Natl Acad Sci U S A*. 2000;97(9):4784-4789.
17. O'Neill HC, Wilson HL, Quah B, et al. Dendritic cell development in long-term spleen stromal cultures. *Stem Cells*. 2004;22(4):475-486.
18. Quah B, Ni K, O'Neill HC. In vitro hematopoiesis produces a distinct class of immature dendritic cells from spleen progenitors with limited T cell stimulation capacity. *Int Immunol*. 2004;16(4):567-577.
19. Wilson HL, O'Neill HC. Dynamics of dendritic cell development from precursors maintained in stroma-dependent long-term cultures. *Immunol Cell Biol*. 2003;81(2):144-151.
20. Despars G, Ni K, Bouchard A, et al. Molecular definition of an *in vitro* niche for dendritic cell development. *Exp Hematol*. 2004;32(3):1182-1193.
21. Despars G, O'Neill HC. Splenic endothelial cell lines support development of dendritic cells from bone marrow. *Stem Cells*. 2006;24:1496-1504.
22. Despars G, Periasamy P, Tan J, et al. Gene signature of stromal cells which support dendritic cell development. *Stem Cells Dev*. 2008;17(5):917-927.
23. Despars G, Tan J, Periasamy P, et al. The role of stroma in hematopoiesis and dendritic cell development. *Curr Stem Cell Res Ther*. 2007;2(1):23-29.
24. Periasamy P, Tan JKH, Griffiths KL, et al. Splenic stromal niches support hematopoiesis of dendritic-like cells from precursors in bone marrow and spleen. *Exp Hematol*. 2009;37:1060-1071.
25. Ni K, O'Neill H. Spleen stromal cells support hematopoiesis and *in vitro* growth of dendritic cells from bone marrow. *Br J Haematol*. 1999;105(1):58-67.
26. Ni K, O'Neill HC. Long-term stromal cultures produce dendritic-like cells. *Br J Haematol*. 1997;97(4):710-725.
27. Ni K, O'Neill HC. Hemopoiesis in long-term stroma-dependent cultures from lymphoid tissue: production of cells with myeloid/dendritic characteristics. *In Vitro Cell Dev Biol Anim*. 1998;34(4):298-307.
28. Hogquist KA, Jameson SC, Heath WR, et al. T cell receptor antagonist peptides induce positive selection. *Cell*. 1994;76(1):17-27.
29. Liu K, Vitoria GD, Schwickert TA, et al. In vivo analysis of dendritic cell development and homeostasis. *Science*. 2009;324:392-397.
30. Lai AY, Lin SM, Kondo M. Heterogeneity of Flt3-expressing multipotent progenitors in mouse bone marrow. *J Immunol*. 2005;175(8):5016-5023.
31. Wolber FM, Leonard E, Michael S, et al. Roles of spleen and liver in development of the murine hematopoietic system. *Exp Hematol*. 2002;30(9):1010-1019.
32. Puza B, O'Neill TJ. Generalised Clopper-Pearson confidence intervals for the binomial proportion. *J Statist Comput Simulation*. 2006;76:489-508.
33. Bedoui S, Prato S, Mintern L, et al. Characterization of an immediate splenic precursor of CD8⁺ dendritic cells capable of inducing antiviral T cell responses. *J Immunol*. 2009;182(7):4200-4207.
34. O'Keeffe M, Hochrein H, Vremec D, et al. Dendritic cell precursor populations of mouse blood: identification of the murine homologues of human blood plasmacytoid pre-DC2 and CD11c(+) DC1 precursors. *Blood*. 2003;101(4):1453-1459.
35. Fogg DK, Sibon C, Miled C, et al. A clonogenic bone marrow progenitor specific for macrophages and dendritic cells. *Science*. 2006;311(5757):83-87.
36. Despars G, Ni K, Bouchard A, et al. Molecular definition of an *in vitro* niche for dendritic cell development. *Exp Hematol*. 2004;32(12):1182-1193.
37. Naik SH, Corcoran LM, Wu L. Development of murine plasmacytoid dendritic cell subsets. *Immunol Cell Biol*. 2005;83(5):563-570.
38. Svensson M, Maroof A, Ato M, et al. Stromal cells direct local differentiation of regulatory dendritic cells. *Immunity*. 2004;21(6):805-816.
39. Tang H, Guo Z, Zhang M, et al. Endothelial stroma programs hematopoietic stem cells to differentiate into regulatory dendritic cells through IL-10. *Blood*. 2006;108(4):1189-1197.
40. Wilson HL, O'Neill HC. Identification of differentially expressed genes representing dendritic cell precursors and their progeny. *Blood*. 2003;102(5):1661-1669.
41. Christensen JL, Weissman IL. Flk-2 is a marker in hematopoietic stem cell differentiation: a simple method to isolate long-term stem cells. *Proc Natl Acad Sci U S A*. 2001;98(25):14541-14546.
42. Kiel MJ, Yilmaz OH, Iwashita T, et al. SLAM family receptors distinguish hematopoietic stem and progenitor cells and reveal endothelial niches for stem cells. *Cell*. 2005;121(7):1109-1121.

This work was supported by ANU institutional funding (H.O.). J.K.H.T. and P.P. were supported by ANU Graduate Scholarships.

Authorship

Contribution: J.K.H.T. designed, implemented, and analyzed the experiments and prepared the manuscript; P.P. designed, performed, and analyzed the experiments; and H.C.O. supervised the project and contributed to design and analysis of experiments and preparation of the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Helen C. O'Neill, Research School of Biology, Bldg No. 41, Linnaeus Way, Australian National University, Canberra, ACT, Australia, 0200; e-mail: Helen.ONeill@anu.edu.au.