6 RNA Viruses and RNA-Based Drugs

*A Perfect Match for
RNA Delivery and the
Identification of Candidate
Therapeutic Target
Inflammatory Molecules*

Brett A. Lidbury, Cristina M. Musso, Jasjit Johal,
Nestor E. Rulli, and Suresh Mahalingam
Virus and Inflammation Research Group,
Centre for Biomolecular and Chemical
Sciences, University of Canberra

Mark T. Heise
The Carolina Vaccine Institute, Department of Genetics,
and Department of Microbiology and Immunology,
University of North Carolina at Chapel Hill

CONTENTS

6.1	Introduction.....	116
6.2	RNA Virus Vectors and Replicon Technologies.....	116
6.2.1	Replication-Competent Viral Vectors.....	117
6.2.2	Propagation-Defective Viral Vectors.....	118
6.3	RNA Virus Evasion of the Interferon System.....	119
6.3.1	Human Immunodeficiency Virus.....	120
6.3.2	Influenza Virus	122
6.3.3	Ross River Virus (RRV)	123
6.4	Concluding Remarks.....	125
	Acknowledgments.....	126
	References.....	126

6.1 INTRODUCTION

Viruses have an unmatched genius at infiltrating cells and manipulating the expression of host genes that influence viral replication and survival. This complexity of virus–host interaction presents significant challenges to contemporary medical science, with viral diseases continuing to cause mortality and morbidity worldwide. Viruses of current major concern to human health show the majority to be those comprising an RNA genome (e.g., HIV, influenza, dengue) [1]. Such RNA virus capacity in terms of the manipulation of host-cell function, in terms of potential RNA-based gene therapies and vaccines, does have positive dimensions:

1. The ability of such viruses to infiltrate cell membranes and deliver genetic material to the cell interior, and thereafter exploit the cell's molecular machinery to replicate viral genes
2. The complexities of the virus–host relationship that emphasize the crucial host molecules/pathways responsible for antiviral defense, as well as inflammation control.

This chapter focuses on two aspects learned from RNA viruses:

1. A review of RNA viral vectors (e.g., alphaviral “replicons”) and their potential to deliver therapeutic RNA molecules to individuals suffering from a disease
2. The identification of host inflammatory pathways and immune responses that, because of their potent antiviral activities, are targeted by viruses for disruption, evasion, or sabotage.

The combination of knowledge in gene delivery and in RNA virus-manipulated host defense responses provides an exciting future arena for RNA-based drug targets, as well as for gene-therapy vector designs derived from engineering RNA virus genomes.

6.2 RNA VIRUS VECTORS AND REPLICON TECHNOLOGIES

Since the demonstration by Racaniello and Baltimore in 1981 that RNA from a molecularly cloned poliovirus was infectious [2], there has been significant interest in utilizing RNA viruses as vectors for vaccination or gene therapy. For many years, much of this work was limited to positive-sense RNA viruses, such as the picornaviruses, togaviruses, and flaviviruses, due to the development of infectious cDNA clones for many of these viruses [2–4]. However, in recent years, infectious cDNA clones have been generated for negative-sense RNA viruses, as well as the large coronaviruses, raising the possibility of generating expression vectors based on many different virus types.

RNA virus vectors, which for the purpose of this discussion are viruses or their derivatives that express heterologous genetic material that has been engineered into the viral genome, can be classified either as replication-competent viral vectors or replication-defective “replicon vectors.” Replication-competent vectors are

RNA Viruses and RNA-Based

fully functional viruses that i virions that will infect subsec which can be as small as the c nucleotides in length, is also e petent vectors are generally t the production of progeny vi of the missing viral sequenc genes, which encode the prot the genes encoding the viral r remain intact. Therefore, if t permissive cell, viral RNA sy will prevent the production o *vitro* by simply introducing t it is often possible to supply t defective genome to be packa of only a single round of repl

6.2.1 REPLICATION-COMPE

Replication-competent viral ber of virus families, includ use, or intended use, of these expressed heterologous nucle epitopes to entire open readin express heterologous genetic involves the insertion of short additional sequence. This ap where antigenic sequences fr the glycoproteins of Sindbis that is nonessential for viral for coronaviruses, the transn engineered to express heterol reading frames [8], while in t 4 can be replaced with a het expressed from the viral ge expression of the heterologo with alphaviruses, where ad introduced at the beginning tional subgenomic promoter sequence in infected cells [10

There are several advant expression vectors. One maj they can be propagated in c greatly decreases the cost of tor to spread and replicate w tion of more robust immu

manipulating the expression. This complexity of contemporary medical science and morbidity worldwide, the majority to be those [1]. Such RNA virus, in terms of potential dimensions:

es and deliver genetic the cell's molecular

it emphasize the crucial defense, as well as

viruses:

ions") and their potentials suffering from a

nd immune responses targeted by viruses for

A virus-manipulated host RNA-based drug targets, engineering RNA virus

TECHNOLOGIES

1981 that RNA from a has been significant inter-gene therapy. For many NA viruses, such as the development of infectious in recent years, infectious A viruses, as well as the expression vectors based on

ssion are viruses or their that has been engineered ion-competent viral vec-m-competent vectors are

fully functional viruses that are capable of infecting a cell and producing progeny virions that will infect subsequent cells. However, a heterologous genetic sequence, which can be as small as the coding sequence of a T-cell epitope or several thousand nucleotides in length, is also expressed from the viral genome. Nonreplication-competent vectors are generally lacking one or more viral genes that are essential for the production of progeny virions, with the heterologous gene expressed in place of the missing viral sequences. Commonly, the missing genes are viral structural genes, which encode the proteins that physically make up the virion particle, while the genes encoding the viral replicase proteins, which mediate viral RNA synthesis, remain intact. Therefore, if the RNA for the defective vector is introduced into a permissive cell, viral RNA synthesis will commence, but the lack of structural genes will prevent the production of progeny virions. Though these vectors can be used *in vitro* by simply introducing the defective viral genome RNA into a permissive cell, it is often possible to supply the viral structural genes *in trans*, which will allow the defective genome to be packaged, with the resulting defective viral particles capable of only a single round of replication.

6.2.1 REPLICATION-COMPETENT VIRAL VECTORS

Replication-competent viral expression vectors have been constructed for a number of virus families, including togaviruses and coronaviruses [5, 6]. The major use, or intended use, of these vectors is in vaccination, where the engineered virus expressed heterologous nucleic acid sequences ranging in size from T- and B-cell epitopes to entire open reading frames. Three basic strategies have been employed to express heterologous genetic material from replication-competent vectors. The first involves the insertion of short genetic sequences into viral genes that will tolerate the additional sequence. This approach has been successfully applied to alphaviruses, where antigenic sequences from heterologous pathogens have been engineered into the glycoproteins of Sindbis virus [7]. A second approach is to replace a viral gene that is nonessential for viral replication with a heterologous sequence. For example, for coronaviruses, the transmissible gastroenteritis virus (TGEV) genome has been engineered to express heterologous genes in place of the nonessential 3a and 3b open reading frames [8], while in the mouse hepatitis virus genome, the nonessential gene 4 can be replaced with a heterologous gene [9]. Finally, a genetic sequence can be expressed from the viral genome by engineering in a new viral promoter to drive expression of the heterologous sequence. This approach has been used extensively with alphaviruses, where additional copies of the 26S subgenomic promoter can be introduced at the beginning of the viral 3' UTR (untranslated region). This additional subgenomic promoter will then drive high-level expression of the heterologous sequence in infected cells [10].

There are several advantages and disadvantages to using replication-competent expression vectors. One major advantage with replication-competent vectors is that they can be propagated in culture by simply allowing the vector to grow, which greatly decreases the cost of producing/packaging the vector. The ability of the vector to spread and replicate within the inoculated host may also result in the generation of more robust immune responses against the vector and the protein derived

from the heterologous sequence. However, though many virus-based expression vectors are derived from existing live attenuated vaccines, such as the 17D strain of yellow fever virus [4], there are obviously some safety concerns whenever a replication-competent vector is used, especially in highly susceptible populations, such as immunocompromised individuals. Many vectors also have limitations with respect to how large a sequence can be inserted, with some vectors limited to the coding sequence for one or a few B- or T-cell epitopes. Finally, vector stability is also an issue, where many recombinant viruses lose transgene expression after a few rounds of replication [11].

6.2.2 PROPAGATION-DEFECTIVE VIRAL VECTORS

In addition to replication-competent viral vectors, propagation-defective viral vectors, or "replicons," have been developed for a number of RNA viruses. Replicon vectors are generally made by removing one or more of the viral structural genes from the viral genome, which results in a virus that can initiate viral RNA synthesis and gene expression, but is unable to produce infectious progeny virions. The missing gene or genes can then be provided in *trans* to package the viral RNA, with the resulting "single-hit" particle (replicon particle) able to infect a new target cell and initiate viral RNA synthesis and gene expression, but not produce progeny virion due to the lack of virally encoded structural genes. This approach has been used effectively with a number of positive-sense RNA viruses, including picornaviruses, togaviruses, flaviviruses, and coronaviruses, as well as negative-sense RNA viruses such as vesicular stomatitis virus (VSV) [12–15].

Alphavirus vectors provide an excellent example of the power of replicon vectors for gene delivery. Though the genome is a single-stranded RNA of positive-sense polarity, the genome can readily be broken down into two functional units. The 5' two-thirds of the viral genome encodes the viral nonstructural genes (nSP), which are essential for viral RNA synthesis, while the 3' one-third of the viral genome encodes the viral structural genes, which are expressed from a highly active subgenomic promoter. Therefore, the structural genes, which are made up of a capsid gene, two major glycoproteins (E2 and E1), as well as two additional proteins (E3 and 6K) that serve as signal sequences, can be removed from the genome without affecting viral RNA synthesis. A heterologous gene can then be expressed from the subgenomic promoter in place of the structural genes. The resulting replicon RNA will initiate viral RNA synthesis if introduced into a permissive cell, and drive high-level expression of the heterologous gene, though no progeny virions will be produced [16]. The replicon RNA can be packaged by providing helper RNA, which contains the alphavirus 5' and 3' ends, as well as the subgenomic promoter driving expression of the viral structural genes, but which have deletion of the viral nonstructural genes [17]. If this helper RNA is introduced into the same cell as the replicon RNA, the replicon-derived nonstructural proteins will amplify the helper RNA in *trans*, and drive expression of the viral structural genes from the helper. The structural proteins will then package the replicon RNA to produce replicon particles, while the helper RNA, which lacks an RNA packaging signal [16], will not be packaged, or will be packaged inefficiently. In order to decrease the likelihood of a recombination event

RNA Viruses and RNA-Based

between the helper and the re-
tent virus, a split helper system
RNA and the viral glycoprotein
used [17]. Variations on this theme
the alphavirus structural gene
lison particles. Vectors based
vaccine delivery or in short-term

Single-hit vectors have also
viruses, including VSV. Defective
viral G protein, which is essential
erologous genes can be inserted
G protein then provided in *trans*.
malian expression vector driving
the fact that the defective VSV
wide range of viruses, which vary
depending on which viral glycoprotein

6.3 RNA VIRUS EVASION

It is widely published that DNA
antiviral immune responses can
"steal" genes from the host that
fewer examples exist for small
evasion strategies employed by
tumor immune evasion strategies
the human immunodeficiency
(RRV), with a special focus on
are crucial in exerting antiviral
ous IFN evasion mechanisms
ment of appropriate drugs or
other virus pathogens.

The activation of host proteins
to correlate with early defense
RNA viruses. Because of this
or neutralize their antiviral proteins
through various intracellular
receptors, which are present
tion factors such as STAT (sig-
become phosphorylated followed
by the translocation of
and STAT-2 heterodimers) to
several antiviral genes (IFN- γ -
inducible synthetase [OAS], nitric
oxide synthase [iNOS], and
IRF-7, which become active
various other downstream an-

Profiles in Drug Design

virus-based expression such as the 17D strain of rns whenever a replicable populations, such as imitations with respect s limited to the coding ctor stability is also an ssion after a few rounds

ion-defective viral vec- RNA viruses. Replicon e viral structural genes iate viral RNA synthe- s progeny virions. The ackage the viral RNA, e to infect a new target ut not produce progeny This approach has been s, including picornavi- as negative-sense RNA

ower of replicon vectors RNA of positive-sense unctional units. The 5' ral genes (nSP), which rd of the viral genome a highly active subge- ade up of a capsid gene, al proteins (E3 and 6K) ome without affecting ressed from the subge- ing replicon RNA will ell, and drive high-level ions will be produced r RNA, which contains oter driving expression ral nonstructural genes the replicon RNA, the per RNA in *trans*, and The structural proteins ticles, while the helper e packaged, or will be a recombination event

between the helper and the replicon RNA that would generate a replication-competent virus, a split helper system, where the viral capsid is expressed from one helper RNA and the viral glycoproteins are expressed from a second helper, can also be used [17]. Variations on this theme, including packaging cell lines that stably express the alphavirus structural genes [17], have also been used to produce alphavirus replicon particles. Vectors based on several different alphaviruses have been used for vaccine delivery or in short-term gene therapy applications.

Single-hit vectors have also been constructed for a number of negative-sense RNA viruses, including VSV. Defective VSV vectors can be generated by deletion of the viral G protein, which is essential for the production of infectious viruses [15]. Heterologous genes can be inserted in place of G within the VSV genome, and the viral G protein then provided in *trans* using a packaging cell line transduced with a mammalian expression vector driving VSV G expression. One advantage of these vectors is the fact that the defective VSV particles can be pseudotyped with glycoproteins from a wide range of viruses, which will allow the vectors to be directed to specific cell types, depending on which viral glycoprotein is used for packaging [15].

6.3 RNA VIRUS EVASION OF THE INTERFERON SYSTEM

It is widely published that DNA viruses utilize a number of strategies to evade the antiviral immune responses of the host [18]. DNA viruses are thought to have "stolen" genes from the host that were modified to benefit virus survival [18]. However, fewer examples exist for smaller-genome RNA viruses, and information on immune evasion strategies employed by this class of viruses is emerging [1]. Several important immune evasion strategies employed by RNA viruses are discussed below for the human immunodeficiency virus (HIV), influenza virus, and Ross River virus (RRV), with a special focus on the interferon (IFN) system. The type I interferons are crucial in exerting antiviral effects against RNA viruses. Understanding the various IFN evasion mechanisms employed by these viruses will assist in the development of appropriate drugs or other therapeutic strategies in combating these and other virus pathogens.

The activation of host proteins such as IFNs, following viral infection, appears to correlate with early defense mechanisms, since IFNs act to inhibit both DNA and RNA viruses. Because of this, viruses often target IFNs in their attempt to modify or neutralize their antiviral potency [19]. The antiviral effects of IFNs are mediated through various intracellular pathways and are initiated by their binding to cellular receptors, which are present on most cells [20] (Figure 6.1). For instance, transcription factors such as STAT (signal transducer and activator of transcription) complexes become phosphorylated following binding of type I IFNs to its receptor. This is followed by the translocation of phosphorylated STAT complexes (consisting of STAT-1 and STAT-2 heterodimers) to the nucleus, where they bind to transcription elements of several antiviral genes (IFN- β , RNA-dependent protein kinase [PKR]), 2' 5' oligoadenylate synthetase [OAS], nitric oxide [NO], as well as secondary transcription factors [interferon-regulatory factor 1 [IRF-1], IRF-3, IRF-7]. Genes such as IRF-1, IRF-3, and IRF-7, which become activated, also play important roles in the transcription of various other downstream antiviral genes. The production of these antiviral proteins

FIGURE 6.1 Virus induction of Type I IFNs. A and B represent primary events. C represents host response. Arrows (→) indicate stimulation or induction. Dashed lines (---) indicate suppression or inhibition. Virus infection and double-stranded RNA (dsRNA) will trigger the early activation of different cell proteins (PKR, NF-κB, 2'5' OAS, IRF3) that mediate the primary expression of IFN alpha/beta. Type I IFN interaction with the specific cell receptor stimulates Jak1 and Tyk2 proteins. The activation of STAT-1 and STAT-2 is followed by the formation of an ISGF3 (interferon-stimulated gene factor-3) complex. ISGF3 will bind to the interferon receptor ISRE, promoting IFN synthesis and IRF-7 induction.

can directly disrupt the regulation of viral and cellular macromolecular synthesis and degradation. In recent years, many human RNA viruses have been shown to interrupt/disrupt the IFN system by targeting various proteins in the signaling cascade, such as STAT complexes, IRF proteins, 2'5' OAS, and PKR [1, 19, 20].

6.3.1 HUMAN IMMUNODEFICIENCY VIRUS

Human Immunodeficiency virus (HIV) is of major concern worldwide, with devastating health impacts seen particularly in the developing world. HIV infection leads to the condition known as acquired immunodeficiency syndrome (AIDS), and is transmitted primarily through infected blood or semen. HIV is extremely successful in modulating various compartments of the host immune responses to its benefit. In the past 10 years, studies have shown that several HIV proteins, such as Nef, Env, and Vpu, can cripple the host immune responses. These proteins are shown to abolish CD4 surface protein from infected cells [21], resulting in the inability of T cells to engage with antigen-presenting cells and mount an immune response. Additionally, the Nef protein can trigger the down-regulation of MHC-I molecules, suppress apoptosis of infected cells by inhibiting the apoptosis-associated kinase 1 (ASK 1), as well as stimulate the expression of Fas ligands on T cells, inducing their death [22].

In addition, immune re- HIV-1 Tat protein, a poten cycle of this virus [23]. Tat of mononuclear cells toward conserved amino acid sequ Furthermore, it is possible class I and II molecules, a erature [24–26].

The binding of the vi fusion, and dsRNA activati of several early viral genes. tion response) RNA bindin binding proteins were show RNA forms an inactive hel phosphorylation of PKR at

HIV

FIGURE 6.2 Early event of HIV infection. HIV-1 Tat protein, a potential cycle of this virus [23]. Tat of mononuclear cells toward conserved amino acid sequences. Furthermore, it is possible class I and II molecules, a erature [24–26]. The binding of the virus fusion, and dsRNA activation of several early viral genes. tion response) RNA binding proteins were shown RNA forms an inactive hel phosphorylation of PKR at

it primary events. C represents. Dashed lines (–) indicate A (dsRNA) will trigger the S, IRF3) that mediate the h the specific cell receptor STAT-2 is followed by the lex. ISGF3 will bind to the action.

macromolecular synthesis have been shown to interfere in the signaling cascade, [1, 19, 20].

worldwide, with devastating HIV infection leads to the AIDS), and is transmitted by successful in modulating its benefit. In the past 10 s Nef, Env, and Vpu, can abolish CD4 surface of T cells to engage with Additionally, the Nef protein induces apoptosis of infected cells, as well as stimulate the J.

In addition, immune regulatory processes may also be manipulated through the HIV-1 Tat protein, a potent monocyte chemoattractant expressed early in the life cycle of this virus [23]. Tat has been shown to facilitate infection by the recruitment of mononuclear cells toward HIV-producing cells, possible through its expression of conserved amino acid sequences that correspond to critical chemokine sequences. Furthermore, it is possible that Tat plays a role in the down-regulation of both HLA class I and II molecules, although there is much controversy present within the literature [24–26].

The binding of the virus to a specific cell receptor, viral cellular membrane fusion, and dsRNA activates NF- κ B, resulting in a strong transcriptional stimulation of several early viral genes. Tat has also been shown to inhibit HIV-1 Tar (transactivation response) RNA binding and activation of 2'-5' OAS. The HIV-1 Tat and Tar RNA binding proteins were shown to inhibit dsRNA-mediated activation of PKR [27]. Tar-RNA forms an inactive heterodimer with PKR. Tat, on the other hand, inhibits autophosphorylation of PKR and competes with eIF2 α [28, 29] (Figure 6.2).

FIGURE 6.2 Early events after HIV infection of a host cell. After internalization and uncoating, reverse transcription results in the production of double-stranded DNA. The viral DNA is integrated into the host genome. Transcription results in various viral mRNAs encoding the regulatory and structural proteins to be translated in the cytosol (steps 1–5). The binding of the virus to a specific cell receptor, viral cellular membrane fusion, and dsRNA activates NF- κ B, resulting in a strong transcriptional stimulation of several early viral genes. TAR and TAT are known to down-regulate PKR function. TAR-RNA forms an inactive heterodimer with PKR. TAT inhibits autophosphorylation of PKR and competes with eIF2 α . NEF induces down-regulation of MHC I molecules and inhibits apoptosis of the infected cells by inhibiting the apoptosis-associated kinase 1 (ASK 1), but stimulates the expression of Fas ligands on the surface of T cells, inducing their death.

Other research has demonstrated the ability of HIV to manipulate the chemokine system. It was shown that chemokine receptors were utilized as co-receptors to facilitate entry into CD4⁺ cells [30]. Activity of cells expressing CD4 as well as CCR5 and CXCR4 may be influenced by the HIV structural protein, gp120, and benefited by its interaction with cellular receptors for viral entry. Such binding activity by gp120 can elicit not only apoptosis in CD4⁺ T cells and endothelial cells, but other outcomes affecting function, including neurodegenerative effects and dysregulated lymphocyte homing [31].

6.3.2 INFLUENZA VIRUS

The influenza virus continues to pose significant global health problems, causing morbidity and mortality worldwide. In the United States, mortality rates over the last 30 years have ranged between 30,000 to 70,000 cases per year, with the elderly and those who are immunocompromised most at risk of death after influenza virus infection [32].

Insights into influenza pathogenesis have been helped over the past decade by studies into the modulation of the host type I IFN response via influenza proteins postinfection. The nonstructural protein of influenza A, NS1, has been shown to have a role in inhibiting the host activation of type I IFNs [33, 34], associated with the disruption of PKR activation by NS1 binding to dsRNA [35]. NF- κ B and other transcription factors play a role in the transactivation of the IFN- β promoter, and in addition to the capacity for influenza to block the PKR response, NS1 can inhibit the activation of both NF- κ B and IRF-3 [36, 37], thus compromising transcriptional activation pathways associated with IFN expression (see Figure 6.3). Therefore, it is evident that NS1 has an immune evasion role in addition to its polymerase activity, suggesting that viral proteins are capable of executing diverse functions within the viral life cycle. For these and other studies, an influenza virus was engineered with a deleted NS1 gene (delNS1), and early results showed that the delNS1 influenza virus could only replicate in IFN-deficient cells [33]. In IFN-intact cells, delNS1 influenza virus infection stimulated increased type I IFN mRNA transcription [33, 34], as well as high levels of NF- κ B activation, in comparison to the NS1-competent parent virus [37], which suggests that the growth inhibition of delNS1 influenza was a result of the robust IFN response postinfection.

Subsequent work on influenza virus interaction with the type I IFN response has also revealed some of the subtle mechanisms employed by the virus to disrupt the IFN antiviral effect. As discussed earlier, influenza infection disrupts PKR activity in the infected host cell via the viral NS1 protein, allowing enhanced influenza virus replication [35]. The early IFN response to influenza infection has found a critical role for the host PKR inhibitory protein, P58^{IPK}, which, through binding to the PKR kinase domain, leads to the suppression of PKR-mediated phosphorylation of eIF2 α [38, 39], a key intracellular protein required for both host and viral protein translation. P58^{IPK} is bound to I-P58^{IPK} as an inactive complex under conditions free of cell stress, such as infection. However, following infection with influenza virus, this inactive P58^{IPK}/I-P58^{IPK} complex becomes disrupted, and P58^{IPK} is released to interact with PKR and inhibit its kinase activity. Therefore, with decreased phosphorylation

FIGURE 6.3 Interference of influenza virus infection results in expression of IRF3/7, ATF2/cJUN, NF- κ B, as virus mobilizes a cellular protein to repress PKR activity. Full lines

of eIF2 α by PKR, the translation of viral proteins is enhanced, leading to the enhanced production of viral proteins.

6.3.3 ROSS RIVER VIRUS (RRV)

Ross River virus (RRV) is a member of the *Togaviridae* family. It is a sible for the greatest incidence of infection is not fatal, but causes arthritis, myalgia, lethargy, and fever [40]. RRV is closely related to the *Alphavirus* family, which includes the *Semliki Forest virus* and the *Borna disease virus*. RRV is found in southern India [42]. RRV infections in humans (arthritis, lethargy, and fever) are similar to those of the *Alphavirus* family. RRV is a "bone virus" [40]. Macrophage and monocyte cellular agent of severe myelitis in the mouse model have been observed in RRV-infected

FIGURE 6.3 Interference of the interferon pathway by the influenza virus. Influenza virus infection results in expression of NS1 protein. NS1-dsRNA binding has the ability to prevent IRF3/7, ATF2/cJUN, NF- κ B, and PKR activation, resulting in the inhibition of IFN α/β . The virus mobilizes a cellular protein (p58^{IPK}), which is released from its own inhibitor I-p58^{IPK} to repress PKR activity. Full lines indicate induction; dotted lines indicate inhibition.

of eIF2 α by PKR, the translation of influenza virus proteins is increased, leading to the enhanced production of progeny influenza virus by the infected cell [38].

6.3.3 ROSS RIVER VIRUS (RRV)

Ross River virus (RRV) is an endemic Australian alphavirus and the agent responsible for the greatest incidence of arboviral disease in Australia. Disease resulting from infection is not fatal, but involves a syndrome of symptoms that includes arthritis/arthralgia, myalgia, lethargy, and/or rash, which can persist for several weeks to months [40]. RRV is closely related to the reemergent Chikungunya virus (CHIK; *Alphaviridae*, Semliki Forest virus clade) [41], which has recently caused dramatic disease outbreaks in the French territories of the West Indian Ocean, as well as in southern India [42]. RRV and CHIK infections have similar disease manifestations in humans (arthritis, lethargy, etc.), and along with other relevant members of the Alphavirus family are broadly referred to as "arthrogenic/arthritisogenic alphaviruses" [40]. Macrophage and monocyte infiltrates have been associated with human disease [43], and monocyte/macrophage (F4/80⁺) cells have been identified as the cellular agent of severe muscle damage in RRV-infected mice [44]. Further studies in the mouse model have also detected inflammatory infiltrates in the joint and bone tissue of RRV-infected mice. Again, the infiltrate was predominantly F4/80⁺

cells, with significant populations of CD4⁺ T cells and NK cells detected via flow cytometric analysis [45].

RRV replicates in human and murine macrophages after cell uptake directly via a "natural" cellular receptor, or through Fc- γ receptor involving the "antibody-dependent enhancement" (ADE) mechanism of infection [46]. Since its discovery and implication as an important infection strategy, ADE has become a subject of significant scrutiny by many interested in human disease viruses. Essentially, the mechanism of ADE involves antibody-facilitated virus entry into macrophages or monocytes via Fc- γ receptors, resulting in a subsequent enhanced growth of RRV or other ADE viruses [46–48]. The first observation of RRV-ADE was reported in 1996 by Linn and colleagues [46], with the RRV-ADE and macrophage infection model leading to novel insights into the viral control of early immune and inflammatory host-gene expression postinfection.

An intriguing observation associated with RRV-ADE was the early suppression or ablation of macrophage antiviral responses, mediated by the disruption of TNF and NOS2 expression (mRNA and protein) [49]. In experiments involving pre-infection lipopolysaccharide (LPS)-stimulated antiviral activity for macrophage cultures (through enhanced pro-inflammatory cytokine and IFN expression), it was found that RRV was able to undergo unrestricted replication where infection was facilitated by polyclonal anti-RRV antibodies at a subneutralizing titer [49]. Conditions of considerable antiviral activity induced by LPS in macrophages did not allow RRV to grow when infection occurs through non-ADE (direct) infection [49]. The decrease in expression of TNF and NOS2 after RRV-ADE infection was associated with the disruption of the transcription factors interferon regulatory factor-1 (IRF-1) and nuclear factor- κ B (NF- κ B) [49, 50]. Significantly, the transcription and translation of non-antiviral *de novo* cellular proteins were unaffected by RRV-ADE [49]. Additional investigations showed that RRV-ADE infection stimulated increased interleukin-10 (IL-10) expression in macrophage cultures [50], explaining the global suppression of inflammatory/antiviral proteins after RRV-ADE infection (see Figure 6.4). This study also showed the suppression of IFN-associated STAT transcriptional protein complexes (e.g., ISGF-3), while the IL-10 associated transcription factor, Sp-1, was not affected by RRV-ADE infection [50].

The increased expression of IL-10 after RRV-ADE infection, at the same time as the suppression/ablation of TNF, NOS2, and type I IFN, suggests that viral infection mediated by ADE enables a sophisticated manipulation of host defense responses involving antiviral cytokines and immune protein expression. It is not simply the case that infection results in either a general suppression or general enhancement of host-gene expression postinfection. This ability of viruses to manipulate the cellular response through their interaction with transcriptional pathways that regulate key defense gene expression will provide important clues to the future design of antiviral and antiinflammatory genetic drugs, for example, the identification of gene candidates for targeting, either to augment or ablate downstream expression, during a given virus infection. The desired outcome for such a strategy is, on one hand, to enhance host immunity for optimum virus clearance, while also reducing host pathology and associated patient side effects. Also, with the above example of TNF/IL-10 expression for RRV-ADE, future antiviral genetic drug design will benefit

FIGURE 6.4 Suppression of replication results in the suppression of the production of IFN α/β is induced in response through IL-10 mediated regulation of SOCS3.

from transcriptional profiling of individual patients giving clues to the essential therapeutic targeting.

6.4 CONCLUDING RE

With the exciting developments and anti-sense technologies, it is essential that less questions associated with new drug development. The delivery and the development of disease protection, do not mean new gene-based drugs. The above challenges can be addressed in section 6.1, viruses, through their penetrating cells and delivery. Furthermore, if constraining the virus will effectively de

cells detected via flow

ter cell uptake directly involving the "antibody-46]. Since its discovery has become a subject of viruses. Essentially, the ry into macrophages or nced growth of RRV or DE was reported in 1996 ophage infection model nune and inflammatory

as the early suppression the disruption of TNF nts involving pre-infec- or macrophage cultures (pression), it was found re infection was facili- titer [49]. Conditions of es did not allow RRV to ion [49]. The decrease in associated with the dis- or-1 (IRF-1) and nuclear and translation of non- /-ADE [49]. Additional ncreased interleukin-10 the global suppression n (see Figure 6.4). This ' transcriptional protein iption factor, Sp-1, was

ion, at the same time as ests that viral infection host defense responses on. It is not simply the r general enhancement s to manipulate the cel- pathways that regulate to the future design of re identification of gene eam expression, during ategy is, on one hand, hile also reducing host above example of TNF/ rug design will benefit

FIGURE 6.4 Suppression of antiviral response by RRV-ADE. RRV-ADE infection and replication results in the suppression of STAT-1 and NF- κ B, but not Sp-1. As a consequence, the production of IFN α/β is inhibited while IL-10 is increased. The suppression of antiviral response through IL-10 mediated activity in monocyte cells is most likely mediated via up-regulation of SOCS3.

from transcriptional profiling of host genes during infection, allowing both the profiling of individual patients to determine disease severity and outcome, while also giving clues to the essential defense gene/protein interactions during infection for therapeutic targeting.

6.4 CONCLUDING REMARKS

With the exciting development of RNA-based gene technologies (for example, RNAi and anti-sense technologies) promising a revolution in drug and vaccine development, it is essential that leaders in such fields take pause to consider fundamental questions associated with pathogen-host interaction and disease mechanism with new drug development. The key questions involve the challenges of effective drug delivery and the development of drugs that, while curing a health disorder or providing disease protection, do not induce deleterious side effects in the patient. In this realm of new gene-based drug technologies, RNA viruses offer enormous hope that the above challenges can be met and overcome. As outlined in the introduction in section 6.1, viruses, through their fundamental nature and life cycle, have a genius in penetrating cells and delivering genetic material to the cell interior for expression. Furthermore, if constructed correctly through recombinant DNA technology, the virus will effectively deliver, replicate, and express therapeutic genes in target

cells, leading to the subsequent modification of a biological response to ameliorate disease. In addition to these properties, and in the context of broader disease processes like inflammation, RNA viruses have proven to be outstanding tools in the elucidation of key host intracellular pathways that control aberrant immune and/or inflammatory response. In this regard, we have focused on the lessons learned from HIV and influenza virus, as well as Ross River virus antibody-dependent enhancement (RRV-ADE), which has exposed how the involvement of host-immunity can assist in the viral manipulation of the early response to infection.

In terms of the first challenge suggested above, viruses as natural gene delivery vectors provide a way to specifically target cells and tissues associated with a disease process. In terms of unwanted side effects, our knowledge of virus-host interaction will assist in the identification of key molecules and intracellular pathways that contribute to nonspecific inflammation and overzealous immune activity that can result from infection or other biological stimuli, and thereafter enhance or exacerbate disease. The delivery of exogenous genes via therapeutic virus infection will be used to block the overexpression of "disease" genes while also, where necessary, reconstituting critical host-gene activation pathways after specific suppression by viral proteins or nucleic acids.

The application of knowledge from the discipline of virology, both engineering RNA virus genomes and the molecular biology of virus-host interaction, is tailor-made to accelerate developments in modern drug design using gene profiles from both the infected eukaryotic cell and the infecting virus itself.

ACKNOWLEDGMENTS

Suresh Mahalingam is an Australian NHMRC R.D. Wright Fellow.

REFERENCES

1. Mahalingam, S., Meanger, J., Foster, P. S., and Lidbury, B. A. 2002. The viral manipulation of the host cellular and immune environments to enhance propagation and survival: a focus on RNA viruses. *J. Leukocyte Biol.* 72: 429.
2. Racaniello, V. R., and Baltimore, D. 1981. Cloned poliovirus complementary DNA is infectious in mammalian cells. *Science* 214: 916.
3. Xiong, C., Levis, R., Shen, P., Schlesinger, S., Rice, C. M., and Huang, H. V. 1989. Sindbis virus: an efficient, broad host range vector for gene expression in animal cells. *Science* 243: 1188.
4. Rice, C. M., Grakoui, A., Galler, R., and Chambers, T. J. 1989. Transcription of infectious yellow fever RNA from full-length cDNA templates produced by in vitro ligation. *New Biol.* 1: 285.
5. Smerdou, C., and Liljestrom, P. 2000. Alphavirus vectors: from protein production to gene therapy. *Gene Therapy and Regulation* 1: 33.
6. Enjuanes, L., Sola, I., Alonso, S., Escors, D., and Zuniga, S. 2005. Coronavirus reverse genetics and development of vectors for gene expression. *Curr. Top. Microbiol. Immunol.* 287: 161.
7. London, S. D., Schmaljohn, A. L., Dalrymple, J. M., and Rice, C. M. 1992. Infectious enveloped RNA virus antigenic chimeras. *Proc. Natl. Acad. Sci. USA* 89: 207.
8. Sola, I., Alonso, S., Zuniga, S. 2005. Engineering the transmissibility of RNA viruses inducing lactogenic immunity. *Proc. Natl. Acad. Sci. USA* 102: 1207.
9. Fischer, F., Stegen, C. F., and Rice, C. M. 1996. A mouse hepatitis virus recombinant expressing a foreign gene. *J. Virol.* 70: 291.
10. Frolov, I., Hoffman, T. A., and Rice, C. M. 1996. Altered host range of RNA viruses. *Proc. Natl. Acad. Sci. USA* 93: 1207.
11. Caley, I. J., Betts, M. R., and Rice, C. M. 1999. Venezuelan equine encephalitis virus vector design strategies for gene delivery. *J. Virol.* 73: 1207.
12. Kaplan, G., and Racaniello, V. R. 1981. Subgenomic replicons. *J. Virol.* 38: 1207.
13. Bredenbeek, P. J., Frolov, I., and Rice, C. M. 1996. Packaging of RNA viruses. *Proc. Natl. Acad. Sci. USA* 93: 6439.
14. Khromykh, A. A., and Wertz, G. E. 1997. Kunjin: construction and use as a vector for gene delivery. *J. Virol.* 71: 1207.
15. Roberts, A., Buonocore, L., and Rice, C. M. 1996. Vesicular stomatitis virus as a vector for gene delivery. *J. Virol.* 70: 1207.
16. Pushko, P., Parker, M., and Rice, C. M. 1997. Replicon-helper system for expression of heterologous genes in vivo. *Virology* 230: 1207.
17. Polo, J. M., Belli, B. A., and Rice, C. M. 1996. Virus packaging cell lines. *Proc. Natl. Acad. Sci. USA* 93: 1207.
18. Alcamí, A., and Kozminski, A. 1997. Immunol. Today 21: 447.
19. Vilcek, J., and Sen, I. C. 1997. B. N. Fields, D. N. Knipe, B. Roizman, and S. E. Strickland. 1996. Str.
20. Samuel, C.E. 2001. Antiviral drug development. *Antiviral Res.* 50: 1207.
21. Piguet, V., Schwartz, O., and MHC-I by primate 1. *Immunol. Rev.* 170: 1207.
22. Xu, X. N., Laffert, B., and Rice, C. M. 1996. Induction of Fas ligand expression by zeta chain. *J. Exp. Med.* 183: 1207.
23. Albini, A., Ferrini, S., Bazzani, A. E., et al. 1998. *Proc. Natl. Acad. Sci. USA* 95: 13153.
24. Howcroft, T. K., Strebel, P., and Rice, C. M. 1996. class I gene promoter activity. *J. Virol.* 70: 1207.
25. Tosi, G., De Lema, B., and Rice, C. M. 2000. HLA class II expression in vivo. *J. Virol.* 74: 19.
26. Matsui, M., Warburton, J. R., and Rice, C. M. 1996. Effects of HIV-1 TAT on gene expression. *Defic. Syndr. Hum. Retrov.* 13: 1207.

response to ameliorate of broader disease pro-outstanding tools in the aberrant immune and/or the lessons learned from study-dependent enhance-ment of host-immunity can-tion.

as natural gene delivery associated with a disease of virus-host interac-racellular pathways that immune activity that can after enhance or exacer-atric virus infection will e also, where necessary, specific suppression by

ology, both engineering ost interaction, is tailor-sing gene profiles from lf.

Fellow.

v. 2002. The viral manipu-ance propagation and sur-

is complementary DNA is

., and Huang, H. V. 1989. expression in animal cells.

89. Transcription of infec-duced by in vitro ligation.

from protein production to

2005. Coronavirus reverse urre. *Top. Microbiol. Immu-*

ice, C. M. 1992. Infectious . *Sci. USA* 89: 207.

8. Sola, I., Alonso, S., Zuniga, S., Balasch, M., Plana-Duran, J., and Enjuanes, L. 2003. Engineering the transmissible gastroenteritis virus genome as an expression vector inducing lactogenic immunity. *J. Virol.* 77: 4357.
9. Fischer, F., Stegen, C. F., Koetzner, C. A., and Masters, P. S. 1998. Construction of a mouse hepatitis virus recombinant expressing a foreign gene. *Adv. Exp. Med. Biol.* 440: 291.
10. Frolov, I., Hoffman, T. A., Prágai, B. M., Dryga, S. A., Huang, H. V., Schlesinger, S., and Rice, C. M. 1996. Alphavirus-based expression vectors: strategies and applications. *Proc. Natl. Acad. Sci. USA* 93: 11371.
11. Caley, I. J., Betts, M. R., Davis, N. L., Swanstrom, R., Frelinger, J. A., and Johnston, R. E. 1999. Venezuelan equine encephalitis virus vectors expressing HIV-1 proteins: vector design strategies for improved vaccine efficacy. *Vaccine* 17: 3124.
12. Kaplan, G., and Racaniello, V. R. 1988. Construction and characterization of poliovirus subgenomic replicons. *J. Virol.* 62: 1687.
13. Bredenbeek, P. J., Frolov, I., Rice, C. M., and Schlesinger, S. 1993. Sindbis virus expression vectors: packaging of RNA replicons by using defective helper RNAs. *J. Virol.* 67: 6439.
14. Khromykh, A. A., and Westaway, E. G. 1997. Subgenomic replicons of the flavivirus Kunjin: construction and applications. *J. Virol.* 71: 1497.
15. Roberts, A., Buonocore, L., Price, R., Forman, J., and Rose, J. K. 1999. Attenuated vesicular stomatitis viruses as vaccine vectors. *J. Virol.* 73: 3723.
16. Pushko, P., Parker, M., Ludwig, G. V., Davis, N. L., Johnston, R. E., and Smith, J. F. 1997. Replicon-helper systems from attenuated Venezuelan equine encephalitis virus: expression of heterologous genes in vitro and immunization against heterologous pathogens in vivo. *Virology* 239: 389.
17. Polo, J. M., Belli, B. A., Driver, D. A., Frolov, I., Sherrill, S., et al. 1999. Stable alpha-virus packaging cell lines for Sindbis virus and Semliki Forest virus-derived vectors. *Proc. Natl. Acad. Sci. USA*, 96: 4598.
18. Alcamí, A., and Koszinowski, U. H. 2000. Viral mechanisms of immune evasion. *Immunol. Today* 21: 447.
19. Vilcek, J., and Sen, I. C. 1996. Interferons and other cytokines. In *Field's Virology*, eds. B. N. Fields, D. N. Knipe, P. M. Howley, R. M. Chanock, J. L. Melnick, T. P. Monath, B. Roizman, and S. E. Straus. Philadelphia: Lippincott-Raven, p. 375.
20. Samuel, C. E. 2001. Antiviral actions of interferons. *Clin. Microbiol. Rev.* 14: 778.
21. Piguet, V., Schwartz, O., Le Gall, S., and Trono, D. 1999. The downregulation of CD4 and MHC-I by primate lentiviruses: a paradigm for the modulation of cell surface receptors. *Immunol. Rev.* 168: 51.
22. Xu, X. N., Laffert, B., Screaton, G. R., Kraft, M., Wolf, D., Kolanus, W., et al. 1999. Induction of Fas ligand expression by HIV involves the interaction of Nef with T cell receptor zeta chain. *J. Exp. Med.* 189: 1489.
23. Albin, A., Ferrini, S., Benelli, R., Sforzini, S., Giunciuglio, D., Aluigi, M. G., Proudfoot, A. E., et al. 1998. HIV-1 Tat protein mimicry of chemokines. *Proc. Natl. Acad. Sci. USA* 95: 13153.
24. Howcroft, T. K., Strebel, K., Martin, M. A., and Singer, D. S. 1993. Repression of MHC class I gene promoter activity by two-exon Tat of HIV. *Science* 260: 1320.
25. Tosi, G., De Lerma Barbaro, A., D'Agostino, A., Valle, M. T., Megiovanni, A. M., Manca, F., et al. 2000. HIV-1 Tat mutants in the cysteine-rich region downregulate HLA class II expression in T lymphocytic and macrophage cell lines. *Eur. J. Immunol.* 30: 19.
26. Matsui, M., Warburton, R. J., Cogswell, P. C., Baldwin, A. S., Jr., and Frelinger, J. A. 1996. Effects of HIV-1 Tat on expression of HLA class I molecules. *J. Acquir. Immune Defic. Syndr. Hum. Retrovirol.* 11: 233.

27. Park, H., Davies, M. V., Langland, J. O., Chang, H. W., Nam, Y. S., Tartaglia, J., Paoletti, E., et al. 1994. TAR RNA-binding protein is an inhibitor of the interferon-induced protein kinase PKR. *Proc. Natl. Acad. Sci. USA* 91: 4713.
28. Hiscott, J., Kwon, H., and Genin, P. 2001. Hostile takeovers: viral appropriation of the NF- κ B pathway. *J. Clin. Invest.* 107: 143.
29. Demarch, F., Gutierrez, M. I., and Giacca, M. 1999. Human Immunodeficiency Virus Type 1 Tat protein activates transcription factor NF- κ B through the cellular interferon-inducible, double-stranded RNA-dependent protein kinase, PKR. *J. Virol.* 73: 7080.
30. Berger, E. A., Murphy, P. M., and Farber, J. M. 1999. Chemokine receptors as HIV-1 coreceptors: roles in viral entry, tropism, and disease. *Annu. Rev. Immunol.* 17: 657.
31. Huang, M. B., Hunter, M., and Bond, V. C. 1999. Effect of extracellular human immunodeficiency virus type 1 glycoprotein 120 on primary human vascular endothelial cell cultures. *AIDS Res. Hum. Retroviruses* 15: 1265.
32. Thompson, W. W., Comanor, L., and Shay, D. K. 2006. Epidemiology of seasonal influenza: use of surveillance data and statistical models to estimate the burden of disease. *J. Infect. Dis.* 194 (Suppl. 2): S82.
33. Garcia-Sastre, A., Egorov, A., Matassov, D., Brandt, S., Levy, D. E., Durbin, J. E., et al. 1998. Influenza A virus lacking the NS1 gene replicates in interferon-deficient systems. *Virology* 252: 324.
34. Garcia-Sastre, A. 2001. Inhibition of interferon-mediated antiviral responses by influenza A viruses and other negative-strand RNA viruses. *Virology* 279: 375.
35. Bergmann, M., Garcia-Sastre, A., Carnero, E., Pehamberger, H., Wolff, K., et al. 2000. Influenza virus NS1 protein counteracts PKR-mediated inhibition of replication. *J. Virol.* 74: 6203.
36. Talon, J., Horvath, C. M., Polley, R., Basler, C. F., Muster, T., Palese, P., and Garcia-Sastre, A. 2000. Activation of interferon regulatory factor 3 is inhibited by the influenza A virus NS1 protein. *J. Virol.* 74: 7989.
37. Wang, X., Li, M., Zheng, H., Muster, T., Palese, P., Beg, A. A., and Garcia-Sastre, A. 2000. Influenza A virus NS1 protein prevents activation of NF- κ B and induction of alpha/beta interferon. *J. Virol.* 74: 11566.
38. Goodman, A. G., Smith, J. A., Balachandran, S., Perwitasari, O., Proll, S. C., Thomas, M. J., Korth, M. J., Barber, G. N., Schiff, L. A., and Katze, M. G. 2007. The cellular protein P58^{IPK} regulates influenza virus mRNA translation and replication through a PKR-mediated mechanism. *J. Virol.* 81: 2221.
39. Yan, W., Frank, C. L., Korth, M. J., Sopher, B. L., Novoa, I., Ron, D., and Katze, M. G. 2002. Control of PERK eIF2 α kinase activity by the endoplasmic reticulum stress-induced molecular chaperone P58^{IPK}. *Proc. Natl. Acad. Sci. USA* 99: 15920.
40. Suhrbier, A., and La Linn, M. 2004. Clinical and pathologic aspects of arthritis due to Ross River virus and other alphaviruses. *Curr. Opin. Rheumatol.* 16: 374.
41. Luers, A. J., Adams, S. D., Smalley, J. V., and Campanella, J. J. 2005. A phylogenomic study of the genus Alphavirus employing whole genome comparison. *Comp. Funct. Genom.* 6: 217.
42. Brisse, S., Itean, I., and Schuffenecker, I. 2007. Chikungunya outbreaks. *N. Engl. J. Med.* 356: 2650.
43. Soden, M., Vasudevan, H., Roberts, B., Coelen, R., Hamlin, G., Vasudevan, S., and La Brooy, J. 2000. Detection of viral ribonucleic acid and histologic analysis of inflamed synovium in Ross River virus infection. *Arthritis Rheum.* 43: 365.
44. Lidbury, B. A., Simeonovic, C., Maxwell, G. E., Marshall, I. D., and Hapel, A. J. 2000. Macrophage-induced muscle pathology results in morbidity and mortality for Ross River virus-infected mice. *J. Infect. Dis.* 181: 27.
45. Morrison, T. E., Whitmon, and Heise, M. T. 2006. Chikungunya virus-induced inflammation in a mouse model. *J. Virol.* 80: 11566.
46. Linn, M. L., Aaskov, J. G., and Linn, M. 2004. Persistence of chikungunya virus in macrophages. *J. Virol.* 78: 407.
47. Porterfield, J. S. 1986. Anti-chikungunya virus. *Antiviral Res.* 31: 335.
48. Suhrbier, A., and Linn, M. 2004. Macrophage-dependent enhancement of chikungunya virus infection. *J. Virol.* 78: 11566.
49. Lidbury, B. A., and Mahalingam, S. 2000. Expression of chikungunya virus in macrophage infection. *J. Virol.* 74: 837.
50. Mahalingam, S., and Lidbury, B. A. 2000. Antiviral transcription factor enhancement of macrophage infection. *J. Virol.* 74: 13819.

Profiles in Drug Design

- Y. S., Tartaglia, J., Paoletti, he interferon-induced pro-
- viral appropriation of the
- Immunodeficiency Virus
gh the cellular interferon-
PKR. *J. Virol.* 73: 7080.
- okine receptors as HIV-1
Rev. Immunol. 17: 657.
- xtracellular human immu-
n vascular endothelial cell
- miology of seasonal influ-
ate the burden of disease.
- , D. E., Durbin, J. E., et al.
erferon-deficient systems.
- tiviral responses by influ-
logy 279: 375.
- H., Wolff, K., et al. 2000.
hibition of replication. *J.*
- I., Palese, P., and Garcia-
is inhibited by the influ-
- A., and Garcia-Sastre, A.
NF- κ B and induction of
- , O., Proll, S. C., Thomas,
M. G. 2007. The cellular
and replication through a
- Ron, D., and Katze, M. G.
oplasmic reticulum stress-
/SA 99: 15920.
- aspects of arthritis due to
itol. 16: 374.
- J. 2005. A phylogenomic
mparison. *Comp. Funct.*
- ya outbreaks. *N. Engl. J.*
- J., Vasudevan, S., and La
ogic analysis of inflamed
365.
- J., and Hapel, A. J. 2000.
and mortality for Ross
45. Morrison, T. E., Whitmore, A. C., Shabman, R. S., Lidbury, B. A., Mahalingam, S., and Heise, M. T. 2006. Characterization of Ross River virus tropism and virus-induced inflammation in a mouse model of viral arthritis and myositis. *J. Virol.* 80: 737.
46. Linn, M. L., Aaskov, J. G., and Suhrbier, A. 1996. Antibody-dependent enhancement and persistence in macrophages of an arbovirus associated with arthritis. *J. Gen. Virol.* 77: 407.
47. Porterfield, J. S. 1986. Antibody-dependent enhancement of viral infectivity. *Adv. Virus Research* 31: 335.
48. Suhrbier, A., and Linn, M. L. 2003. Suppression of antiviral responses by antibody-dependent enhancement of macrophage infection. *Trends Immunol.* 24: 165.
49. Lidbury, B. A., and Mahalingam, S. 2000. The specific ablation of antiviral gene expression in macrophages by antibody-dependent enhancement of Ross River virus infection. *J. Virol.* 74: 8376.
50. Mahalingam, S., and Lidbury, B. A. 2002. Suppression of lipopolysaccharide induced antiviral transcription factor (STAT-1 and NF- κ B) complexes by antibody-dependent enhancement of macrophage infection by Ross River virus. *Proc. Natl. Acad. Sci. USA* 99: 13819.