1 This is the accepted version of the paper after review. The formatted version is available from
2 <http://onlinelibrary.wiley.com/doi/10.1111/aec.12024/abstract>

3

4 **Detecting invertebrate responses to fire depends on sampling method and**
5 **taxonomic resolution.**

6

7 Luisa C. Teasdale¹, Annabel L. Smith^{1,2*}, Mailyn Thomas¹, Catherine A. Whitehead¹ and Don
8 A. Driscoll^{1,2}

9

10 ¹ Fenner School of Environment & Society, Frank Fenner Building 141, Australian National
11 University, Canberra ACT 0200, Australia.

12 ² ARC Centre of Excellence for Environmental Decisions and the NERP Environmental
13 Decisions Hub, Frank Fenner Building 141, Australian National University, Canberra ACT
14 0200, Australia.

15

16 * Author for correspondence: E: annabel.smith@anu.edu.au, P: +612 6125 9339, F: +612
17 6125 0757

18

19

20 **Running title:** Detecting invertebrate responses to fire

21

22

23 **Abstract**

24 New knowledge about the responses of species to fire is needed to plan for
25 biodiversity conservation in the face of changing fire regimes. However, the knowledge that
26 is acquired may be influenced by the sampling method and the taxonomic resolution of a
27 study. To investigate these potential sampling biases, we examined invertebrate responses
28 to time since fire in mallee woodlands of southern Australia. Using a large-scale replicated
29 study system we sampled over 60,000 invertebrates with large pitfall traps, wet pitfall traps,
30 and sweep nets, and undertook analyses at morphospecies and order level. Large pitfalls
31 and sweep nets detected several strong fire effects whereas wet pitfall traps detected few
32 effects. Invertebrate abundance in sweep nets was highest shortly after fire due to
33 grasshopper outbreaks. Several additional morphospecies showed strong preferences for
34 different stages in the post-fire succession. In contrast with morphospecies effects, analyses
35 at order level either failed to detect fire effects or were driven by the most abundant species.
36 For fire research to produce credible results with the potential to guide management, it must
37 use a range of sampling techniques and undertake analyses at (morpho) species level.
38 Homogeneous fire management, such as fire suppression in fragmented landscapes or
39 widespread frequent burning for asset-protection, is likely to cause declines in fire-affected
40 invertebrates.

41

42 **Key words:** fire management, time since fire, succession, trapping method, morphospecies

43

44 **Introduction**

45 Fire plays an integral role in shaping and maintaining many ecosystems around the
46 world (Bond & Keeley 2005). However, land-clearing, extreme management practices such
47 as complete fire suppression and climate change are altering the way that fire affects
48 ecosystems (Williams *et al.* 2001; Whitlock *et al.* 2003; Brennan *et al.* 2009). There is

49 currently debate about the best way to manage fire to balance built asset protection with
50 conservation of native vegetation and wildlife (Morrison *et al.* 1996; Parr & Andersen 2006;
51 Driscoll *et al.* 2010a). Fire suppression could threaten species reliant on recently burnt
52 habitat (Wikars 2002; Woinarski *et al.* 2004) but burning too frequently can have a negative
53 impact on species which rely on mature vegetation (Andersen *et al.* 2005; Moretti *et al.*
54 2006). Understanding the response of a wide range of species to fire is therefore a
55 prerequisite for planning effective fire management (Driscoll *et al.* 2010a, b).

56 As a community recovers after fire the vegetation undergoes succession, changing in
57 species composition and structure. This pattern of recovery means that the suitability of the
58 habitat for animal species may also change over time (Fox 1982; Letnic *et al.* 2004). Such
59 change is the basic premise of the habitat accommodation model of succession (Fox *et al.*
60 2003). The model predicts that a species abundance will be highest at its optimum
61 successional stage of vegetation recovery. In partial support of this model, preferences for a
62 particular stage of regeneration have been shown in a number of taxa including reptiles
63 (Singh *et al.* 2002; Fenner & Bull 2007; Driscoll & Henderson 2008), mammals (Fox 1982;
64 Letnic *et al.* 2004), birds (Smucker *et al.* 2005) and invertebrates (Moretti *et al.* 2004; Paquin
65 2008; Rodrigo *et al.* 2008; Driscoll *et al.* 2010c). These preferences mean that managing fire
66 to maintain habitat at different successional stages (i.e. a fire mosaic) could enable the
67 maintenance of biodiversity (Richards *et al.* 1999; Moretti *et al.* 2004). However, a better
68 understanding of wildlife ecology in fire prone ecosystems is needed to define acceptable
69 bounds for the spatial and temporal scales of fire mosaics (Bradstock *et al.* 2005; Parr &
70 Andersen 2006; Clarke 2008). A first step towards understanding the potential importance of
71 fire mosaics for conservation is to discover if species show a strong successional response
72 and only occur at a particular time since fire (Driscoll & Henderson 2008). This will help
73 determine the extent to which a fire mosaic may be important.

74 Invertebrates are the largest component of global biodiversity, play a major role in
75 herbivory, nutrient cycling and maintaining soil structure (Lavelle *et al.* 1997) and are an

important food source for many vertebrate species (Losey & Vaughan 2006). Incorporating information about invertebrates into fire management plans should therefore be a priority, but invertebrates are often ignored in fire ecology research (New *et al.* 2010). Most previous studies of invertebrate fire ecology have focused on coarse taxonomic groups or functional groups (Bailey & Whitham 2002; Moretti *et al.* 2006; Engle *et al.* 2008; Fattorini 2010; Radford & Andersen 2012). While functional groups provide a way to simplify responses to fire in an ecologically meaningful way (Langlands *et al.* 2011) understanding species level responses is essential in order to quantify extinction risk under changing fire regimes (Driscoll *et al.* 2010b). Many of the studies that do look at species level responses also only examine a small number of taxa (e.g. Formicidae, Andersen 1991; Coleoptera, Gandhi *et al.* 2001, Driscoll & Weir 2005; Araneae, Langlands *et al.* 2006) limiting the scope of inference. It is also typical for such studies to use only one sampling method, usually pit-fall traps of one size. Different methods sample biased subsets of the fauna, so results based on a single approach will not represent the response of invertebrates across the community (Abensperg-Traun & Steven 1995). Management recommendations should be based on knowledge of fire responses for many species within a community rather than on a narrow subset (Clarke 2008; Driscoll *et al.* 2010b; Pyrke & Samways 2012). Using a range of methods and morphospecies classifications is an efficient way to achieve this (Oliver & Beattie 1996; Derraik *et al.* 2002; Pyrke & Samways 2012).

To avoid the limitations of many previous invertebrate-fire studies we examined the fire response of a mallee woodland invertebrate community at the morphospecies level using multiple sampling methods. The aims of this study were to 1) determine which morphospecies had significant changes in abundance with time since fire, 2) determine the influence of sampling method and classification level on the ability to detect ecological effects of fire and 3) compare the time investment and outcome of different sampling methods. Addressing these issues is essential to manage fire in a way that conserves biodiversity.

103 **Methods**

104 *Study sites*

105 We sampled invertebrates at Hincks Wilderness Area (66,658 ha; 33° 46' 10" S, 136°
106 03' 24" E) and Pinkawillinie Conservation Park (130,148 ha; 32° 54' 30" S, 135° 53' 23" E)
107 on the Eyre Peninsula, South Australia (Fig. 1) during summer 2010. Both reserves contain
108 large areas of mallee vegetation (multi-stemmed *Eucalyptus* spp. up to 6m tall in our study
109 region) surrounded by land cleared for agriculture. The landscape consists of parabolic and
110 longitudinal siliceous sand dunes over solid limestone, calcrete bedrock (Twidale *et al.*
111 1985). The annual average rainfall is approximately 300 mm. Mallee typically experiences
112 fire on a decadal (10-100 yr) time scale (Bradstock & Cohn 2002) but can remain unburnt for
113 over 100 years (Clarke *et al.* 2010). Both of our study reserves have a documented history of
114 planned and unplanned fires dating back to the 1950s.

115 *Survey Design*

116 We used a chronosequence survey design to examine the effect of time since fire on
117 invertebrate abundance (Driscoll *et al.* 2010b). At each reserve, two 1 ha sites were sampled
118 in each of three fire categories: burnt 4-5, 9-11, and over 40 years ago (Fig. 1). Replicate
119 sites in each category were separated by approximately 1 km. To minimise edge effects, we
120 placed all sites at least 200 m from the fire edge and/or the park boundary (Driscoll &
121 Henderson 2008).

122 All sampling sites straddled sand dunes and incorporated dune ridges, slopes, and
123 dune bases which, at some sites descended to the hard, clayish swale. All sites were
124 selected for their sandy soil, presence of *Tridodia irritans* (an important habitat for many
125 animal species) and for their similar topography and vegetation (dominated by *Eucalyptus*
126 *spp.* and *Melaleuca uncinata*). The 4-5 yr sites were characterised by a low (< 1 m), sparse
127 canopy with very little leaf litter and large areas of bare sand. The 9-11 yr sites had a 1-2 m
128 canopy with moderate leaf litter and bare ground and the >40 yr sites had a high canopy (> 2

m), many shrubs, dense leaf litter and little bare ground (Smith *et al.* 2012). At each location we sampled two sites within each fire category meaning that conditions were similar for replicate sites within fire categories. This form of pseudoreplication is often unavoidable in fire ecology because of limited fire histories within landscapes (Whelan *et al.* 2002) but we accommodated this in our analysis using mixed-effects models (see *Analysis*). At each site we used three sampling methods to collect invertebrates: large pitfall traps, sweep netting and wet pitfall traps.

Large pitfall traps

The large pitfall traps were used concurrently for a related reptile study (Smith *et al.* 2012) and consisted of 20 L plastic buckets (28.5cm diameter) buried flush with the ground and placed midway along a 10 m plastic drift fence (black plastic, 30cm high). Traps were arranged in 5 x 5 grids with 25 m between each bucket and the direction of the fences alternating at right angles. Samples were collected over a six day period in summer, January 2010 (Hincks 7th-12th, Pinkawillinie 21st-26th). During this time daily minimum and maximum temperatures averaged 20°C and 38°C at Hincks and 17°C and 33°C at Pinkawillinie. A total of 300 large pitfall traps (100 in each treatment) were sampled during the survey. Invertebrates smaller than 3mm in length and ants were not collected from the large pitfall traps because of time constraints. This trapping method therefore had a collection filter in addition to the bucket size limiting the invertebrates sampled.

Sweep netting

At each site invertebrates were sampled along four 100 m transects (within the large pitfall trap grids) using butterfly nets approximately 40 cm in diameter and 1 m in length. To incorporate potential time of day effects associated with diel activity patterns in invertebrates we sampled each site twice, once before 10am or between 3pm and 7pm and once between 10am and 3pm. Each transect was sampled by two people walking approximately 15 m apart at a steady pace for 10 min. All vegetation including tree canopies, shrubs, understory

and bare ground was swept with the nets. All invertebrates were collected from the nets at the end of each transect and preserved.

Wet pitfall traps

Wet pitfall traps consisted of plastic jars 10 cm deep and 4 cm in diameter containing approximately 60 mL of 9 % salt water and a drop of detergent to reduce surface tension. Twenty traps were set at each of the twelve sites giving a total of 240 traps with 80 in each treatment. Wet pitfalls were placed approximately five meters away from, and perpendicular to the fence of the large traps. The traps were open for seven days during January 2010 (Hincks 6th – 13th, Pinkawillinie 20th – 27th), corresponding closely with the large pitfall trap collection dates.

After collection, invertebrates from all three survey methods were stored in 70 % ethanol. Samples were identified using the entomology collections at CSIRO, Canberra and the South Australian Museum, Adelaide. Where accurate identification was not possible even after consultation with museum staff, individuals were assigned to a morphospecies.

Data Analysis

To determine whether the mean abundance of each morphospecies varied among fire categories we used Poisson generalised linear mixed models (GLMM) with log link functions. We fitted time since fire, location and their interactions as fixed effects. To account for our pseudoreplicated design we fitted fire (a factor delimited areas burnt in separate fire events) as a random effect. The two sites in each fire category within reserves were thus treated as subsamples rather than true replicates for statistical analysis (Bolker *et al.* 2008). To account for over-dispersion in residual variation we also fitted an observation level random effect that modelled extra-Poisson variation (Maindonald & Braun 2010). GLMMs were fitted using the glmer function in the lme4 package (Bates *et al.* 2011) for R (R Development Core Team, 2009). We obtained predicted values and standard errors using the predictSE.mer function in the AICcmodavg package (Mazerolle 2011).

We obtained P -values for multi-level terms (time since fire and its interaction with location) using Wald tests (Harrell 2001). P -values for location (a two-level term) were derived from z-scores (fixed effect/SE: Crawley 2002). Given the large number of statistical tests in our analysis we calculated Q-values using the R package qvalue (Storey 2002). Q-values estimate the number of false positive results obtained, thus controlling the false-discovery rate (Storey 2002). Q-values less than 0.05 were taken as significant meaning that 5% of our significant results could be false positives. Only morphospecies found at both locations were analysed to incorporate the replication in our study design and ensure that our focus was on time since fire and not local abundance variation. We analysed data from morphospecies with an equal or greater number of captures than sample sites (12) following Didham *et al.* (1998) (large pitfall traps = 34 morphospecies out of a total of 184; sweep netting = 42/249; wet pitfall traps = 63/144). To determine how taxonomic resolution influenced our results, the analyses were repeated at the order level for each sampling method using all morphospecies within each order. Centipedes in the order Scolopendromorpha could not be assigned confidently to morphospecies as they had desiccated. These were analysed at the order level only.

A Venn diagram was constructed to compare the number of morphospecies sampled with each method and to examine the degree of overlap among methods. A time investment and outcomes table was also compiled to compare the costs and benefits of each method.

Results

A total of 61150 invertebrates was captured during this study: 3343 in the large pitfall traps, 8034 by sweep netting and 49773 in the wet pitfall traps (mostly ants). We identified 461 morphospecies. There was no significant difference in the number of morphospecies detected among the three fire categories by any sampling method. There was no significant difference in total invertebrate abundance among the fire categories using the large pitfall traps ($P = 0.398$) or the wet pitfall traps ($P = 0.079$). The total abundance of invertebrates

207 caught by sweep netting was significantly higher in the 4-5 yr sites than in the 9-11 yr or >40
 208 yr sites ($P = 0.014$).

209 *Large pitfall traps*

210 Of 34 morphospecies analysed from the large pitfall traps the abundance of five was
 211 significantly affected by time since fire (Table 1). Lycosidae sp. 1 (Araneae) was most
 212 abundant in the 4-5 yr sites while Endacusta sp. 1 (Orthoptera) had lowest abundance in 4-5
 213 yr sites (Fig. 2). Zoridae sp. 1 (Araneae) was more abundant in 9-11 yr sites and Lycosidae
 214 sp. 2 (Araneae) and *Platyzosteria* sp. 1 (Blattodea) were more abundant in >40 yr sites (Fig.
 215 2).

216 A total of 15 invertebrate orders were detected in the large pitfall trap sample.
 217 Fourteen of these had no significant response to time since fire. Blattodea was significantly
 218 more abundant in long unburnt vegetation ($P = 0.004$, $Q = 0.029$). This result was not
 219 significant when *Platyzosteria* sp. 1 was excluded from the data set ($P = 0.107$, $Q = 0.438$)
 220 indicating the result was driven by the strong response in this species.

221 *Sweep netting*

222 Of the 42 species analysed from the sweep net sample, nine showed a significant
 223 response to time since fire (Table 1). *Warramunga* sp. 1 (Orthoptera) was the most
 224 commonly captured species and was significantly more abundant in the 4-5 yr sites (Fig. 3).
 225 One morphospecies was more abundant in the 9-11 yr sites (Psyllidae sp. 1), one was more
 226 abundant in 9-11 and >40 yrs (Lepidoptera sp. 2), and three morphospecies were more
 227 abundant in the >40 yr sites (Fig. 3). Two morphospecies were more abundant in the 4-5 yr
 228 and 9-11 yr sites than in the >40 yr sites (Fig. 3). Cicadellidae sp. 1 (Hemiptera) showed
 229 different peaks in abundance at different locations (Fig 3).

230 A total of 11 invertebrate orders were detected during the sweep net survey.
 231 Orthoptera had a significant response to time since fire with higher abundance in 4-5 yr sites

($P = 0.007$, $Q = 0.040$). Diptera had a significant interaction between fire and location ($P = 0.008$, $Q = 0.040$), being more abundant in the >40 yr sites at Pinkawillinie but having no fire response Hincks. These results were not significant when the most abundant species was excluded from the data set (*Warramunga* sp. 1 (Orthoptera) and Culicidae sp. 1 (Diptera)) indicating that the order-level results were driven by the most abundant species.

Wet pitfall traps

Of the 63 morphospecies analysed from the wet pitfall traps, three significantly varied in abundance among the three fire categories (Table 1). Formicidae sp. 1 (Hymenoptera) and Zoridae sp. 1 (Araneae) were more abundant in the 9-11 yr sites and *Melophorus* sp. 1 (Hymenoptera) was more abundant in the 4-5 yr sites (Fig. 4). A total of 14 orders were detected in the wet pitfall trap survey and none varied significantly in abundance with time since fire. There were also no significant results when the most abundant species in each order was excluded from the analysis.

Comparison of sampling methods

We observed little overlap between the three survey methods in the morphospecies detected (Fig. 5). Of the three methods the wet pitfall traps required the least field effort and captured most invertebrates (Table 2). However, wet pitfall traps required the most time to sort the samples and detected the fewest fire responses (Table 2). Weighted by hours of effort, sweep netting was six times more efficient at detecting invertebrate responses to fire than wet pitfall traps (Table 2).

Discussion

Effect of fire on invertebrate abundance

In our study 23 morphospecies showed a significant response to time since fire. Of these, five species were captured most often in the 4-5 yr or 4-5 yr and 9-11yr sites, four for the 9-11 yr sites only, and seven for the >40 yr or 9-11 yr and >40 yr sites. All of these

species had very low numbers in one or two of the fire categories. Our results demonstrate that several invertebrate species specialise on a post-fire successional stage. These species may be at risk of local extinction if fire is not managed at appropriate temporal or spatial scales (Fahrig 1997; York 1999; Driscoll *et al.* 2012; Pyrke & Samways 2012). This might not have substantial consequences in very large patches of mallee woodland where fires are unlikely to affect the whole patch. In such areas, re-colonisation would be possible from adjacent areas when the optimal time since fire returned (assuming dispersal is not limiting which is currently unknown). The consequences of local extinction may be more severe in fragmented landscapes because entire fragments can be burnt by a single fire and source populations for recolonisation may not be nearby. We do not know if the fire-affected species identified in our study also occur in the surrounding matrix of cleared agricultural land. However, in a previous study, only about a quarter of mallee beetle fauna were found in the agricultural matrix (Driscoll & Weir 2005). The role of the matrix in providing alternative habitat or in limiting dispersal in this system needs to be explored further.

Our results revealed extreme opposite responses to time since fire in species pairs from the same family (Lycosidae, Fig. 2a,b). This finding could be driven by competitive exclusion or niche differentiation where closely related species specialise on different resources (Hardin 1960; Pfennig 2009), facilitating coexistence (Schluter 2000). This pattern is consistent with the habitat accommodation model where one species can competitively exclude another when their key habitat element becomes available (Fox 1982). It also means that predicting the response of animal species to fire based on simple morphological, or family-level traits may not be possible.

Many vertebrates in this ecosystem rely on invertebrates as a source of prey. Insectivorous vertebrates often show abundance differences with time since fire (Fox 1982; Letnic *et al.* 2004; Fenner & Bull 2007; Driscoll & Henderson 2008; Smith *et al.* 2012; Driscoll *et al.* 2012) and these changes may be driven by variation in the fire response of their prey. Fire regimes have the potential to affect communities at a number of trophic

levels, but interactions between fire responses of animal species from different trophic levels have rarely been investigated. Using sweep netting, we found that invertebrates were more abundant in the 4-5 yr sites primarily due to outbreaks of the grasshopper *Warramunga* sp. 1. Insectivorous vertebrates that are able to forage above the ground, such as arboreal reptiles, birds and mammals, may have increased abundance after fire in response to this increase in prey availability (Radford and Andersen 2012). Detailed information about preferred prey across a range of vertebrate species is now needed to examine this possibility.

The mean abundance of most invertebrate morphospecies did not vary significantly with time since fire in this study. There are two potential reasons for this. First, many species may not be affected by time since fire (Herrando *et al.* 2002; Driscoll & Henderson 2008). The resources required for many ground-dwelling invertebrates including spiders, scorpions, centipedes and predatory beetles may be consistently available across post-fire succession stages. If species can survive fire (e.g. by sheltering underground) then the abundance of many species could remain unchanged. Second, many species may have had responses to time since fire that we did not detect. Statistical power of our study was likely to be low, with only two sites in each fire category within each reserve. Large-scale natural experiments have proximate factors such as local rainfall that can increase variation and reduce power (Hargrove & Pickering 1992; Driscoll *et al.* 2010b). Fire severity and frequency also have the potential to affect fauna abundance (Pardon *et al.* 2003; Smucker *et al.* 2005) but were not examined in this study. The invertebrate assemblage can also be highly seasonal and this study was a snapshot of a particular time of year. Sampling throughout the year may give a broader insight into the invertebrate community's response to fire.

Effect of taxonomic level on results

We found few significant responses when we compared changes in invertebrate abundance with time since fire at the order level. Only Blattodea and Orthoptera showed

significant responses, but these results were driven by the dominant species in that group. It is not surprising that many studies investigate ecological processes at the invertebrate order level (e.g. Bailey & Whitham 2002; Moretti *et al.* 2006; Engle *et al.* 2008; Radford & Andersen 2012) given the complexity of invertebrate taxonomy. However, our results showed that important ecological responses may not be detected using coarser taxonomic groupings. Erroneous management conclusions are likely to be drawn from studies that undertake analyses using higher taxonomic levels. When used carefully, morphospecies can be a valuable tool in broad scale invertebrate studies (Oliver & Beattie 1996; Derraik *et al.* 2002) and our study has demonstrated the benefit of this approach. However, as discussed previously, statistical power was low so small effects at the order level may not have been detected.

Methods influence interpretation of time since fire impacts

We found little overlap in the morphospecies detected using the three different sampling methods. This is not surprising as the sweep netting samples were collected predominantly from above-ground vegetation whereas the other two methods were sampling mainly ground-dwelling invertebrates. It is well established that different methods will sample different components of the habitat (e.g. dry vs. moist microhabitats: Prasifka *et al.* 2007) but our study demonstrates that different methods reveal different perspectives on the influence of fire within the same vegetation type. If we had only sampled using wet pitfall traps we would not have detected any species that prefer mallee unburnt for > 40 yrs. This could lead to management recommendations that increase the amount of fire in the landscape, with negative consequences for invertebrate species that were more common in long-unburnt mallee (5 out of 17 species with significant responses to fire). Although some studies have shown fire responses in invertebrates using small wet pitfall traps (e.g. Andersen 1991), our study highlights the importance of using a range of methods to gain a broad understanding of invertebrate fire ecology.

Each of our sampling methods included a range of "filters", and these are likely to apply in other studies that use these trapping techniques. For example, our large pitfall traps were dry, so probably captured fewer flying beetles than wet pitfall traps might. There was also some risk that invertebrates in large pitfall traps were predated upon by captured vertebrates, although the risk would have applied in all fire categories. Small (<3mm) invertebrates are very difficult to detect in large pitfall traps because these traps must have some soil in them to help protect captured vertebrates. Each method had different sized filters and sampled different subsets of the invertebrate population (excluding ants meant the large pitfall traps also had a collection filter). Our comparative research approach gave us different perspectives and showed that combining multiple methods can give a more complete representation of fire responses in the invertebrate community.

The results from our study have important implications for fire ecology, field methods and fire management. The contrasting responses of individual species to fire imply that landscapes with a range of seral stages are needed to maintain biodiversity. In fragmented landscapes with small mallee remnants, such a range has not been achieved in the past and local extinctions are expected (Driscoll & Henderson 2008). To reliably determine when there is no response to fire, research must focus on analyses at the species level rather than the order level. Robust management decisions are only possible when based on results arising from a range of trapping methods with data analysed at the species level. Predators that can eat grasshoppers and forage in low shrubs may have a response to fire that is mediated by invertebrate prey. For most generalist predators of arthropods however, prey availability remains unchanged with time since fire.

Current fire management protocols are based on the requirements of a very narrow range of taxa (Clarke 2008). Typically these are based on the fire interval requirements of vascular plants (Keith *et al.* 2002; Menges 2007). Our study has demonstrated that invertebrate species can have strong and contrasting responses to fire. To manage fire in a

way that conserves invertebrate biodiversity, a range of successional stages in a landscape are needed, and this may be broader than the range implied by models of plant succession.

Acknowledgements

This research was supported by funds from the Australian Research Council, Department of Environment and Natural Resources South Australia, Native Vegetation Council South Australia, Department for Environment and Climate Change New South Wales and the Australian National University. Thanks to Beth Mantle, Cate Lemann, Steve Shattuck, Rolf Oberprieler, Tom Weir and Matthew Colloff from CSIRO Entomology and to Peter Hudson and staff at the Terrestrial Invertebrate Collection, South Australian Museum for their help with identification. Thanks also to Kevin Mayes, Sally South, Steve Morley and Handoko Wahjudi without whose help much of these data could not have been collected. This research was conducted under permits from the South Australian Department for Environment and Heritage (S25589-2) and the Flinders University Animal Welfare Committee (E256).

References

- Abensperg-Traun M. & Steven D. (1995) The effects of pitfall trap diameter on ant species richness (Hymenoptera: Formicidae) and species composition of the catch in a semi-arid eucalypt woodland. *Aust. J. Ecol.* **20**, 282-287.
- Andersen A. N. (1991) Responses of ground-foraging ant communities to 3 experimental fire regimes in a savanna forest of tropical Australia. *Biotropica*. **23**, 575-585.
- Andersen A. N., Cook G. D., Corbett L. K., Douglas M. M., Eager R. W., Russell-Smith J., Setterfield S. A., Williams R. J. & Woinarski J. C. Z. (2005) Fire frequency and biodiversity

- 386 conservation in Australian tropical savannas: implications from the Kapalga fire experiment.
387 *Austral Ecol.* **30**, 155-167.
- 388 Bailey J. K. & Whitham T. G. (2002) Interactions among fire, aspen, and elk affect insect
389 diversity: Reversal of a community response. *Ecology*. **83**, 1701-1712.
- 390 Bates D., Maechler M., Bolker B. (2011) lme4: linear mixed-effects models using S4 classes.
391 R package, version 0.999375-40. <http://lme4.r-forge.r-project.org/> (accessed September
392 2011).
- 393 Bolker B. M., Brooks M. E., Clark C. J., Geange S. W., Poulsen J. R., Stevens M. H. H. &
394 White J. S. (2008) Generalized linear mixed models: a practical guide for ecology and
395 evolution. *Trends Ecol. Evol.* **24**, 127-135.
- 396 Bond W. J. & Keeley J. E. (2005) Fire as a global 'herbivore': the ecology and evolution of
397 flammable ecosystems. *Trends Ecol. Evol.* **20**, 387-394.
- 398 Bradstock R. A. & Cohn J. S. (2002) Demographic characteristics of mallee pine (*Callitris*
399 *verrucosa*) in fire-prone mallee communities of central New South Wales. *Aust. J. Bot.* **50**,
400 653-665.
- 401 Bradstock R. A., Bedward M., Gill A. M. & Cohn J. S. (2005) Which mosaic? A landscape
402 ecological approach for evaluating interactions between fire regimes, habitat and animals.
403 *Wildlife Res.* **32**, 409-423.
- 404 Brennan K. E. C., Christie F. J. & York A. (2009) Global climate change and litter
405 decomposition: more frequent fire slows decomposition and increases the functional
406 importance of invertebrates. *Glob. Change Biol.* **15**, 2958-2971.
- 407 Clarke M. F. (2008) Catering for the needs of fauna in fire management: science or just
408 wishful thinking? *Wildlife Res.* **35**, 385-394.

- 409 Clarke M.F., Avitabile S.C., Brown L., Callister K.E., Haslem A., Holland G.J., Kelly L.T.,
 410 Kenny S.A., Nimmo D.G., Spence-Bailey L.M., Taylor R.S., Watson S.J., Bennett A.F.
 411 (2010) Ageing mallee eucalypt vegetation after fire: insights for successional trajectories in
 412 semi-arid mallee ecosystems. *Aust. J. Bot.* **58**, 363-372.
- 413 Crawley M.J. (2002) Statistical Computing. An Introduction to Data Analysis using S-Plus.
 414 Wiley, West Sussex.
- 415 Derraik J. G. B., Closs G. P., Dickinson K. J. M., Sirvid P., Barratt B. I. P. & Patrick B. H.
 416 (2002) Arthropod morphospecies versus taxonomic species: a case study with Araneae,
 417 Coleoptera, and Lepidoptera. *Conserv. Biol.* **16**, 1015-1023.
- 418 Didham R.K., Hammond P.M., Lawton J.H., Eggleton P. & Stork, N.E. (1998) Beetle species
 419 responses to tropical forest fragmentation. *Ecol. Monogr.* **68**, 295-323.
- 420 Driscoll D. A. & Weir T. (2005) Beetle responses to habitat fragmentation depend on
 421 ecological traits, habitat condition, and remnant size. *Conserv. Biol.* **19**, 182-194.
- 422 Driscoll D. A. & Henderson M. K. (2008) How many common reptile species are fire
 423 specialists? A replicated natural experiment highlights the predictive weakness of a fire
 424 succession model. *Biol. Conserv.* **141**, 460-471.
- 425 Driscoll D. A., Lindenmayer D. B., Bennett A. F., Bode M., Bradstock R. A., Cary G. J.,
 426 Clarke M. F., Dexter N., Fensham R., Friend G., Gill M., James S., Kay G., Keith D. A.,
 427 MacGregor C., Possingham H. P., Russell-Smith J., Salt D., Watson J. E. M., Williams R. J.
 428 & York A. (2010a) Resolving conflicts in fire management using decision theory: asset-
 429 protection versus biodiversity conservation. *Conserv. Lett.* **3**, 215-223.
- 430 Driscoll D. A., Lindenmayer D. B., Bennett A. F., Bode M., Bradstock R. A., Cary G. J.,
 431 Clarke M. F., Dexter N., Fensham R., Friend G., Gill M., James S., Kay G., Keith D. A.,
 432 MacGregor C., Russell-Smith J., Salt D., Watson J. E. M., Williams R. J. & York A. (2010b)

- 433 Fire management for biodiversity conservation: Key research questions and our capacity to
434 answer them. *Biol. Conserv.* **143**, 1928-1939.
- 435 Driscoll D. A., Kirkpatrick J. B., McQuillan P. B., Bonham K. (2010c) Classic metapopulations
436 are rare among beetle species from a naturally fragmented landscape. *J. Anim. Ecol.* **79**,
437 294-303.
- 438 Driscoll D. A., Smith A. L., Blight S. & Maindonald J. (2012) Reptile responses to fire and the
439 risk of post-disturbance sampling bias. *Biodivers. Conserv.* **21**, 1607-1625.
- 440 Engle D. M., Fuhlendorf S. D., Roper A., Leslie D. M. (2008) Invertebrate community
441 response to a shifting mosaic of habitat. *Rangeland Ecol. Manag.* **61**, 55-62.
- 442 Fattorini S. (2010) Effects of fire on tenebrionid communities of a *Pinus pinea* plantation: a
443 case study in a Mediterranean site. *Biodivers. Conserv.* **19**, 1237-1250.
- 444 Fahrig L. (1997) Relative effects of habitat loss and fragmentation on population extinction.
445 *J. Wildl. Manage.* **61**, 603-610.
- 446 Fenner A. L. & Bull C. M. (2007) Short-term impact of grassland fire on the endangered
447 pygmy bluetongue lizard. *J. Zool.* **272**, 444-450.
- 448 Fox B. J. (1982) Fire and mammalian secondary succession in an Australian coastal heath.
449 *Ecology.* **63**, 1332-1341.
- 450 Fox B. J., Taylor J. E. & Thompson P. T. (2003) Experimental manipulation of habitat
451 structure: a retrogression of the small mammal succession. *J. Anim. Ecol.* **72**, 927-940.
- 452 Gandhi K. J. K., Spence J. R., Langor D. W. & Morgantini L. E. (2001) Fire residuals as
453 habitat reserves for epigaeic beetles (Coleoptera : Carabidae and Staphylinidae). *Biol.*
454 *Conserv.* **102**, 131-141.
- 455 Hardin G. (1960) Competitive exclusion principle. *Science.* **131**, 1292-1297.

- 456 Hargrove W. W. & Pickering J. (1992) Pseudoreplication: a *sine qua non* for regional
457 ecology. *Landscape Ecol.* **6**, 251-258.
- 458 Harrell F.E. (2001) Regression modelling strategies. Springer, New York.
- 459 Herrando S., Brotons L., Del Amo R. & Llacuna S. (2002) Bird community succession after
460 fire in a dry Mediterranean shrubland. *Ardea*. **90**, 303-310.
- 461 Keith D. A., Williams J. E. & Woinarski J. C. W. (2002) Fire management and biodiversity
462 conservation: key approaches and principles. In: *Flammable Australia: the fire regimes and*
463 *biodiversity of a continent* (eds R. A. Bradstock, A. M. Gill & J. E. Williams) pp. 401-425.
464 Cambridge University Press, Cambridge.
- 465 Langlands P. R., Brennan K. E. C. & Pearson D. J. (2006) Spiders, spinifex, rainfall and fire:
466 Long-term changes in an arid spider assemblage. *J. Arid Environ.* **67**, 36-59.
- 467 Langlands P. R., Brennan K. E. C., Framenau, V. W. & Main B. Y. (2011) Predicting the
468 post-fire responses of animal assemblages: testing a trait-based approach using spiders. *J.*
469 *Anim. Ecol.* **80**, 558-568.
- 470 Lavelle P., Bignell D., Lepage M., Wolters V., Roger P., Ineson P., Heal O. W. & Dhillon S.
471 (1997) Soil function in a changing world: the role of invertebrate ecosystem engineers. *Eur.*
472 *J. Soil Biol.* **33**, 159-193.
- 473 Letnic M., Dickman C. R., Tischler M. K., Tamayo B. & Beh C. L. (2004) The responses of
474 small mammals and lizards to post-fire succession and rainfall in arid Australia. *J. Arid*
475 *Environ.* **59**, 85-114.
- 476 Losey J. E. & Vaughan M. (2006) The economic value of ecological services provided by
477 insects. *Bioscience*. **56**, 311-323.
- 478 Maindonald J.H., Braun W.J. (2010) Data analysis and graphics using R: an example-based
479 approach, 3rd edn, Cambridge University Press.

- 480 Mazerolle M.J. (2011). AICcmodavg. R package, version 1.2. [http://cran.r-](http://cran.r-project.org/web/packages/AICcmodavg/index.html)
481 [project.org/web/packages/AICcmodavg/index.html](http://cran.r-project.org/web/packages/AICcmodavg/index.html) (accessed September 2011).
- 482 Menges E. S. (2007) Integrating demography and fire management: an example from
483 Florida scrub. *Aust. J. Bot.* **55**, 261-272.
- 484 Moretti M., Obrist M. K. & Duelli P. (2004) Arthropod biodiversity after forest fires: winners
485 and losers in the winter fire regime of the southern Alps. *Ecography*. **27**, 173-186.
- 486 Moretti M., Duelli P. & Obrist M. K. (2006) Biodiversity and resilience of arthropod
487 communities after fire disturbance in temperate forests. *Oecologia*. **149**, 312-327.
- 488 Morrison D. A., Buckney R. T., Bewick B. J. & Cary G. J. (1996) Conservation conflicts over
489 burning bush in south-eastern Australia. *Biol. Conserv.* **76**, 167-175.
- 490 New T. R., Yen A. L., Sands D. P. A., Greenslade P., Neville P. J., York A. & Collett N. G.
491 (2010) Planned fires and invertebrate conservation in south east Australian. *J. Insect*
492 *Conserv.* **14**, 567-574.
- 493 Oliver I. & Beattie A. J. (1996) Invertebrate morphospecies as surrogates for species: A case
494 study. *Conserv. Biol.* **10**, 99-109.
- 495 Paquin P. (2008) Carabid beetle (Coleoptera: Carabidae) diversity in the black spruce
496 succession of eastern Canada. *Biol. Conserv.* **141**, 261-275.
- 497 Pardon L. G., Brook B. W., Griffiths A. D. & Braithwaite R. W. (2003) Determinants of
498 survival for the northern brown bandicoot under a landscape-scale fire experiment. *J. Anim.*
499 *Ecol.* **72**, 106-115.
- 500 Parr C. L. & Andersen A. N. (2006) Patch mosaic burning for biodiversity conservation: a
501 critique of the pyrodiversity paradigm. *Conserv. Biol.* **20**, 1610-1619.
- 502 Pfennig D. W. (2009) Character displacement: Ecological and reproductive responses to a
503 common evolutionary problem. *Q. Rev. Biol.* **84**, 253-276.

- 504 Prasifka J. R., Lopez M. D., Hellmich R. L., Lewis L. C. & Dively G. P. (2007) Comparison of
505 pitfall traps and litter bags for sampling ground-dwelling arthropods. *J. Appl. Entomol.* **131**,
506 115-120.
- 507 Pryke J. S. & Samways M. J. (2012) Differential resilience of invertebrates to fire. *Austral*
508 *Ecol.* **37**, 460-469.
- 509 R Development Core Team (2009) R: A language and environment for statistical computing.
510 R Foundation for Statistical Computing, Vienna, Austria.
- 511 Radford I.J., Andersen A.N. (2012) Effects of fire on grass-layer savanna macroinvertebrates
512 as key food resources for insectivorous vertebrates in northern Australia. *Austral Ecol.* **37**,
513 733-742.
- 514 Richards S. A., Possingham H. P., Tizard J. (1999) Optimal fire management for maintaining
515 community diversity. *Ecol. Appl.* **9**, 880-892.
- 516 Rodrigo A., Sarda-Palomera F., Bosch J. & Retana J. (2008) Changes of dominant ground
517 beetles in black pine forests with fire severity and successional age. *Ecoscience.* **15**, 442-
518 452.
- 519 Schluter D. (2000) Ecological character displacement in adaptive radiation. *Amer. Nat.* **156**,
520 S4-S16.
- 521 Singh S., Smyth A. K. & Blomberg S. P. (2002) Effect of a control burn on lizards and their
522 structural environment in a eucalypt open-forest. *Wildlife Res.* **29**, 447-454.
- 523 Smith A. L., Bull C. M. & Driscoll D. A. (2012) Post-fire succession affects abundance and
524 survival but not detectability in a knob-tailed gecko. *Biol. Conserv.* **145**, 139–147.
- 525 Smucker K. M., Hutto R. L. & Steele B. M. (2005) Changes in bird abundance after wildfire:
526 Importance of fire severity and time since fire. *Ecol. Appl.* **15**, 1535-1549.

- 527 Storey J.D. (2002) A direct approach to false discovery rates. *J. Roy. Stat. Soc. B.* **64**, 479-
528 498.
- 529 Twidale C. R., Horwitz R. C. & Campbell E. M. (1985) Hamersley landscapes of the
530 northwest of Western Australia. *Rev. Geol. Dyn. Geogr.* **26**, 173-186.
- 531 Whelan R.J., Rodgerson L., Dickman C.R., Sutherland E.F. (2002) Critical life cycles of
532 plants and animals: developing a process-based understanding of population changes in
533 fire-prone landscapes. In *Flammable Australia: the fire regimes and biodiversity of a*
534 *continent.* eds R.A. Bradstock, J.E. Williams, A.M. Gill. Cambridge University Press,
535 Cambridge.
- 536 Whitlock C., Shafer S. L. & Marlon J. (2003) The role of climate and vegetation change in
537 shaping past and future fire regimes in the northwestern US and the implications for
538 ecosystem management. *Forest Ecol. Manag.* **178**, 5-21.
- 539 Wikars L. O. (2002) Dependence on fire in wood-living insects: An experiment with burned
540 and unburned spruce and birch logs. *J. Insect Conserv.* **6**, 1-12.
- 541 Williams A. A. J., Karoly D. J. & Tapper N. (2001) The sensitivity of Australian fire danger to
542 climate change. *Climatic Change.* **49**, 171-191.
- 543 Woinarski J. C. Z., Risler J. & Kean L. (2004) Response of vegetation and vertebrate fauna
544 to 23 years of fire exclusion in a tropical Eucalyptus open forest, Northern Territory,
545 Australia. *Austral Ecol.* **29**, 156-176.
- 546 York A. (1999) Long-term effects of frequent low-intensity burning on the abundance of litter-
547 dwelling invertebrates in coastal blackbutt forests of southeastern Australia. *J. Insect*
548 *Conserv.* **3**, 191-199.
- 549
- 550

Figure legends

Fig. 1. Invertebrate study locations at A) Hincks Wilderness Area and Pinkawillinie Conservation Park on the Eyre Peninsula South Australia. B & C) at each reserve two sites were sampled in three fire treatments.

Fig. 2. The abundance of five invertebrate morphospecies from large pitfall traps had significant responses to time since fire: a) *Lycosidae sp. 1* (Araneae) b) *Lycosidae sp. 2* (Araneae) c) *Zoridae sp. 1* (Araneae) d) *Platyzosteria sp. 1* (Blattodea) and e) *Endacusta sp. 1* (Orthoptera). Error bars are 95% confidence limits.

Fig. 3. The abundance of nine species caught in sweep nets had significant responses to time since fire: a) *Warramunga sp. 1* (Orthoptera) b) *Cicadellidae sp. 1* (Hemiptera) c) *Polyphrades marmoratus* (Coleoptera) d) *Cicadellidae sp. 2* (Hemiptera) e) *Psyllidae sp. 1* (Hemiptera) f) *Lepidoptera sp. 1* (Lepidoptera) g) *Lepidoptera sp. 2* (Lepidoptera) h) *Dicranolaius sp. 1* (Coleoptera) and i) *Mantodea sp. 1* (Mantodea). Error bars are 95% confidence limits. (H = Hincks Wilderness Area, P = Pinkawillinie Conservation Park).

Fig. 4. The abundance of three species caught in wet pitfall trap samples had significant responses to time since fire: a) *Formicidae sp. 1* (Hymenoptera) b) *Melophorus sp. 1* (Hymenoptera) and c) *Zoridae sp. 1* (Araneae). Error bars are 95% confidence limits.

Fig. 5. Number of morphospecies detected using large pitfall traps, wet pitfall traps and sweep netting, and the overlap between the three methods. Brackets refer to the number of morphospecies where only one or two individuals were detected.

574 **Table 1.** The abundance of 17 invertebrate morphospecies varied significantly with time
 575 since fire in mallee vegetation of South Australia.

Sampling method	Species	Order	Fire category where most abundant	P-value	Q-value	Figure reference
Large pitfall traps	Lycosidae sp. 1	Araneae	4-5yr	<0.001	<0.001	2a
	Lycosidae sp.2	Araneae	>40yr	<0.001	<0.001	2b
	Zoridae sp. 1	Araneae	9-11yr	<0.001	<0.001	2c
	Platyzosteria sp.	Blattodea	>40yr	<0.001	<0.001	2d
	Endacusta sp.1	Orthoptera	9-11yr and >40yr	<0.001	0.012	2e
Sweep netting	Warramunga sp.	Orthoptera	4-5yr	<0.001	<0.001	3a
	Cicadellidae sp.	Hemiptera	Interaction	<0.001	<0.001	3b
	<i>Polyphrades marmoratus</i>	Coleoptera	>40yr	<0.001	<0.001	3c
	Cicadellidae sp.	Hemiptera	4-5yr and 9-11yr	<0.001	<0.001	3d
	Psyllidae sp. 1	Hemiptera	9-11yr	<0.001	<0.001	3e
	Lepidoptera sp.	Lepidoptera	>40yr	<0.001	<0.001	3f
	Lepidoptera sp.	Lepidoptera	9-11yr and >40yr	<0.001	<0.001	3g
	<i>Dicranolaius</i> sp.	Coleoptera	>40yr	<0.001	<0.001	3h
	Mantodea sp. 1	Mantodea	4-5yr and 9-11yr	0.002	0.012	3i
Wet pitfall traps	Formicidae sp. 1	Hymenoptera	9-11yr	<0.001	<0.001	4a
	<i>Melophorus</i> sp.	Hymenoptera	4-5yr	0.002	0.041	4b
	Zoridae sp. 1	Araneae	9-11yr	0.001	0.023	4c

576

577

578

579 **Table 2.** Time investment and outcomes of three sampling methods used to detect
 580 invertebrate responses to time since fire.

Method	Field effort (hrs)	Sorting effort (hrs)	No. indiv.s	No. species / No. of species analysed	No. morphospecies with a fire response	No. responses / hr of effort
Large pitfall traps	60	24	3343	177/34	5	0.059
Sweep netting	32	24	8034	240/42	9	0.161
Wet pitfall traps	28	85	49773	150/63	3	0.027

581

582

583

584

585

586

Figure 1

591 **Figure 2**

592

593

594

595

Figure 3

Figure 4

a) Formacidae sp. 1

b) *Melophorus* sp. 1

c) Zoridae sp. 1

606 **Figure 5**

607

608

609

610

611