



---

Maturing Arc Signatures Monitored by Trace Element and Hf Isotope Systematics in the Early Cretaceous Zacatecas Volcanic Field, Mexico

Author(s): Felipe de Jesús Escalona-Alcázar, Luis A. Delgado-Argote, Oliver Nebel, Fernando Velasco-Tapia, Bodo Weber, and Ernesto Patricio Núñez-Peña

Source: *The Journal of Geology*, Vol. 122, No. 5 (September 2014), pp. 549-566

Published by: [The University of Chicago Press](#)

Stable URL: <http://www.jstor.org/stable/10.1086/677045>

Accessed: 17/06/2015 23:18

---

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



*The University of Chicago Press* is collaborating with JSTOR to digitize, preserve and extend access to *The Journal of Geology*.

<http://www.jstor.org>

# Maturing Arc Signatures Monitored by Trace Element and Hf Isotope Systematics in the Early Cretaceous Zacatecas Volcanic Field, Mexico

Felipe de Jesús Escalona-Alcázar,<sup>1,2,\*</sup> Luis A. Delgado-Argote,<sup>2</sup> Oliver Nebel,<sup>3</sup>  
Fernando Velasco-Tapia,<sup>4</sup> Bodo Weber,<sup>2</sup> and Ernesto Patricio Núñez-Peña<sup>1</sup>

1. Unidad Académica de Ciencias de la Tierra, Universidad Autónoma de Zacatecas, Calzada de la Universidad #108, C. P. 98058 Zacatecas, Zacatecas, México; 2. Departamento de Geología, Centro de Investigación Científica y Educación Superior de Ensenada, Carretera Ensenada-Tijuana 3918, Zona Playitas, C. P. 22860 Ensenada, Baja California, México; 3. Research School of Earth Sciences, Australian National University, Bldg. 61, Mills Road, Acton ACT 0200, Canberra, Australia; 4. Facultad de Ciencias de la Tierra, Universidad Autónoma de Nuevo León, Carretera a Cerro Prieto km 8, Ex-Hacienda de Guadalupe, C. P. 67700 Linares, Nuevo León, México

## ABSTRACT

Mesozoic growth of continental crust along the southwestern margin of North America and its southern extension in Mexico has been partly explained by the accretion of terranes. These terranes have been considered to be fragments of exotic, intraoceanic island arcs that approached mainland Mexico after the Early Cretaceous. Trace elements and Lu-Hf isotopic systematics for primitive arc successions of the Zacatecas Volcanic Field indicate a close relationship with parts of the northern Guerrero superterrane. Major and trace element systematics of lava flows and dioritic rocks from laccoliths suggest a cogenetic origin of the Zacatecas Formation and Las Pilas Complex rocks, here combined in the Zacatecas Group. This group represents a single arc succession that evolves from a primitive to mature arc. Initial  $^{176}\text{Hf}/^{177}\text{Hf}$  (age corrected to 130 Ma) ranges from 0.28296 to 0.28307, corresponding to  $\varepsilon\text{Hf}(t) = +9.3$  to  $+13.4$ , indicating a source related to a depleted mantle wedge with a superimposed subducted sediment contribution. Based on combined field and geochemical evidence, we propose an arc model and suggest a spatial extension of paleoarc spreading north-south from Baja California beyond the present-day Trans-Mexican Volcanic Belt in the Early Cretaceous.

**Online enhancements:** supplementary tables.

## Introduction

The study of modern island arc systems can help to reconstruct paleotectonic environments and, with this, contribute to a better understanding of modes of formation, growth, and evolution of active continental margins. Studies of intraoceanic arc systems show that subduction zones evolve from infant stages shortly after subduction initiation toward mature arc systems (e.g., Hergt and Woodhead 2007). This transition is best reflected in the geochemical nature of the mantle wedge by an ultradepletion history and long-term overprint by slab-derived fluids (Tatsumi et al. 1992; Miller et al. 1994; Pearce and Peate 1995; Hergt and Wood-


head 2007; Escuder-Viruete et al. 2009). In maturing arcs, the geochemistry of both island arc basalts (IAB) and associated sediments show increasing crustal influence deviant from a mid-ocean ridge basalt (MORB) source. However, complications may arise from crustal assimilation superimposed on source characteristics so that geochemical tracing of arc sources is not straightforward. In addition, accreted arc systems have often been subject to low or intermediate metamorphic overprints, which may promote some geochemical changes and obliterate the original chemical signatures.

In this contribution, we study the major and trace element geochemistry and Hf isotope composition of a series of basaltic rocks and sediments from an accreted terrane in central Mexico that are inter-

Manuscript received January 19, 2013; accepted April 15, 2014; electronically published August 20, 2014.

\* Author for correspondence; e-mail: papiasca@yahoo.com.

[The Journal of Geology, 2014, volume 122, p. 549–566] © 2014 by The University of Chicago.  
All rights reserved. 0022-1376/2014/12205-0004\$15.00. DOI: 10.1086/677045



**Figure 1.** a, Map of Mexico showing study area and location of the Guerrero superterrane (boldface dashed line) and subterrane (names in italics; modified from Campa and Coney 1983; Dickinson and Lawton 2001; Centeno-García et al. 2008). The Sierra Madre Occidental is based on Ferrari et al. (2005). The underlined names are from massive sulfides. The  $\epsilon\text{Nd}(t)$  values were taken from Verma (1984, 1993), Schaaf (1990), Böhnelt et al. (1992), Lapierre et al. (1992), Centeno-García et al. (1993), Tardy et al. (1994), Schaaf et al. (1995), Centeno-García and Silva-Romo (1997), Freydier et al. (1997, 2000), and Valdez-Moreno et al. (2006). The  $\epsilon\text{Nd}(t)$  contour values were performed following the nearest neighbor procedure. Values of  $\epsilon\text{Nd}(t) < +2.0$  were not plotted because they merge with the zero values where there are no available data. b, Simplified geologic map of the study area (modified from Escalona-Alcázar et al. 2009) showing the location of the samples for geochemistry and Hf isotopic analysis. Cities are in boldface, subterrane names are in italics, and massive sulfides are underlined.

preted to be fragments of island arc sequences from the Late Jurassic–Early Cretaceous Guerrero superterrane. The Guerrero terrane is divided into subterrane, each of which represents accreted successions of supposed outboard island arcs associated with ancient subduction. Thus, the Guerrero terrane has the potential to allow for the study of time-integrated geochemical changes in maturing arc systems. In this study, we employ parameters that are least susceptible to metamorphic disturbance, including the Lu-Hf isotope system, to iden-

tify mantle sources, tectonic environment, and maturing arc signatures.

### Regional Geology

The Guerrero superterrane is exposed at the southeastern and southern ends of the Sierra Madre Occidental and divided into northern and southern successions by the Trans-Mexican Volcanic Belt (TMVB; fig. 1). Its location and mafic nature have been key in studying the accretionary history of

western Mexico, leading to the development of a fringing arc model for its northern (Busby et al. 1998; Busby 2004) and southern exposures (Centeno-García et al. 2011). The Guerrero superterrane is considered an association of island-arc, back-arc, and ocean-floor assemblages of Late Jurassic–Early Cretaceous age (Campa and Coney 1983; Centeno-García et al. 1993). However, its paleolocation and relationship with mainland Mexico, either as an exotic terrane or as an island arc, is still a matter of debate (Campa and Coney 1983; Lapierre et al. 1992; Sedlock et al. 1993; Dickinson and Lawton 2001; Centeno-García et al. 2008; Mortensen et al. 2008; Martini et al. 2013). Traditionally, the assemblage of arcs in various terranes in the southwest United States and in Mexico were described as exotic terranes representing outboard arcs that were accreted to the continental margin of North America (Campa and Coney 1983; Centeno-García et al. 1993; Sedlock et al. 1993; Dickinson and Lawton 2001; Umhoefer 2003; Talavera-Mendoza et al. 2007). A fringing arc evolution as an alternative explanation for the juxtaposition of terranes has been proposed for the northwestern part of the superterrane, i.e., in the Alisitos subterrane of the Baja California peninsula (fig. 1a; Busby et al. 1998; Busby 2004). In this model, a Triassic–Jurassic island arc was separated from the continent by a possibly narrow back-arc basin. During subsequent arc evolution, this basin suffered an extensional phase, followed by converging forces, prior to accretion to mainland Mexico (Busby et al. 1998; Busby 2004; Alsleben et al. 2008). Whereas this explanation seems most plausible for the Baja California succession, the evolution of its southern extension is yet to be established. If this is indeed the case, the fringing arc model envisages fundamental processes to explain the growth of the continental crust at the margin of the southwestern Cordillera.

South of the TMVB, the tectonic setting and petrogenetic models of the Guerrero superterrane have been defined mostly on the basis of field data, supported by some trace element abundances and  $^{147}\text{Sm}$ – $^{143}\text{Nd}$ ,  $^{87}\text{Rb}$ – $^{87}\text{Sr}$ , and, to a lesser extent,  $^{207}\text{Pb}$ – $^{206}\text{Pb}$  isotope systematics. Based on these data, up to three independent island arcs have been proposed: Zihuatanejo, Teloloapan, and Arcelia (fig. 1a; Lapierre et al. 1992; Centeno-García et al. 1993, 2003, 2011; Tardy et al. 1994; Freydier et al. 2000; Talavera-Mendoza et al. 2007; Martini et al. 2009, 2010; Vega-Granillo et al. 2012). In particular, MORB-like Nd isotope signatures, i.e.,  $\epsilon\text{Nd}(t) > +8$ , in lithologic units from central Mexico have been used to favor an exotic terrane assemblage, likely

representing an intraoceanic arc system (Lapierre et al. 1992). Moreover, isotope systematics (Rb–Sr and Sm–Nd) of Late Cretaceous plutons exposed along the southwest Mexican continental margin (e.g., Puerto Vallarta, Manzanillo, Jilotlán) record the southeast waning of arc magmatism, evidenced by increasing crustal assimilation (Verma and Luhr 1993; Valdez-Moreno et al. 2006; isotope data from Schaaf 1990 were published in Böhnell et al. 1992; Schaaf et al. 1995).

The eastern and northern edges of the Guerrero superterrane included ocean floor–like assemblages (Lapierre et al. 1992; Tardy et al. 1994; Freydier et al. 1997) and ultramafic associations. Ultramafic rocks at the northwestern edge of the superterrane are probably ocean-floor related (Ortega-Gutiérrez et al. 1979), whereas those in the southeast are arc related (Delgado-Argote et al. 1992). North of the TMVB, the Guerrero superterrane is widely covered by Tertiary volcanic rocks of the Sierra Madre Occidental. The documented outcrops in Guanajuato (fig. 1a) were interpreted as nascent intraoceanic arc and oceanic crust (Lapierre et al. 1992; Ortiz-Hernández et al. 1992, 2003; Tardy et al. 1992, 1994). Further to the northwest, a possible back-arc setting developed in San Nicolás (Mortensen et al. 2008) and Francisco I. Madero (Yta et al. 2003), both of which are located in the state of Zacatecas (fig. 1a).

Based on field relationships, Centeno-García et al. (2011) suggested that the southwesternmost arc, the Zihuatanejo subterrane, possibly evolved as a fringing arc, placing the Guerrero superterrane close to the continental margin. Such interpretation implies that the fringing arc model proposed for the Baja California peninsula (Busby et al. 1998; Busby 2004) could be extended southward as far as the Sierra de Zacatecas (Escalona-Alcázar et al. 2009) and Guanajuato (Lapierre et al. 1992; Tardy et al. 1992; Freydier et al. 1996).

**Geology of the Zacatecas Group.** The study area is located in the state of Zacatecas, Mexico (fig. 1). Several investigations on the Sierra de Zacatecas have suggested two main Mesozoic lithologic units in tectonic contact: the lower Zacatecas Formation and the upper Las Pilas Complex (Ponce and Clark 1988; Monod and Calvet 1992; Centeno-García et al. 1993; Centeno-García and Silva-Romo 1997; Bartolini et al. 2001). Recently, Escalona-Alcázar et al. (2009) identified these two units as one continuous sequence that formed in the Early Cretaceous. The older unit is the dominantly sedimentary Zacatecas Formation (ZF) that grades to the younger, dominantly volcanic Las Pilas Complex (LPC).

Contact relationships between the ZF and LPC units are gradual. The ZF is composed of feldspathic wacke, mudstone, chert, and discrete limestone lenses, accompanied by basaltic lava flows, rare dikes, and hydrothermal vent-like structures. The LPC is mainly composed of laccolithic intrusions and basaltic lava flows, interlayered with feldspathic and lithic wacke, mudstone, chert, and rare limestone. This assemblage is here referred to as the Zacatecas Group.

Available geochemical data for the Zacatecas Group are limited. Because these rocks underwent hydrothermal/deuteric alteration and greenschist facies metamorphism, fluid-mobile elements such as K, Na, Rb, or Sr need to be treated with caution in petrogenetic and tectonic interpretation. In the Zacatecas Formation, limited Sm-Nd isotopic studies on three samples indicate a primitive igneous suite, interpreted to be akin to ocean-floor assemblages (Centeno-García and Silva-Romo 1997). For the Las Pilas Complex, Ranson et al. (1982) presented major elements and two trace elements (Rb and Sr) for five igneous rocks. However, due to hydrothermal/deuteric alteration, these results may be compromised by secondary processes rather than reflect primary igneous signatures. Verma (1984) analyzed Rb-Sr and Sm-Nd isotopes in one basaltic sample from the LPC and inferred that oceanic crust was involved in its magma generation. Subsequently, Tardy et al. (1992, 1994) and Centeno-García and Silva-Romo (1997), based on two igneous rock samples, defined an island arc tectonic setting. It is noteworthy that only Verma (1984) mentioned the sample location; all the other samples are from unknown localities. In summary, all previous analyses suggest a primitive, mantle-derived origin for the volcanic rocks but with opposing tectonic interpretations (IAB vs. MORB) and based on rather limited data. Hence, our systematic approach of sampling volcanic, plutonic, and sedimentary rocks in conjunction with the robustness—i.e., dominantly fluid immobile nature—of the Lu-Hf isotope system during alteration offers a promising way to reevaluate the tectonic framework of this terrane.

### Analytical Procedure

Fifteen samples comprising six plutonic and nine volcanic rocks were collected from the Mesozoic sequence of the Sierra de Zacatecas for geochemical analyses (table S1; tables S1–S3 are available online). Of these, four volcanic and five plutonic rock samples, along with four sedimentary rock samples, were analyzed for Lu-Hf isotope systematics.

Coarse rock samples were processed for geochemical analyses in the facilities of the Department of Geology, Centro de Investigación Científica y Educación Superior de Ensenada. Approximately 1 kg of each sample was crushed using a jaw crusher and subsequently powdered in a tungsten carbide mill. Major and trace element analyses were performed in the ActLabs Laboratories, Canada, on fused lithium tetraborate discs (1 : 4 dilution). Rock powders were mixed with internal standards prior to the melting procedures, and fusion discs were subsequently digested in nitric acid. Major elements were measured with an ENVIRO II ICP-OES (Thermo Jarrell), and trace elements were analyzed with a SCIEX ELAN 600 ICP-MS (Perkin-Elmer). Data quality was monitored with an in-house standard (sample 34/XII/05) that was cross-calibrated against standard reference materials, and obtained results are within previous results for the same sample. The geochemical analyses are listed in table S1.

Taking into account the hydrothermal alteration, we employed immobile elements Y, Zr, Ti, and Nb for classification and petrogenetic interpretation (Winchester and Floyd 1977; MacLean and Barret 1993). The normalizing values of rare earth elements (REEs) and MORB (N-MORB) are taken from Sun and McDonough (1989).

For Lu-Hf isotopic analyses, samples were powdered in an agate mortar to avoid contamination from tungsten carbide. Approximately 100 mg of each sample was spiked with a  $^{176}\text{Lu}$ - $^{178}\text{Hf}$  isotope tracer and dissolved in 3-mL Savillex teflon vials in high-pressure autoclaves (bombs) in an acid solution of ~1 : 1 HF : HNO<sub>3</sub>. Once evaporated, the samples were dried three times with 2 mL of  $\text{HNO}_3$  and traces of HF (<0.05 M) and then equilibrated with 6 M HCl. The Lu-Hf separation from the rock matrix and details of the isotope tracer are listed in Morel et al. (2008).

Isotope measurements were performed on an MC-ICPMS Thermo Fisher Neptune at Vrije Universiteit Amsterdam, The Netherlands, following the protocol listed in Nebel et al. (2009). The JMC-475 Hf isotope standard gave a mean of  $^{176}\text{Hf}/^{177}\text{Hf} = 0.282159 \pm 8$ , and all samples were corrected relative to a value of 0.282160 for international data comparison. Procedural blanks of 10 pg and 25 pg for Lu and Hf, respectively, were in the range of the long-term laboratory blanks (Nebel et al. 2009) and negligible.

### Results

**Petrography.** The lava flows of the Zacatecas Formation have porphyritic textures with plagioclase

( $\leq 1$  mm), clinopyroxene (augite), and rare olivine occurring as phenocrysts (table S2). Chlorite and epidote are secondary minerals as a result of hydrothermal and/or deuteric alteration. In the Las Pilas Complex, lava flows are composed of plagioclase (0%–39% of  $An_{20}$ - $An_{45}$ ), clinopyroxene (augite; 0%–48%), olivine ( $\leq 8\%$ ), and rare quartz ( $\leq 2\%$ ; table S2). Plagioclase and clinopyroxene are sometimes found as glomerophytic aggregates. Mesostasis varies from micro- to cryptocrystalline. As in the ZF, alteration developed chlorite, epidote, sericite, and calcite together with quartz and calcite veinlets.


The laccolithic bodies are composed by rare quartz ( $\leq 1\%$ ), plagioclase (25%–50% of  $An_{20}$ - $An_{45}$ ), clinopyroxene (3%–25% of augite), and scarce olivine ( $\leq 3\%$ ; table S2). In accordance with their mineralogical content, intrusive bodies show increasing quartz content from diorite to quartzdiorite (Le Maitre 2002; fig. 2c).

**Geochemical Results and Sample Classification.** All the igneous rocks were analyzed for their major and trace element compositions (table S1). Samples yield loss-on-ignition values of 1.7%–11.2%, indicating a significant proportion of hydrous minerals, which are likely secondary alteration products. We thus focus on elements considered relatively im-

mobile in hydrous fluids as, e.g., high field strength elements (HFSE) and REE.

The volcanic samples of the Zacatecas Group are subalkaline basalts, most with a tholeiitic affinity (fig. 2a, 2b). The  $Zr/TiO_2$  is considered indicative of melt evolution (Winchester and Floyd 1977; Pearce 1996), and increasing values in the Zacatecas Group reflect increasing magmatic differentiation (fig. 2a). The tholeiitic basalts from the Zacatecas Formation have  $TiO_2$  concentrations of  $\sim 1.6$  wt%, while those from the Las Pilas Complex vary from 1.1 wt% to 0.7 wt%, indicating fractionation of titanomagnetite in the more evolved andesites. The lava flows from the Zacatecas Formation have Zr values of  $>100$  ppm, while Las Pilas Complex samples have  $<100$  ppm.

To circumvent employment of the classical total alkaline versus silica content diagram for volcanic rock classification (which may be compromised by alteration), an additional parameter to identify the original alkaline nature of the sample is Nb/Y (Pearce 1996). Strong alkaline characters are often associated with low degrees of melting: incompatible Nb will be elevated for a given Y concentration (Pearce 1996) so that the Nb/Y ratio is a useful discriminator between alkaline and subalkaline series. The rocks from the Zacatecas Group are all



**Figure 2.** a, Classification diagrams for altered igneous extrusive rocks (Pearce 1996). b, Diagram showing the calc-alkaline, transitional, and tholeiitic affinities (MacLean and Barret 1993). c, Modal classification of the intrusive bodies from Las Pilas Complex (after Le Maitre 2002).


subalkaline (fig. 2a). The tholeiitic nature and magmatic differentiation can also be seen in figure 2b, where rocks with higher  $Zr/TiO_2$  plot in the transitional field.

The magmatic differentiation was further evaluated by using Mg# versus  $FeO^*$  (fig. 3a) and  $Al_2O_3$  (fig. 3b). In terms of Mg#, the volcanic rocks from the Zacatecas Formation plot on the tholeiitic trend, with a slight increase in  $FeO^*$  as the Mg# decreases. Las Pilas Complex samples tend to cluster in the edge between the tholeiitic and calc-alkaline trends (fig. 3a, 3b). The plutonic rocks have a tendency toward the calc-alkaline trend (fig. 3a, 3b). The  $Al_2O_3$  increases as the Mg# decreases (fig. 3b); this is related to the delay in plagioclase fractionation, as is typical for tholeiitic series. Crystal fractionation was further evaluated by using distributions of selected trace elements indicative for key phases, i.e., Ni for olivine and Sr for plagioclase. For this purpose, samples from lava flows are placed in their stratigraphic order (fig. 3c). Figure 3c shows

that upward in the stratigraphic section, Ni has a slight negative slope, indicative for a decrease in olivine, whereas the Sr slope is positive, indicating plagioclase fractionation.

The tholeiitic character of the sequence is also observed in figure 3d in  $(Ba/La)_N$  versus  $(La/Sm)_N$  (Arculus and Powell 1986). In this diagram, the island arc affinity of the sequence is confirmed by an increase in  $(Ba/La)_N$ , reflecting alkali enrichment by slab-derived fluids (Arculus and Powell 1986; Woodhead et al. 2001).

**Rare Earth Element and Multielement Patterns.** The two lava flows of the Zacatecas Formation (fig. 4a) show almost flat REE patterns with slightly depleted light rare earth elements (LREEs;  $[La/Sm]_N = 0.8\text{--}0.94$ ) and positive  $Eu/Eu^*$  anomalies (1.15–1.24). The REE patterns with 20–30 times chondrite values are in the range of other island arc basalts (Wilson 1989), which is likely caused by crystal fractionation. Such fractionation is more evident from the patterns of the volcanic and plutonic rocks




**Figure 3.** a, In the Mg# versus  $FeO^*$  diagram, the volcanic rocks of the Zacatecas Formation are closer to the tholeiitic trend compared with the volcanic rocks of the Las Pilas Complex. In the same diagram, the plutonic rocks show a tendency toward the calc-alkaline trend. b, Most of the volcanic and plutonic rocks follow the calc-alkaline trend in the Mg# versus  $Al_2O_3$  diagram. c, Toward the top of the stratigraphic section, the Ni content decreases and the Sr increases, indicating an upward decrease in olivine and plagioclase fractionation, respectively. d, The island arc tholeiitic character is well defined in the  $(Ba/La)_N$  versus  $(La/Sm)_N$  diagram (Arculus and Powell 1986).

of the Las Pilas Complex, where total concentration increases and LREE exhibit an increment with respect to chondrite 10–40 times in volcanic rocks and 20–60 times in the laccolithic bodies (fig. 4a, 4b). The slight increase in LREE and the persistent positive Eu anomaly in the two sequences correlate with the enrichment in incompatible elements (Gill 2010).

The positive Eu/Eu\* anomaly suggests plagioclase accumulation, and the flat to fractionated trend for the middle rare earth elements and heavy rare earth element (HREEs) indicates neither residual garnet during melting nor the presence of refractory minerals such as monazite, rutile, or sphene.


In the normalized trace element diagrams of the volcanic rocks from the Zacatecas Group, the volcanic and plutonic rocks show similar patterns (fig. 5). There is a slight negative anomaly in Nb, as is typical for IAB. The Zr and Ti have small positive spikes, whereas the Sr is more enriched. These values are typical from ocean island arc tholeiitic basalts, with melts derived from a depleted mantle that is becoming enriched by adding fluids from the subducted slab (Wilson 1989). The similarity in the patterns shown in figure 5 confirms a comagmatic origin of the entire Zacatecas Group.

**Lu-Hf Isotope Systematics.** The Lu and Hf isotope results, as well as initial Hf isotope compositions corrected to an assumed age of 130 Ma and  $TDM_{(Hf)}$  model ages of the igneous and sediments suites, are listed in table S3. The  $^{176}Hf/^{177}Hf$  ratios range from 0.28302 to 0.28314 for basalts, from 0.28299 to 0.28305 for diorites, and from 0.28276 to 0.28297 for sediments. The Lu/Hf ratios of all volcanic rocks of the Zacatecas Group are in the range of




**Figure 5.** Trace element concentrations of the igneous rocks of the Zacatecas Group normalized to mid-ocean ridge basalt (MORB; Sun and McDonough 1989). *a*, Volcanic rocks. *b*, Plutonic rocks.

global island arc rocks (Vervoort and Blichet-Toft 1999). One basalt sample (70) has a  $^{176}Lu/^{177}Hf$  of 0.0538 that exceeds by far those of the average depleted mantle of  $^{176}Lu/^{177}Hf = 0.03836$  (Weber et al. 2012), indicating a cumulus effect. A similar but minor effect is also observed in basalt sample 79, which has a  $^{176}Lu/^{177}Hf$  of 0.0340, close to the depleted MORB mantle (DMM), yielding an overestimated  $TDM_{(Hf)}$  of 1091 Ma (table S3). The  $\epsilon_{Hf}(130Ma)$



**Figure 4.** Rare earth element diagrams of the igneous rocks from the Zacatecas Volcanic Field normalized to chondrite (Sun and McDonough 1989). The thick black lines are the samples from the Zacatecas Formation, whereas the thin black lines are the samples from the Las Pilas Complex. *a*, Volcanic rocks. *b*, Plutonic rocks.





**Figure 6.** Comparison of immobile elements ratios from the Zacatecas Group with present-day tectonic settings: mid-ocean ridge basalt (MORB; Jenner et al. 2012), the Aleutians, and the Trans-Mexican Volcanic Belt (TMVB; both from the georoc database, <http://georoc.mpchmainz.gwdg.de/georoc/>). Abbreviations, in alphabetic order: LPC-E = Las Pilas Complex lava flows, LPC-I = Las Pilas Complex laccolithic bodies, ZF = Zacatecas Formation. The Zacatecas Group tends from a MORB source toward the present-day TMVB, indicating increasing island arc signatures.

values range from +10.2 to +13.4 in basalts and from +9.3 to +11.4 in diorites, all of which are clearly below the DMM composition of  $\varepsilon\text{Hf}_{(130\text{Ma})} \approx +17.5$ . Sedimentary rocks yield  $\varepsilon\text{Hf}_{(130\text{Ma})}$  values between +1.5 and +8.6, suggesting crustal sources admixed to a dominantly arc-derived detritus (cf. Vervoort and Blichert-Toft 1999) that is also reflected in  $\text{TDM}_{(\text{Hf})}$  from 931 to 579 Ma.

## Discussion

**An Arc Origin of the Zacatecas Group.** To evaluate the nature of the mantle source of the primitive igneous rocks and to test if the samples are indeed arc related, we use ratios of elements with different compatibility during melting and different affinities during slab fluid transport. A comparison with global MORB (Jenner et al. 2012), the Aleutian arc (a fluid-dominated arc setting), and the present-day TMVB (which is dominated by evolved crustal compositions data taken from the georoc database, <http://georoc.mpchmainz.gwdg.de/georoc/>) is used to investigate the geochemical setting, which formed the Zacatecas Group.

Ratios of immobile elements usually reflect the contribution from the subducted slab (Pearce and Peate 1995), either in the form of sediment melts or a fluid component with elevated Th/Yb. Likewise, in fluid-dominated arcs, the Nb/Yb indicates the source fertility with exhaustion of Nb over Yb in maturing arcs. Hence, in an island arc, setting

the Yb-normalized ratios of Th-Nb is an effective way to distinguish between arc settings deviant from the MORB array and paleo-ridge segments (Pearce and Peate 1995). In figure 6, it is demonstrated that Zacatecas samples clearly plot off the MORB array, indicating an island arc setting.

The La/Nb ratio support this view, with values ranging from 1.4 to 2.8, which are elevated compared to MORB (Rollinson 1993). These variations possibly reflect a subducted slab component.

Zirconium is an HFSE and considered immobile in arc fluids (e.g., Woodhead et al. 2001). When normalized to Yb, Zr can reflect enrichment of the mantle source in trace elements by sediment melts or indicate the crustal character of arc rocks. In figure 6a, we show that all samples of the Zacatecas Group overlap with samples from the Aleutian arc, which is an intraoceanic arc dominated by slab fluids. However, some samples of the Las Pilas Complex laccoliths tend toward more evolved compositions and approach compositions of the present-day TMVB. These chemical features, together with field evidence, indicate a younging succession from the lower and older Zacatecas Formation toward the overlying, younger Las Pilas Complex and suggest that the arc suite evolves from primitive to mature and/or more crustal compositions.

Escalona-Alcázar et al. (2009) argued that the Zacatecas Group is composed of two conformable units. These two units exhibit two trends with depletion and enrichment in LREE. Based on a limited

number of samples, Centeno-García and Silva-Romo (1997) obtained depleted LREE patterns from two volcanic samples of the older Zacatecas Formation, which were interpreted in terms of an ocean-floor association. Opposing this view, in the younger Las Pilas Complex, Tardy et al. (1992, 1994) obtained enriched LREE patterns in two samples and proposed an island arc setting. The same contrasting results were also used as argument to support the existence of two distinct stratigraphic units. By using our systematic sampling approach for the Sierra de Zacatecas and considering that the stratigraphic sequence appears continuous, our sequential sampling shows an almost flat REE pattern. Abundances in the Zacatecas Formation are almost twice as enriched compared to the lower part of the Las Pilas Complex. The LREEs in the Las Pilas Complex show a gradual increase from the bottom to the top of the complex (fig. 7a). These evolving patterns obtained in both units not only support a cogenetic relationship but also likely reflect the evolution of the island arc tholeiite as the arc becomes mature.


The REE patterns obtained for other island arc subterranean of the Guerrero superterrane also support this interpretation. In particular, the intrusive rocks of the Las Pilas Complex overlap with patterns of the Tuna Mansa Diorite and the Santa Ana Dike Complex, located about 200 km southeast within the Guanajuato region, with the exception of the low LREE observed in the Guanajuato units (figs. 1, 7b; Ortiz-Hernández et al. 1992). The Early Cretaceous magmatic sequence in Guanajuato has been interpreted as having a common magmatic origin (Ortiz-Hernández et al. 1992; Tardy et al. 1992, 1994; Mendoza and Suastegui 2000).

In addition to LREE enrichment, the spider diagrams of the igneous rocks of the Zacatecas Group show negative Nb anomalies that are a typical feature of many island arcs (fig. 5; Wilson 1989; Miller et al. 1994; Clift and Lee 1998; Falloon et al. 2007; Hergt and Woodhead 2007), including those of the genetically akin Guerrero superterrane.

In summary, the geochemical arguments, together with field relationships, reinforce the hypothesis that the Zacatecas Formation and the Las Pilas Complex form a group that developed in an arc environment, notably geochemically similar to other subterranean of the Guerrero superterrane.

We thus conclude that the Zacatecas Group is the remnant of an arc sequences that was fluid dominated, possibly growing in crustal components in a maturing arc.


**Constraints from Hf Isotope Compositions.** The initial Hf isotope compositions of the igneous rocks



**Figure 7.** Rare earth element patterns normalized to chondrite (Sun and McDonough 1989) from the Zacatecas Group compared with other Early Cretaceous Mexican subterranean. The continuous lines are the results presented in this study. *a*, Data for the Zacatecas Formation (Centeno-García and Silva-Romo 1997), Las Pilas Complex (Tardy et al. 1992, 1994), Chilitos Formation (Tardy et al. 1992, 1994), La Luz Basalts (Ortiz-Hernández et al. 1992), and Arperos basin (Freydier et al. 1996, 2000). *b*, Intrusive rocks from the Las Pilas Complex compared with the Santa Ana Dyke Complex and Tuna Mansa Diorite (Lapierre et al. 1992; Ortiz-Hernández et al. 1992, 2003; Tardy et al. 1994).

are deviant from age-corrected DMM. This indicates a crustal component in the rocks, which in island arc settings can be due to contribution of sediments from the subducted slab (fig. 8a; Pearce et al. 1999; Yogodzinski 2010; Nebel et al. 2011).

The igneous rocks of the study area carry an  $\epsilon_{\text{Hf}}(t)$  signal that is intermediate between the mantle wedge source (assumed to be similar to DMM) and the sediments, with values ranging from +9.3 (diorite from the Las Pilas Complex) to +13.4 (basalt from the Zacatecas Formation). Hence, the ra-



**Figure 8.** *a*,  $\epsilon\text{Hf}(t)$  versus  $\text{Hf}/\text{Yb}$  fields taken from Pearce et al. (1999). *b*,  $\text{Hf}/\text{Hf}^*$  anomalies versus  $\epsilon\text{Hf}(t)$  mixing lines proposed by Tollstrup and Gill (2005) for the Marianas arc for depleted mantle with volcanoclastic sediments including marine clays with (1) and without (2) zircons. MORB = mid-ocean ridge basalt.


diogenic Hf isotope composition is generally consistent with a source with time-integrated superchondritic  $\text{Lu}/\text{Hf}$ , i.e., a depleted mantle origin (fig. 8; Pearce et al. 1999). Likewise, even though the magmas were presumably derived from a depleted mantle source, their model ages (listed in table S3) show a wide range that suggests the mixing of components with different residence times in different proportions. Given the above-mentioned constraints in an intraoceanic arc setting in the absence of evolved continental crust, such bias in the isotope composition is likely related to sediment contribution from the slab (Nebel et al. 2011).

In order to explore the contribution of sedimentary Hf, we compared our Hf isotope systematics with a modeled origin of positive/negative anomalies (expressed as  $\text{Hf}/\text{Hf}^*$ ) in an extended REE plot. The  $\text{Hf}/\text{Hf}^*$  represent the extrapolated space for Hf in REE compared to its actual position, and anomalies were calculated according to the Tollstrup and Gill (2005) model to define the crustal contribution

in the magma generation. The model uses a mix of (i) a depleted mantle component and (ii) clay and volcanoclastics, the latter either with the dissolution of zircon (fig. 8*b*, line 1) or without a zircon contribution (fig. 8*b*, line 2), and assuming a maximum 25% melting (batch-melting model).

Igneous rocks from the Zacatecas Group plot close to the mixing line 1, which argues for a sediment melt. Notably,  $\text{Hf}/\text{Hf}^*$  have recently been attributed to crystal fractionation, in addition to source parameters of arc rocks (Handley et al. 2011; Woodhead et al. 2011). All samples cluster around a value of  $\text{Hf}/\text{Hf}^*$  close to 1 (0.81–1.15), so that both source variability consequent to sediment melting and crystal fractionation may account for the observed effect. However, the covariations with isotope compositions argue for the former rather than the latter.

An intriguing fact is the correlation of  $^{176}\text{Hf}/^{177}\text{Hf}$  of the igneous rocks with LREE (expressed as  $\text{La}/\text{Sm}$ ; fig. 9). Sediment melts show high  $(\text{La}/\text{Sm})_N$  accompanied with low  $^{176}\text{Hf}/^{177}\text{Hf}$  ratios and represent extension of the trend defined by igneous rocks. This indicates that local sediments are indeed possible end members for a mixing relationship. Assimilation of existing arc crust during magma ascent can have large effects on isotope and elemental abundances of new arc rocks, which, however, can be similar to effects caused by subducted continental material contribution to the magma source (Arculus and Powell 1986; McCulloch and Gamble 1991; Hawkesworth et al. 1993; Thirlwall et al. 1996; Woodhead et al. 1998). Crustal assimilation



**Figure 9.**  $(\text{La}/\text{Sm})_N$  versus  $^{176}\text{Hf}/^{177}\text{Hf}$ . Chondrite normalizing values from Sun and McDonough (1989).

and partial sediment melts from the top of the slab can be expected to result in distinct contamination paths in  $(\text{La}/\text{Sm})_N$  versus Hf isotopes (where subscript N denotes the normalization to chondritic abundances). Observed variations in  $(\text{La}/\text{Sm})_N$  in fluid-dominated shallow arc settings are minor (0.3–3; e.g., Pearce et al. 1999; Woodhead et al. 2001), including amphibole cumulates and rocks that experienced amphibole fractionation (Nebel et al. 2007). In contrast, deep island arc melts that have incorporated water-bearing-sediment-derived liquids show elevated  $(\text{La}/\text{Sm})_N$  of  $\gg 3$  (White and Dupré 1986; Thirlwall et al. 1996). This is a consequence of higher liquid-solid residue partitioning LREE (e.g., La, Ce) over HREE (e.g., Yb, Lu) in hydrous melts or supercritical fluids from subducted sediment and the presence of residual HREE-retaining garnet (van Westrenen et al. 2001; Kessel et al. 2005; Rubatto and Hermann 2007; Klimm et al. 2008). The igneous rocks of the current study (fig. 9) show a trend of  $^{176}\text{Hf}/^{177}\text{Hf}$  from volcanic to plutonic rocks toward sediments, here interpreted to indicate a mixing of mafic magma with sedimentary melts.


Notably, this observation is consistent with previous suggestions of sedimentary contribution in the magma genesis of the greater Guerrero and Caribbean island arc chains (Tardy et al. 1992, 1994; Freydier et al. 1997; Kerr et al. 1999; Mendoza and Suastegui 2000; Centeno-García 2005; Marchesi et al. 2007; Centeno-García et al. 2011).

Our preferred scenario of sediment melt contribution can also explain the Nd isotope signatures in previous isotope studies: reported values for basalts from the Zacatecas Group range from  $\epsilon\text{Nd}(t) = +1.4$  to  $\epsilon\text{Nd}(t) = +7.2$  (Verma 1984; Centeno-García and Silva-Romo 1997), values that are in general agreement with those reported south of the TMVB, where  $\epsilon\text{Nd}(t)$  ranges from +1.6 to +9.1 (Freydier et al. 1997, 2000). Whereas these values generally support a mantle origin for these rocks (Verma 1984; Centeno-García and Silva-Romo 1997), it seems obvious that their deviation from a pure DMM source requires slab fluids contribution, sediment melts, or crustal contamination. In light of our Hf isotope values and in conjunction with trace element constraints, sediment melts appear a feasible scenario for the Hf-Nd isotope compositions.

**The Guerrero Superterrane in a Fringing Arc Model.** Interpretations of the origin of the Zacatecas Group have considered two or three lithologic units of different age, each deposited or emplaced in different tectonic settings (Gutiérrez-Amador 1908; Monod and Calvet 1992; Centeno-García et al. 1993; Centeno-García and Silva-Romo 1997;

Bartolini et al. 2001; Centeno-García 2005). The oldest is the Zacatecas Formation, which has been interpreted to be deposited in an oceanic basin during the Late Triassic. The Nazas Formation of the Middle-Upper Jurassic is interpreted to be emplaced in a continental arc setting, whereas the youngest, the Las Pilas Complex, is interpreted as a Lower Cretaceous island arc. The Mesozoic tectonic evolution of western Mexico is related to an active continental margin developed in northern and western Mexico from the Permian to the Middle Jurassic (DeCserna et al. 1994; Barboza-Gudiño et al. 1998, 2008, 2010; Bartolini et al. 2001; Centeno-García et al. 2008). The magmatism from the Late Jurassic to Early Cretaceous has been explained by two hypotheses: the first considers the arc magmatism to have been continuous in time but not space (Damon et al. 1981). In this scenario, during the Late Jurassic, the active arc may have migrated toward the east, perhaps close to the actual west coast of Mexico (Lapierre et al. 1992; Tardy et al. 1992; Freydier et al. 1996; Barboza-Gudiño et al. 1998; Mendoza and Suastegui 2000; Umhoefer 2003; Talavera-Mendoza et al. 2007; Martini et al. 2010; Centeno-García et al. 2011). In addition to this hypothesis, Martini et al. (2011) suggested a para-autochthonous arc model in which, initiated by slab rollback, the arc migrated to the west, where it formed an island arc that was separated from mainland Mexico by a back-arc basin (Arperos basin). During the Late Cretaceous, the island arc accreted to the continent, thereby closing the Arperos basin (Martini et al. 2013). The second hypothesis suggests that the arc developed simultaneously from the Late Jurassic to Early Cretaceous in two or three parallel island arcs. However, this model fails to explain necessary distances between the arcs and the continent (Mendoza and Suastegui, 2000; Dickinson and Lawton 2001; Talavera-Mendoza et al. 2007; Centeno-García et al. 2008; Escuder-Virueite et al. 2013). However, Centeno-García et al. (2011), in a detailed study of the stratigraphy and U-Pb ages of detrital zircons of the Zihuatanejo subterrane, concluded that this subterrane evolved as an extensional arc. In such a model, the Zihuatanejo subterrane evolved as a fringing arc close to the continent, and the Zacatecas Formation is still considered an exotic block.

Our new interpretation of the igneous suites of the Zacatecas Group supports the first hypothesis for the tectonic evolution of central Mexico during the Mesozoic. Sampling of the Mesozoic sequence of the Sierra de Zacatecas reported in the current study indicates a magmatic affinity for the igneous units, even though they were probably generated



**Figure 10.** *a*, Paleogeographic model of the tectonic setting and evolution of the Zacatecas Group during the Hauterivian. 1 = Pico de Teyra, 2 = Sierra de Catorce, 3 = Cañon de la Peregrina, 4 = Sierra de Salinas, 5 = Guanajuato, 6 = Arcelia, 7 = Huetamo, 8 = Oaxaca. The stippled pattern is oceanic crust. Modified from Pindell and Kennan (2009). *b*, A possible scenario of the development of the Zacatecas Group.

at different stages during the evolution of the island arc. Hence, based on the field evidence and our geochemical interpretation, we argue for a single arc origin for all these units that evolved from a juvenile to a more mature stage.

We thus support an evolutionary scenario of a continuous arc rather than the existence of differ-

ent arcs. In a paleotectonic context, the presence of continental crust has broader implications for the arc development. Our data support a fringing arc model similar to that described in the northern part of the Baja California peninsula (Busby et al. 1998, 2004), rather than an exotic arc model. Hafnium isotopes have been successfully applied to

exotic outboard arcs, providing a true mantle signature, even in mature arc sequences (e.g., Nebel et al. 2007).

Our preferred model for the development of the Zacatecas Group in the Guerrero superterrane is illustrated in figure 10. In this model, the arc developed parallel and close to the continental margin (Busby et al. 2006; Alsleben et al. 2008; Martini et al. 2010, 2011; Centeno-García et al. 2011), and Zacatecas is in the back side of the Alisitos-Teloloapan arc.

In our model, the Arperos basin possibly developed an oceanic-like crust in a back-arc basin behind the Guerrero arc, as earlier proposed (Tardy et al. 1994). In Zacatecas, ~10 km west of our study area, the Francisco I. Madero massive sulfide deposits possibly developed in an island arc setting as well (Yta et al. 2003). As the island arc evolves, the Las Pilas Complex was emplaced, forming a volcanic field in the W margin of the Arperos basin. The details of the distribution of the lithologic units are shown in figure 10*b*. This model is similar to that proposed by Escuder-Viruete et al. (2013) for the Caribbean arc and their extension in Mexico for the Guerrero superterrane. Our model further supports an arc development on oceanic-like crust covered by sediments of the Potosí fan along the western margin of Mexico. If there was a slab roll-back, then the trench migrated to the west and the sediments could be subducted together with the oceanic Farallon plate (Martini et al. 2011). In this scenario, the mixing of sedimentary melt contribution with depleted mantle is the origin of the Hf isotope signatures for the magma generation.

We propose a fringing arc model for the evolution of central Mexico similar to that of the Guerrero superterrane at the western margin of North America. In this model, the island-arc, back-arc, and ocean-floor assemblages of the Late Jurassic–Early Cretaceous developed close to the continent, extending from the border of the United States with Mexico to south of the TMVB.

### Conclusions

New Lu-Hf-isotope data from the Zacatecas Group coupled with trace element systematics indicate an island arc origin of the igneous rocks. On the basis

of its geochemistry and age distributions, the island arc of the Guerrero superterrane represents different stages of the same arc section in various evolutionary stages. With respect to the continental hinterland of the outboard Guerrero arc, the Zacatecas-Teloloapan subterrane was likely in close proximity to a continent.

During magma genesis, a depleted mantle wedge experienced sedimentary melt contribution, resulting in the offset of isotope and trace elements from a pure DMM source. The sediment contribution may have been related to the proximity of the paleoarc to the North American continent. Hence, a fringing arc model that was first suggested for the Alisitos subterrane in Baja California can be extended south- and eastwards to the Zacatecas Group in Central Mexico. In this model, the arc could be separated from the continent by the Arperos basin, whose extension is unknown.

We thus conclude that the island arc of the Guerrero terrane shows a continuous evolution in time but with significant spatial variations that promoted the development of transitional arc crust in a maturing arc system.

### ACKNOWLEDGMENTS

This research was mainly supported by Consejo Nacional de Ciencia y Tecnología project 45817, "Emplacement Environments of the Southern Plutons of Baja California: Petrologic, Structural, Paleomagnetic and Geophysical Indicators," granted to L. A. Delgado-Argote. Partial support was also given by the Programa de Apoyo a la Investigación Científica y Tecnológica–Universidad Autónoma de Nuevo León project "Mineralogy, Geochemistry and Geochronology of the Altered Tuffs of the San Felipe Formation (Upper Cretaceous), Northeastern Mexico" granted to F. Velasco-Tapia. F. Escalona-Alcázar thanks J. Fernández-Ávalos, G. Mercado-Sánchez, and J. P. Tavizón-García for their support. We also thank P. Avilés, P. Juárez, V. Pérez, and S. Rosas for their help with the sample preparation. Comments from M. Corver, A. Esparza-Martínez, P. García-Sandoval, J. C. García y Barragán, D. Müller, F. Piñero-Fernández, and L. Solari, as well as comments from the reviewers L. Ferrari and P. Schaaf, greatly helped to improve this article.

### REFERENCES CITED

- Alsleben, H.; Wetmore, P. H.; Schmidt, K. L.; Paterson, S. R.; and Melis, E. A. 2008. Complex deformation during arc-continent collision: quantifying finite strain in the accreted Alisitos arc, Peninsular Ranges batholith. *J. Struct. Geol.* 30:220–236.
- Arculus, R. J., and Powell, R. 1986. Source component

- mixing in the regions of arc magma generation. *J. Geophys. Res.* 91:5913–5926.
- Barboza-Gudiño, J. R.; Orozco-Esquivel, M. T.; Gómez-Anguiano, M.; and Zavala-Monsiváis, A. 2008. The Early Mesozoic volcanic arc from western North America in northeastern Mexico. *J. S. Am. Earth Sci.* 25:49–63.
- Barboza-Gudiño, J. R.; Tristán-González, M.; and Torres-Hernández, J. R. 1998. The Late Triassic–Early Jurassic active continental margin of western North America in northeastern Mexico. *Geofis. Int.* 37:283–292.
- Barboza-Gudiño, R.; Zavala-Monsiváis, A.; Venegas-Rodríguez, G.; and Barajas-Nigoche, L. D. 2010. Late Triassic stratigraphy and facies from northeastern Mexico: tectonic setting and provenance. *Geosphere* 6: 621–640.
- Bartolini, C.; Lang, H.; Cantú-Chapa, A.; and Barboza-Gudiño, J. 2001. The Triassic Zacatecas formation in central Mexico: paleotectonic, paleogeographic and paleobiogeographic implications. *In* Bartolini, C.; Buffer, R. T.; and Cantú-Chapa, A., eds. The western Gulf of Mexico basin: tectonics, sedimentary basins, and petroleum systems. *Am. Assoc. Pet. Geol. Mem.* 75:295–315.
- Böhnell, H.; Morán-Zenteno, D.; Schaaf, P.; and Urrutia-Fucugauchi, J. 1992. Paleomagnetic and isotope data from southern Mexico and the controversy over the pre-neogene position of Baja California. *Geofis. Int.* 31:253–261.
- Busby, C. 2004. Continental growth at convergent margins facing large ocean basins: a case study from Mesozoic convergent-margin basins of Baja California, Mexico. *Tectonophysics* 392:241–277.
- Busby, C.; Fackler-Adams, B.; Mattinson, J.; and Deoreo, J. 2006. View of an intact oceanic arc, from surficial to mesozonal levels: Cretaceous Alisitos arc, Baja California. *J. Volcanol. Geotherm. Res.* 149:1–46.
- Busby, C.; Smith, D.; Morris, W.; and Fackler-Adams, B. 1998. Evolutionary model for convergent margins facing large ocean basins: Mesozoic Baja California, Mexico. *Geology* 26:227–230.
- Campa, M. F., and Coney, P. 1983. Tectonostratigraphic terranes and mineral resource distributions in Mexico. *Can. J. Earth Sci.* 20:1040–1051.
- Centeno-García, E. 2005. Review of Upper Paleozoic and Lower Mesozoic stratigraphy and depositional environments of central and west Mexico: constraints on terrane analysis and paleogeography. *In* Anderson, T. H.; Nourse, J. A.; McKee, J. W.; and Steiner, M. B., eds. The Mojave-Sonora Megashear hypothesis: development, assessment, and alternatives. *Geol. Soc. Am. Spec. Pap.* 393:233–258.
- Centeno-García, E.; Busby, C.; Busby, M.; and Gehrels, G. 2011. Evolution of the Guerrero composite terrane along the Mexican margin, from extensional fringing arc to contractional continental arc. *Geol. Soc. Am. Bull.* 123:1776–1797.
- Centeno-García, E.; Corona-Chávez, P.; Talavera-Mendoza, O.; and Iriondo A. 2003. Geology and tectonic evolution of the western Guerrero terrane: a transect from Puerto Vallarta to Zihuatanejo, Mexico. *In* Alcayde, M., and Gómez-Caballero, A., eds. *Geologic transects across cordilleran Mexico. Guidebook for the field trips of the 99th Geological Society of America Annual Meeting (Puerto Vallarta, Jalisco, Mexico, April 4–7, 2003).* Univ. Nac. Auton. Mex. Inst. Geol. Spec. Pap. 1:201–228.
- Centeno-García, E.; Guerrero-Suastegui, M.; and Talavera-Mendoza, O. 2008. The Guerrero composite terrane of western Mexico: collision and subsequent rifting in a supra-subduction zone. *In* Draut, A.; Clift, P. D.; and Scholl, D. W., eds. *Formation and applications of sedimentary record in arc collision zones.* *Geol. Soc. Am. Spec. Pap.* 436:279–308.
- Centeno-García, E.; Ruiz, J.; Coney, P.; Patchett, P. J.; and Ortega-Gutiérrez, F. 1993. Guerrero terrane of Mexico: its role in the Southern Cordillera from new geochemical data. *Geology* 21:419–422.
- Centeno-García, E., and Silva-Romo, G. 1997. Petrogenesis and tectonic evolution of central Mexico during Triassic-Jurassic time. *Rev. Mex. Cienc. Geol.* 14:244–260.
- Clift, P. D., and Lee, J. 1998. Temporal evolution of the Mariana arc during rifting of the Mariana Trough traced through the volcanoclastic record. *Island Arc* 7:496–512.
- Damon, P. E.; Shafiqullah, M.; and Clark, K. F. 1981. Evolución de los arcos magmáticos en México y su relación con la metalogénesis. *Inst. Geol. Rev.* 5:223–238.
- DeCserna, Z.; Pantoja-Alor, J.; Elías-Herrera, M.; and Sánchez-Zavala, J. L. 1994. Calc-alkaline volcanic rocks in the pre-Jurassic basement of south-central Mexico: a Late Paleozoic–Early Mesozoic convergent plate margin? *Rev. Mex. Cienc. Geol.* 11:253–256.
- Delgado-Argote, L. A.; López-Martínez, M.; York, D.; and Hall, C. M. 1992. Geologic framework and geochronology of ultramafic complexes in southern Mexico. *Can. J. Earth Sci.* 29:1590–1604.
- Dickinson, W. R., and Lawton, T. F. 2001. Carboniferous to Cretaceous assembly and fragmentation of Mexico. *Geol. Soc. Am. Bull.* 113:1142–1160.
- Escalona-Alcázar, F. J.; Delgado-Argote, L. A.; Weber, B.; Núñez-Peña, E. P.; Valencia, V. A.; and Ortiz-Acevedo, O. 2009. Kinematics and U-Pb dating of detrital zircons from the Sierra de Zacatecas, Mexico. *Rev. Mex. Cienc. Geol.* 26:48–64.
- Escuder-Viruete, J.; Pérez-Estaún, A.; and Weis, D. 2009. Geochemical constraints on the origin of the late Jurassic proto-Caribbean oceanic crust in Hispaniola. *Int. J. Earth Sci.* 98:407–425.
- Escuder-Viruete, J.; Valverde-Vaquero, P.; Rojas-Agramonte, Y.; Jabites, J.; and Pérez-Estaún, A. 2013. From intra-oceanic subduction to arc accretion and arc-continent collision: insights from the structural evolution of the Río San Juan metamorphic complex, northern Hispaniola. *J. Struct. Geol.* 46:34–56.
- Falloon, J. T.; Danyushevsky, L. V.; Crawford, T. J.; Maas, R.; Woodhead, J. D.; Eggins, S. M.; Bloomer, S. H.; Wright, D. J.; Zlobin, S. K.; and Stacey, A. R. 2007.

- Multiple mantle plume components involved in the petrogenesis of subduction-related lavas from the northern termination of the Tonga Arc and northern Lau Basin: evidence from the geochemistry of arc and backarc submarine volcanics. *Geochem. Geophys. Geosyst.* doi:10.1029/2007GC001619.
- Ferrari, L.; Valencia-Moreno, M.; and Bryan, S. 2005. Magmatismo y tectónica en la Sierra Madre Occidental y su relación con la evolución de la margen occidental de Norteamérica. *Bol. Soc. Geol. Mex.* 57:343–378.
- Freydier, C.; Lapiere, H.; Briquieu, L.; Tardy, M.; Coulon, C.; and Reyes-Martínez, J. 1997. Volcaniclastic sequences with continental affinities within the Late Jurassic–Early Cretaceous Guerrero intra-oceanic Arc terrane (western Mexico). *J. Geol.* 5:483–502.
- Freydier, C.; Lapiere, H.; Ruiz, J.; Tardy, M.; Martínez, R.; and Coulon, C. 2000. The Early Cretaceous Arperos basin: an oceanic domain diving the Guerrero arc from nuclear Mexico evidenced by the geochemistry of lavas and sediments. *J. S. Am. Earth Sci.* 13: 325–336.
- Freydier, C.; Martinez, A.; Lapiere, H.; and Coulon, C. 1996. The Early Cretaceous Arperos basin (western Mexico), geochemical evidence for an aseismic ridge formed near a spreading center. *Tectonophysics* 259: 343–367.
- Gill, R. 2010. *Igneous rocks and processes: a practical guide.* Wiley-Blackwell, Chichester, 428 p.
- Gutiérrez-Amador, M. 1908. Las capas Cárnicas de Zacatecas. *Bol. Soc. Geol. Mex.* 4:28–35.
- Handley, H. K.; Turner, S.; MacPherson, C. G.; Gertisser, R.; and Davidson, J. P. 2011. Hf-Nd isotope and trace element constraints on subduction inputs at island arcs: limitations of Hf anomalies as sediment input indicators. *Earth Planet. Sci. Lett.* 304:212–223.
- Hawkesworth, C. J.; Gallagher, K.; Hergt, J. M.; and McDermott, F. 1993. Mantle and slab contribution to arc magma. *Ann. Rev. Earth Planet. Sci.* 21:175–204.
- Hergt, J. M., and Woodhead, J. D. 2007. A critical evaluation of recent models for Tau-Tonga arc-backarc basin magmatic evolution. *Chem. Geol.* 245:9–44.
- Jenner, F. E.; Hugh, S.; and O'Neill, C. 2012. Analysis of 60 elements in 616 ocean floor basaltic glasses. *Geochem. Geophys. Geosyst.* doi:10.1029/2011GC004009.
- Kerr, A. C.; Iturralde-Vinent, M. A.; Saunders, A. D.; Babbs, T. L.; and Tarney, J. 1999. A new plate tectonic model of the Caribbean: implications from a geochemical reconnaissance of Cuban Mesozoic volcanic rocks. *Geol. Soc. Am. Bull.* 111:1581–1599.
- Kessel, R.; Schmidt, M. W.; Ulmer, P.; and Pettke, T. 2005. Trace element signature of subduction-zone fluids, melts and supercritical liquids at 120–180 km depth. *Nature* 437:724–727.
- Klimm, K.; Blundy, J. D.; and Green, T. H. 2008. Trace element partitioning and accessory phase saturation during H<sub>2</sub>O-saturated melting of basalt with implications for subduction zone chemical fluxes. *J. Petrol.* 49:523–553.
- Lapiere, H.; Ortiz, L. E.; Abouchami, W.; Monod, O.; Coulon, C.; and Zimmermann, J. L. 1992. A crustal section of an intra-oceanic island arc: the Late Jurassic–Early Cretaceous Guanajuato magmatic sequence, central Mexico. *Earth Planet. Sci. Lett.* 108:61–77.
- Le Maitre, R. W., ed. 2002. *Igneous rocks: a classification and glossary of terms.* 2nd edition. Cambridge University Press, New York, 236 p.
- MacLean, W. H., and Barret, T. J. 1993. Lithochemical techniques using immobile elements. *J. Geochem. Explorat.* 48:109–133.
- Marchesi, C.; Garrido, C. J.; Bosch, D.; Proenza, J. A.; Gervilla, F.; Monié, P.; and Rodríguez-Vega, A. 2007. Geochemistry of Cretaceous magmatism in Eastern Cuba: recycling of North American continental sediments and implications for subduction polarity in the Greater Antilles paleo-arc. *J. Petrol.* 48:1813–1840.
- Martini, M.; Ferrari, L.; López-Martínez, M.; Cerca-Martínez, M.; Valencia, V. A.; and Serrano-Durán, L. 2009. Cretaceous-Eocene magmatism and Laramide deformation in southwestern Mexico: no role for terrane accretion. *In* Kay, S. M.; Ramos, V. A.; and Dickinson, W. R., eds. *Backbone of the Americas: shallow subduction, plateau uplift, and ridge and terrane collision.* *Geol. Soc. Am. Mem.* 204:151–182.
- Martini, M.; Ferrari, L.; López-Martínez, M.; and Valencia, V. 2010. Stratigraphic redefinition of the Zihuatanejo area, southwestern Mexico. *Rev. Mex. Cienc. Geol.* 27:412–430.
- Martini, M.; Mori, L.; Solari, L.; and Centeno-García, E. 2011. Sandstone provenance of the Arperos Basin (Sierra de Guanajuato, Central Mexico): Late Jurassic–Early Cretaceous back-arc spreading as the foundation of the Guerrero terrane. *J. Geol.* 119:597–617.
- Martini, M.; Solari, L.; and Camprubí, A. 2013. Kinematics of the Guerrero terrane accretion in the Sierra de Guanajuato, central Mexico: new insights for the structural evolution of arc-continent collisional zones. *Int. Geol. Rev.* 55:574–589.
- McCulloch, M. T.; and Gamble, J. A. 1991. Geochemical and geodynamical constraints on subduction zone magmatism. *Earth Planet. Sci. Lett.* 102:358–374.
- Mendoza, O., and Suastegui, M. 2000. Geochemistry and isotopic composition of the Guerrero terrane (western Mexico): implications for the tectono-magmatic evolution of southwestern North America during Late Mesozoic. *J. S. Am. Earth Sci.* 13:297–324.
- Miller, C. A.; Barton, M.; Hanson, R. E.; and Fleming, T. H. 1994. An Early Cretaceous volcanic arc/marginal basin transition zone, Peninsula Hardy, southernmost Chile. *J. Volcanol. Geotherm. Res.* 63:33–58.
- Miyashiro, A. 1974. Volcanic rock series in island arcs and active continental margins. *Am. J. Sci.* 274:321–355.
- Monod, O., and Calvet, Ph. 1992. Structural and stratigraphic re-interpretation of the Triassic units near Zacatecas (Zac.), Central Mexico: evidence of a Laramide nappe pile. *Zentralbl. Geol. Paläontol.* I 6:1533–1544.
- Morel, M. L. A.; Nebel, O.; Nebel-Jacobsen, Y. J.; Miller, J. S.; and Vroon, P. Z. 2008. Hafnium isotope characterization of the GJ-1 zircon reference material by so-



- lution and laser ablation MC-ICPMS. *Chem. Geol.* 255:231–235.
- Mortensen, J. K.; Hall, B. V.; Bissig, T.; Friedman, R. M.; Danielson, T.; Oliver, J.; Rhys, D. A.; Ross, K. V.; and Gabites, J. E. 2008. Age and paleotectonic setting of volcanogenic massive sulphide deposits in the Guerrero terrane of Central Mexico: constraints from U-Pb age and Pb isotope studies. *Econ. Geol.* 103:117–140.
- Nebel, O.; Morel, M. L. A.; and Vroon, P. Z. 2009. Isotope dilution analyses of Lu, Hf, Zr, Ta and W, and Hf-isotope compositions of NIST SRM-612 glass wafers. *Geostand. Geoanal. Res.* 33:487–499.
- Nebel, O.; Münker, C.; Nebel-Jacobsen, Y. J.; Kleine, T.; Mezger, K.; and Mortimer, N. 2007. Hf-Nd-Pb isotope evidence from Permian arc rocks for the long-term presence of the Indian-Pacific mantle boundary in the SW Pacific. *Earth Planet. Sci. Lett.* 254:377–392.
- Nebel, O.; Vroon, P. Z.; van Westrenen, W.; Iizuka, T.; and Davies, G. R. 2011. The effect of sediment recycling in subduction zones on the Hf isotope character of new arc crust, Banda arc, Indonesia. *Earth Planet. Sci. Lett.* 303:240–250.
- Ortega-Gutiérrez, F.; Prieto-Vélez, R.; Zúñiga, Y.; and Flores, S. 1979. Una secuencia volcano-plutónica-sedimentaria Cretácica en el norte de Sinaloa: ¿un complejo ofiolítico? *Univ. Nac. Auton. Mex. Inst. Geol. Rev.* 3:1–8.
- Ortiz-Hernández, L. E.; Acevedo-Sandoval, O. A.; and Flores-Castro, K. 2003. Early Cretaceous intraplate seamounts from Guanajuato, central Mexico: geochemical and mineralogical data. *Rev. Mex. Cienc. Geol.* 20:27–40.
- Ortiz-Hernández, L. E.; Chiodi, M.; Lapierre, H.; Monod, O.; and Calvet, P. 1992. El arco intraoceánico alóctono (Cretácico Inferior) de Guanajuato, características petrográficas, geoquímicas, estructurales e isotópicas del complejo filoniano y de las lavas basálticas asociadas; implicaciones geodinámicas. *Univ. Nac. Auton. Mex. Inst. Geol. Rev.* 9:126–145.
- Pearce, J. A. 1996. A user's guide to basalt discrimination diagrams. In Wyman, D. A., ed. *Trace element geochemistry of volcanic rocks: applications for massive sulphide exploration*. *Geol. Assoc. Can. Short Course Notes* 12:79–113.
- Pearce, J. A.; Kempton, P. D.; Nowell, G. M.; and Noble, S. R. 1999. Hf-Nd Element and isotope perspective on the nature and provenance of mantle and subduction components in Western Pacific Arc-Basin systems. *J. Petrol.* 40:1579–1611.
- Pearce, J. A., and Peate, D. W. 1995. Tectonic implications of the composition of volcanic arc magmas. *Ann. Rev. Earth Planet. Sci.* 23:251–283.
- Pindell, J., and Kennan, L. 2009. Tectonic evolution of the Gulf of Mexico, Caribbean and northern South America in the mantle reference frame: an update. In James, K. H.; Lorente, M. A.; and Pindell, J., eds. *The origin and evolution of the Caribbean Plate*. *Geol. Soc. Lond. Spec. Pub.* 328:1–55.
- Ponce, B. F., and Clark, K. F. 1988. The Zacatecas mining district: a tertiary caldera complex associated with precious and base metal mineralization. *Econ. Geol.* 83:1668–1682.
- Ranson, W. A.; Fernández, L. A.; Simmons Jr., W. B.; and Enciso-de la Vega, S. 1982. Petrology of the metamorphic rocks of Zacatecas, Zac., Mexico. *Bol. Soc. Geol. Mex.* 42:37–59.
- Rollinson, H. R. 1993. *Using geochemical data: evaluation, presentation, interpretation*. Longman, Essex, 352 p.
- Rubatto, D., and Hermann, J. 2007. Experimental zircon/melt and zircon/garnet trace element partitioning and implications for the geochronology of crustal rocks. *Chem. Geol.* 241:38–61.
- Schaaf, P. 1990. *Isotopengeochemische Untersuchungen an Granitoiden Gesteinen eines aktiven Kontinentalrandes. Alter und Herkunft der Tiefengesteinskomplexe an der Pazifikküste Mexikos zwischen Puerto Vallarta und Acapulco*. PhD thesis, Ludwig-Maximilians Universität, München.
- Schaaf, P.; Morán-Zenteno, D.; Hernández-Bernal, M. S.; Solís-Pichardo, G.; Tolson, G.; and Köhler, H. 1995. Paleogene continental margin truncation in southwestern Mexico: geochronological evidence. *Tectonics* 14:1339–1350.
- Sedlock, R. L.; Ortega-Gutiérrez, F.; and Speed, R. C. 1993. Tectonostratigraphic terranes and tectonic evolution of Mexico. *Geol. Soc. Am. Spec. Pap.* 278, 153 p.
- Sun, S. S., and McDonough, W. F. 1989. Chemical and isotopic systematics of oceanic basalts: implications for mantle compositions and processes. In Saunders, A. D., and Norry, N. J., eds. *Magmatism in the ocean basins*. *Geol. Soc. Lond. Spec. Pub.* 42:313–345.
- Talavera-Mendoza, O.; Ruiz, J.; Gehrels, G. E.; Valencia, V. A.; and Centeno-García, E. 2007. Detrital zircon U/PB geochronology of southern Guerrero and western Mixteca arc successions (southern México): new insights for the tectonic evolution of southwestern North America during late Mesozoic. *Geol. Soc. Am. Bull.* 119:1052–1065.
- Tardy, M.; Lapierre, H.; Bourdier, J. L.; Coulon, C.; Ortiz-Hernández, L. E.; and Yta, M. 1992. Intraoceanic setting of the western Mexico Guerrero Terrane: implications for the Pacific-Thetys geodynamic relationships during the Cretaceous. *Univ. Nac. Auton. Mex. Inst. Geol. Rev.* 10:118–128.
- Tardy, M.; Lapierre, H.; Freydier, C.; Coulon, C.; Gill, J. B.; Mercier-de Lepinay, B.; Beck, C.; Martínez, J.; Talavera, O.; Ortiz, E.; Stein, G.; Bourdier, J. L.; and Yta, M. 1994. The Guerrero suspect terrane (western Mexico) and coeval arc terranes (the Greater Antilles and the Western Cordillera of Colombia): a Late Mesozoic intra-oceanic arc accreted to cratonic America during the Cretaceous. *Tectonophysics* 230:49–73.
- Tatsumi, Y.; Murasaki, M.; and Nohda, S. 1992. Arc-arc variation of lava chemistry in the Izu-Bonin Arc: identification of subduction components. *J. Volcanol. Geotherm. Res.* 49:179–190.
- Thirlwall, M. F.; Graham, A. M.; Arculus, R. J.; Harmon,

- R. S.; and Macpherson, C. G. 1996. Resolution of the effects of crustal assimilation, sediment subduction, and fluid transport in island arc magmas: Pb-Sr-Nd-O isotope geochemistry of Grenada, Lesser Antilles. *Geochim. Cosmochim. Acta* 60:4785–4810.
- Tollstrup, D. L., and Gill, J. B. 2005. Hafnium systematics of the Mariana Arc: evidence for sediment melt and residual phases. *Geology* 33:737–740.
- Umhoefer, P. 2003. A speculative model on the North American Cordillera in the Early Cretaceous: tectonic escape related to arc collision of the Guerrero terrane and a change in North America plate motion. In Johnson, S.; Paterson, S.; Kimbrough, D.; Martin-Barajas, A.; and Girty, G., eds. *The tectonic evolution of Northwestern Mexico and Southwestern USA*. *Geol. Soc. Am. Spec. Pap.* 374:117–134.
- Valdez-Moreno, G.; Schaaf, P.; Macias, J. L.; and Kuskabe, M. 2006. New Sr-Nd-Pb-O isotope data for Colima volcano and evidence for the nature of local basement. In Siebe, C.; Macias, J. L.; and Aguirre-Díaz, G. J., eds. *Neogene-Quaternary continental margin volcanism: a perspective from Mexico*. *Geol. Soc. Am. Spec. Pap.* 402:45–63.
- van Westrenen, W.; Blundy, J. D.; and Wood, B. J. 2001. High field strength element/rare earth element fractionation during partial melting in the presence of garnet: implications for identification of mantle heterogeneities. *Geochim. Geophys. Geosyst.* 2:2000GC000133.
- Vega-Granillo, R.; Vidal-Solano, J. R.; and Herrera-Urbina, S. 2012. Island arc tholeiites of Early Silurian, Late Jurassic and Late Cretaceous ages in the El Fuerte region, northwestern Mexico. *Rev. Mex. Cienc. Geol.* 29:492–513.
- Verma, S. P. 1984. Sr and Nd isotopic evidence for petrogenesis of Mid-Tertiary felsic volcanism in the mineral district of Zacatecas, Zac. (Sierra Madre Occidental), Mexico. *Isotope Geosci.* 2:37–53.
- Verma, S. P., and Luhr, J. F. 1993. Sr-Nd-Pb isotope and trace element geochemistry of calc-alkaline andesite from Volcán Colima, Mexico. *Geof. Int.* 32:617–631.
- Vervoort, J. D., and Blichert-Toft, J. 1999. Evolution of the depleted mantle: Hf isotope evidence from juvenile rocks through time. *Geochim. Cosmochim. Acta* 63:533–556.
- Weber, B.; Scherer, E.; Martens, U.; and Mezger, K. 2012. Where did the lower Paleozoic rocks of Yucatán come from? A U-Pb, Lu-Hf, and Sm-Nd isotope study. *Chem. Geol.* 312–313:1–17.
- White, W. M., and Dupré, B. 1986. Sediment subduction and magma genesis in the Lesser Antilles: isotopic and trace element constraints. *J. Geophys. Res.* 91:5927–5941.
- Wilson, M. 1989. *Igneous petrogenesis*. Unwin Hyman, London, 466 p.
- Winchester, J. A., and Floyd, P. A. 1977. Geochemical discrimination of different magma series and their differentiation products using immobile elements. *Chem. Geol.* 20:325–343.
- Woodhead, J.; Hergt, J.; Greig, A.; and Edwards, L. 2011. Subduction zone Hf-anomalies: mantle messenger, melting artifact or crustal process?. *Earth Planet. Sci. Lett.* 304:231–239.
- Woodhead, J. D.; Eggins, S. M.; and Johnson, R. W. 1998. Magma genesis in the New Britain island arc: further insights into melting and mass transfer processes. *J. Petrol.* 39:1641–1668.
- Woodhead, J. D.; Hergt, J. M.; Davidson, J. P.; and Eggins, S. M. 2001. Hafnium isotope evidence for “conservative” element mobility during subduction zone processes. *Earth Planet. Sci. Lett.* 192:331–346.
- Yogodzinski, G. M. 2010. Subduction controls of Hf and Nd isotopes in lavas of the Aleutian island arc. *Earth Planet. Sci. Lett.* 300:226–238.
- Yta, M.; Moreno-Tovar, F.; Cardona-Patiño, F.; and Córdoba-Méndez, D. A. 2003. Contribución a la definición de superposición de eventos metalogénicos en el yacimiento de Francisco I. Madero. *Rev. Mex. Cienc. Geol.* 20:124–132.