1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

**The halogen (F, Cl, Br, I) and H₂O systematics of Samoan lavas:
assimilated-seawater, EM2 and high-³He/⁴He components**

Mark A. Kendrick^{1,2}, Matthew G. Jackson³, Erik H. Hauri⁴, David Phillips¹

- 1 – School of Earth Sciences, University of Melbourne, Victoria 3010, Australia.
- 2 – Research School of Earth Sciences, Australian National University, Canberra, ACT 0200,
Australia.
- 3 – Department of Earth Science, University of California Santa Barbara, CA 93106-9630,
USA
- 4 - Department of Terrestrial Magnetism, Carnegie Institute of Washington, DC 20015, USA

Abstract. The Samoan mantle plume samples two or more mantle components including: an extreme EM2 composition with $^{87}\text{Sr}/^{86}\text{Sr} > 0.720$, and a primitive component with high $^3\text{He}/^4\text{He}$. The high $^{87}\text{Sr}/^{86}\text{Sr}$ melts have a unique potential to constrain the composition of the EM2 mantle end-member that is commonly attributed to subduction recycling. However, a previous study of H_2O , CO_2 , S, F and Cl in Samoan glasses was hampered by the presence of unresolved assimilated sea water. The current study builds on the earlier work by extending the volatile database to include the trace halogens Br and I, and reporting new volatile data for additional glasses with $^{87}\text{Sr}/^{86}\text{Sr}$ up to 0.7125 and $^3\text{He}/^4\text{He}$ of up to 15 Ra (Ra is the atmospheric $^3\text{He}/^4\text{He}$ ratio of 1.39×10^{-6}).

The selected glasses with MgO of 4.0 to 6.5 wt. % have CO_2 concentrations of 4 to 200 ppm that reflect degassing of CO_2 on the seafloor. In contrast, the glasses contain 0.7-1.9 wt. % H_2O , 880-1870 ppm F, 470-1700 ppm Cl, 1.8-6.9 ppm Br, 18-130 ppb I and 0.6-1.5 wt. % K. Correlations between the concentrations of these elements suggest the melts retain H_2O concentrations close to pre-degassing values and demonstrate the melts have been variably affected by assimilation of seawater-derived brines. The brines are indicated to have had salinities of 55 ± 15 wt. % salt, F/Cl ratios close to seawater (e.g. < 0.0001), Br/Cl ratios ~ 40 % higher than seawater, and I/Cl ratios ten times the seawater value. It is calculated the melts assimilated brine fractions of 0 to 0.45 wt. %, which contributed up to ~ 30 % of the total H_2O and up to ~ 70 % of the total Cl in the melts.

After accounting for the effects of brine assimilation, the Samoan melts are suggested to have a fairly constant magmatic Cl/K of 0.05 ± 0.2 , which is lower than the median MORB value. Assimilation-corrected $\text{H}_2\text{O}/\text{Ce}$ values are not correlated with $^{87}\text{Sr}/^{86}\text{Sr}$, but the sample with the highest $^{87}\text{Sr}/^{86}\text{Sr}$ of 0.7125 has an assimilation-corrected $\text{H}_2\text{O}/\text{Ce}$ of 83 ± 3 , that is significantly lower than typical MORB values of 150-300. These data confirm that the Samoan EM2 source is depleted in H_2O , and perhaps Cl, relative to lithophile elements of similar compatibility.

The glasses deemed free of seawater components, including the least enriched sample with $^3\text{He}/^4\text{He}$ of 15 Ra and the most enriched sample with $^{87}\text{Sr}/^{86}\text{Sr}$ of 0.7125, have indistinguishable Br/Cl and I/Cl that are within the narrow range determined for EM1 and EM2 glasses from Pitcairn and Society and mid-ocean ridge basalts (e.g. $\text{Br}/\text{Cl} = 0.0028 \pm 0.0006$; $\text{I}/\text{Cl} = 0.00006 \pm 0.00003$; 2σ). The uniformity of mantle Br/Cl and I/Cl could be explained if halogens are not significantly subducted into the mantle. However, an alternative possibility is that the subduction zone 'filter' controls the abundance ratios of halogens recycled into the mantle, and recycled halogens with associated H_2O , have been circulated throughout the mantle.

1. Introduction

The enriched trace element and isotope signatures of Oceanic Island Basalts (OIB) that sample the EM1, EM2 and HIMU mantle reservoirs are commonly attributed to the involvement of recycled sediments and/or altered oceanic crust/lithospheric materials introduced by subduction-recycling (e.g. Eiler et al., 2000; Hofmann and White, 1982; Hofmann, 2003; Jackson et al., 2007; White and Hofmann, 1982; Weaver, 1991; Workman et al., 2008; Zindler and Hart, 1986). OIB melts commonly have high concentrations of H₂O and other volatiles compared to MORB (e.g. Moore, 1970; Schilling et al., 1980); however, it is unclear if the high volatile content of EM1 and EM2 melts can be explained by subduction recycling. EM1 and EM2 melts are typically depleted in H₂O relative to trace elements of similar compatibility (e.g. Ce, La) that are inferred to have a recycled origin (e.g. Dixon and Clague, 2001; Dixon et al., 2002; Wallace, 2002; Workman et al., 2006; Kendrick et al., 2014a). Furthermore, a high proportion of seawater-derived volatiles (e.g. H₂O, Cl, noble gases) entering subduction zones are lost as slab fluids into the sub-arc mantle, and the extent to which these volatiles can be subducted into the deeper mantle is unclear (Hilton et al., 2002; Ito et al., 1983; Parai and Mukhopadhyay, 2012; Rüpke et al., 2004). The relative abundances of magmatic volatiles and trace elements in OIB melts that sample deeply recycled components can therefore provide a better understanding of global volatile recycling processes (Dixon et al., 2002; Wallace, 2002; Workman et al., 2006), with implications for the relative importance of subducted versus primordial volatile components in the Earth's mantle (cf. Holland and Ballentine, 2006; Mukhopadhyay, 2012; Staudacher and Allègre, 1988).

Samoan OIB are of particular interest for constraining the composition of the EM2 mantle end-member because they record the strongest ⁸⁷Sr/⁸⁶Sr enrichment of any known mantle component (>0.720; Jackson et al., 2007; 2009). Furthermore, submarine glasses

86 which are required for measuring the magmas volatile contents have been recovered with
87 $^{87}\text{Sr}/^{86}\text{Sr}$ up to 0.7125, compared to a maxima of 0.706 in EM2 glasses from the Society
88 seamounts (Fig 1; Devey et al., 1990). The Samoan glasses can be used to place potentially
89 unique constraints on the volatile characteristics of the EM2 end-member, provided the
90 geochemical complexity in melts is sufficiently well understood. Current data indicate that
91 isotopic variation in Samoan melts result from mixing an EM2 component with a primitive
92 component characterised by lower $^{87}\text{Sr}/^{86}\text{Sr}$ and high $^3\text{He}/^4\text{He}$ (e.g. Fig 1; Farley et al., 1992;
93 Jackson et al., 2007). However, additional mantle components could also be present, and a
94 previous study of volatiles (CO_2 , H_2O , S, F and Cl) in Samoan lavas was hampered by the
95 unresolved presence of assimilated seawater (Workman et al., 2006).

96 The aim of the current study is to further elucidate the origins of halogens and water
97 in Samoan melts. This is achieved by analysing the scarcest halogens Br and I, in addition to
98 Cl and F, and extending the volatile database to include the most enriched glasses now
99 available (Jackson et al., 2007). The new data are of particular interest because iodine is an
100 essential element for life that has a high abundance in organic-rich marine sediments and has
101 previously been suggested as a possible marker for subducted sedimentary components in the
102 mantle (Deruelle et al. 1992). We have already shown that the enriched mantle reservoirs
103 sampled by the Society and Pitcairn seamounts have MORB-like I/Cl (Kendrick et al., 2012a;
104 2014a); however, we deemed the current study necessary to test if the more strongly enriched
105 Samoan melts with exceptionally high $^{87}\text{Sr}/^{86}\text{Sr}$ (Fig 1) might be characterised by more
106 elevated I/Cl ratios. In addition, multi-element correlations between F, Cl, Br, I, H_2O and
107 incompatible trace elements such as K enable us to rigorously evaluate possible seawater
108 assimilation processes (Kendrick et al., 2013a). We demonstrate that Samoan melts
109 assimilated variable amounts of high salinity brine(s), and we use a quantitative mixing

110 model to correct the measured H₂O and Cl abundances for assimilation, thus providing
111 improved constraints on mantle-derived H₂O and Cl in all of the samples investigated.

112

113 2. Samples and methodology

114 The samples selected for the current study include: i) nineteen pristine glasses from
115 Samoan volcanoes in the SW Pacific, including fourteen samples previously characterised by
116 Workman et al. (2006); ii) palagonite separated from a Samoan glass, and iii) two pristine
117 glasses from Baffin Bay, Canada.

118 The Samoan glasses were all dredged from water depths of between 780 and 4400m.
119 The majority of samples were recovered during the 1999 AVON2/3 cruise of *RV Melville*
120 from the Vailulu'u, Malumalu and Ta'u volcanoes at the eastern end of the Samoan chain
121 (Fig 2; Workman et al., 2004; 2006). The Vailulu'u volcano represents the current
122 expression of the Samoan hotspot and U-series and ⁴⁰Ar-³⁹Ar dating indicate the samples
123 from Vailulu'u, Malumalu and Ta'u have ages of between ~100 yrs and ~0.3 Ma
124 (McDougall, 2010; Hart et al., 2000; Sims et al., 2008; Workman et al., 2006). Additional
125 samples recovered during the AVON2/3 cruise and the 2005 ALIA cruise of *RV Kilo Moana*
126 (Jackson et al., 2007; Koppers et al., 2008; 2011), were selected from the more westerly Muli
127 and Taumatau volcanos for this study. Sample 128-21 recovered from Taumatau, at the
128 western termination of the Samoan chain, has an age of 4.8 Ma (Fig 2; Koppers et al., 2008),
129 and the highest ⁸⁷Sr/⁸⁶Sr of any glass yet recovered from Samoa (Jackson et al., 2007).

130 The palagonite alteration separated from the surface of ALIA-104-04 glass was
131 analysed to test if significant halogens are incorporated into altered glass and therefore if

132 undetected alteration of magmatic glasses could significantly influence our results (cf.
133 Kendrick et al., 2012b; 2013a).

134 The Baffin Bay glasses which sample the proto-Icelandic plume, are ~60 Ma and
135 were selected from coastal outcrops, their eruption depths are unknown but pillow textures
136 suggest subaqueous conditions (Jackson et al., 2010; Robillard et al., 1992; Kent et al., 2004).
137 These glasses are related to the picrites that preserve the highest measured terrestrial $^3\text{He}/^4\text{He}$
138 ratio of ~50 Ra in olivine phenocrysts (Starkey et al., 2009; Stuart et al., 2003). They have
139 primitive Pb isotope signatures and primitive $^3\text{He}/^4\text{He}$ up to 24 Ra, that are probably lower
140 than the picrites because of their low He concentrations and radiogenic ingrowth of ^4He over
141 the ~60 Ma since eruption (discussion in Jackson et al., 2010). These samples were selected
142 to test if high $^3\text{He}/^4\text{He}$ glasses from Baffin Bay and Samoa have similar and unique Br/Cl and
143 I/Cl ratios characteristic of high $^3\text{He}/^4\text{He}$ reservoirs within Earth's mantle (see also Kendrick
144 et al., 2013a).

145 Analyses undertaken for this study include: i) Cl, Br, I and K measurements on all
146 Samoan and Baffin samples conducted using the noble gas method utilising irradiation-
147 produced noble gas proxy isotopes ($^{38}\text{Ar}_{\text{Cl}}$, $^{80}\text{Kr}_{\text{Br}}$, $^{128}\text{Xe}_{\text{I}}$ and $^{39}\text{Ar}_{\text{K}}$; Kendrick, 2012).
148 Pristine glass fragments (0.4-2 mm in size) were hand-picked under a binocular microscope
149 and cleaned using distilled water and acetone. 10-30 mg of each sample was then irradiated
150 in either position 5c of the McMaster nuclear reactor, Canada (irradiation UM#48; 15th
151 December 2011; 42 hours; total neutron fluence of 1.2×10^{19} neutrons cm^{-2} ; thermal/fast =
152 2.7); or in the Central Thimble facility of the USGS Triga reactor, Denver, USA (irradiation
153 UM#53; 28th November 2012; 80 hours; total neutron fluence of 6.4×10^{18} neutrons cm^{-2} ;
154 thermal/fast = 0.8).

155 Irradiation produced noble gas proxy isotopes were extracted from the samples by
156 heating in an ultra-high vacuum tantalum resistance furnace. Palagonite was analysed in 20
157 minute heating steps at temperatures of 300 °C and 1500 °C, because up to 50% of the
158 halogen-derived noble gases in palagonite are released at 300 °C. In comparison, noble gases
159 are only released from glasses at >500 °C (Kendrick et al., 2012a) and the glasses were
160 preheated to 300 °C before being fused in single 20 minute heating step at 1500 °C. The
161 extracted gases were purified by standard gettering procedures and analysed with the MAP-
162 215 noble gas mass spectrometer at the University of Melbourne (see Kendrick, 2012 or
163 Kendrick et al., 2013a for details).

164 Noble gas proxy isotopes were converted to Cl, Br, I, K and Ca abundances on the
165 basis of $^{38}\text{Ar}_{\text{Cl}}/\text{Cl}$, $^{80}\text{Kr}_{\text{Br}}/\text{Br}$, $^{128}\text{Xe}_{\text{I}}/\text{I}$, $^{39}\text{Ar}_{\text{K}}/\text{K}$ and $^{37}\text{Ar}_{\text{Ca}}/\text{Ca}$ production ratios monitored with
166 the Hb3gr ^{40}Ar - ^{39}Ar flux monitor (1072 Ma; Roddick, 1983) and three scapolite standards
167 (Kendrick, 2012; Kendrick et al., 2013a). The reported concentrations were cross-checked
168 with electron microprobe Cl, K and Ca determinations undertaken at the University of
169 Melbourne (see Table S1; Kendrick et al., 2014a). Halogen concentration and ratio
170 measurements have internal precision as high as 1-2% (2σ ; Table S1). However,
171 standardisation limits 2σ reproducibility between irradiations to the 5% level for Br and 10 %
172 for I (Fig S2), and the external precision is estimated at 10% for all elements (see Kendrick,
173 2012 and Kendrick et al., 2013a for detailed discussion).

174 SIMS measurements of H_2O , CO_2 , S, F and Cl in three new samples (77-09, 104-04
175 and 128-21) were undertaken at the Department of Terrestrial Magnetism, Carnegie Institute
176 of Washington, using a Cameca IMS 6F ion microprobe, following procedures comparable to
177 those of Workman et al. (2006). Several glasses from dredges 71, 77, 78, 79, 104 and 128
178 were analysed by electron microprobe and laser ablation inductively coupled plasma mass

179 spectrometry at the University of Melbourne. The operating conditions of the Cameca SX-50
180 electron microprobe included an accelerating voltage of 15 keV, current of 30 nA and 20 sec
181 counting times, which gave a detection limit of 260 ppm for Cl. The Agilent 7700x ICP-MS
182 is coupled to a 193 nm excimer laser. The glasses were analysed using a beam diameter of
183 ~80 µm and BHVO2G as the calibration standard: BCR2G, analysed as an unknown, gave
184 reproducible results close to recommended values (Table S1). Typical analytical conditions
185 have been described in detail elsewhere (Woodhead et al., 2007).

186

187 3. Results

188 The new volatile, trace and major element data for Samoan and Baffin glasses are
189 summarised together with representative data from the literature in Table 1 and presented
190 with the standard data in the electronic supplement (Table S1; Fig S2).

191 The Samoan glasses represent alkali basalts and trachy-basalts, they contain 45.2 to
192 48.2 wt. % SiO₂, 3.4 to 6.1 wt. % total alkalis (Na₂O + K₂O), and 6.5 to 4.0 wt. % MgO. The
193 Baffin glasses are basalts with ~49 wt. % SiO₂, ~2.2 wt. % alkalis (Na₂O + K₂O), and ~8.5 wt.
194 % MgO (Table S1; Kent et al., 2004; Workman et al. 2006).

195 The Samoan glasses have 4 to 200 ppm CO₂ but are strongly enriched in incompatible
196 trace elements with 0.6 to 1.7 wt. % K. They have water concentrations of 0.67-1.9 wt. %
197 H₂O, and halogen concentrations of 490-1790 ppm Cl, 1.9-7.2 ppm Br, 19-130 ppb I and
198 900-1900 ppm F (Fig 3; Table 1). In contrast, the Baffin glasses have much lower
199 concentrations of incompatible trace elements and volatiles with ~0.1 wt. % K, 70-80 ppm
200 Cl, 130-240 ppb Br and 3-6 ppb I (Fig 3; Table 1). The low halogen content of the Baffin
201 glasses is unlikely to be explained by degassing because the Cl concentrations are at the

Commented [MK1]: Standard data for BCR2G are in Table 1, halogen standards are shown in Fig S2

202 upper end of the range determined for melt inclusions in related samples (Maisonneuve,
203 2012) and the glasses have normal K/Cl ratios of 13-20 ($\text{Cl/K} = 0.05\text{-}0.08$) that do not
204 suggest Cl loss.

205 The concentrations of Cl, Br and I vary over more than two orders of magnitude
206 between the Baffin and Samoan samples (Fig 3). The majority of samples have I/Cl within
207 the range of $[60 \pm 30] \times 10^{-6}$ that is typical of OIB and MORB glasses (Fig 3c; see Kendrick et
208 al., 2013a). In contrast, the Samoan samples with the highest Cl concentrations have
209 unusually high Br/Cl ratios of up to 4.3×10^{-3} ; these values exceed the range of $[2.8 \pm 0.6] \times 10^{-3}$
210 (2σ) defined by 55 MORB and OIB glasses from other locations analysed by the same
211 technique (Kendrick et al., 2013a), and 45 glasses analysed in other laboratories (Jambon et
212 al., 1995; Schilling et al., 1980), that are collectively considered typical of the mantle (Fig
213 3b). The highest K/Cl of 18 ($\text{Cl/K} = 0.06$) determined for the Samoan samples is fairly
214 typical of mantle values defined by a global database of 566 submarine glasses (Fig S3;
215 Michael and Cornell, 1998; Kendrick et al., 2012b), but the samples with the highest Cl
216 concentrations have low K/Cl of 5.7 ($\text{Cl/K} = 0.17$) and the lowest measured F/Cl of 0.5 (Figs
217 3ad).

218 The palagonite separated from sample 104-04 has concentrations of 35 ppm Cl, 200
219 ppb Br, 85 ppb I and 0.09 wt. % K, which indicate that it is depleted in Cl, Br and K by ~7-30
220 times, and enriched in I by ~5 times, compared to pristine glass from this sample (Table 1).
221 As a result, the palagonite has an I/Cl ratio of 2400×10^{-6} that is 140 times greater than the
222 unaltered glass (Table S1). The high concentration of I in palagonite confirms that the I/Cl
223 ratio of a glass separate could be influenced by the presence of undetected palagonite
224 contaminants, or incipient alteration (Kendrick et al., 2013a); however, it is important to note
225 that minor contamination is unlikely to significantly influence the Br/Cl or K/Cl ratios.

Commented [MK2]: This section makes it clearer than the basis for 'mantle' halogen ratios is 55 MORB and OIB glasses

Consistent with many more from Jambon and Schilling – but these studies had different standards.

Commented [MK3]: This paper included a compilation of published K/Cl data

226 The purity of our glass separates was tested by performing duplicate analyses of glass
227 separated from samples 76-08 and 78-01, that were both strongly affected by palagonite
228 alteration of the rims. Similar results were obtained for 76-08-a and 76-08-b (Tables 1 and
229 S1), confirming efficient removal of contaminating material and demonstrating that the
230 iodine data are reliable for the majority of samples that were more easily prepared as high
231 purity glass separates (Table 1). In contrast, duplicates of 78-01 gave significantly different
232 results for I/Cl, which confirms the sensitivity of this measurement to minor contamination.
233 In this case 78-01-a with an I/Cl ratio of 90×10^{-6} is considered most representative of the melt
234 (Table 1), and the higher I/Cl of 270×10^{-6} obtained for 78-01-b is ascribed to contamination
235 and has been omitted from the figures for clarity (Table 1).

Commented [MK4]: This paragraph has been shortened and moved forward from the discussion – we did perform some duplicate analyses

236 237 4. Discussion

238 The H₂O and halogen content of the Samoan glasses is partly controlled by the mantle
239 source; however, before evaluating the volatile systematics of the source we must first
240 determine the extent to which these volatiles have been modified by shallow level processes
241 including fractional crystallisation, degassing and seawater assimilation.

242 4.1 Magma degassing and assimilation of seawater components

243 The Samoan glasses have vesicularities of 1 to 30 vol. % and low CO₂ concentrations
244 of 4-230 ppm that provide evidence for degassing (see Fig 4; Table 1; Workman et al., 2006).
245 The combined H₂O and CO₂ data indicate that many of the melts were in equilibrium with a
246 H₂O-rich vapour phase (shown by the isopleths in Fig 4a). However, the glasses are
247 suggested to preserve water concentrations close to their pre-degassing values because: i)
248 very little H₂O is lost during open system degassing (Fig 4a); and ii) the equilibrium vapour
249 phase in samples 68-03 and 73-03 (erupted at the shallowest depths of ~1000 m, with

250 vesicularities of 5-20 vol. %), is calculated to contain only 0.01-0.05 wt % H₂O, thus
251 demonstrating that the low water content of these samples (0.7 wt. %; Fig 4) cannot be
252 explained by closed system degassing (Workman et al., 2006; this study). Complex multi-
253 stage degassing scenarios are possible; however, we demonstrate below that the majority of
254 Samoan melts were more strongly influenced by H₂O added through assimilation processes
255 than by H₂O lost during degassing.

256 In the absence of significant H₂O degassing, the concentration of H₂O, as well as
257 incompatible trace elements such as K and Cl, are expected to vary in the Samoan melts as a
258 result of partial melting, mixing isotopically distinct components in the mantle source, and
259 subsequent fractional crystallisation (Fig 5). The liquid lines of descent (LLD) in Fig 5 were
260 modelled with Petrolog3 (Danyushevsky and Plechov, 2011) by progressively removing
261 olivine, pyroxene, plagioclase, ilmenite and magnetite, assuming K, Cl and H₂O are all
262 perfectly incompatible (Fig 5). The Samoan volcanoes sample multiple magma batches and
263 the concentrations of K, H₂O and Cl in the mantle end-members are not known. However, it
264 is significant that, in contrast to K and H₂O, Cl does not follow the expected broad trend
265 toward higher concentrations in the more evolved melts (Fig 5). The lack of a relationship
266 between Cl and MgO, together with the low K/Cl of Cl-rich glasses (Fig 2a and 4; K/Cl of
267 <10 or Cl/K of >0.1), could be explained if some of the melts assimilated seawater-derived Cl
268 (e.g. Coombs et al., 2004; Kendrick et al., 2013a; Kent et al., 1999; 2002; le Roux et al.,
269 2006; Michael and Cornell, 1998; Michael and Schilling, 1989).

270 Possible assimilation processes are evaluated in Fig 6 by examining F, Cl, Br, I, H₂O
271 and K co-variation in three element diagrams that use Cl as a common denominator. This
272 method of plotting the data is advantageous for this purpose because mixing trends are
273 revealed as straight lines (Fig 6). The binary mixing trends in Fig 6 are interpreted to result

Commented [MK5]: This caveat is added to address reviewer 1 comments – we have not fitted a curve to the data the LLD are to illustrate melt evolution which applies regardless of batch.

274 from mixing mantle-derived halogens and halogens introduced by an assimilated brine
275 because:

- 276 i) Extrapolation of the data suggests that one end-member has the
277 characteristics of a brine including: 1) low $\text{H}_2\text{O}/\text{Cl}$ that indicates a high
278 salinity; 2) low K/Cl of <0.2 that is required to maintain charge balance in a
279 Cl^- dominated, Na^+ , Ca^{++} , Fe^{++} , K^+ brine; and 3) low F/Cl ratio that would
280 result from the low solubility of F in a seawater-derived Ca- ($\pm$ P-) bearing
281 aqueous solution (Fig 6; Seyfried and Ding, 1995). In contrast, the second
282 end-member has K/Cl , $\text{H}_2\text{O}/\text{Cl}$, Br/Cl and I/Cl ratios similar to those defined
283 as representative of the mantle in several previous studies of submarine
284 MORB and OIB glasses (Jambon et al., 1995; Schilling et al., 1978, 1980;
285 Michael and Cornell, 1998; Kendrick et al., 2012ab; 2013a; 2014a).
- 286 ii) The variations in Br/Cl , K/Cl , F/Cl and $\text{H}_2\text{O}/\text{Cl}$ are not correlated with
287 $^{87}\text{Sr}/^{86}\text{Sr}$ (or $^3\text{He}/^4\text{He}$), which is the expected result of assimilating brines
288 with very high concentrations of Cl, and low concentrations of Sr, relative to
289 the melts (Kendrick et al., 2013a). In contrast, correlations between K/Cl
290 and $^{87}\text{Sr}/^{86}\text{Sr}$, or Br/Cl and $^{87}\text{Sr}/^{86}\text{Sr}$, might be expected if the variation was
291 intrinsic to the mantle or related to subduction recycling (e.g. Stroncik and
292 Haase, 2004; Kendrick et al., 2012a; 2014a).
- 293 iii) Similar mixing trends have been observed in MORB previously and are
294 therefore unrelated to subduction but can be explained by brine assimilation
295 (Kendrick et al., 2013a).
- 296 iv) The samples closest to the suggested brine end-member come from the
297 Vailulu'u volcano which is the most hydrothermally active of the Samoan

Commented [MK6]: This section is organised differently, the brine characteristics are listed prominently.

volcanos investigated (Staudigel et al., 2004), and therefore a likely location for magmas to assimilate hydrothermal brines.

v) The lack of a correlation between Cl and MgO provides independent evidence that the melts have assimilated seawater-derived Cl in some form (Fig 5).

Commented [MK7]: This is important supporting evidence assimilation has occurred.

The good fit of the mixing model to halogen data for glasses from 5 different volcanoes that erupted from ~100 yrs to 5 Ma is extremely significant (Fig 6). It suggests that over the last 5 Ma, all the Samoan volcanoes have had sources characterised by similar Br/Cl, I/Cl, F/Cl, H₂O/Cl and K/Cl ratios and that the magmas have been affected by remarkably uniform assimilation processes (section 4.1.1). This observation is consistent with the uniformly high salinity of brines (55 ± 15 wt. % salts) assimilated by magmas exhibiting anomalously high Cl concentrations at other mid-ocean ridge and oceanic island settings investigated for Cl assimilation processes (see Fig 8 of Kendrick et al., 2013a; data of Coombs et al., 2004; Kent et al., 1999; 2002; le Roux et al., 2006; Wanless et al., 2011). As a point of comparison, assimilation of Cl at both Samoan and Hawaiian volcanoes appears to be related to location (Dixon and Clague, 2001), and in contrast to some previous studies, there is no evidence that the most evolved melts have assimilated the highest proportion of Cl (Fig 5; cf. Wanless et al., 2011).

4.1.1 Brine characteristics and origin

Hydrothermal vent fluids typically have K/Cl of <0.05 and F/Cl of <0.0001 that are close to the seawater values of 0.02 and 0.00007, respectively (Li and Schoonmaker, 2003; Mottl et al., 2011). As already mentioned, a low K/Cl ratio is required to maintain charge balance in a seawater-derived Na-Ca brine (Vanko, 1988), and a low F/Cl ratio is expected

Commented [MK8]: Data for vents are summarised

322 because of fluorine's uptake by minerals and low solubility in aqueous fluids (e.g. Frohlich et
323 al., 1983; Seyfried and Ding, 1995). The salinity of vent fluids is variable, and typically in
324 the range 1-8 wt. % salts (Oosting and von Damm, 1996; Fontaine et al., 2007), but much
325 higher salinity fluids with up to 60 wt. % salt are preserved in fluid inclusions from deeper
326 greenschist-amphibolite facies environments in the oceanic crust (Vanko, 1986; 1988; Kelley
327 et al., 1993). Therefore, the uniformly high salinity of the brines (55 ± 15 wt % salt)
328 assimilated by magmas from several different Samoan volcanoes (Fig 6), and other seafloor
329 locations (see Kendrick et al., 2013a data of Wanless et al. 2011, Coombs et al., 2004; Kent
330 et al, 1999; 2002; Le Roux et al., 2006), suggests that: i) assimilation may be restricted to
331 high salinity brines by the relative solubilities of Cl and H₂O in silicate melts; and ii)
332 assimilation is more likely to occur at depths of >2-3 km in the crust rather than on the
333 seafloor (see also Coombs et al., 2004; Kendrick et al., 2013a; le Roux et al., 2006).

334 The Br/Cl and I/Cl ratios of the melts do not fit the binary mixing model in Fig 6 as
335 well as the K/Cl, F/Cl and H₂O/Cl ratios. This is not explained by analytical uncertainty
336 which is at the 2-8 % level (2σ) for both Br/Cl and I/Cl (Table S1). Palagonite contamination
337 (or incipient alteration of the glasses) has the potential to influence the glasses' I/Cl ratios;
338 however, the low concentrations of Cl and Br in palagonite indicate that its presence would
339 have a negligible effect on Br/Cl ratios (section 3; Table 1). Therefore we suggest that the
340 quality of fit for both Br/Cl and I/Cl in our mixing model (Fig 6) is influenced by variation of
341 these ratios in the assimilated brines. This is possible because whereas K/Cl, F/Cl and H₂O/Cl
342 are all fixed by requirements of solubility and charge balance, Br and I are highly soluble
343 trace constituents. As a result the Br/Cl and I/Cl ratios of brines behave as essentially free
344 variables. The mixing model indicates that, on average, the Samoan melts assimilated brines
345 with Br/Cl of ~0.005, that were about 40 % higher than seawater, and I/Cl ratios that were on
346 average about ten times seawater (Fig 6c,d and e).

A previous study of magmatic glasses demonstrated that, as in Samoa, brines assimilated by melts from the Galapagos Spreading Centre and the NW part of the Lau Basin had Br/Cl ratios 10's of percent higher than seawater (Kendrick et al., 2013a). However, in contrast to Samoa, these glasses preserved I/Cl ratios closer to the seawater value (Kendrick et al., 2013a). The data available for vent fluids demonstrate that fluid interaction with I-rich organic matter present in marine sediments elevates the I/Cl ratios, and to a lesser extent Br/Cl ratios, of some vent fluids above seawater values (Campbell and Edmund, 1989; You et al., 1994; Kawagucci et al., 2011). The majority of vent fluids preserve seawater-like Br/Cl (e.g. von Damm et al., 1997; Seyfried et al., 2003); however, lower than seawater Br/Cl ratios are known from condensed vapour phases (~1 wt % salt) emitted at 9-10° N on the East Pacific Rise (Oosting and von Damm, 1996).

It is concluded that the elevated I/Cl ratios inferred for the Samoan brines are consistent with a fluid history involving interaction with sediments. A number of processes may have contributed to the elevated Br/Cl ratios of the brines including fluid interaction with sediments, and partitioning of $H_2O > Cl > Br$ into hydrous alteration minerals such as amphibole (Fig 6; Kendrick et al., 2013a). It remains unclear if high salinity brines formed by phase separation under specific conditions might also be enriched in Br/Cl relative to the vapour phase (Oosting and von Damm, 1996; Berndt and Seyfried, 1997; von Damm et al., 1997; Liebscher et al., 2006; Foustoukos and Seyfried, 2007). However, sub-critical seawater would boil when heated to magmatic temperatures during assimilation, with the likely result that vapour phases would be preferentially vented on the seafloor, and the dense residual brines would be preferentially assimilated by the magma (Fig 6; Kendrick et al., 2013a) and/or retained in the lower crust (e.g. Bischoff and Rosenbauer, 1989).

If brine assimilation occurred at depths of >2 km beneath the seafloor as suggested, then the preservation of the mixing trends in Fig 6 is consistent with the suggestion that very

372 little H₂O was lost during degassing of CO₂ from these melts (Fig 4a; Table 1). Sample 128-
 373 21 with only 4 ppm CO₂ (Fig 4a), lies on the mixing lines in Figs 6b and 6d, suggesting
 374 minimal H₂O loss occurred (Fig 4a). The Vailulu'u samples show scatter in H₂O/Cl;
 375 however, the samples recovered from ~1000 m in dredges 70 and 73 (Table 1) include both
 376 the lowest and highest H₂O/Cl ratios, suggesting the scatter is unrelated to degassing (Fig 6a).
 377 The preservation of the mixing trends also suggests ratios of incompatible elements (e.g.
 378 Br/Cl, I/Cl, F/Cl, H₂O/Cl) were not significantly altered by fractional crystallisation of the
 379 Samoan magmas after assimilation (Fig 6).

380

381 4.2 Samoan mantle source characteristics

382 The mixing model in Fig 6 enables the amount of H₂O and Cl introduced into the
 383 Samoan magmas by brine assimilation to be quantified. The calculations can be undertaken
 384 using any ratio that differs between the mantle and brine and 10% increments of the total Cl
 385 are shown on each of the mixing lines in Fig 6 with the associated uncertainties summarised
 386 for individual samples in Table 1. The Br/Cl data are important because this ratio exhibits
 387 limited variation in the mantle (0.0028 ± 0.0006 (2σ); Kendrick et al., 2013a; 2014a),
 388 suggesting the least contaminated Samoan samples with Br/Cl of 0.0029 lie close to the
 389 mantle end-member (Fig 6). In comparison, the Br/Cl of the brine obtained from the
 390 regression of data in Fig 6c (0.0046 ± 0.004 ; 2σ), is about twice the mantle value. In contrast,
 391 hydrothermal fluids have F/Cl of $<<0.01$ (Seyfried and Ding, 1995; Li and Schoonmaker,
 392 2003), which is many times lower than the maximum F/Cl of 2.3 measured in a Samoan
 393 glass (Fig 6a). As a result, mixing calculations based the F/Cl ratio, using conservative
 394 estimates for the brine (0.01000 ± 0.00999) and mantle (2.4 ± 0.5) end-members (Fig 6a),

395 give results that are more precise but indistinguishable from similar calculations undertaken
396 with the Br/Cl data (cf. samples 76-08-a and 76-08-b in Table 1).

397 The mixing calculations indicate that sample 75-10 with the highest measured F/Cl
398 ratio assimilated between zero and 26% of its total Cl, and the other T'au samples from
399 dredges 74 and 75 probably also assimilated very little Cl (Table 1). In comparison, samples
400 68-03, 71-02, 71-11, 71-13, 71-22 and 76-11 are estimated to have assimilated about 70% of
401 their total Cl (Table 1). The quantitative mixing model developed here indicates that water
402 assimilation is more significant than was previously recognised on a qualitative basis (cf.
403 Workman et al., 2006). However, the proportion of the H₂O assimilated by the T'au magmas
404 is close to zero and probably accounts for a maximum of 20-30 % of the total H₂O in samples
405 68-03 and 73-03 (Table 1). In absolute terms, the T'au melts are estimated to have
406 assimilated up to a few hundred parts per million of brine and sample 71-02 is estimated to
407 have assimilated 0.45 ± 13 wt. % brine (Table 1). Based on reasonable estimates for the
408 densities of brine (1.3-1.4 g cm³) and melt (2.6-2.7 g cm³), this is equivalent to a maximum of
409 8 or 9 cm³ of brine being assimilated by a litre of magma (Fig 6; Table 1).

410 The assimilation-corrected H₂O/Ce and Cl/K ratios of Samoan melts are plotted
411 together with the measured F/Nd ratio, which is not influenced by brine assimilation, as a
412 function of ⁸⁷Sr/⁸⁶Sr in Fig 7. Note that each ratio represents an element pair of similar
413 compatibility, and that Cl/K is used in preference to K/Cl to enable comparison of
414 volatile/lithophile pairs expressed in the same format (whereas K/Cl provides a common
415 denominator in Fig 6). The MORB mantle has H₂O/Ce of 150-300 (e.g. Michael, 1995) and
416 F/Nd of 20 ± 6 (Workman et al., 2006), but as a result of unresolved seawater assimilation, its
417 Cl/K is less well defined (Michael and Cornell, 1998): we adopt a Cl/K of 0.05-0.1 that
418 encompasses the median value of previous MORB analyses (0.08 ± 0.01), but it should be
419 noted that MORB can have much lower Cl/K values (see Fig S3).

420 The assimilation-correction reduces the measured H₂O/Ce ratio by 0-30 %, which has
421 a minor impact on the interpretation of the H₂O data (Fig 7a; Table 1; Workman et al., 2006).
422 In contrast, the assimilation correction has a profound effect on the Cl/K data reducing
423 measured Cl/K ratios from maxima of 0.17-0.18 to assimilation-corrected values of ~0.05
424 (Figs 7b; Table 1). The mean assimilation-corrected Cl/K of 0.05 ± 0.02 (2σ) is controlled by
425 the binary mixing model used to correct the data (Fig 6), and the scatter of the data around
426 the mean corrected Cl/K value therefore reflects the fit quality of the model (Fig 6). The
427 modelling indicates that most of the variation in Cl/K (or K/Cl) measured in Samoan glasses
428 results from brine assimilation (Fig 6); however, the fit quality means that minor systematic
429 variation in the Cl/K of the Samoan source is not precluded (Fig 7b).

430 In general, the model indicates the Samoan source has lower H₂O/Ce and Cl/K than
431 median MORB, but F/Nd at the high end of the MORB range (Fig 7). Therefore, despite
432 their overall trace element and volatile enrichment, the Samoan, Society and Pitcairn EM
433 magmas are all depleted in H₂O and Cl, but not F, compared to median MORB and lithophile
434 elements of similar compatibility (Kendrick et al., 2014a). A relative depletion of Cl as well
435 as H₂O may therefore be a characteristic of EM reservoirs (Fig 7; Dixon et al., 2002, Wallace,
436 2002; Stroncik and Haase, 2004; Kendrick et al., 2014a). However, pooling the data from
437 Society, Pitcairn and Samoa which extends to the highest ⁸⁷Sr/⁸⁶Sr (Fig 1), suggests that the
438 H₂O/Ce and Cl/K ratios are not strongly correlated with ⁸⁷Sr/⁸⁶Sr (Fig 7; cf. Workman et al.,
439 2006), and/or that part of the Samoan source probably has higher H₂O/Ce and Cl/K at any
440 given ⁸⁷Sr/⁸⁶Sr than either the Pitcairn or Society sources (Fig 7).

441 Finally, the five Samoan glasses least affected by brine assimilation (samples 74-02;
442 75-02; 75-10; 77-09 and 128-21) include sample 128-21 with the highest ⁸⁷Sr/⁸⁶Sr of 0.7125,
443 and sample 75-02 with a ³He/⁴He of 15 R/Ra (Table 1) that represent the EM2 and primitive
444 components of the Samoan plume, respectively (Fig 1; Jackson et al., 2007). The Br/Cl and

I/Cl ratios of these samples are similar to the Baffin Bay sample with $^3\text{He}/^4\text{He}$ of 24 R/Ra (Table 1) and other MORB and OIB samples, suggesting that all these mantle reservoirs have Br/Cl and I/Cl ratios similar to the MORB range (Fig 8; Kendrick et al., 2013a; 2014a). This conclusion is further supported by the mixing systematics of Br/Cl and I/Cl in glasses with $^3\text{He}/^4\text{He}$ of 16-28 Ra from the NW part of the Lau Basin (Kendrick et al., 2013a).

4.3 Implications for the global volatile cycling

Commented [MK9]: This section has been simplified and is more circumspect

The apparent uniformity of Br/Cl and I/Cl in the Earth's mantle (Fig 8; $\text{I/Cl} = 0.00006 \pm 0.00003$) reflects the similar compatibilities of these elements in the mantle (e.g. Schilling et al., 1980; Balcone-Boissard et al., 2010; Kendrick et al., 2012b), and is in stark contrast to the behaviour of these elements in Earth's surface reservoirs and subduction zones. The Br/Cl ratio varies by more than an order of magnitude in the surface reservoir and I/Cl is even more variable (e.g. Holser, 1979; Muramatsu and Wedepohl, 1998; Kendrick et al., 2013b). Seawater has an I/Cl ratio of 0.000003 that is $\sim 10,000$ times lower than the maximum values of ~ 0.03 found in some organic-rich marine sediments, sedimentary marine pore fluids and serpentinites (Kendrick et al., 2013b; John et al., 2011; Muramatsu and Wedepohl, 1998; Muramatsu et al., 2001; Snyder et al., 2005).

The uniformly low I/Cl of the mantle (0.00006 ± 0.00003) compared to the maximum of ~ 0.03 in sedimentary rocks and serpentinites, could be simply explained if halogens are not significantly recycled into the mantle, and halogens have a dominantly primordial origin in the mantle (Schilling et al., 1978). However, it is increasingly recognised that a significant portion of the volatiles entering subduction zones could be deeply subducted into the mantle (e.g. Jacobsen and van der Lee, 2006; Walter et al., 2011). Furthermore, the non-radiogenic noble gas isotopes in the mantle are now believed to have a dominantly subducted atmospheric origin (cf. Porcelli and Wasserburg, 1995; Sarda et al., 1999; Sarda 2004;

469 Tieloff et al., 2000; Holland and Ballentine, 2006), implying that subduction of other
470 seawater-derived volatiles is likely.

471 Estimated rates of Cl subduction in sediments, altered ocean crust and serpentinites
472 are more poorly defined than Cl outputs through volcanism and it is unclear from the existing
473 data if there is a net flow of Cl into the mantle, or a net flow of Cl out of the mantle, at the
474 present day (cf. Table 2; Schilling et al., 1978; Ito et al., 1983; Sharp and Barnes, 2004; Sano
475 et al., 2008; Barnes and Cisneros, 2012). The scenario in Table 2 demonstrates that contrary
476 to early assumptions, net subduction of Cl is possible, if Cl and other volatiles are
477 inefficiently lost through magmatic arcs. In this scenario, we envisage that deeply subducted
478 Cl would be stored in cold deep reservoirs within the slab, such as serpentinites, that can be
479 subducted beyond magmatic arcs (Schmidt and Poli, 1998; Green II et al., 2010). The
480 subducted Cl would then be incrementally lost into mantle reservoirs beyond the arc, with
481 perhaps ~1-10 % of the initial Cl concentrated with other trace elements in the EM source
482 (Kendrick et al. 2014b). This figure is comparable to the proportion of subducted H₂O
483 estimated to reach the EM source (Dixon et al., 2002). Higher efficiencies of Cl loss through
484 the arc are of course possible, if balanced by significant reductions in the input flux, or
485 increases in the arc magma Cl flux (cf. Table 2). However, the net flow of Cl is more likely
486 to have been into the mantle, in the past, if primordial heavy halogens (like noble gases), had
487 very low abundances as a result of catastrophic outgassing early in Earth's history (Tolstikhin
488 et al., 2014), or if volatiles were introduced to Earth in a late veneer post-dating accretion of
489 the mantle (Wänke, 1981; Deruelle et al., 1992; Javoy, 1997; Bonifacie et al., 2008; Holland
490 et al., 2009).

491 The relative uniformity of mantle Br/Cl and I/Cl (Fig 8) can be reconciled with
492 significant Cl subduction in the following ways: one possibility is that the subduction of Cl in
493 sedimentary rocks and serpentinites with high I/Cl is balanced by subduction of altered ocean

494 crust, which has an unknown halogen signature but could include much lower I/Cl.
495 Alternatively, if the altered ocean crust has Br/Cl and I/Cl overlapping the MORB range
496 subduction of altered ocean crust could dominate the Cl subduction budget (cf. Table 2;
497 Kendrick et al., 2014b). A second possibility is that halogens present in sediments and
498 serpentinites with initially high I/Cl might be fractionated during subduction such that I is
499 preferentially lost in fluids, and the minerals, fluid inclusions and grain boundaries hosting
500 halogens in deeper slab environments acquire MORB-like Br/Cl and I/Cl ratios (cf. Kendrick
501 et al., 2011). A combination of these processes seems plausible because studies of eclogite
502 facies serpentinites have shown that Cl is preferentially subducted to greater depths than Br
503 or I (Kendrick, et al., 2011; John et al., 2011). Furthermore, halogens in subduction related
504 fluids have I/Cl ratios that decrease across the arc from maximum values of ~0.01 in the
505 forearc, to ~0.001 in the arc and to lower MORB-like ratios of <0.0001 in some backarc
506 basins (Kendrick et al., 2013b; 2014b). The alternative possibility that halogens were not
507 subducted into the sources of the investigated back arc basin basalts (BABB) with MORB-
508 like Br/Cl and I/Cl, is not favoured by the elevated Cl/K and H₂O/Ce ratios of these BABB,
509 that demonstrate up to 90 % of their total Cl is related to subduction (Kendrick et al., 2014b).

Commented [MK10]: Eclogite facies rocks are mentioned

511 5. Summary and conclusions

512 Submarine glasses from Samoa contain halogens derived from the mantle and introduced by
513 assimilation of high salinity brines (e.g. 55 ± 15 wt % salts), with F/Cl of close to seawater,
514 Br/Cl of ~0.005 that is ~40% higher than seawater, and I/Cl of ten times the seawater value.
515 Quantitative corrections for brine assimilation show 0-70 % of the total Cl and 0-30 % of the
516 total H₂O in the Samoan melts investigated was introduced by brines accounting for up to
517 0.45 % of the melt mass. The preservation of the mixing trends in melts that have

518 experienced some degassing favours the dominance of open system degassing with minimal
519 H₂O loss.

520 Assimilation-corrected data confirm that, despite an overall volatile enrichment, the
521 Samoan EM2 end-member with Cl/K of 0.05 ± 0.02 (2σ) and H₂O/Ce of 83 ± 3 is modestly
522 depleted in Cl relative to K as well as H₂O relative to Ce, compared to median MORB.
523 Furthermore, the samples least affected by brine assimilation, with $^{87}\text{Sr}/^{86}\text{Sr}$ of 0.7045-
524 0.7125, have indistinguishable Br/Cl and I/Cl, within the MORB range that is currently
525 estimated as $(2.8 \pm 0.6) \times 10^{-3}$ for Br/Cl and $(60 \pm 30) \times 10^{-6}$ for I/Cl (2σ ; Kendrick et al.,
526 2013a). The uniformity of mantle Br/Cl and I/Cl can be explained if the subduction zone
527 ‘filter’ controls the relative abundances of Cl, Br and I in deeply subducted slabs and recycled
528 halogens have been circulated throughout the entire mantle. Differences in the Cl/K and
529 H₂O/Ce of different mantle components then reflect the efficiency and timing of slab
530 dehydration.

531

532 **Words = 6417**

533

534 **Acknowledgements**

535 Stanislav Szczepanski is thanked for technical assistance in the University of Melbourne
536 noble gas laboratory. Dr Mark Kendrick has been the recipient of an Australian Research
537 Council QEII Fellowship and Future Fellowship (project number FT 130100141).
538 Jackson acknowledges support from NSF grants EAR-1145202 and OCE-1153894. This
539 research used samples recovered by the 1999 AVON2/3 cruise of *R/V Melville* and the 2005
540 Alia cruise of the *R/V Kilo Moana*. We thank Ray Burgess and John Lassiter for

constructive reviews, and Tim Elliot for editorial suggestions, that improved the focus of this manuscript.

543

544 **References**

- 545 Balcone-Boissard, H., Villemant, B., Boudon, G., 2010. Behavior of halogens during the
546 degassing of felsic magmas. *Geochem. Geophys. Geosyst.* 11, 22.
- 547 Barnes, J. D. and Cisneros, M., 2012. Mineralogical control on the chlorine isotope
548 composition of altered oceanic crust. *Chemical Geology* **326–327**, 51-60.
- 549 Berndt, M. E. and Seyfried, W. E., 1997. Calibration of Br/Cl fractionation during subcritical
550 phase separation of seawater: Possible halite at 9 to 10 degrees N East Pacific Rise.
551 *Geochimica et Cosmochimica Acta* **61**, 2849-2854.
- 552 Bischoff, J.L., Rosenbauer, R.J., 1989. Salinity variations in submarine hydrothermal systems
553 by layered double diffusive convection. *Journal of Geology* 97, 613-623.
- 554 Bonifacie, M., Jendrzewski, N., Agrinier, P., Humler, E., Coleman, M., and Javoy, M.,
555 2008. The chlorine isotope composition of Earth's mantle. *Science* **319**, 1518-1520.
- 556 Campbell, A.C., Edmond, J.M., 1989. Halide systematics of submarine hydrothermal vents.
557 *Nature* 342, 168-170.
- 558 Coombs, M.L., Sisson, T.W., and Kimura, J.I., 2004, Ultra-high chlorine in submarine
559 Kilauea glasses: evidence for direct assimilation of brine by magma: *Earth and*
560 *Planetary Science Letters*, v. 217, p. 297-313.
- 561 Danyushevsky, L.V., Plechov, P., 2011. Petrolog3: Integrated software for modeling
562 crystallization processes. *Geochemistry, Geophysics, Geosystems* 12, Q07021.
- 563 Deruelle, B., Dreibus, G., and Jambon, A., 1992, Iodine abundances in oceanic basalts:
564 implications for Earth dynamics: *Earth and Planetary Science Letters*, v. 108, p. 217-
565 227.

566 Devey, C.W., Albarede, F., Cheminee, J.L., Mischard, A., Muhe, R., Stoffers, P., 1990.
 567 Active Submarine Volcanism on the Society Hotspot Swell (West Pacific): A
 568 Geochemical Study. *Journal of Geophysical Research* 95, 5049-5066.
 569 Dixon, J.E., and Clague, D.A., 2001, Volatiles in basaltic glasses from Loihi seamount,
 570 Hawaii: Evidence for a relatively dry plume component: *Journal of Petrology*, v. 42,
 571 p. 627-654.
 572 Dixon, J.E., Leist, L., Langmuir, C., and Schilling, J.-G., 2002, Recycled dehydrated
 573 lithosphere observed in plume-influenced mid-ocean-ridge basalt: *Nature*, v. 420, p.
 574 385-389.
 575 Eiler, J.M., Schiano, P., Kitchen, N., and Stolper, E.M., 2000, Oxygen-isotope evidence for
 576 recycled crust in the sources of mid-ocean-ridge basalts: *Nature*, v. 403, p. 530-534.
 577 Farley, K.A., Natland, J.H., and Craig, H., 1992, Binary mixing of enriched and undegassed
 578 (primitive?) mantle components (He, Sr, Nd, Pb) in Samoan lavas: *Earth and*
 579 *Planetary Science Letters*, v. 111, p. 183-199.
 580 Frohlich, P.N., Kim, K.H., Jahnke, R., Burnett, W.C., Soutar, A., Deakin, M., 1983. Pore
 581 water fluoride in Peru continental margin sediments: Uptake from seawater.
 582 *Geochimica et Cosmochimica Acta* 47, 1605-1612.
 583 Fontaine, F.J., Wilcock, W.S.D., Butterfield, D.A., 2007. Physical controls on the salinity of
 584 mid-ocean ridge hydrothermal vent fluids. *Earth and Planetary Science Letters* 257,
 585 132-145.
 586 Foustoukos, D.I., Seyfried, W.E., 2007. Trace element partitioning between vapor, brine and
 587 halite under extreme phase separation conditions. *Geochimica et Cosmochimica Acta*
 588 71, 2056-2071.
 589 Green II, H. W., Chen, W.-P., and Brudzinski, M. R., 2010. Seismic evidence of negligible
 590 water carried below 400-km depth in subducting lithosphere. *Nature* **467**, 828-831.

591 Hart, S.R., Staudigel, H., Koppers, A.A.P., Blusztajn, J., Baker, E.T., Workman, R., Jackson,
592 M., Hauri, E., Kurz, M., Sims, K., Fornari, D., Saal, A., Lyons, S., 2000. Vailulu'u
593 undersea volcano: The New Samoa. *Geochem. Geophys. Geosyst.* 1.

594 Hilton, D.R., Fischer, T.P., and Marty, B., 2002, Noble Gases and Volatile Recycling at
595 Subduction Zones, in Porcelli, D., Ballentine, C., and Wieler, R., eds., Noble Gases in
596 Geochemistry and Cosmochemistry, Volume 47: Washington D.C., Mineralogical
597 Society of America, p. 319-370.

598 Hofmann, A.W., White, W.M., 1982. Mantle Plumes from Ancient Oceanic-Crust. *Earth and*
599 *Planetary Science Letters* 57, 421-436.

600 Hofmann, A.W., 2003, Sampling Mantle Heterogeneity through Oceanic Basalts: Isotopes
601 and Trace Elements, in Carlson, R.L., ed., Treatise of Geochemistry Volume 2: The
602 Core and Mantle, Volume 2, Elsevier Ltd., p. 61-101.

603 Holland, G., and Ballentine, C.J., 2006, Seawater subduction controls the heavy noble gas
604 composition of the mantle: *Nature*, v. 441, p. 186-191.

605 Holland, G., Cassidy, M., and Ballentine, C. J., 2009. Meteorite Kr in Earth's Mantle
606 Suggests a Late Accretionary Source for the Atmosphere. *Science* **326**, 1522-1525.

607 Holser, W.T., 1979. Trace elements and isotopes in evaporites, in: Burns, R.G. (Ed.), Marine
608 minerals: Mineralogical Society of America Short Course Notes, pp. 295-346.

609 Ito, E., Harris, D.M., and Anderson, A.T., 1983, Alteration of Oceanic-Crust and Geologic
610 Cycling of Chlorine and Water: *Geochimica et Cosmochimica Acta*, v. 47, p. 1613-
611 1624.

612 Jackson, M.G., Carlson, R.W., Kurz, M.D., Kempton, P.D., Francis, D., and Blusztajn, J.,
613 2010, Evidence for the survival of the oldest terrestrial mantle reservoir: *Nature*, v.
614 466, p. 853-856.

615 Jackson, M.G., Hart, S.R., Koppers, A.A.P., Staudigel, H., Konter, J., Blusztajn, J., Kurz, M.,
 616 and Russell, J.A., 2007, The return of subducted continental crust in Samoan lavas:
 617 *Nature*, v. 448, p. 684-687.

618 Jackson, M.G., S.R. Hart, N. Shimizu, and J. Blusztajn (2009). Pervasive cpx-whole rock
 619 isotopic disequilibrium in Polynesian hotspot lavas: Evidence supporting isotopic
 620 variability in olivine and clinopyroxene-hosted melt inclusions. *Geochem. Geophys.*
 621 *Geosys. (G-cubed)* 10, Q03006, doi:10.1029/2008GC002324

622 Jacobsen, S. D. and van der Lee, S., 2006. *Earth's Deep Water Cycle*. AGU, Washington,
 623 DC.

624 Jambon, A., Deruelle, B., Dreibus, G., Pineau, F., 1995. Chlorine and bromine abundance in
 625 MORB: The contrasting behaviour of the Mid-Atlantic Ridge and East Pacific Rise
 626 and implications for chlorine geodynamic cycle. *Chemical Geology* 126, 101-117.

627 Javoy, M., 1997. The major volatile elements of the Earth: Their origin, behavior, and fate.
 628 *Geophysical Research Letters* **24**, 177-180.

629 John, T., Scambelluri, M., Frische, M., Barnes, J.D., Bach, W., 2011. Dehydration of
 630 subducting serpentinite: Implications for halogen mobility in subduction zones and
 631 the deep halogen cycle. *Earth and Planetary Science Letters* 308, 65-76.

632 Kawagucci, S., Chiba, H., Shibash, J.-i., Yamanaka, T., Toki, T., Muramatsu, Y., Ueno, Y.,
 633 Makabe, A., Inoue, K., Yoshida, N., Nakagawa, S., Nunoura, T., Takai, K., Takahata,
 634 N., Sano, Y., Narita, T., Teranishi, G., Obata, H., and Gamo, T., 2011. Hydrothermal
 635 fluid geochemistry at the Iheya North field in the mid-Okinawa Trough: Implication
 636 for origin of methane in subseafloor fluid circulation systems. *Geochem. J.* **45**, 109-
 637 124.

638 Kelley, D.S., Gillis, K.M., Thompson, G., 1993. Fluid evolution in submarine magma-
 639 hydrothermal systems at the Mid-Atlantic Ridge. *J. Geophys. Res.-Solid Earth* 98,
 640 19579-19596.

641 Kendrick, M.A., 2012, High precision Cl, Br and I determination in mineral standards using
 642 the noble gas method: *Chemical Geology*, v. 292-293, p. 116-126.

643 Kendrick, M.A., Arculus, R.J., Burnard, P., Honda, M., 2013a. Quantifying brine assimilation
 644 by submarine magmas: Examples from the Galápagos Spreading Centre and Lau
 645 Basin. *Geochimica et Cosmochimica Acta* 123, 150-165.

646 Kendrick, M. A., Arculus, R. J., Danyushevsky, L., Kamenetsky, V. S., Woodhead, J., and
 647 Honda, M., 2014. Subduction-related halogens (Cl, Br and I) and H₂O in magmatic
 648 glasses from Southwest Pacific Backarc Basins. *Earth and Planetary Science Letters*
 649 400, 165-176.

650 Kendrick, M.A., Honda, M., Pettke, T., Scambelluri, M., Phillips, D., and Giuliani, A.,
 651 2013b, Subduction zone fluxes of halogens and noble gases in seafloor and forearc
 652 serpentinites: *Earth and Planetary Science Letters*, v. 365, p. 86-96.

653 Kendrick, M.A., Jackson, M., Kent, A.J.R., Hauri, E., Wallace, P., and Woodhead, J., 2014a,
 654 Contrasting behaviours of CO₂, S, H₂O and halogens (F, Cl, Br and I) in the
 655 enriched-mantle melts of the Pitcairn and Society seamounts: *Chemical Geology*, 370,
 656 69-81.

657 Kendrick, M.A., Kamenetsky, V.S., Phillips, D., and Honda, M., 2012b, Halogen (Cl, Br, I)
 658 systematics of mid-ocean ridge basalts: a Macquarie Island case study: *Geochimica et*
 659 *Cosmochimica Acta*, v. 81, p. 82-93.

660 Kendrick, M.A., Scambelluri, M., Honda, M., Phillips, D., 2011. High abundances of noble
 661 gas and chlorine delivered to the mantle by serpentinite subduction. *Nat. Geosci.* 4,
 662 807-812.

663 Kendrick, M.A., Woodhead, J.D., and Kamenetsky, V.S., 2012a, Tracking halogens through
 664 the subduction cycle: *Geology*, v. 40, p. 1075-1078.
 665 Kent, A.J.R., Clague, D.A., Honda, M., Stolper, E.M., Hutcheon, I.D., and Norman, M.D.,
 666 1999, Widespread assimilation of a seawater-derived component at Loihi Seamount,
 667 Hawaii: *Geochimica et Cosmochimica Acta*, v. 63, p. 2749-2761.
 668 Kent, A.J.R., Peate, D.W., Newman, S., Stolper, E.M., and Pearce, J.A., 2002, Chlorine in
 669 submarine glasses from the Lau Basin: seawater contamination and constraints on the
 670 composition of slab-derived fluids: *Earth and Planetary Science Letters*, v. 202, p.
 671 361-377.
 672 Kent, A.J.R., Stolper, E.M., Francis, D., Woodhead, J., Frei, R., Eiler, J., 2004. Mantle
 673 heterogeneity during the formation of the North Atlantic Igneous Province:
 674 Constraints from trace element and Sr-Nd-Os-O isotope systematics of Baffin Island
 675 picrites. *Geochemistry, Geophysics, Geosystems* 5, Q11004.
 676 Koppers, A.A.P., Russell, J.A., Jackson, M.G., Konter, J., Staudigel, H., Hart, S.R., 2008.
 677 Samoa reinstated as a primary hotspot trail. *Geology* 36, 435-438.
 678 Koppers, A.A.P., Russell, J.A., Roberts, J., Jackson, M.G., Konter, J.G., Wright, D.J.,
 679 Staudigel, H., Hart, S.R., 2011. Age systematics of two young en echelon Samoan
 680 volcanic trails. *Geochem. Geophys. Geosyst.* 12.
 681 le Roux, P.J., Shirey, S.B., Hauri, E.H., Perfit, M.R., and Bender, J.F., 2006, The effects of
 682 variable sources, processes and contaminants on the composition of northern EPR
 683 MORB (8-10 degrees N and 12-14 degrees N): Evidence from volatiles (H₂O, CO₂, S)
 684 and halogens (F, Cl): *Earth and Planetary Science Letters*, v. 251, p. 209-231.
 685 Li, Y.-H., Schoonmaker, J.E., 2003. Chemical composition and mineralogy of marine
 686 sediments, *Treatise on Geochemistry* pp. 1-35.

687 Liebscher, A., Luders, V., Heinrich, W., and Schettler, G., 2006. Br/Cl signature of
 688 hydrothermal fluids: liquid-vapour fractionation of bromine revisited. *Geofluids* 6,
 689 113-121.

690 Ludwig, K.R., 2009, User's manual for Isoplot 3.7, Berkeley Geochronology Center Special
 691 Publication No. 4.

692 Maisonneuve, M., 2012. Constraining the nature of E- and N-type components in the Baffin
 693 Island picrites using olivine-hosted melt inclusions. McGill University, Montreal,
 694 Quebec, Canada, p. 139.

695 McDougall, I., 2010. Age of volcanism and its migration in the Samoa Islands. *Geological*
 696 *Magazine* 147, 705-717.

697 Michael, P., 1995, Regionally distinctive sources of depleted MORB - evidence from trace
 698 elements and H₂O: *Earth and Planetary Science Letters*, v. 131, p. 301-320.

699 Michael, P.J., and Cornell, W.C., 1998, Influence of spreading rate and magma supply on
 700 crystallization and assimilation beneath mid-ocean ridges: Evidence from chlorine
 701 and major element chemistry of mid-ocean ridge basalts: *Journal of Geophysical*
 702 *Research-Solid Earth*, v. 103, p. 18325-18356.

703 Michael, P.J., and Schilling, J.-G., 1989, Chlorine in mid-ocean ridge magmas: Evidence for
 704 assimilation of seawater-influenced components: *Geochimica et Cosmochimica Acta*,
 705 v. 53, p. 3131-3143.

706 Moore, J., 1970. Water Content of Basalt Erupted on the ocean floor. *Contributions to*
 707 *Mineralogy and Petrology* 28, 272-279.

708 Mottl, M. J., Seewald, J. S., Wheat, C. G., Tivey, M. K., Michael, P. J., Proskurowski, G.,
 709 McCollom, T. M., Reeves, E., Sharkey, J., You, C. F., Chan, L. H., and Pichler, T.,
 710 2011. Chemistry of hot springs along the Eastern Lau Spreading Center. *Geochimica*
 711 *et Cosmochimica Acta* **75**, 1013-1038.

712 Mukhopadhyay, S., 2012, Early differentiation and volatile accretion recorded in deep-mantle
 713 neon and xenon: *Nature*, v. 486, p. 101-104.
 714 Muramatsu, Y., Fehn, U., and Yoshida, S., 2001, Recycling of iodine in fore-arc areas:
 715 evidence from the iodine brines in Chiba, Japan: *Earth and Planetary Science Letters*,
 716 v. 192, p. 583-593.
 717 Muramatsu, Y., Wedepohl, K.H., 1998. The distribution of iodine in the earth's crust.
 718 *Chemical Geology* 147, 201-216.
 719 Oosting, S.E., Von Damm, K.L., 1996. Bromide/chloride fractionation in seafloor
 720 hydrothermal fluids from 9–10°N East Pacific Rise. *Earth and Planetary Science*
 721 *Letters* 144, 133-145.
 722 Parai, R., and Mukhopadhyay, S., 2012, How large is the subducted water flux? New
 723 constraints on mantle regassing rates: *Earth and Planetary Science Letters*, v. 317, p.
 724 396-406.
 725 Porcelli, D. and Wasserburg, G. J., 1995. Mass transfer of helium, neon, argon, and xenon
 726 through a steady-state upper mantle. *Geochimica et Cosmochimica Acta* **59**, 4921-
 727 4937.
 728 Ranero, C.R., Phipps Morgan, J., McIntosh, K., Reichert, C., 2003. Bending-related faulting
 729 and mantle serpentinization at the Middle America trench. *Nature* 425, 367-373.
 730 Robillard, I., Francis, D., and Ludden, J.N., 1992, The Relationship Between E-type and N-
 731 type Magmas in the Baffin-Bay Lavas: *Contributions to Mineralogy and Petrology*, v.
 732 112, p. 230-241.
 733 Roddick, J.C., 1983. High precision intercalibration of ⁴⁰Ar-³⁹Ar standards. *Geochimica et*
 734 *Cosmochimica Acta* 47, 887-898.
 735 Rüpke, L.H., Morgan, J.P., Hort, M., and Connolly, J.A.D., 2004, Serpentine and the
 736 subduction zone water cycle: *Earth and Planetary Science Letters*, v. 223, p. 17-34.

737 Sano, T., Miyoshi, M., Ingle, S., Banerjee, N. R., Ishimoto, M., and Fukuoka, T., 2008.
738 Boron and chlorine contents of upper oceanic crust: Basement samples from IODP
739 Hole 1256D. *Geochemistry, Geophysics, Geosystems* **9**, Q12O15.

740 Sarda, P., 2004. Surface noble gas recycling to the terrestrial mantle. *Earth and Planetary*
741 *Science Letters* **228**, 49-63.

742 Sarda, P., Moreira, M., and Staudacher, T., 1999. Argon-lead isotopic correlation in Mid-
743 Atlantic Ridge basalts. *Science* **283**, 666-668.

744 Schilling, J.G., Unni, C.K., Bender, M.L., 1978. Origin of Chlorine and Bromine in the
745 oceans. *Nature* **273**, 631-636.

746 Schilling, J.G., Bergeron, M.B., and Evans, R., 1980, Halogens in the mantle beneath the
747 North Atlantic: *Philosophical Transactions of the Royal Society of London Series a-*
748 *Mathematical Physical and Engineering Sciences*, v. 297, p. 147-178.

749 Schmidt, M. W. and Poli, S., 1998. Experimentally based water budgets for dehydrating slabs
750 and consequences for arc magma generation. *Earth and Planetary Science Letters*
751 **163**, 361-379.

752 Seyfried, W. E. and Ding, K., 1995. The hydrothermal chemistry of fluoride in seawater.
753 *Geochimica et Cosmochimica Acta* **59**, 1063-1071.

754 Seyfried, W.E., Jr., Seewald, J.S., Berndt, M.E., Ding, K., Foustoukos, D.I., 2003. Chemistry
755 of hydrothermal vent fluids from the Main Endeavour Field, northern Juan de Fuca
756 Ridge: Geochemical controls in the aftermath of June 1999 seismic events. *J.*
757 *Geophys. Res.* **108**, 2429.

758 Sims, K.W.W., Hart, S.R., Reagan, M.K., Blusztajn, J., Staudigel, H., Sohn, R.A., Layne,
759 G.D., Ball, L.A., Andrews, J., 2008. 238U-230Th-226Ra-210Pb-210Po, 232Th-
760 228Ra, and 235U-231Pa constraints on the ages and petrogenesis of Vailulu'u and
761 Malumalu Lavas, Samoa. *Geochemistry, Geophysics, Geosystems* **9**, Q04003.

762 Snyder, G., Savov, I.P., Muramatsu, Y., 2005. 5. Iodine and Boron in Mariana Serpentinite
 763 Mud Volcanoes (ODP legs 125 and 195): Implications for Forearc Processes and
 764 Subduction Recycling, in: Sinohara, M., Salisbury, M.H., Richter, C. (Eds.),
 765 Proceedings of the Ocean Drilling Program, Scientific Results <[http://www-
 767 odp.tamu.edu/publications/195_SR/VOLUME/CHAPTERS/102.PDF](http://www-

 766 odp.tamu.edu/publications/195_SR/VOLUME/CHAPTERS/102.PDF)>, pp. 1-18.
 768 Starkey, N.A., Stuart, F.M., Ellam, R.M., Fitton, J.G., Basu, S., and Larsen, L.M., 2009,
 769 Helium isotopes in early Iceland plume picrites: Constraints on the composition of
 770 high $^3\text{He}/^4\text{He}$ mantle: *Earth and Planetary Science Letters*, v. 277, p. 91-100.
 771 Staudacher, T., and Allègre, C.J., 1988, Recycling of oceanic crust and sediments: the noble
 772 gas subduction barrier: *Earth and Planetary Science Letters*, v. 89, p. 173-183.
 773 Staudigel, H., Hart, S.R., Koppers, A.A.P., Constable, C., Workman, R., Kurz, M., and
 774 Baker, E.T., 2004, Hydrothermal venting at Vailulu'u Seamount: The smoking end of
 775 the Samoan chain: *Geochemistry Geophysics Geosystems*, v. 5.
 776 Stroncik, N.A., and Haase, K.M., 2004, Chlorine in oceanic intraplate basalts: Constraints on
 777 mantle sources and recycling processes: *Geology*, v. 32, p. 945-948.
 778 Stuart, F.M., Lass-Evans, S., Fitton, J.G., and Ellam, R.M., 2003, High He-3/He-4 ratios in
 779 picritic basalts from Baffin Island and the role of a mixed reservoir in mantle plumes:
 780 *Nature*, v. 424, p. 57-59.
 781 Tieloff, M., Kunz, J., Clague, D.A., Harrison, D., Allegre, C.J., 2000. The nature of pristine
 782 noble gases in mantle plumes. *Science* 288, 1036-1038.
 783 Tolstikhin, I., Marty, B., Porcelli, D., and Hofmann, A., 2014. Evolution of volatile species in
 784 the earth's mantle: A view from xenology. *Geochimica et Cosmochimica Acta* **136**,
 785 229-246.
 786 Vanko, D.A., 1986. High-chlorine amphiboles from oceanic rocks: product of highly-saline
 hydrothermal fluids? *Am. Miner.* 71, 51-59

787 Vanko, D.A., 1988, Temperature, pressure, and composition of hydrothermal fluids, with
 788 their bearing on the magnitude of tectonic uplift at mid-ocean ridges, inferred from
 789 fluid inclusions in oceanic layer 3 rocks: *Journal of Geophysical Research: Solid*
 790 *Earth*, v. 93, p. 4595-4611.

791 Von Damm, K. L., Buttermore, L. G., Oosting, S. E., Bray, A. M., Fornari, D. J., Lilley, M.
 792 D., and Shanks Iii, W. C., 1997. Direct observation of the evolution of a seafloor
 793 'black smoker' from vapor to brine. *Earth and Planetary Science Letters* **149**, 101-
 794 111.

795 Wallace, P.J., 2002. Volatiles in submarine basaltic glasses from the Northern Kerguelen
 796 Plateau (ODP Site 1140): implications for source region compositions, magmatic
 797 processes and plateau subsidence. *J. Petrol.* 43, 1311-1326.

798 Walter, M. J., Kohn, S. C., Araujo, D., Bulanova, G. P., Smith, C. B., Gaillou, E., Wang, J.,
 799 Steele, A., and Shirey, S. B., 2011. Deep Mantle Cycling of Oceanic Crust: Evidence
 800 from Diamonds and Their Mineral Inclusions. *Science* 334, 54-57

801 Wanless, V.D., Perfit, M.R., Ridley, W.I., Wallace, P.J., Grimes, C.B., and Klein, E.M.,
 802 2011, Volatile abundances and oxygen isotopes in basaltic to dacitic lavas on mid-
 803 ocean ridges: The role of assimilation at spreading centers: *Chemical Geology*, v. 287,
 804 p. 54-65.

805 Wänke, H., 1981. Constitution of Terrestrial Planets [and Discussion]. *Philosophical*
 806 *Transactions of the Royal Society of London. Series A, Mathematical and Physical*
 807 *Sciences* **303**, 287-302

808 Weaver, B.L., 1991. The origin of ocean island basalt end-member compositions - trace
 809 element and isotopic constraints. *Earth and Planetary Science Letters* 104, 381-397.

810 White, W.M., and Hofmann, A.W., 1982, Sr and Nd isotope geochemistry of oceanic basalts
 811 and mantle evolution: *Nature*, v. 296, p. 821-825.

812 Workman, R.K., Hart, S.R., Jackson, M., Regelous, M., Farley, K.A., Blusztajn, J., Kurz, M.,
813 Staudigel, H., 2004. Recycled metasomatized lithosphere as the origin of the Enriched
814 Mantle II (EM2) end-member: Evidence from the Samoan Volcanic Chain. *Geochem.*
815 *Geophys. Geosyst.* 5, 4.

816 Workman, R.K., Eiler, J.M., Hart, S.R., and Jackson, M.G., 2008, Oxygen isotopes in
817 Samoan lavas: Confirmation of continent recycling: *Geology*, v. 36, p. 551-554.

818 Workman, R.K., Hauri, E., Hart, S.R., Wang, J., and Blusztajn, J., 2006, Volatile and trace
819 elements in basaltic glasses from Samoa: Implications for water distribution in the
820 mantle: *Earth and Planetary Science Letters*, v. 241, p. 932-951.

821 Woodhead, J.D., Hellstrom, J., Hergt, J.M., Greig, A., Maas, R., 2007. Isotopic and elemental
822 imaging of geological materials by laser ablation inductively coupled plasma mass
823 spectrometry. *Geostandards and Geoanalytical Research* 31, 331-343.

824 You, C.F., Butterfield, D.A., Spivack, A.J., Gieskes, J.M., Gamo, T., Campbell, A.J., 1994.
825 Boron and halide systematics in submarine hydrothermal systems: Effects of phase
826 separation and sedimentary contributions. *Earth and Planetary Science Letters* 123,
827 227-238.

828 Zindler, A., Hart, S., 1986. Chemical Geodynamics. *Annual Review of Earth and Planetary*
829 *Sciences* 14, 493-571.

830

Table 1. Summary of measured and assimilation-corrected volatile data for Samoan and Baffin samples

Volcano/Location	<i>Vailulu'u</i>								<i>Ta'u</i>		<i>Malumalu</i>	
Sample number	68-03	70-01	71-02	71-11	71-13	71-22	73-03	73-12	74-02	75-02	75-10	76-03
Depth (bars)	100-60	150-70	440-400	440-400	440-400	440-400	100-80	100-80	250-190	270-190	270-190	280-200
<i>Selected volatiles¹</i>												
CO ₂ ppm	23	7	179	150	150	200	13	12	46	21	35	52
H ₂ O wt %	0.67	1.0	1.2	1.3	1.1	1.2	0.71	0.81	1.4	1.5	1.2	1.3
F ppm	930	920	900	920	880	1030	1000	980	1870	1370	1230	1180
Cl ppm	1570	1010	1790	1610	1520	1530	1600	1360	870	720	650	1610
Br ppb	6290	3830	7230	5480	6510	6580	6600	5420	2480	2120	1890	5630
I ppb	59	47	82	56	74	71	75	59	41	46	40	72
<i>Selected major and trace elements</i>												
MgO wt %	6.0	6.3	6.5	6.2	6.2	5.5	5.9	6.0	4.0	4.9	5.2	5.5
K wt %	0.98	0.81	1.06	0.99	0.97	0.84	1.30	1.07	1.37	1.13	1.07	1.56
Ce ppm	91	72	96	88	86	90	97	109	161	112	110	123
<i>Assimilation-correction model (2σ uncertainties)²</i>												
F/Cl *s	0.61 ± 0.7	1.0 ± 0.1	0.60 ± 0.07	0.61 ± 0.07	0.64 ± 0.07	0.61 ± 0.07	0.91 ± 0.10	0.89 ± 0.10	1.9 ± 0.2	1.9 ± 0.2	2.3 ± 0.3	1.1 ± 0.1
Br/Cl ×10 ⁻³												
Cl _{assim} ppm	1170 ± 90	590 ± 100	1350 ± 110	1210 ± 100	1120 ± 100	1150 ± 90	1000 ± 140	860 ± 120	180 ± 160	160 ± 130	20 ± 150	870 ± 180
Cl _{mantle} ppm	390 ± 90	420 ± 100	450 ± 110	400 ± 100	400 ± 100	380 ± 90	600 ± 140	500 ± 120	690 ± 160	560 ± 130	630 ± 150	740 ± 180
H ₂ O _{assim} wt. %	0.17 ± 0.08	0.09 ± 0.04	0.20 ± 0.09	0.18 ± 0.08	0.17 ± 0.08	0.17 ± 0.07	0.15 ± 0.07	0.13 ± 0.06	0.03 ± 0.03	0.02 ± 0.02	0.00 ± 0.02	0.13 ± 0.06
H ₂ O _{mantle} wt. %	0.50 ± 0.08	0.91 ± 0.04	0.98 ± 0.09	1.12 ± 0.08	0.93 ± 0.08	1.03 ± 0.08	0.56 ± 0.07	0.68 ± 0.06	1.37 ± 0.03	1.48 ± 0.02	1.20 ± 0.02	1.17 ± 0.06
Brine wt. %	0.39 ± 0.11	0.19 ± 0.06	0.45 ± 0.13	0.40 ± 0.11	0.37 ± 0.11	0.38 ± 0.11	0.33 ± 0.10	0.28 ± 0.09	0.06 ± 0.06	0.05 ± 0.05	0.01 ± 0.05	0.29 ± 0.10
% assim. Cl	75 ± 6	59 ± 10	75 ± 6	75 ± 6	74 ± 6	75 ± 6	62 ± 9	63 ± 9	21 ± 19	22 ± 18	3 ± 23	54 ± 11
% assim H ₂ O	26 ± 11	9 ± 4	17 ± 7	14 ± 6	15 ± 7	14 ± 6	21 ± 9	16 ± 7	2 ± 2	2 ± 1	0 ± 2	10 ± 5
Cl/K _{measured}	0.16	0.12	0.17	0.16	0.16	0.18	0.12	0.13	0.06	0.06	0.06	0.10
Cl/K _{corrected}	0.04 ± 0.01	0.05 ± 0.01	0.04 ± 0.01	0.04 ± 0.01	0.04 ± 0.01	0.05 ± 0.01	0.05 ± 0.01	0.05 ± 0.01	0.05 ± 0.01	0.05 ± 0.01	0.06 ± 0.01	0.05 ± 0.01
H ₂ O/Ce _{measured}	74	139	123	148	128	133	73	74	87	134	109	106
H ₂ O/Ce _{corrected}	55 ± 8	127 ± 6	102 ± 9	127 ± 9	109 ± 9	114 ± 8	58 ± 7	63 ± 5	85 ± 2	132 ± 2	109 ± 2	95 ± 5
<i>Selected isotopes¹</i>												
⁸⁷ Sr/ ⁸⁶ Sr	0.70540	0.70537	0.70594	0.70539		0.70547	0.70562	0.70665		0.70452	0.70453	0.70641
³ He/ ⁴ He (R/Ra)	10.0		9.5	9.9		9.6				15.1		

1 – CO₂, H₂O and F data by SIMS, Cl, Br and I data by the noble gas method (see Table S1 for the full dataset). Legacy data are shown in bold italics are from Workman et al. (2004; 2006); Jackson et al.(2007; 2010); Kent et al. (2004); the ³He/⁴He ratios measured by crushing are reported as R/Ra where Ra is the atmospheric ³He/⁴He value of 1.39×10⁻⁶.

2 - The model parameters are derived from the measured H₂O, F, Cl and Br together with the following formula:

% assim Cl = (X/Cl_{meas}-X/Cl_{mantle})/(X/Cl_{brine}-X/Cl_{mantle})×100, where X = Br or F; Br/Cl_{Mantle} = (2.8 ± 0.6)×10⁻³ (Kendrick et al., 2013a); Br/Cl_{brine} = (4.6 ± 0.4)×10⁻³(Fig 6c); F/Cl_{Mantle} = 2.4 ± 0.5; and F/Cl_{brine} = 0.01 ± 0.00999. H₂O_{assim} = Cl_{assim}/brine salinity/0.55, where brine salinity = 55 ± 15 wt. % salts and salts have seawater composition with 55 wt. % Cl.

Table 1. Continued

Volcano/Location	Malumalu						Muli			Taumatau	Baffin Bay	
Sample number	76-08-a	76-08-b	76-11	77-09	78-01-a	78-01-b	79-04	104-04 glass	104-04 palagonite	128-21	100-24	100-21
Depth (bars)	280-200	280-200	280-200	360-290	230-220	230-220	350-280	250	250	260	Submarine	Submarine
<i>Selected volatiles¹</i>												
CO ₂ ppm	37	As for a	59	108	nd	nd	nd	77	nd	4	nd	nd
H ₂ O wt %	1.2	As for a	1.2	1.9	nd	nd	nd	1.7	nd	1.2	nd	nd
F ppm	1240	As for a	1230	1600	nd	nd	nd	1030	nd	1270	nd	nd
Cl ppm	1370	1420	1620	1070	960	490	1060	1090	35	1050	79	67
Br ppb	4890	5200	6220	3550	3470	1920	2010	3810	200	3000	240	130
I ppb	69	62	67	70	83	133	29	19	85	93	6	3
<i>Selected major and trace elements</i>												
MgO wt %	5.6	As for a	4.9	5.0	4.7	As for a	4.2	6.2	nd	6.0	8.4	8.6
K wt %	1.34	1.39	1.40	1.49	1.73	0.92	1.61	0.62	0.09	1.60	0.10	0.14
Ce ppm	116	As for a	120	144	186	As for a	324	91	nd	135	13	12
<i>Assimilation-correction model (2σ uncertainties)²</i>												
F/Cl	0.89 ± 0.10		0.76 ± 0.08	1.7 ± 0.2				0.99 ± 0.11		1.5 ± 0.2		
Br/Cl×10 ⁻³		3.7 ± 0.1			3.6 ± 0.1	3.9 ± 0.3	1.90 ± 0.05				2.99 ± 0.05	1.92 ± 0.04
Cl _{assim} ppm	870 ± 120	690 ± 440	1110 ± 120	320 ± 180	440 ± 300	300 ± 160		640 ± 110		380 ± 160	8 ± 30	
Cl _{mantle} ppm	500 ± 120	730 ± 440	510 ± 120	750 ± 180	510 ± 300	190 ± 160		450 ± 110		670 ± 160	70 ± 30	
H ₂ O _{assim} wt. %	0.13 ± 0.06	0.10 ± 0.08	0.17 ± 0.07	0.05 ± 0.03	0.07 ± 0.05	0.04 ± 0.03		0.10 ± 0.04		0.06 ± 0.03	<0.01	
H ₂ O _{mantle} wt. %	1.07 ± 0.06	1.10 ± 0.08	1.03 ± 0.07	1.85 ± 0.03				1.60 ± 0.04		1.14 ± 0.03		
Brine wt. %	0.29 ± 0.09	0.23 ± 0.16	0.37 ± 0.11	0.11 ± 0.07	0.15 ± 0.11	0.10 ± 0.06		0.21 ± 0.07		0.13 ± 0.06	0.00 ± 0.00	
% assim. Cl	63 ± 9	48 ± 31	69 ± 7	30 ± 17	46 ± 32	61 ± 32		59 ± 10		36 ± 15		
% assim H ₂ O	11 ± 5	9 ± 7	14 ± 6	3 ± 2				6 ± 3		5 ± 3		
Cl/K _{measured}	0.10	0.10	0.12	0.07	0.06	0.05		0.17		0.07	0.08	
Cl/K _{corrected}	0.04 ± 0.01	0.05 ± 0.03	0.04 ± 0.01	0.05 ± 0.01	0.03 ± 0.02	0.02 ± 0.02		0.07 ± 0.02		0.04 ± 0.01	0.07 ± 0.03	
H ₂ O/Ce _{measured}	103	103	100	132				186		89		
H ₂ O/Ce _{corrected}	92 ± 5	95 ± 7	86 ± 6	129 ± 2				176 ± 5		83 ± 3		
<i>Selected isotopes¹</i>												
⁸⁷ Sr/ ⁸⁶ Sr	0.70637	As for a	0.70720	0.70726	0.70890	As for a	0.70490	0.70483		0.71250	0.70322	0.70381
³ He/ ⁴ He (R/Ra)				10.6	8.1	As for a		15.3			23.8	9.0

Table 2. A possible Cl budget favouring net Cl subduction

	Thickness km	Rock 10 ⁹ kg a ⁻¹	Cl ² ppm	Cl 10 ⁹ kg a ⁻¹
<i>Input</i> ¹				
Sediments	0.5	4	600	2
Altered Ocean Crust (AOC)	6	52	150	8
Serpentinite (20 %)	3	4	1200	5
40 % lost during arc magmatism*, supply to arcs (R) ³ =				6
Net input to deeper mantle =				9
<i>Output</i>				
Mid-ocean ridge (MORB)	~6	48	100	5
Oceanic Island (OIB)	-	4	300	1
Volcanic Arc (M + R) ³	-	6	1200	7
Output from mantle through arc (M) =				1
Net output from mantle =				7

Notes: 1) AOC (kg) = MORB + OIB (kg); (cf. Schilling et al., 1978; Ito et al., 1983); serpentinites and sediments (kg) are scaled to AOC based on relative thicknesses; 2) Cl concentrations assume AOC>MORB; Ito et al. 1983; Sharp and Barnes, 2004; Bonifacie et al., 2008; Sano et al., 2008; Barnes and Cisneros, 2012; Kendrick et al., 2013b; 3) Recycled (R) output through volcanic arc is related to input and Cl loss efficiency.

Fig 1. Kendrick et al. (2014)

Fig 1. $^{87}\text{Sr}/^{86}\text{Sr}$ versus $^{206}\text{Pb}/^{204}\text{Pb}$ data for the Samoan and Baffin glasses used in this study and glasses used in previous halogen studies (Kendrick et al., 2012ab; 2013a, 2014a). The mantle components are: EM1= enriched mantle 1; EM2 = enriched mantle 2; Himu = high U/Pb; Fozo = focus zone; and DMM = depleted MORB mantle (e.g. Hofmann, 2003). Helium data are shown in Fig S1.

Fig 2. Kendrick et al. (2014)

Fig 2. Sketch locality map of the Samoan Islands showing the volcanoes from which submarine glasses were selected for this study in bold. Emergent Islands are shaded grey and have uppercase labels, whereas seamounts have lower case labels. Vailulu'u is the youngest volcano in the Samoan chain, ages increase westward to a maximum of 4.8 Ma for sample 128-21 dredged from Taumatau (Koppers et al., 2008).

Fig 3 (Kendrick et al., 2014)

Fig 3. K and halogen concentration plots: a) K versus Cl; b) Br versus Cl; c) I versus Cl; and d) F versus Cl. Note that slopes of K/Cl, Br/Cl and I/Cl given for reference encompass values typical of the mantle (shaded areas; see Kendrick et al., 2012b; 2013a). The mantle range of F/Cl is not well known but it extends up to at least 7 (e.g. Le Roux et al., 2006).

Fig 4. Kendrick et al. (2014)

Fig 4. The volatile content of Samoan glasses: a) H_2O wt % versus CO_2 ppm; and b) dredge depth versus equilibration depth in metres (1 bar = 10 metres) calculated using the VolatileCalc solubility model (Newman and Lowenstern, 2002). The isobars (200 and 400 bars), isopleths of vapours comprising 20, 50 and 90 mol % H_2O and examples of open and closed system degassing paths were calculated for a basalt melt with 48wt % SiO_2 at 1200 °C. The sample equilibration depths are based on the measured H_2O , CO_2 , SiO_2 and a temperature of 1200 °C. Most of the data are from Workman et al. (2006) with the new data for samples 77-09, 104-04 and 128-21 identified by circles.

Fig 5. K, Cl and H₂O versus MgO concentration plots: a) K versus MgO; b) Cl versus MgO, and c) H₂O versus MgO. Liquid lines of descent showing the progress of fractional crystallisation in 10% increments are shown for variably enriched melts (dotted and dashed lines). The initial K concentrations were chosen so that the LLD bracket the data; the initial Cl and H₂O concentrations then depend on the K/Cl and K/H₂O ratios of the melts given in b and c.

Fig 6. Mixing systematics of F, Cl, Br, I, H₂O and K in Samoan glasses: a) F/Cl versus H₂O/Cl; b) K/Cl versus H₂O/Cl; c) Br/Cl versus H₂O/Cl; d) K/Cl versus Br/Cl; e) I/Cl versus H₂O/Cl; and f) I/Cl versus Br/Cl. The composition of seawater, brines with 5, 10, 20, 30 and 50 wt. % salt and a field suggested to be representative of the Samoan mantle are shown for reference. The composition of the brine end-member obtained with model 2 regressions in Isoplot (Ludwig, 2009) shows 2σ uncertainties. Note that parts d and e show data for selective regressions that exclude the circled data points, and data for non-selective regressions in parentheses. Note that the mixing lines show 10% increments of total Cl between the mantle and brine end-members.

Fig 7. Kendrick et al. (2014)

Fig 7. Assimilation-corrected $\text{H}_2\text{O}/\text{Ce}$ (a) and Cl/K (b), and the measured F/Nd (c) of Samoan melts as a function of $^{87}\text{Sr}/^{86}\text{Sr}$. Uncertainty introduced by the assimilation correction controlled by the model fit in Fig 6 is shown in parts a and b (see Table 1). Data for Society and Pitcairn glasses (Kendrick et al., 2014a), and MORB fields are shown for reference.

Fig 8. Kendrick et al. (2014)

Fig 8. The Br/Cl versus I/Cl systematics of the five Samoan glasses least affected by brine assimilation and a high $^3\text{He}/^4\text{He}$ Baffin Bay glass, showing mantle has limited variation in Br/Cl and I/Cl that is not related to $^{87}\text{Sr}/^{86}\text{Sr}$ or $^3\text{He}/^4\text{He}$. Data for glasses from Society and Pitcairn (Kendrick et al., 2014a), various MORB glasses (Kendrick et al., 2013a), seawater and the brine used in our mixing model (Fig 6), are shown for reference.