

ORIGINAL ARTICLE

Plants originating from more extreme biomes have improved leaf thermoregulation

Pieter A. Arnold^{1,*}, Monique J. White¹, Alicia M. Cook², Andy Leigh², Verónica F. Briceño¹ and Adrienne B. Nicotra¹

¹Division of Ecology and Evolution, Research School of Biology, The Australian National University, Canberra, ACT 2600, Australia and ²University of Technology Sydney, School of Life Sciences, Sydney, NSW 2007, Australia

*For correspondence. E-mail pieter.arnold@anu.edu.au

Received: 22 January 2025 Returned for revision: 7 April 2025 Editorial decision: 23 April 2025 Accepted: 26 April 2025

• **Background and Aims** Many plants have some capacity for leaf thermoregulation via stomatal conductance (g_{sw}), such that leaf temperature (T_{leaf}) is rarely coupled with air temperature (T_{air}). The difference between leaf and air temperature (thermal offset, ΔT) and the slope (thermal coupling strength, β) is mediated by interactions between the immediate environment of the plant and its leaf traits. The aim of this study was to determine whether species originating from biomes with contrasting environmental conditions (alpine, desert and coastal temperate) would differ in their tendency to thermoregulate in a common environment.

• **Methods** Using benign-temperature (25 °C) and high-temperature (38 °C) glasshouse treatments, we measured paired canopy T_{air} and T_{leaf} for 15 diverse species, 5 from each biome, in a common garden experiment. Instantaneous stomatal conductance and a suite of leaf traits were measured and calculated to test for associations with leaf thermoregulation.

• **Key Results** We found clear evidence for greater leaf cooling occurring during high-temperature exposure, especially in alpine and desert species. The leaves of temperate species were largely warmer than air in both treatments. Thicker leaves with higher water content and high stomatal conductance clearly were more effective at cooling. Species originating from different biomes displayed divergent responses of thermal offset and thermal coupling with leaf traits.

• **Conclusions** Our findings suggest that plants originating from more extreme biomes have innately greater scope for thermoregulation, especially desert plants, which could better counter the risk of reaching excess temperatures at the cost of higher water loss. Leaf thermoregulation is a complex plant–environment interaction, and our work contributes to the development of more accurate predictions of leaf temperature during heat exposure across diverse species and biomes.

Key words: Alpine, climate warming, desert, heatwave, leaf temperature, limited homeothermy, stomatal conductance, temperate, thermal coupling, thermal offset, thermal sensitivity, thermoregulation.

INTRODUCTION

Extreme climatic events are major contemporary challenges to terrestrial plants (Perkins-Kirkpatrick *et al.*, 2024). Pulse events that include periods of extremely high temperatures, such as heatwaves, are increasing in frequency, intensity and duration in Australia and are expected to worsen in future decades (Cowan *et al.*, 2014; Perkins-Kirkpatrick and Lewis, 2020). Against the backdrop of accelerated climate warming, heat pulses will expose plants to acute high temperatures that far exceed their typical range (Harris *et al.*, 2018). High temperature affects many physiological and biochemical processes in plants, potentially inflicting injury to tissues and membranes that maintain homeostasis (Goraya *et al.*, 2017). Plants have therefore developed an arsenal of mechanisms to help avoid, tolerate or acclimatize to high temperature to reduce the impact

of heat on plant function (Goraya *et al.*, 2017; Nievola *et al.*, 2017; Deva *et al.*, 2020; Geange *et al.*, 2021).

Leaf temperature (T_{leaf}) is central to the maintenance of photosynthetic performance and metabolic homeostasis (Gates, 1968; Jones, 2014). It is now well established that plants are not necessarily poikilotherms that conform to air temperatures (T_{air}) of their environment (Mahan and Upchurch, 1988; Michaletz *et al.*, 2015), which is apparent from individual leaves (Tserej and Feeley, 2021) to ecosystem canopies (Guo *et al.*, 2023). T_{leaf} can decouple markedly from T_{air} in a range of environmental conditions but is typically exacerbated during periods with high sun exposure and low wind and during heat pulses and heatwaves (Leigh *et al.*, 2012, 2017; Hüve *et al.*, 2019; Slot *et al.*, 2021; Kitudom *et al.*, 2022; Kullberg *et al.*, 2023; Manzi *et al.*, 2024). Leaves are often warmer than air when T_{air} is cold and there is sufficient insolation, whereas leaves can be cooler than air

when T_{air} is warm and water is available to the plant for transpiration (Michaletz *et al.*, 2015), i.e. plants can exhibit limited homeothermy.

The limited homeothermy hypothesis posits that plants can maintain an operative temperature by reducing T_{leaf} through active transpiration (Mahan and Upchurch, 1988). Decoupling of T_{leaf} from T_{air} occurs owing to structural properties of the leaf and thermoregulatory behaviour (Michaletz *et al.*, 2015; Lin *et al.*, 2017; Tserej and Feeley, 2021). Mechanisms of thermoregulation in plants can be described simply as either passive or active via structural and physiological means (Drake, 2023). Intrinsic leaf structural traits allow plants to thermoregulate passively (e.g. leaf lamina area or width; Leigh *et al.*, 2017) and avoid rapid excursions to temperature extremes by slowing heat transfer (e.g. leaf thickness and water content; Vogel, 2009; Leigh *et al.*, 2012). Differences in leaf structural traits are driven by differences in biomes or environmental conditions (Gibson, 1998; Lusk *et al.*, 2018); in a common environment, leaf trait differences might be less pronounced among species (Reich *et al.*, 2003). In contrast to passive influences of leaf structural traits, plants can thermoregulate actively by dynamically adjusting stomatal conductance of water vapour (hereafter, g_{sw}) (Michaletz *et al.*, 2015; Guo *et al.*, 2022). Stomata can be regulated finely between closed and fully open states to optimize gas exchange and water loss and to regulate T_{leaf} in the absence of photosynthesis (Gates, 1968; Matthews *et al.*, 2017; Drake *et al.*, 2018).

During drought stress, species differ in their stomatal behaviour and water-management strategies along a continuum from avoidance (stomatal closure to limit water loss) to tolerance (stomatal opening, which allows dehydration), sometimes referred to as isohydric and anisohydric (Klein, 2014; but see Hochberg *et al.*, 2018). In high-temperature conditions, regulation of T_{leaf} depends strongly on active evaporative cooling via transpiration (Drake *et al.*, 2018; Marchin *et al.*, 2022). During a record heatwave that exceeded 48 °C, Posch *et al.* (2024) found dynamic patterns of T_{leaf} during a common garden experiment. T_{leaf} was typically lower than T_{air} when water was readily available, which enabled g_{sw} to be relatively high. Thereafter, a water-stress treatment applied during extreme heat led to T_{leaf} exceeding T_{air} (disrupting homeothermy) when water availability was low and g_{sw} was near zero (Posch *et al.*, 2024). If high temperatures coincide with water limitation, many plant species are unable to transpire to dissipate heat, hence leaves can reach damaging temperatures (Cook *et al.*, 2021; Marchin *et al.*, 2022; Posch *et al.*, 2024). In contrast, other species have recently been observed to maintain partly open stomata in high temperatures, even in droughted plants (Marchin *et al.*, 2022).

There is a clear trade-off between water use and active thermoregulation (Fauset *et al.*, 2018). However, species that originate from distinct biomes and/or that have different leaf traits will differ in their thresholds for when and how much stomata are opened based on their relative position on the avoidance–tolerance spectrum (Marchin *et al.*, 2022). Leaf thermoregulation therefore involves more than the biophysical effects of structural leaf traits; stomatal strategy makes a substantive difference to leaf temperature. We therefore expect that species originating from contrasting environments would have developed divergent leaf thermoregulation tendencies or different thermal coupling responses (Blonder and Michaletz,

2018). Cooling via stomatal behaviour can be more effective than the mediating effects of passive leaf traits when sufficient water is available (Lin *et al.*, 2017), although both contribute to thermoregulation strategy.

Two simple temperature metrics encapsulate $T_{\text{leaf}}-T_{\text{air}}$ coupling relationships. The thermal offset (ΔT) describes the magnitude of difference between T_{leaf} and T_{air} , and the thermal coupling strength (β) describes the slope of the relationship between T_{leaf} and T_{air} (Blonder and Michaletz, 2018; Blonder *et al.*, 2020). In nature, leaf thermal offsets can exceed ± 15 °C (Salisbury and Spomer, 1964; Leuzinger and Körner, 2007; Blonder and Michaletz, 2018; Fauset *et al.*, 2018). Thermal coupling strength classifies plant thermoregulatory state into three categories: poikilothermy ($\beta \approx 1$), limited homeothermy ($\beta < 1$) and megathermy ($\beta > 1$) (Blonder *et al.*, 2020; Cavaleri, 2020). Blonder *et al.* (2020) demonstrated that both ΔT and β can differ with environment across a range of T_{air} values in plant species from contrasting North American biomes. Specifically, at cool T_{air} , species from temperate forests and meadows exhibit limited homeothermy [they have T_{leaf} warmer than T_{air} (negative ΔT)], but at warm T_{air} , T_{leaf} is cooler than T_{air} and $\beta < 1$. In contrast, those from subalpine meadows were often poikilothermic, but sometimes exhibited megathermy with positive ΔT when T_{air} was high. High desert species were more variable but frequently exhibited megathermy with generally large positive ΔT , especially when T_{air} was high.

Plants from hot, arid environments, such as deserts, are frequently exposed to very high T_{air} and may not have water available to transpire freely to reduce T_{leaf} (Cook *et al.*, 2021), such that many desert plants tolerate rather than avoid high T_{leaf} (Curtis *et al.*, 2016). A common adaptation in desert plants is small leaf area to minimize overheating, reduce transpiration and increase water-use efficiency, but some large-leafed desert plants can maintain much higher transpiration rates and relatively low T_{leaf} (Smith, 1978). Many leaf traits contribute to mediating large thermal offsets (Guo *et al.*, 2022). For example, in tropical plants, T_{leaf} readily exceeds T_{air} (Manzi *et al.*, 2024); however, structural leaf traits are not necessarily individually related to ΔT . For example, in tropical shrubs and herbs, no relationship was found between ΔT and leaf area, leaf mass per area or leaf thickness (Pedraza, 2024). Data from dry temperate and tropical trees support the idea that transpirational cooling can be a strategy used to improve net carbon gain by avoiding leaf mortality or by maintaining temperature homeostasis near the optimal temperatures for photosynthesis (Slot and Winter, 2017; Drake *et al.*, 2018). Alpine plants tend to have strategies that aim to retain heat, because their environment is typically limited by cold temperatures, and T_{leaf} can exceed T_{air} by 15 °C or more, especially in short-statured plants (Salisbury and Spomer, 1964). Thus, high temperatures that occur during heatwaves and extremely hot days will result in unequal thermal exposure among different plant species, especially those with different thermoregulation strategies.

Determining the drivers of variation in thermal coupling in high-temperature conditions should therefore be a priority for understanding impacts to plant performance in the context of global change. Although theoretical predictions of how leaf thermoregulation should vary with environments have been established for decades, empirical studies addressing this question are rare. A recent field study along a temperature and

TABLE 1. List of the 15 species studied, including their biome of origin, taxonomic family, general growth form and origin of plant material used in the experiment.

Species	Biome	Family	Growth form	Plant material
<i>Eucalyptus pauciflora</i>	Alpine	Myrtaceae	Tree	Nursery
<i>Leptorhynchos squamatus</i>	Alpine	Asteraceae	Forb	Nursery
<i>Oxylobium ellipticum</i>	Alpine	Fabaceae	Shrub	Seedbank
<i>Ranunculus graniticola</i>	Alpine	Ranunculaceae	Forb	Nursery
<i>Xerochrysum subundulatum</i>	Alpine	Asteraceae	Forb	Nursery
<i>Acacia binervata</i>	Temperate	Fabaceae	Tree–shrub	Seedbank
<i>Acacia longifolia</i>	Temperate	Fabaceae	Tree–shrub	Seedbank
<i>Backhousia myrtifolia</i>	Temperate	Myrtaceae	Tree	Seedbank
<i>Melaleuca hypericifolia</i>	Temperate	Myrtaceae	Tree–shrub	Nursery
<i>Pittosporum undulatum</i>	Temperate	Pittosporaceae	Tree–shrub	Nursery
<i>Acacia aneura</i>	Desert	Fabaceae	Tree–shrub	Seedbank
<i>Acacia salicina</i>	Desert	Fabaceae	Tree–shrub	Seedbank
<i>Dodonaea viscosa</i>	Desert	Sapindaceae	Shrub	Seedbank
<i>Eucalyptus largiflorens</i>	Desert	Myrtaceae	Tree	Seedbank
<i>Flindersia maculosa</i>	Desert	Rutaceae	Tree	Seedbank

precipitation gradient showed that plants from hotter sites showed greater transpirational cooling and that physical leaf traits were important for maintaining thermoregulation (Zhou *et al.*, 2023). To our knowledge, there have not been empirical studies in controlled environments that explore how common-grown species adapted to very different biomes vary in their leaf thermodynamic properties, and the structural or physiological drivers of leaf thermoregulation.

Our overarching goal was to determine how leaf characteristics facilitate or constrain leaf thermoregulation via thermal coupling. Here, we determined ΔT and β in 15 plant species, 5 from each of three contrasting biomes (alpine, desert and coastal temperate) in benign and high air temperatures in a controlled-environment glasshouse experiment. We then tested whether leaf structural traits and stomatal conductance were associated with leaf thermoregulation. We hypothesized that species originating from biomes with more extreme climates (alpine and desert) would have greater thermoregulatory capacity than those from more benign climates (coastal temperate). This difference would reflect varying combinations of leaf traits with stomatal strategy. We expected that plants with relatively small and less succulent leaves (i.e. low water content, thinner) might be closer to T_{air} and that plants with conservative (i.e. lower and/or less dynamic) g_{sw} would be most limited in their ability to thermoregulate. Assessment of the proximal causes of variation in plant thermoregulation in diverse species in controlled conditions will contribute to an improvement in our understanding of plant thermal sensitivity and vulnerability during heat extremes in nature.

MATERIALS AND METHODS

Information on species, growth conditions and origin biome

Five native Australian plant species that each originated from one of three contrasting biomes were chosen to be grown in

common conditions in glasshouses at The Australian National University, Canberra, ACT, Australia. The 15 species cover seven families and four growth forms (Table 1). A simple phylogenetic tree of the study species is shown in Supplementary Data Fig. S1.

Plants used in the experiment were germinated between August and December 2020 from seed accessions obtained from the Australian National Botanic Gardens Seed Bank and the Australian Botanic Gardens Australian PlantBank. Seed accessions were collected originally within a 50 km radius within three distinct biomes (temperate: Wollongong, NSW; alpine: Kosciuszko National Park, NSW; and desert: Bourke, NSW) and were stored in these facilities for <20 years. Mean climatic parameters of these origin biomes are provided in Table 2. Some species had poor seed germination rates and were purchased as seedlings from Monaro Native Tree Nursery, NSW and Bodalla Nursery, NSW at ~3 months old, which were then acclimated and grown in the same conditions as plants grown from seed (Table 1). Additional information is provided by Harris *et al.* (2024). The plants were grown in common garden well-watered conditions (watered to field capacity daily) in shade houses. Plants were transplanted in August 2021 to large pots (150–200 mm in diameter and ≥ 200 mm in depth) based on their individual size. The plants had grown for ~12–18 months before being moved to glasshouse conditions for this experiment in January–February 2022 (Austral summer) and ranged in size from 0.15 to 1.5 m in height at the time of the experiment. We used five replicate plants of each species for the temperature experiment. The plants were watered to saturation in the morning, before applying the temperature treatments to plants in controlled glasshouse rooms from 12.00 to 15.00 h, where the initial 30-min period from 12.00 to 12.30 h was considered temperature equilibration time. Plants did not show visual signs of water stress (i.e. they did not run out of water during the treatment phase) and were re-watered after the treatments.

TABLE 2. Environmental conditions of biomes of origin based on averages of downsampled long-term (1981–2010) climate data from CHELSA v2.1 database (Karger et al., 2017) using field locations for these alpine, temperate and desert biomes (Briceño et al., 2024).

Biome	MAT (°C)	MinT (°C)	MaxT (°C)	T_{range} (°C)	MAP (mm)
Alpine	4.5	−5.2	16.5	22.7	1764
Temperate	16.5	7.4	24.6	17.2	1285
Desert	20.2	4.5	36.0	31.5	332

Abbreviations: MAP, mean annual precipitation; MAT, mean annual temperature; MaxT, mean maximum temperature of the warmest month; MinT, mean minimum temperature of the coldest month; T_{range} , MaxT – MinT).

Temperature treatments

Two temperature treatments referred to as ‘benign’ and ‘high temperature’ were applied sequentially using controlled-temperature glasshouse rooms. The benign glasshouse room was set to 25 °C (06.00–20.00 h) during the day, and the high-temperature glasshouse room was set to 38 °C. The high temperature of 38 °C was chosen as a temperature that would be sufficiently stressful, but not lethal, for all species (Harris et al., 2024). Both treatment glasshouses were set to and 16 °C overnight (20.00–06.00 h). All plants ($n = 75$) were moved from their shade house to the benign room 14 days before the experiment began, to allow for acclimation to the higher-light environment. Preliminary tests of high-temperature treatment duration effects on plant temperatures showed that ΔT (calculated as $T_{\text{leaf}} - T_{\text{air}}$) of ten test plants averaged over 2.5 h was not different from longer periods of 4 or 6 h of high-temperature exposure, hence the 2.5 h duration (i.e. 12.30–15.00 h) was used. The experiment was conducted over six separate days (three for each treatment), where 30 plants were measured at a time. T_{air} at canopy level averaged across each of the plants during the treatments over 2.5 h was ~23.2 °C in the benign treatment and 35.7 °C in the high-temperature treatment (Fig. 1). Glasshouse conditions during the treatments were as follows for benign: temperature (T_{gh}) = 26.3 ± 0.6 °C, relative humidity (RH) = 30.5 ± 4.5 % and vapour pressure deficit (VPD_{air}) = 2.4 ± 0.5 kPa; and for high temperature: T_{gh} = 38.5 ± 0.4 °C, RH = 23.8 ± 3.4 % and VPD_{air} = 6.3 ± 0.3 kPa (full details are in Supplementary Data Table S1).

Leaf temperature measurements

Leaf temperature (T_{leaf}) measurements were taken using data loggers (Onset HOBO UX120-014M; Onset Computer Co., Bourne, MA, USA) and type-T thermocouples on mature, fully expanded, sun leaves emerging from the main stem or central part of each plant. Each thermocouple for measuring T_{leaf} was attached firmly to the underside of a leaf using porous surgical tape, and the thermocouple wire was supported by malleable wire on the stem to hold it in position without altering the natural leaf position. A second thermocouple was also anchored to the main stem of each plant, with the thermocouple tip open to air shielded from direct sunlight. This pairing enabled us to measure T_{air} immediately adjacent to the thermocouple measuring T_{leaf} . The data loggers recorded temperature at 1-min intervals from 12.00 to 15.00 h.

Plants that were wired for temperature measurements in benign conditions were then transferred to the high-temperature conditions 2–4 days later, at ~11.00 h. Wherever possible, after the benign treatment the thermocouples were left in position, meaning that the T_{leaf} and T_{air} measurements were taken from the same location in both treatments. If a leaf began to discolour or if the thermocouple detached and could not be reattached easily, the thermocouple was moved to the nearest healthy, mature leaf to capture a similar microclimate. Logged measurements were trimmed to above 16 °C for the benign treatment and 31 °C for the high-temperature treatment, to exclude data when glasshouse evaporative coolers were active, because air circulation patterns during the active heating/cooling cycles introduced high variance and did not address our scientific questions (~10 % of the data; Fig. 1A, B). We calculated the thermal offset (ΔT), as $T_{\text{leaf}} - T_{\text{air}}$ (in degrees Celsius) between 12.30 and 15.00 h to allow for temperatures to equilibrate. Negative values of ΔT occur when leaves are cooler than air and positive values of ΔT occur when leaves are warmer than air (Fig. 1C, D). We also calculated thermal coupling strength (β) as the slope of the relationship between T_{leaf} and T_{air} at 30-min intervals, following Blonder et al. (2020). Mean temperature responses per species are shown in Supplementary Data Table S2.

Stomatal conductance

The stomatal conductance to water (g_{sw} ; in moles per metre squared per second) of light-adapted leaves was measured using a porometer–fluorometer (LI-600; LI-COR Biosciences, Lincoln, NE, USA). Transpiration (E) correlated strongly with g_{sw} (Pearson’s $r = 0.90$); therefore, we report only the g_{sw} results. The g_{sw} was measured on the same leaf that had the T_{leaf} thermocouple attached wherever possible, and species with small or compact leaves had g_{sw} measured on the closest mature, similar leaf. There were 18 (of 75) plants for which g_{sw} could not be measured owing to small leaf size; therefore, there were $n = 57$ plants in each temperature treatment for which there were a complete set of leaf traits for principal components analysis. Measurements of g_{sw} were taken twice between 13.30 and 14.30 h, after the plants had been exposed to the treatments for ≥ 1.5 h, and the average of both measurements was used.

Leaf structural traits

After completing T_{leaf} and T_{air} measurements for both benign and high-temperature treatments, the same leaves that were measured for temperature were excised carefully from the plant to measure structural traits. Leaf wet mass (in milligrams) was measured with a precision balance (ML203T; Mettler-Toledo, Columbus, OH, USA), then leaf area (LA; in centimetres squared) using the *leafscan* app (Anderson and Rosas-Anderson, 2017), and leaf width (LW; in millimetres) and leaf thickness (LT; in millimetres) with precision callipers. The leaves were then placed in an oven at 60 °C for ≥ 72 h to dry completely. Dried leaves were then weighed for dry mass (in milligrams), allowing the calculation of leaf water content [LWC; (wet mass – dry mass)/wet mass], leaf density [LD; dry mass/(LA $\times$ LT); in grams per centimetre cubed], leaf mass




FIG. 1. Temperature profiles and relationships between air temperature (T_{air}) and leaf temperature (T_{leaf}) in the glasshouse experiment. (A, B) Canopy T_{air} (A) and T_{leaf} (B) profiles over time across all plants for the benign- and high-temperature treatments during the experiment. Coloured dashed lines represent the glasshouse set temperatures for each treatment. Data shown are means $\pm$ 95 % confidence intervals across 6 days of measurement. (C, D) Relationship between T_{air} and T_{leaf} in the benign treatment (C) and the high-temperature treatment (D). Black dashed lines represent an isometric relationship, and coloured solid lines are simple linear regressions $\pm$ 95 % confidence intervals.

per area (LMA; dry mass/leaf area; in kilograms per metre squared), and leaf dry matter content (LDMC; dry mass/wet mass; in kilograms per kilogram). Summary statistics for individual traits are shown in Supplementary Data Table S3.

Thermal time constant

We calculated the theoretical leaf thermal time constant (τ ; in seconds) as a mechanistic composite trait that links leaf traits to

time-dependent decoupling of T_{leaf} from ambient conditions in the absence of thermoregulation via latent heat flux (Michaletz *et al.*, 2015, 2016; Bison and Michaletz, 2024).

$$\tau = \varphi \times \text{LMA} \times \left(\frac{c_{p,w}}{\text{LDMC} \times h} + \frac{c_{p,d} - c_{p,w}}{h} \right)$$

Values for parameters (φ , $c_{p,w}$ and $c_{p,d}$) were as defined by Bison and Michaletz (2024), i.e. φ (the ratio of projected to total leaf area) was taken to be 0.5, specific heat capacities

$c_{p,w}$ and $c_{p,d}$ were taken as 4181 and 2814 J kg⁻¹ K⁻¹, respectively, and h is a heat transfer coefficient (in watts per metre squared per kelvin) that depends on leaf width (Michaletz *et al.*, 2016). Small values of τ represent leaves that change temperature rapidly in response to environmental temperature changes, and large values correspond to leaves that respond slowly. For additional information, see Supplementary Data Appendix S1.

Statistical analyses

To test the nature of thermal decoupling (ΔT and β) in benign and high-temperature treatments across different species originating from the three biomes, we fitted linear mixed-effects regression (LMER) models. The temperate biome species and benign-temperature treatment were used as reference levels, and all models contained random effect (intercept) terms for growth form, species nested within taxonomic family, and plant identity to account for repeated measures on the same plants. The LMER models were fitted with either ΔT or β as the response variable, with treatment, biome and their interaction as categorical fixed effects.

To determine the effects of the combined leaf traits and their interaction with biome on ΔT and β , LMER models were fitted initially to the benign and high-temperature treatments separately. We generated composite leaf traits in two ways: principal components analysis and the thermal time constant (τ). For principal components analysis, we included the five passive leaf traits (LA, LW, LT, LWC and LD) and the active leaf trait (g_{sw}), which generated two major axes of variation (PC1 and PC2; Supplementary Data Table S4). For these models, the random effects of species and growth form explained near-zero variance owing to redundancy with the leaf traits; therefore, simplified linear models were fitted to the benign and high-temperature treatments separately to determine the effects of composite leaf traits on ΔT and β . These models included two-way interactions between either PC1 and PC2 or τ with biome. We applied type III ANOVAs with Satterthwaite's degrees of freedom (d.f.) to LMER models, followed by Tukey's honest significant differences *post hoc* tests with Kenward-Roger's d.f. for reporting. *Post hoc* tests compared pairwise differences among combinations of biome and treatment. The 95 % confidence intervals were obtained using non-parametric bootstrapping with the *mean_cl_boot* function from *Hmisc* (Harrell, 2019). All data analyses were conducted in R v.4.3.1 (R Core Team, 2023) using *lme4* (Bates *et al.*, 2015), *performance* (Lüdtke *et al.*, 2021), *emmeans* (Lenth, 2023), *factoextra* (Kassambara and Mundt, 2020) and *tidyverse* R packages (Wickham *et al.*, 2019).

RESULTS

Leaf thermal decoupling depends on both biomes and temperature treatments

We hypothesized that species originating from the more extreme alpine and desert climates would have greater thermoregulation tendency (ΔT differing from zero and β differing from one, exhibiting either megathermy or limited

homeothermy) than those originating from the more benign temperate climate. The overall effect of treatment on ΔT was significant (Table 3), whereby high-temperature conditions resulted in significantly more negative ΔT (cooler leaves) than the benign treatment (Fig. 1). There was substantial variation in T_{leaf} along the T_{air} continuum both within and among biomes (Fig. 2A). On average, ΔT was positive for temperate species in both benign (1.99 ± 1.30 °C) and high-temperature (0.60 ± 0.91 °C) treatments (Fig. 2B). For both alpine and desert species, ΔT was positive in benign (alpine, 0.63 ± 1.01 °C; desert, 0.50 ± 1.05 °C) and negative in high-temperature (alpine, -1.25 ± 0.77 °C; desert, -1.66 ± 0.92 °C) treatments.

The effect of biome on ΔT was significant (Table 3), with alpine and desert species having ~ 1.2 °C cooler leaves than temperate species in both treatments (Fig. 2B). However, there were no significant interactions between biome and treatment (Table 3), such that the magnitude of difference in ΔT across biomes was consistent in both treatments (Fig. 2B). *Post hoc* tests revealed that pairwise temperature treatment differences in ΔT were significant within each biome (Fig. 2B; Supplementary Data Table S5). The temperate species were significantly different from desert species in either benign or high temperature and different from alpine species in high temperature (Fig. 2B; Supplementary Data Table S5).

Thermal coupling strength (β) was significantly higher in the high-temperature treatment, but not significantly different among biomes (Table 3; Fig. 2C). Species from all biomes typically exhibited limited homeothermy ($\beta < 1$) in benign conditions, but at high temperature, on average, temperate species exhibited megathermy ($\beta > 1$), whereas alpine and desert species exhibited poikilothermy ($\beta \approx 1$) (Fig. 2C). β differed significantly between treatments in only the temperate and desert species, and the only other significant contrast was the

TABLE 3. Type III ANOVA outputs from linear mixed-effects regression (LMER) models that test the contributions of temperature treatment and biome on thermal offset (ΔT) and thermal coupling strength (β). *P < 0.05. Random effects are reported from the LMER summary.

Fixed effect	F	d.f.	P-value	Random effect	s.d.	Variance (%)
Response: thermal offset (ΔT)						
Treatment	77.893	1,72	<0.001*	Species	<0.001	0.0
Biome	22.332	2,45	<0.001*	Family	0.430	6.6
Treatment × biome	0.543	2,72	0.584	Growth form	0.665	15.8
				Plant identity	0.667	15.9
				Residual	1.313	61.7
$R^2 = 0.545$						
Response: thermal coupling strength (β)						
Treatment	27.244	1132	<0.001*	Species	0.033	3.5
Biome	2.230	2,7	0.181	Family	0.052	8.7
Treatment × biome	0.205	2132	0.815	Growth form	0.009	0.3
				Plant identity	<0.001	0.0
				Residual	0.165	87.6
$R^2 = 0.190$						




FIG. 2. Canopy leaf and air temperature relationships, thermal coupling strength and thermal offsets among species from three biomes and two temperature treatments. (A) The overall raw data for relationships between T_{leaf} and T_{air} , where linear regressions are fitted to individual plants. (B) Mean thermal offset (ΔT), which is the magnitude of the difference, $T_{\text{leaf}} - T_{\text{air}}$. (C) Mean thermal coupling strengths (β), which is the slope of the relationship between T_{leaf} and T_{air} , calculated at 30-min intervals. $\beta > 1$ indicates megathermy, $\beta < 1$ indicates limited homeothermy, and $\beta \approx 1$ indicates poikilothermy. Data shown are means $\pm$ s.e.m. The grey lines for all panels (isometric, $\beta = 1$, $\Delta T = 0$) indicate when T_{air} and T_{leaf} are equivalent.

temperate species in high temperatures compared with desert species in benign conditions (Fig. 2C; Supplementary Data Table S5).

Taxonomic and growth form differences explained relatively small proportions of variance (in both temperature treatments, for both ΔT and β) beyond that explained by biome (Table 3). Across species, ΔT showed similar patterns in both temperature treatments (Fig. 3), with a few notable exceptions. *Acacia longifolia* (temperate) had the highest ΔT among

temperate species in benign conditions but the lowest ΔT in high-temperature conditions (Fig. 3). *Eucalyptus largiflorens* (desert) also shifted from positive ΔT in benign conditions to a strongly negative ΔT in high-temperature conditions (Fig. 3). The most negative values of ΔT were achieved by two desert *Acacia* species, *Acacia salicina* and *Acacia aneura*. Both these species could cool their leaves below T_{air} by $>3\text{ }^{\circ}\text{C}$ in high-temperature conditions; $>1\text{ }^{\circ}\text{C}$ greater cooling than any other species tested.




FIG. 3. Thermal coupling parameters across species. Thermal offsets (ΔT ; A) and thermal coupling strength (β ; B) in benign (left) and high-temperature (right) treatments for each of 15 species originating from three biomes. Data shown are means $\pm$ bootstrapped 95% confidence intervals. The dashed grey line at $\Delta T = 0$ and $\beta = 1$ indicates when T_{air} and T_{leaf} are equivalent.

TABLE 4. ANOVA outputs from linear regression models that test the contributions of biome and leaf traits as principal components (PC1 and PC2) or as a composite thermal time constant (τ) on thermal offset (ΔT) and thermal coupling strength (β) separately in benign- and high-temperature conditions. * $P < 0.05$.

Fixed effects	Thermal offset (ΔT)				Thermal coupling strength (β)			
	Benign		High temperature		Benign		High temperature	
	<i>F</i>	<i>P</i> -value	<i>F</i>	<i>P</i> -value	<i>F</i>	<i>P</i> -value	<i>F</i>	<i>P</i> -value
Biome	6.178	0.004*	23.349	<0.001*	4.076	0.023*	1.512	0.231
PC1	5.730	0.021*	25.554	<0.001*	1.938	0.170	4.207	0.046*
PC2	0.489	0.488	0.847	0.362	0.787	0.380	0.061	0.807
Biome $\times$ PC1	2.934	0.063	0.115	0.892	1.194	0.312	0.454	0.638
Biome $\times$ PC2	0.442	0.645	1.041	0.361	1.732	0.188	0.019	0.981
Biome	6.000	0.005*	19.845	<0.001*	3.732	0.031*	1.575	0.217
τ	0.225	0.638	0.015	0.902	0.067	0.797	0.006	0.940
Biome $\times$ τ	3.996	0.024*	7.091	0.002*	0.369	0.693	2.215	0.120




FIG. 4. Principal components analysis of six leaf traits: leaf water content (LWC), leaf area (LA), leaf thickness (LT), leaf width (LW), stomatal conductance to water (g_{sw}) and leaf density (LD). (A, B) Contributions of each leaf trait to PC1 (A) and PC2 (B) major axes. Red dashed line corresponds to the expected value if contributions were uniform among traits, where grey bars that are higher than the red line indicate dominant variables to that principal component. (C) Principal component space of dominant PC1 and PC2 axes that together explain 75.7 % of the variance in the leaf traits. Coloured ellipses represent the 95 % confidence space for each biome.

Leaf traits can moderate thermoregulation

Species-level leaf traits are shown in Supplementary Data Fig. S2, and the relationships of individual leaf traits and thermal coupling are shown in Supplementary Data Fig. S3. Given the strong effect of temperature treatment on ΔT and β , we analysed the effects of composite leaf traits on thermal

coupling in each treatment separately. Biome was accounted for in all models and was significant in all cases except β at high temperature, indicating that differences in biome contributed to thermoregulatory differences indirectly (Table 4).

The composite leaf trait major axis (PC1) can be interpreted as an axis from negative values representing thick, less dense leaves with high water content and high stomatal conductance




FIG. 5. Relationships between leaf thermal coupling and composite leaf traits from principal components analysis (PC1 and PC2 axes) and thermal time constant (τ). (A, B) Thermal offset (ΔT ; A) in relationship to leaf traits PC1 and thermal coupling strength (β ; B) in relationship to leaf traits PC1 under benign and high-temperature treatments. (C, D) ΔT (C) and β (D) in relationship to leaf traits PC2. (E, F) ΔT (E) and β (F) in relationship to τ . Raw data are shown, and linear regressions are overlaid where relationships between trait and thermal coupling are significant (solid where $P < 0.05$ and dashed where $P < 0.1$) overall (black) or interact with biome (coloured). Corresponding model outputs are shown in Table 4. The dashed grey horizontal lines at $\Delta T = 0$ and $\beta = 1$ indicate when T_{air} and T_{leaf} are equivalent, and the dashed grey vertical lines in A–D show $PC1 = 0$ and $PC2 = 0$.

to positive values increasing towards thin, dense leaves with low water content and low stomatal conductance (Fig. 4A, C). The other dominant composite leaf trait axis (PC2) can be interpreted as a continuum from negative values representing wider, larger leaves to positive values increasing towards narrower, smaller leaves (Fig. 4B, C). The three biomes formed generally distinct clusters in principal component space. The leaves of alpine species were defined by negative PC1 (thick, less dense, high water content and high g_{sw}) and a narrow range of slightly negative PC2 (Fig. 4C). The leaves of temperate species were defined by positive PC1 (thin, dense, low water content and low g_{sw}) but spanned a wide range along PC2 (from broad and large to narrow and small) (Fig. 4C). The leaves of desert species covered a wide range of PC1, but all were positive along PC2 (small and narrow leaves) (Fig. 4C).

Thermal offset (ΔT) increased significantly as PC1 increased in both treatments (Table 4), i.e. leaves that were thin, dense and had low water content and relatively low g_{sw} characteristics tended to be warmer than air (Fig. 5A). Thermal coupling strength (β) increased significantly as PC1 increased in the high-temperature treatment (Table 4), i.e. leaves that were thicker, less dense and had high water content and high g_{sw} exhibited limited homeothermy, whereas leaves that were thinner, denser and had low water content and low g_{sw} exhibited megathermy (Fig. 5B). There was a marginally non-significant interaction between PC1 and biome (Fig. 5A, C; Table 4). Neither ΔT nor β was significantly related to PC2 (Fig. 5C, D; Table 4). The relationship between ΔT and thermal time constant (τ) was not significant overall, but did differ significantly among biomes (Fig. 5E; Table 4). Desert species had a negative relationship between ΔT and τ , whereas temperate and alpine species had a positive relationship, and these patterns were consistent in both treatments (Fig. 5E; Table 4). Thermal coupling strength (β) showed no significant relationships with τ or the interaction between τ and biome in either treatment (Fig. 5F; Table 4).

DISCUSSION

Here, we have shown that these diverse plant species that originate from contrasting biomes differ clearly in their thermoregulation in both benign and high-temperature conditions. Our hypotheses about the drivers of variation in plant thermoregulation were largely supported: variation in T_{leaf} , ΔT and β during high temperatures was dependent on origin biome and composite leaf traits, especially leaf water content and g_{sw} . Different species exposed to nearly identical conditions will reach different T_{leaf} owing to the unique interactions of their leaf properties with the environment (Perez and Feeley, 2020). Thus, understanding the sources of variation in T_{leaf} is essential: empirical data inform leaf energy budget theory and more accurate predictive models of T_{leaf} (Michaletz et al., 2015; Blonder et al., 2020; Kearney and Leigh, 2024).

Plants from extreme climates can thermoregulate more effectively

Our hypothesis that species originating from biomes with more extreme climates would have greater thermoregulatory tendency than those from more benign climates was generally supported. That is, the adaptations a plant has to the environmental

conditions of its biome of origin explain thermoregulation even in common conditions. Temperate species had leaves that were almost always warmer than air, whereas the leaves of alpine and desert species were equivalent to T_{air} in benign conditions but often much cooler than air at high temperatures. Species from all biomes showed limited homeothermy ($\beta < 1$) in benign conditions, but at high temperature, most exhibited poikilothermy ($\beta \approx 1$).

Leaf thermoregulation is thought to originate from selection on leaf traits to maximize carbon gain in their environment (Michaletz et al., 2016). That is, plants will maintain T_{leaf} within an optimal range for photosynthesis via variable stomatal opening to thermoregulate actively at an approximate cross-over point when T_{leaf} reaches ~ 27 – 30 °C and ΔT becomes negative (Michaletz et al., 2016; Dong et al., 2017). Experimental tests of the limited homeothermy hypothesis found that cotton (*Gossypium hirsutum*) exhibited poikilothermy until T_{air} reached 27 °C, then switched to limited homeothermy when T_{air} was 27–40 °C to maintain $T_{leaf} = 27 \pm 2$ °C, when water was available for transpiration (Upchurch and Mahan, 1988). However, recent large-scale analyses of canopy temperatures generally do not support a hypothesis of universal limited homeothermy; rather, there is evidence for a diverse range of viable thermoregulation strategies (β range = 0.7–1.3) (Still et al., 2022; Guo et al., 2023; Manzi et al., 2024). Our present study provides empirical support for these recent analyses at a smaller scale, where plants (on average) exhibited limited homeothermy in benign conditions, but a wide range of β in high temperatures.

Potential links between leaf thermoregulation and photosystem heat tolerance

A common measure of photosystem heat tolerance is T_{crit} , the critical temperature for photosystem II functional impairment and subsequent damage, which is derived from ramping assays of the temperature-dependent change in chlorophyll *a* fluorescence (Arnold et al., 2021). Many of the species in the present study are known to differ in T_{crit} from field surveys (Briceño et al., 2024) and controlled-environment experiments (Harris et al., 2024). Drawing links between T_{crit} from these studies and thermoregulation from our study returns some unexpected outcomes. Notably, *Dodonaea viscosa* was one of the least heat-tolerant desert species in the aforementioned studies (controlled 25 °C environment, 39.9 ± 1.0 °C; field, 45.4 ± 0.5 °C), and in the present study it had the highest T_{leaf} and $\Delta T \approx 0$ °C in high-temperature conditions. The relatively low heat-tolerance threshold of the alpine *Eucalyptus pauciflora* (controlled 25 °C environment, 42.7 ± 1.7 °C; field, 39.0 ± 0.8 °C) and the temperate *Acacia binervata* (controlled 25 °C environment, 42.0 ± 1.5 °C), which both had positive ΔT values at high temperatures in the present study, is consistent with this pattern. The reverse is true for the desert *Acacia* species, which are both extremely heat tolerant (*A. salicina*: controlled 25 °C environment, 46.9 ± 0.8 °C; field, 49.6 ± 0.8 °C; and *A. aneura*: controlled 25 °C environment, 48.6 ± 0.9 °C; field, 53.0 ± 4.0 °C), and here were found to have the lowest T_{leaf} and negative ΔT values at high temperatures.

We advocate for testing the association of thermal coupling metrics and heat tolerance as a focus of future investigations.

Initially, it appears counter-intuitive that a species should have adaptations to avoid and tolerate high temperatures concurrently. In an extremely hot and dry environment, plants may typically avoid the worst of heat stress through their structural leaf properties and evaporative cooling via transpiration. Yet, sustained dry conditions may render evaporative cooling an unviable option for avoiding heat, and high heat tolerance would become necessary (Gong *et al.*, 2023). A species that does not cool T_{leaf} below T_{air} (or only cools moderately), might indicate limited capacity for cooling or a high heat-tolerance threshold before initiating cooling. As such, we hypothesize that the tendency to cool T_{leaf} below T_{air} could be associated with higher heat tolerance in some species, but that it will probably also depend on water-use strategy.

Composite leaf traits contribute to thermoregulation at high temperature

The dominant axis of variation (PC1) was defined by three structural leaf traits [leaf thickness (LT), leaf density (LD) water content (LWC)] and by the active leaf trait, stomatal conductance (g_{sw}). PC1 correlated strongly with ΔT in benign conditions and with ΔT and β in high-temperature conditions. PC2, which was largely defined by leaf area (LA) and leaf width (LW) had relatively little consistent relationship with ΔT and β in either environment. LWC plays a key role in leaf thermoregulation and leaf economics (Michaletz *et al.*, 2015; Wang *et al.*, 2022). The specific heat capacity is positively correlated with the water content of leaves (Zhang *et al.*, 2025), and there is also a strong positive association between LWC and maximum g_{sw} (Zhou *et al.*, 2023). Water availability to plants is generally linked to the capacity to regulate T_{leaf} (Lambers and Oliveira, 2019; Cook *et al.*, 2021; Manzi *et al.*, 2024), and the transport and storage of water directly in leaf tissues can reduce heat loading while facilitating greater cooling (Zhou *et al.*, 2023). In desert species, thicker leaves (that often also have higher LWC) have slower heating response times relative to thinner leaves, resulting in lower T_{leaf} during temperature extremes, even in the absence of transpiration, but the effect is reduced for large leaves (Leigh *et al.*, 2012). Our study shows that leaf cooling is more effective in plants that have higher LWC and LT and lower LD in both temperature treatments.

Combinations of leaf functional and energy budget traits and environments across 41 species and seven sites along an elevation gradient showed that regression approaches achieved relatively low predictive power for ΔT and especially for β (Blonder *et al.*, 2020). In their study, the site environment played a more substantial role than commonly measured functional traits and energy balance traits, and interactions between traits and environment were relevant. Blonder *et al.* (2020) concluded that the low predictability of thermal coupling and the variation encountered at a given site indicate that a range of strategies will result in viable performance. Our results are consistent with these findings: variation in ΔT was more readily explained than β . The common environment approach we used highlights that both external environment (temperature treatment) and origin biome strongly influence ΔT , whereas only environmental conditions influence β .

The thermal time constant (τ) differed among plants from the different biomes. The alpine plants in this study had relatively large values of τ , indicating that they respond more slowly to environmental changes than the desert or temperate plants, which could be to buffer against the rapid environmental temperature fluctuations that occur naturally in the alpine biome (Körner, 2003). We also found that the relationship between τ and ΔT differed among biomes, being positive for temperate and alpine plants, such that leaves that respond relatively more slowly to environmental changes were warmer than air or at least cooled less effectively. In contrast, the leaves of desert plants that respond relatively more slowly to environmental changes were, nonetheless, far more effective at cooling below T_{air} , especially in comparison to temperate plants. That is, for the same value of τ , ΔT differed by $\leq 4^\circ\text{C}$ between desert and temperate plants, which suggests that g_{sw} was the main driver of these differences, because it is not involved in the calculation of τ . The dynamic fluctuations of T_{air} in glasshouse conditions suggests that T_{leaf} might not frequently reach a steady state within the range of τ , hence delays in both leaf warming and cooling might influence the relationships between traits and thermoregulation. All plants had access to adequate water throughout the heat event and could have transpired freely; however, desert plants transpired far more than temperate plants in both temperature treatments. If the desert plants with larger τ values opened their stomata earlier to achieve high g_{sw} and did so for longer than temperate plants with larger τ values, that could explain why the leaves of these desert plants were much cooler than air and why relationships with ΔT differed between these biomes.

Inherently low stomatal conductance limits evaporative cooling

We predicted that species with inherently low g_{sw} would be most limited in their thermoregulation. Generally, high g_{sw} strongly reduced ΔT , which was consistent across biomes except for temperate species in benign conditions. Stomatal conductance and LWC both play pivotal roles in enabling leaf cooling at high temperature, thus reducing T_{leaf} on acutely hot days and during heatwaves will clearly depend on water availability and water-use strategies (Drake *et al.*, 2018; Aparecido *et al.*, 2020; Cook *et al.*, 2021; Marchin *et al.*, 2022; Manzi *et al.*, 2024). The temperate species originate from a biome that is typically not water limited, yet these species appear to be more limited in their tendency for thermoregulation via regulating stomata. One potential explanation for this is the intricate link between temperature and VPD. Increases in VPD are a major concern with climate change, because it can also limit evapotranspiration by exacerbating water stress and forcing stomatal closure (Grossiord *et al.*, 2020). Although the relative humidity in our glasshouse experiment was generally low enough to enable cooling to take place (Mahan and Upchurch, 1988), VPD increased in the high-temperature treatment. The lower g_{sw} in the species originating from the coastal temperate biome probably responded to the high-temperature (with relatively high VPD) treatment by closing stomata more than alpine and desert species that are adapted to typically drier air.

An alternative explanation is that these coastal temperate species have intrinsically lower g_{sw} or slower stomatal response to high temperatures in comparison to species that originated

from more extreme climates. For example, some desert and alpine species can open their stomata rapidly to optimize the trade-offs between carbon fixation, water loss and leaf thermoregulation during narrower windows of suitable conditions in these challenging environments (Knapp and Smith, 1988, 1991; Fernández-Marín *et al.*, 2020). Glasshouse experiments with plants originating from hot dry and hot wet habitats suggest that transpiration is greater in species from hot dry habitats that have sporadic rain (Lin *et al.*, 2017), supporting the idea that extreme climate is a driver of thermoregulation strategy. Similar to our study, those authors found that cooling via stomatal behaviour was more effective than passive leaf traits when water was sufficient (Lin *et al.*, 2017). We infer that high T_{air} in our high-temperature treatment probably increased T_{leaf} to a point that exceeded the heat load that most of the temperate species could dissipate via transpiration.

Leaf thermoregulation is a complex plant–environment interaction

Plant species are often interpreted as being on a water stress avoidance–tolerance (isohydric–anisohydric) spectrum; however, rather than being a simple plant hydraulic trait, isohydricity is a complex plant–environment interaction (Hochberg *et al.*, 2018). Leaf thermoregulation seems analogous to this complexity. There are many causes for leaf thermoregulation depending on the immediate environment of the plant. For example, thermoregulation can optimize photosynthesis, but it also plays a role in hydraulic maintenance; then at extreme temperatures, thermoregulation facilitates avoidance of heat damage in the absence of photosynthesis (Slot and Winter, 2017; Drake *et al.*, 2018; Fauset *et al.*, 2018; Guo *et al.*, 2022). Therefore, the balance among the available thermoregulation mechanisms depends on these dynamic plant–environment interactions (Guo *et al.*, 2023). Contrasting patterns of leaf thermoregulatory traits and strategies among provenances across tropical trees demonstrates that warm-adapted provenances are not necessarily less vulnerable to heat stress based on their operating temperatures and heat tolerance (Middleby *et al.*, 2024). Elucidating the mechanisms that underlie differences in thermoregulatory strategies of plants across different origin biomes will be essential empirical research for applications to plant breeding and management of wild populations. Determining the physical leaf and stem properties and the underlying genetic markers and mechanisms for stomatal responsiveness that contribute to variation in plant thermoregulation and plasticity therein could be used to identify and select on target traits (Fritz *et al.*, 2018).

CONCLUSION

Advanced tools for rapidly estimating leaf size (Schrader *et al.*, 2021; Leigh, 2022) and predicting leaf temperatures based on biophysical modelling with microclimates and energy budgets are now available (Kearney and Leigh, 2024). The accuracy of predicted leaf temperatures requires capturing and understanding the diversity of functional leaf traits and stomatal conductance behaviour, which can have a large impact on T_{leaf} predictions (Perez and Feeley, 2020; Kearney and Leigh, 2024). Our study provides empirical evidence that species from contrasting biomes that are exposed to common conditions (benign

or high temperature) will respond to the conditions by regulating T_{leaf} to different extents. We also identify that composite leaf traits explain variation in leaf thermoregulation among species. Our findings suggest that, beyond simple expectations of leaf size, species from a coastal temperate biome appear to possess a suite of thermoregulatory traits more likely to increase exposure to heat stress, particularly if combined with dry conditions, than those adapted to more extreme conditions. The increasingly extreme environmental conditions that are occurring during the Anthropocene are exerting significant pressure on plants in many regions to avoid, tolerate and acclimatize to higher temperatures. Further work should evaluate interactive effects among temperature, VPD and water availability to discern the impacts of these major global change factors on leaf thermoregulation.

SUPPLEMENTARY DATA

Supplementary data are available at *Annals of Botany* online and consist of the following.

Appendix S1: thermal time constant notes. Figure S1: phylogenetic tree of the 15 species in the experiment. Figure S2: leaf trait values across species. Figure S3: relationships between leaf thermal coupling and six key leaf traits. Table S1: glasshouse environmental parameters. Table S2: summary statistics for each thermal trait. Table S3: summary statistics for each leaf trait. Table S4: principal components analysis variable loadings for leaf traits. Table S5: pairwise contrasts for ΔT and β among biomes and treatments.

FUNDING

This work was supported by the Australian Research Council through a Linkage Project and Discovery Project (grant numbers LP180100942 and DP240100177).

ACKNOWLEDGEMENTS

We respectfully acknowledge the traditional custodians of the land on which we conducted this work, the Ngannawal and Ngambri people, and we acknowledge the many First Nations peoples from land where our seeds were collected. Rosalie Harris and Philippa Alvarez germinated the seeds, and Freya Brown, Angela Bracken and Alan Vincent assisted with propagating the seedlings. Dmitry Grishin, Jacynnda Bovill, Sarah Woodcock and Isabelle Bennett assisted with experimental set-up and measurements, and Haydon Siiteri provided the Li-COR LI600. Two anonymous reviewers provided excellent feedback that improved an earlier version of this manuscript. We sincerely thank the Plant Services team at The Australian National University for their efforts in maintaining the plants used in this experiment.

AUTHOR CONTRIBUTIONS

V.F.B. and P.A.A. conceptualized the study with input from A.M.C., A.B.N. and A.L.; M.J.W. collected the data; P.A.A.

analysed the data and wrote the manuscript; all authors contributed to editing.

CONFLICT OF INTEREST

None declared.

DATA AVAILABILITY

Data and R code are available from the Figshare digital repository: <https://doi.org/10.6084/m9.figshare.28741331>.

LITERATURE CITED

- Anderson CJR, Rosas-Anderson PJ. 2017. Leafscan app. <https://www.leafscanapp.com/> (7 January 2022, date last accessed).
- Aparecido LMT, Woo S, Suazo C, Hultine KR, Blonder B. 2020. High water use in desert plants exposed to extreme heat. *Ecology Letters* 23: 1189–1200.
- Arnold PA, Briceño VF, Gowland KM, Catling AA, Bravo LA, Nicotra AB. 2021. A high-throughput method for measuring critical thermal limits of leaves by chlorophyll imaging fluorescence. *Functional Plant Biology* 48: 634–646.
- Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67: 1–48.
- Bison NN, Michaletz ST. 2024. Variation in leaf carbon economics, energy balance, and heat tolerance traits highlights differing timescales of adaptation and acclimation. *The New Phytologist* 242: 1919–1931.
- Blonder B, Michaletz ST. 2018. A model for leaf temperature decoupling from air temperature. *Agricultural and Forest Meteorology* 262: 354–360.
- Blonder B, Escobar S, Kapás RE, Michaletz ST. 2020. Low predictability of energy balance traits and leaf temperature metrics in desert, montane and alpine plant communities. *Functional Ecology* 34: 1882–1897.
- Briceño VF, Arnold PA, Cook AM, et al. 2024. Drivers of thermal tolerance breadth of plants across contrasting biomes: do mean or seasonality in climate indices matter more? *bioRxiv*: 2023.10.02.560437 [preprint, not peer reviewed].
- Cavaleri MA. 2020. Cold-blooded forests in a warming world. *The New Phytologist* 228: 1455–1457.
- Cook AM, Berry N, Milner KV, Leigh A. 2021. Water availability influences thermal safety margins for leaves. *Functional Ecology* 35: 2179–2189.
- Cowan T, Purich A, Perkins S, Pezza A, Bosch G, Sadler K. 2014. More frequent, longer, and hotter heat waves for Australia in the twenty-first century. *Journal of Climate* 27: 5851–5871.
- Curtis EM, Gollan J, Murray BR, Leigh A. 2016. Native microhabitats better predict tolerance to warming than latitudinal macro-climatic variables in arid-zone plants. *Journal of Biogeography* 43: 1156–1165.
- Deva CR, Urban MO, Challinor AJ, Falloon P, Svitáková L. 2020. Enhanced leaf cooling is a pathway to heat tolerance in common bean. *Frontiers in Plant Science* 11: 19.
- Dong N, Prentice IC, Harrison SP, Song QH, Zhang YP. 2017. Biophysical homeostasis of leaf temperature: a neglected process for vegetation and land-surface modelling. *Global Ecology and Biogeography* 26: 998–1007.
- Drake JE. 2023. A data-intensive documentation of plant ecosystem thermoregulation across spatial and temporal scales. *The New Phytologist* 238: 921–923.
- Drake JE, Tjoelker MG, Vårhammar A, et al. 2018. Trees tolerate an extreme heatwave via sustained transpirational cooling and increased leaf thermal tolerance. *Global Change Biology* 24: 2390–2402.
- Fauset S, Freitas HC, Galbraith DR, et al. 2018. Differences in leaf thermoregulation and water use strategies between three co-occurring Atlantic forest tree species. *Plant, Cell & Environment* 41: 1618–1631.
- Fernández-Marín B, Gullás J, Figueroa CM, et al. 2020. How do vascular plants perform photosynthesis in extreme environments? An integrative ecophysiological and biochemical story. *The Plant Journal: for Cell and Molecular Biology* 101: 979–1000.
- Fritz MA, Rosa S, Sicard A. 2018. Mechanisms underlying the environmentally induced plasticity of leaf morphology. *Frontiers in Genetics* 9: 478.
- Gates DM. 1968. Transpiration and leaf temperature. *Annual Review of Plant Physiology* 19: 211–238.
- Geange SR, Arnold PA, Catling AA, et al. 2021. The thermal tolerance of photosynthetic tissues: a global systematic review and agenda for future research. *The New Phytologist* 229: 2497–2513.
- Gibson AC. 1998. Photosynthetic organs of desert plants. *BioScience* 48: 911–920.
- Gong X-W, Leigh A, Guo J-J, Fang L-D, Hao G-Y. 2023. Sand dune shrub species prioritize hydraulic integrity over transpirational cooling during an experimental heatwave. *Agricultural and Forest Meteorology* 336: 109483.
- Goraya GK, Kaur B, Asthir B, Bala S, Kaur G, Farooq M. 2017. Rapid injuries of high temperature in plants. *Journal of Plant Biology* 60: 298–305.
- Grossiord C, Buckley TN, Cernusak LA, et al. 2020. Plant responses to rising vapor pressure deficit. *The New Phytologist* 226: 1550–1566.
- Guo Z, Yan Z, Majcher BM, et al. 2022. Dynamic biotic controls of leaf thermoregulation across the diel timescale. *Agricultural and Forest Meteorology* 315: 108827.
- Guo Z, Still CJ, Lee CKF, et al. 2023. Does plant ecosystem thermoregulation occur? An extratropical assessment at different spatial and temporal scales. *The New Phytologist* 238: 1004–1018.
- Harrell FEJ. 2019. *Hmisc: Harrell Miscellaneous. R package, v5.1-0*. <https://CRAN.R-project.org/package=Hmisc> (11 March 2025, date last accessed).
- Harris RMB, Beaumont LJ, Vance TR, et al. 2018. Biological responses to the press and pulse of climate trends and extreme events. *Nature Climate Change* 8: 840.
- Harris RJ, Alvarez PR, Bryant C, et al. 2024. Acclimation of thermal tolerance in juvenile plants from three biomes is suppressed when extremes co-occur. *Conservation Physiology* 12: coae027.
- Hochberg U, Rockwell FE, Holbrook NM, Cochard H. 2018. Iso/anisohydry: a plant–environment interaction rather than a simple hydraulic trait. *Trends in Plant Science* 23: 112–120.
- Hüve K, Bichele I, Kaldmæ H, Rasulov B, Valladares F, Niinemets U. 2019. Responses of aspen leaves to heatflecks: both damaging and non-damaging rapid temperature excursions reduce photosynthesis. *Plants (Basel, Switzerland)* 8: 145.
- Jones HG. 2014. *Plants and microclimate: a quantitative approach to environmental plant physiology*. Cambridge, UK: Cambridge University Press.
- Karger DN, Conrad O, Böhner J, et al. 2017. Climatologies at high resolution for the Earth's land surface areas. *Scientific Data* 4: 170122.
- Kassambara A, Mundt F. 2020. *factoextra: Extract and Visualize the Results of Multivariate Data Analyses. R package, v1.0.7*. <https://CRAN.R-project.org/package=factoextra> (1 March 2025, date last accessed).
- Kearney MR, Leigh A. 2024. Fast, accurate and accessible calculations of leaf temperature and its physiological consequences with NicheMapR. *Methods in Ecology and Evolution* 15: 1516–1531.
- Kitudom N, Fauset S, Zhou Y, et al. 2022. Thermal safety margins of plant leaves across biomes under a heatwave. *The Science of the Total Environment* 806: 150416.
- Klein T. 2014. The variability of stomatal sensitivity to leaf water potential across tree species indicates a continuum between isohydric and anisohydric behaviours. *Functional Ecology* 28: 1313–1320.
- Knapp AK, Smith WK. 1988. Effect of water stress on stomatal and photosynthetic responses in subalpine plants to cloud patterns. *American Journal of Botany* 75: 851–858.
- Knapp AK, Smith WK. 1991. Gas exchange responses to variable sunlight in two Sonoran desert herbs: comparison with subalpine species. *Botanical Gazette* 152: 269–274.
- Körner C. 2003. *Alpine plant life*. Berlin: Springer.
- Kullberg AT, Coombs L, Soria Ahuanari RD, Fortier RP, Feeley KJ. 2023. Leaf thermal safety margins decline at hotter temperatures in a natural warming ‘experiment’ in the Amazon. *New Phytologist* 241: 1447–1463.
- Lambers H, Oliveira RS. 2019. Plant water relations. In: Lambers H, Oliveira RS, eds. *Plant physiological ecology*. Cham: Springer International.
- Leigh A. 2022. Using leaf shape to determine leaf size could be a game-changer. A commentary on: ‘Leaf size estimation based on leaf length, width and shape’. *Annals of Botany* 129: i–ii.
- Leigh A, Sevanto S, Ball MC, et al. 2012. Do thick leaves avoid thermal damage in critically low wind speeds? *The New Phytologist* 194: 477–487.
- Leigh A, Sevanto S, Close JD, Nicotra AB. 2017. The influence of leaf size and shape on leaf thermal dynamics: does theory hold up under natural conditions? *Plant, Cell & Environment* 40: 237–248.
- Lenth RV. 2023. *emmeans: estimated marginal means, aka least-squares means. R package, v1.8.7*. <https://CRAN.R-project.org/package=emmeans> (11 March 2025, date last accessed).

- Leuzinger S, Körner C. 2007. Tree species diversity affects canopy leaf temperatures in a mature temperate forest. *Agricultural and Forest Meteorology* **146**: 29–37.
- Lin H, Chen Y, Zhang H, Fu P, Fan Z. 2017. Stronger cooling effects of transpiration and leaf physical traits of plants from a hot dry habitat than from a hot wet habitat. *Functional Ecology* **31**: 2202–2211.
- Lüdecke D, Ben-Shachar MS, Patil I, Waggoner P, Makowski D. 2021. performance: an R package for assessment, comparison and testing of statistical models. *Journal of Open Source Software* **6**: 3139.
- Lusk CH, Clearwater MJ, Laughlin DC, et al. 2018. Frost and leaf-size gradients in forests: global patterns and experimental evidence. *The New Phytologist* **219**: 565–573.
- Mahan JR, Upchurch DR. 1988. Maintenance of constant leaf temperature by plants—I. Hypothesis-limited homeothermy. *Environmental and Experimental Botany* **28**: 351–357.
- Manzi OJL, Wittemann M, Dusinge ME, et al. 2024. Canopy temperatures strongly overestimate leaf thermal safety margins of tropical trees. *The New Phytologist* **243**: 2115–2129.
- Marchin RM, Backes D, Ossola A, Leishman MR, Tjoelker MG, Ellsworth DS. 2022. Extreme heat increases stomatal conductance and drought-induced mortality risk in vulnerable plant species. *Global Change Biology* **28**: 1133–1146.
- Matthews JSA, Viallet-Chabrand SRM, Lawson T. 2017. Diurnal variation in gas exchange: the balance between carbon fixation and water loss. *Plant Physiology* **174**: 614–623.
- Michaletz ST, Weiser MD, Zhou J, Kaspari M, Helliker BR, Enquist BJ. 2015. Plant thermoregulation: energetics, trait–environment interactions, and carbon economics. *Trends in Ecology & Evolution* **30**: 714–724.
- Michaletz ST, Weiser MD, McDowell NG, et al. 2016. The energetic and carbon economic origins of leaf thermoregulation. *Nature Plants* **2**: 16129.
- Middleby KB, Cheesman AW, Hopkinson R, et al. 2024. Ecotypic variation in leaf thermoregulation and heat tolerance but not thermal safety margins in tropical trees. *Plant, Cell & Environment* **48**: 649–663.
- Nieola CC, Carvalho CP, Carvalho V, Rodrigues E. 2017. Rapid responses of plants to temperature changes. *Temperature (Austin, Tex.)* **4**: 371–405.
- Pedraza S. 2024. Leaf traits and leaf-to-air temperature differences in tropical plants suggest variability in thermoregulatory capacities across elevations. *Biotropica* **56**: e13332.
- Perez TM, Feeley KJ. 2020. Photosynthetic heat tolerances and extreme leaf temperatures. *Functional Ecology* **34**: 2236–2245.
- Perkins-Kirkpatrick SE, Lewis SC. 2020. Increasing trends in regional heat waves. *Nature Communications* **11**: 3357.
- Perkins-Kirkpatrick S, Barriopedro D, Jha R, et al. 2024. Extreme terrestrial heat in 2023. *Nature Reviews Earth & Environment* **5**: 244–246.
- Posch BC, Bush SE, Koepke DF, et al. 2024. Intensive leaf cooling promotes tree survival during a record heatwave. *Proceedings of the National Academy of Sciences of the United States of America* **121**: e2408583121.
- R Core Team. 2023. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Reich PB, Wright IJ, Cavender-Bares J, et al. 2003. The evolution of plant functional variation: traits, spectra, and strategies. *International Journal of Plant Sciences* **164**: S143–S164.
- Salisbury FB, Spomer GG. 1964. Leaf temperatures of alpine plants in the field. *Planta* **60**: 497–505.
- Schrader J, Shi P, Royer DL, et al. 2021. Leaf size estimation based on leaf length, width and shape. *Annals of Botany* **128**: 395–406.
- Slot M, Winter K. 2017. *In situ* temperature response of photosynthesis of 42 tree and liana species in the canopy of two Panamanian lowland tropical forests with contrasting rainfall regimes. *New Phytologist* **214**: 1103–1117.
- Slot M, Cala D, Aranda J, Virgo A, Michaletz ST, Winter K. 2021. Leaf heat tolerance of 147 tropical forest species varies with elevation and leaf functional traits, but not with phylogeny. *Plant, Cell & Environment* **44**: 2414–2427.
- Smith WK. 1978. Temperatures of desert plants: another perspective on the adaptability of leaf size. *Science* **201**: 614–616.
- Still CJ, Page G, Rastogi B, et al. 2022. No evidence of canopy-scale leaf thermoregulation to cool leaves below air temperature across a range of forest ecosystems. *Proceedings of the National Academy of Sciences of the United States of America* **119**: e2205682119.
- Tserej O, Feeley KJ. 2021. Variation in leaf temperatures of tropical and subtropical trees are related to leaf thermoregulatory traits and not geographic distributions. *Biotropica* **53**: 868–878.
- Upchurch DR, Mahan JR. 1988. Maintenance of constant leaf temperature by plants—II. Experimental observations in cotton. *Environmental and Experimental Botany* **28**: 359–366.
- Vogel S. 2009. Leaves in the lowest and highest winds: temperature, force and shape. *The New Phytologist* **183**: 13–26.
- Wang Z, Huang H, Wang H, et al. 2022. Leaf water content contributes to global leaf trait relationships. *Nature Communications* **13**: 5525.
- Wickham H, Averick M, Bryan J, et al. 2019. Welcome to the tidyverse. *Journal of Open Source Software* **4**: 1686.
- Zhang J, Kaiser E, Zhang H, Marcelis LFM, Viallet-Chabrand S. 2025. A simple new method to determine leaf specific heat capacity. *Plant Methods* **21**: 6.
- Zhou Y, Kitudom N, Fauset S, et al. 2023. Leaf thermal regulation strategies of canopy species across four vegetation types along a temperature and precipitation gradient. *Agricultural and Forest Meteorology* **343**: 109766.