

# Selective Distance Measurements Using Triple Spin Labeling with $\text{Gd}^{3+}$ , $\text{Mn}^{2+}$ , and a Nitroxide

*Zuyan Wu, Akiva Feintuch<sup>2</sup>, Alberto Collauto<sup>2</sup>, Luke A. Adams<sup>3</sup>, Luigi Aurelio<sup>3</sup>, Bim Graham<sup>3</sup>, Gottfried Otting<sup>1\*</sup>, and Daniella Goldfarb<sup>2\*</sup>.*

<sup>1</sup> Research School of Chemistry, Australian National University, Canberra, ACT 2601, Australia

<sup>2</sup> Department of Chemical Physics, Weizmann Institute of Science, Rehovot 76100, Israel

<sup>3</sup> Monash Institute of Pharmaceutical Sciences, Monash University, Parkville VIC 3052, Australia


## **Corresponding Author**

Gottfried Otting, e-mail [gottfried.otting@anu.edu.au](mailto:gottfried.otting@anu.edu.au), Daniella Goldfarb,  
[Daniella.goldfarb@weizmann.ac.il](mailto:Daniella.goldfarb@weizmann.ac.il)

## ABSTRACT

Distance measurements by pulse EPR techniques, such as double electron-electron resonance (DEER, also called PELDOR), have become an established tool to explore structural properties of bio-macromolecules and their assemblies. In such measurements a pair of spin labels provides a single distance constraint. Here we show that by employing three different types of spin labels that differ in their spectroscopic and spin dynamics properties it is possible to extract three independent distances from a single sample. We demonstrate this using the *Antennapedia* homeodomain orthogonally labeled with  $Gd^{3+}$  and  $Mn^{2+}$  tags in complex with its cognate DNA binding site labeled with a nitroxide.

## TOC GRAPHICS



Double electron-electron resonance (DEER, also called PELDOR) experiments provide valuable long-range distance restraints to probe the structure of biological macromolecules and their assemblies by measuring the dipolar interaction between unpaired electron spins at different sites.<sup>1</sup> For diamagnetic systems, DEER experiments are commonly performed following the covalent attachment of two, usually identical, spin labels at well defined, strategically chosen locations within the biological macromolecule. While the majority of DEER applications have been carried out using the well-established nitroxide spin label, other options for spin labelling have emerged in the last decade, which are based on  $Gd^{3+}$ <sup>2-3</sup>,  $Mn^{2+}$ <sup>4,5,6</sup> and  $Cu^{2+}$  ions<sup>7-8</sup> as well as trityl radicals.<sup>9-11</sup>  $Cu^{2+}$  has the advantage of being a simple  $S=1/2$  and is attractive at common X-band frequencies where its spectra width is limited. Trityl radicals are unique as they allow for room temperature distance measurements.<sup>11</sup> Among the metal ions  $Gd^{3+}$  and  $Mn^{2+}$  labels are particularly attractive for measurements at a high field, such as W-band (95 GHz, ~3.5 T), though Q-band (~34 GHz, ~1.2 T) is also possible.<sup>12</sup> The use of  $Gd^{3+}$  complexes as spin labels at W-band has been shown to be advantageous over the commonly used nitroxide label due to the absence of orientation selection<sup>13-14</sup> and their stability in living cells.<sup>15-16</sup>

The most popular method for introducing  $Gd^{3+}$  and  $Mn^{2+}$  spin labels is via the thiol group of a native or genetically engineered cysteine residue as is commonly done with nitroxide spin labels. An alternative method is to employ an unnatural amino acid that has been genetically encoded to be incorporated in response to an amber stop codon.<sup>17-18</sup> While DEER applications using  $Gd^{3+}$  and  $Mn^{2+}$  have so far involved mostly  $Gd^{3+}$ – $Gd^{3+}$  and  $Mn^{2+}$ – $Mn^{2+}$  distance measurements, the possibility to use different chemistries to attach different types of spin labels triggered the exploration of distance measurements between different types of spin labels. So far,  $Gd^{3+}$ –nitroxide<sup>19,22</sup> and  $Mn^{2+}$ –nitroxide<sup>23-25</sup> distance measurements have been reported both in model compounds and in bio-

macromolecules. The advantage of this approach is the increase in sensitivity as compared to the “homogenous” metal-metal spin labeling scheme, although orientation selection is reintroduced to a limited extent.<sup>21</sup> Trityl-nitroxide distance measurements,<sup>26,27</sup> as well as distance measurements between a paramagnetic metal ion that is an integral part of a metalloprotein, such as Cu<sup>2+</sup> and Fe<sup>3+</sup>, and a nitroxide<sup>28</sup> or a trityl spin label,<sup>29</sup> have also been reported.


When using two spin labels (similar or different), the EPR-based distance measurements produce a single distance (except for oligomeric proteins); this implies that  $n$  different samples need to be prepared, if structural information is to be obtained from  $n$  distances. This is labor intensive. The possibility of performing spectroscopically selective DEER measurements using two types of paramagnetic centers in a three spin system and obtaining information beyond a single distance has been demonstrated on model systems, nanoparticles and trans-membrane model peptides.<sup>23,30-33</sup> In addition, measurements were carried out on a homodimeric protein, produced by mixing monomers that were each labeled with a different spin label (Gd<sup>3+</sup> and nitroxide), and spectroscopic selection of Gd<sup>3+</sup>-nitroxide, nitroxide-nitroxide and Gd<sup>3+</sup>-Gd<sup>3+</sup> distances was demonstrated.<sup>22</sup> However, as the two labels were attached to the same site of the protein, the different distances extracted reflected the different chemical structures of the spin labels rather than giving three independent structural constraints on the protein. An alternative approach for introducing two different spin labels takes advantage of two orthogonal labeling chemistries by using a cysteine to label one site and an unnatural amino acid to label the other site.<sup>9</sup>

In the present work we show that by using three different spin labels based on Gd<sup>3+</sup>, Mn<sup>2+</sup>, and nitroxide tags it is possible to extract three independent distances, Gd<sup>3+</sup>-Mn<sup>2+</sup>, Gd<sup>3+</sup>-nitroxide and Mn<sup>2+</sup>-nitroxide, from a single sample.<sup>34</sup> We demonstrate this using the *Antennapedia* homeodomain labeled with Gd<sup>3+</sup> and Mn<sup>2+</sup> tags, introduced via orthogonal labeling, in complex


with its cognate DNA binding site<sup>35</sup>, which was labeled with a nitroxide spin label (see Fig. 1a). More specifically, the unnatural amino-acid *p*-azido-L-phenylalanine (AzF) was introduced at position 22 and labeled with a Gd<sup>3+</sup> chelate, C3,<sup>17</sup> and the Mn<sup>2+</sup> ion was introduced via a MTS-EDTA tag attached to Cys39. Finally, the nitroxide was introduced through a reaction of the oligonucleotide d(CTCTAATGGCTT\*TC), containing a phosphorothioate (PT) group between the second and third last nucleotides (indicated by a star), with IAM-PROXYL. PT groups in DNA are known to be amenable to selective alkylation.<sup>36-37</sup> The complementary strand was not labeled (see Fig. 1b-d for the reaction schemes). Experimental details are given in the Supplementary Information, SI.

The selection and assignment of the three different distances was based on: (i) The different spectra of the three spins, particularly at W-band, which provide increased spectral resolution. (ii) The different nutation frequencies of nitroxide ( $S=1/2$ ), Mn<sup>2+</sup> ( $S=5/2$ ) and Gd<sup>3+</sup> ( $S=7/2$ ) which, for the central transitions, follow the ratio 1:3:4. (iii) The longer electron spin-lattice relaxation time,  $T_1$ , of nitroxides relative to Gd<sup>3+</sup> and Mn<sup>2+</sup>. Initially we performed a series of W-band echo-detected EPR (ED-EPR) measurements at different conditions to optimize the EPR spectra of the different paramagnetic centers, which have very different spectroscopic properties. The conditions found for maximizing the ED-EPR intensity of each of the spin labels are summarized in the SI and the spectra are shown in Figure 2. The nitroxide spectrum (Fig. 2a) is best observed while optimizing the pulse lengths for a  $S=1/2$  system, which will lead to too long pulses for the Mn<sup>2+</sup> and Gd<sup>3+</sup> central transition. For example, for the same power setting a  $\pi/2$  pulse of 30 ns for a nitroxide corresponds to  $3\pi/2$  and  $2\pi$  pulses for Mn<sup>2+</sup> and Gd<sup>3+</sup>, respectively. Similarly, for optimized observation of Mn<sup>2+</sup> and Gd<sup>3+</sup> (Fig. 2b,c), in addition to the too short pulses for the nitroxide, a fast repetition rate, suitable for the short  $T_1$  values of Mn<sup>2+</sup> ( $\sim 121 \mu\text{s}$ )

and  $Gd^{3+}$  (115  $\mu s$ ) will saturate the nitroxide ( $T_1 = 45$  ms) signals (see Fig. S1). These concepts were used in the past to select between  $Gd^{3+}$  and nitroxides.<sup>20, 22, 31</sup> The  $Mn^{2+}$  spectrum reveals the presence of a small contribution from  $Mn^{2+}$  ions that are not bound to the EDTA tag and display much narrower peaks. We attribute this to migration of a small amount of  $Mn^{2+}$  ions from the protein to the DNA (see Fig. S2).




**Figure 1.** Labeling schemes used. (a) Cartoon representation of the complex between the *Antennapedia* homeodomain and its cognate DNA. The three different tags used in the present work were modeled on the NMR structure of the complex (PDB ID: 1AHD)<sup>36</sup> showing the tags in red and the paramagnetic centers in magenta. (b) Reaction scheme of a PT group in DNA with IAM-PROXYL. (c) Reaction scheme of a cysteine residue with EDTA-MTS. (d) Cu(I)-catalyzed click reaction of an AzF residue with the C3-Gd tag.



**Figure 2.** W-band ED-EPR spectra of the *Antennapedia* homeodomain-DNA complex recorded under different conditions (see SI) suited to highlight each one of the labels separately. Measurements were optimized for the observation of (a) nitroxide (blue), (b) Mn<sup>2+</sup> (red), and (c) Gd<sup>3+</sup> (black). The positions of the pump and observe pulses for the selection of different distances are also indicated. The top trace shows in blue the set up for Gd<sup>3+</sup>-NO and Mn<sup>2+</sup>-NO distances and in green the set-up for Gd<sup>3+</sup>-NO distance selection using a dual-mode cavity. The bottom trace shows the pulse position for Mn<sup>2+</sup>-Gd<sup>3+</sup> selection.

To determine the distances between the three labels we performed a series of DEER measurements under different conditions optimized for selective detection of the different distances (Fig. 3). The full experimental details of the different measurements are given in the SI. The pulse frequencies for selecting individual distance distributions are also summarized in Fig. 2. All DEER traces were background-corrected and analyzed using the DeerAnalysis software<sup>38</sup> and for each trace the regularization parameter recommended by DeerAnalysis was used for obtaining the distance distributions. To measure the Gd<sup>3+</sup>-Mn<sup>2+</sup> distance we positioned the pump pulse on the Gd<sup>3+</sup> peak and the observe frequency 100 MHz up-field, close to the last peak of the

$Mn^{2+}$  multiplet to maximize the  $Mn^{2+}$  signal (Fig. 3, black trace). These conditions ensured that there was no overlap with the nitroxide spectrum, thus avoiding any contributions from distances involving the nitroxide ( $Gd^{3+}$ -nitroxide,  $Mn^{2+}$ -nitroxide). Furthermore, the repetition rate used was such that the nitroxide is saturated. The distance distribution consists of a major peak at 3 nm and two a shoulder at 3.6 and a smaller peak at 4.5 nm. To examine the impact of DNA binding on this distance we also measured the  $Mn^{2+}$ - $Gd^{3+}$  distance in the homeodomain in the absence of DNA under the same conditions (black trace in Fig. 3). Here the distance distribution gave a strong peak at 2.8 nm along with a weak peak at 4.5 nm, indicating that the binding caused a slight shift of the maximum of the distance distribution and a broadening of the distribution. The broadening can also include a small contribution from  $Mn^{2+}$  ions that migrated to the DNA.




**Figure 3.** (a) Series of DEER traces (after background removal) and fits obtained for the homeodomain/DNA complex designed to select the three different distances in the sample along with DEER traces of control samples (see text). The traces were shifted vertically relative to each other for improved visibility. The pulse set up is indicated next to the traces in (b) and as indicated in Fig. 2 and the Table S2. The blue trace was obtained with a dual-mode cavity. The black trace corresponds to the homeodomain alone and the magenta trace corresponds to the homeodomain/DNA complex where the protein was labeled only with  $Mn^{2+}$  only. (b) Distance distributions obtained from these traces with the DeerAnalysis<sup>®</sup> software. The primary DEER data are shown in Fig. S4. Dashed lines indicate the distance distributions calculated for the three labels modeled on the NMR structure of the *Antennapedia* homeodomain-DNA complex (PDB ID: 1AHD).<sup>39</sup>

The distance between the nitroxide label on the DNA and the  $Gd^{3+}$  label on the homeodomain was measured with a dual-mode cavity (blue trace in Fig. 3).<sup>21, 40</sup> This allowed positioning the

pump pulse on the maximum of the nitroxide spectrum and the detection frequency on the peak of the  $\text{Gd}^{3+}$  spectrum ( $\Delta\nu = 695$  MHz). At this observe position the contribution of the signal from  $\text{Mn}^{2+}$  is negligible compared to that from  $\text{Gd}^{3+}$  (see Fig. 2c) and therefore any contributions from the  $\text{Mn}^{2+}$ -nitroxide distance are equally negligible. The ED-EPR spectrum was measured under these conditions also for a  $\tau$  value of  $4.5 \mu\text{s}$  (see Fig. S3), relevant for the DEER measurements, showing that the relative contribution of the  $\text{Mn}^{2+}$  remained low, namely its phase memory is similar to that of the  $\text{Gd}^{3+}$ . Under this set-up the nitroxide and  $\text{Mn}^{2+}$  contribute to the spectrum at the pump pulse position, but because the pump  $\pi$  pulse is optimized for the nitroxide ( $S = 1/2$ ) it is very inefficient for  $\text{Mn}^{2+}$  and therefore the contribution from the  $\text{Mn}^{2+}$ - $\text{Gd}^{3+}$  distances to the DEER trace is expected to be negligible as well. This is confirmed by the large modulation depth of 5.5%, which is much larger than the depth expected for  $\text{Mn}^{2+}$ - $\text{Gd}^{3+}$  DEER when the pump pulse frequency is set outside the central transition of the  $\text{Gd}^{3+}$  spectrum. For this optimized setup we obtained a  $\text{Gd}^{3+}$ -nitroxide distance distribution with a maximum at 2.8 nm. Although this distance is closely similar to the  $\text{Mn}^{2+}$ - $\text{Gd}^{3+}$  distance, our experimental conditions allow unambiguous discrimination between the two. Measurements with the pump pulse set to different positions within the nitroxide EPR spectrum to test for orientation selection are shown in Fig. S5.

To measure the  $\text{Mn}^{2+}$ -nitroxide distance distribution we positioned the pump pulse on the maximum of the nitroxide spectrum and placed the observe pulses 140 MHz upfield (See Fig. 2a). At this observe position we expect the signal to have contributions from all three spin labels. Therefore, the distance distribution should have contributions from both  $\text{Mn}^{2+}$ -nitroxide and  $\text{Gd}^{3+}$ -nitroxide dipolar interactions, but we can exclude contributions from the  $\text{Mn}^{2+}$ - $\text{Gd}^{3+}$  interactions because the pump pulse length was optimized for the nitroxide. Indeed, the distance distribution

shows two distances, the first at about 2.8 nm, which is very similar to the Gd<sup>3+</sup>–nitroxide distance measured using the dual-mode cavity, and the second distance at 4.4 nm, which we attribute to the Mn<sup>2+</sup>–nitroxide distance (see green traces in Fig. 3).



**Figure 4.** DEER measurements on the *Antennapedia* homeodomain-DNA complex using different observe pulse durations and same microwave power. The pump pulse was set on the nitroxide maximum and the observe pulses on the overlapping signals from Gd<sup>3+</sup> and Mn<sup>2+</sup> (see Fig. 2a) (a) DEER traces after background subtraction, including the fit based on the distance distributions shown in (b). (b) Distance distributions obtained from the data in (a).

To substantiate this assignment we repeated the DEER measurements with different observe pulse lengths (same microwave power) to differentiate between Gd<sup>3+</sup> and Mn<sup>2+</sup>. At this observe frequency the major contribution to the Gd<sup>3+</sup> spectrum comes from the  $M_s = -3/2$  to  $M_s = -1/2$  transition, while that of the Mn<sup>2+</sup> spectrum is mostly from the central transition,  $M_s = -1/2$  to  $M_s = 1/2$ . Accordingly, the optimal nutation frequency of Gd<sup>3+</sup> is expected to be larger by a factor of at least  $\sqrt{15/9} = 1.29$  due to its higher spin. For a constant power setup with observed pulses of 15 and 30 ns, optimized for the central transition of Gd<sup>3+</sup>, which has a nutation frequency very similar to that of the Gd<sup>3+</sup> observe transition ( $\sqrt{15/16} = 1.03$ ), we thus expect longer observe

pulses to reduce the  $Gd^{3+}$  contribution and increase the  $Mn^{2+}$  contribution to the echo and, consequently, increase the relative contribution of the  $Mn^{2+}$ -nitroxide distances to the distance distribution. Figure 4 shows the result of such an experiment. While for the 15/30 and 20/40 ns set up the observed difference in the relative intensity of the peaks is within experimental error, for the 25/50 ns the relative weight of the 4.4 nm peak clearly grows. Here we observe a shift of the maxima of two peaks from 4.4-4.45 nm to 4.2 nm and from 2.7 to 2.9 nm, the origin of which is currently unknown. (See Fig. S6 for the primary time domain DEER data along with the ED-EPR spectra recorded with the observe pulses indicated in Fig. 4, and Figs. S7 and S8 for uncertainties in the distance distribution as obtained from the validation process using DeerAnalysis for the data shown in Figs. 3 and 4). To confirm this assignment we repeated the nitroxide- $Mn^{2+}$  DEER experiment on a sample where the homeodomain was labeled only with  $Mn^{2+}$ . As expected, this experiment yielded only one peak in the distance distribution, at 4.3 nm (magenta curve in Figure 3).

Finally, we compared the distances we obtained from the DEER experiments to distances predicted from the structure of the homeodomain-DNA complex.<sup>39</sup> The results are shown as dashed traces in Figure 3b. The experimentally determined  $Mn^{2+}$ -nitroxide and  $Gd^{3+}$ - $Mn^{2+}$  distance distributions agree well with the calculated ones except for the additional small peak at 4.5 nm in the  $Gd^{3+}$ - $Mn^{2+}$  distribution, which may arise from a small amount of  $Mn^{2+}$  that migrated to the DNA (see Fig. S1). The calculated  $Gd^{3+}$ -nitroxide distance is shifted by  $\sim 0.5$  nm towards a longer distance.

To conclude, we have demonstrated for the first time DEER distance measurements between three different spin labels in a single biological sample and obtained three independent distance constraints. Specifically, a protein was labeled orthogonally with a  $Gd^{3+}$  tag inserted through the

use of an unnatural amino acid and a Mn<sup>2+</sup> tag attached to a cysteine residue. The third spin label, a nitroxide, was introduced in the DNA part of the complex. This approach reduced the number of samples needed to obtain the distance information from three to one. This new approach is particularly useful for studying complex formation in bio-macromolecules.

**Supporting Information.** Experimental details, T<sub>1</sub> measurements, ED-EPR spectra of Antennapedia homeodomain with and without DNA, primary DEER data, Gd<sup>3+</sup>-nitroxide DEER, validation of distance distributions.

#### AUTHOR INFORMATION

The authors declare no competing financial interests.

#### ACKNOWLEDGMENT

D. G. and G. O. acknowledge the support of the Israel Science Foundation (ISF grant No. 334/14) and the Australian Research Council, respectively. This research was made possible in part by the historic generosity of the Harold Perlman Family (D. G.). D. G. holds the Erich Klieger Professorial Chair in Chemical Physics.

#### REFERENCES

1. Jeschke, G.; Polyhach, Y., Distance Measurements on Spin-labelled Biomacromolecules By pulsed Electron Paramagnetic Resonance. *Phys. Chem. Chem. Phys.* **2007**, *9*, 1895-1910.
2. Feintuch, A.; Otting, G.; Goldfarb, D., Gd<sup>3+</sup> Spin Labeling for Measuring Distances in Biomacromolecules: Why and How? *Methods Enzymol.* **2015**, *563*, 415-457.
3. Goldfarb, D., Gd<sup>3+</sup> Spin Labeling for Distance Measurements by Pulse EPR Spectroscopy. *Phys. Chem. Chem. Phys.* **2014**, *16*, 9685-9699.
4. Banerjee, D.; Yagi, H.; Huber, T.; Otting, G.; Goldfarb, D., Nanometer-Range Distance Measurement in a Protein Using Mn<sup>2+</sup> Tags. *J. Phys. Chem. Lett.* **2012**, *3*, 157-160.
5. Martorana, A.; Yang, Y.; Zhao, Y.; Li, Q. F.; Su, X. C.; Goldfarb, D., Mn(II) Tags for DEER Distance Measurements in Proteins via C-S Attachment. *Dalton Trans.* **2015**, *44*, 20812-20816.
6. Ching, H. Y. V.; Mascali, F. C.; Bertrand, H. C.; Bruch, E. M.; Demay-Drouhard, P.; Rasia, R. M.; Policar, C.; Tabares, L. C.; Un, S., The Use of Mn(II) Bound to His-Tags as Genetically Encodable Spin-Label for Nanometric Distance Determination in Proteins. *J. Phys. Chem. Lett.* **2016**, *7*, 1072-1076.

7. Cunningham, T. F.; Putterman, M. R.; Desai, A.; Horne, W. S.; Saxena, S., The Double-histidine Cu<sup>2+</sup>-Binding Motif: a Highly Rigid, Site-Specific Spin Probe for Electron Spin Resonance Distance Measurements. *Angew. Chem. Int. Ed.* **2015**, *54*, 6330-6334.
8. Yang, Z. Y.; Kise, D.; Saxena, S., An Approach towards the Measurement of Nanometer Range Distances Based on Cu<sup>2+</sup> Ions and ESR. *J. Phys. Chem. B* **2010**, *114*, 6165-6174.
9. Reginsson, G. W.; Kunjir, N. C.; Sigurdsson, S. T.; Schiemann, O., Trityl Radicals: Spin Labels for Nanometer-Distance Measurements. *Chem. Eur. J.* **2012**, *18*, 13580-13584.
10. Shevelev, G. Y.; Krunkacheva, O. A.; Lomzov, A. A.; Kuzhelev, A. A.; Rogozhnikova, O. Y.; Trukhin, D. V.; Troitskaya, T. I.; Tormyshev, V. M.; Fedin, M. V.; Pyshnyi, D. V.; Bagryanskaya, E. G., Physiological-Temperature Distance Measurement in Nucleic Acid using Triarylmethyl-Based Spin Labels and Pulsed Dipolar EPR Spectroscopy. *J. Am. Chem. Soc.* **2014**, *136*, 9874-9877.
11. Yang, Z.; Liu, Y.; Borbat, P.; Zweier, J. L.; Freed, J. H.; Hubbell, W. L., Pulsed ESR Dipolar Spectroscopy for Distance Measurements in Immobilized Spin Labeled Proteins in Liquid Solution. *J. Am. Chem. Soc.* **2012**, *134*, 9950-9952.
12. Raitsimring, A. M.; Astashkin, A. V.; Enemark, J. H.; Kaminker, I.; Goldfarb, D.; Walter, E. D.; Song, Y.; Meade, T. J., Optimization of Pulsed-DEER Measurements for Gd-Based Labels: Choice of Operational Frequencies, Pulse Durations and Positions, and Temperature. *Appl. Magn. Reson.* **2013**, *44*, 649-670.
13. Potapov, A.; Yagi, H.; Huber, T.; Jergic, S.; Dixon, N. E.; Otting, G.; Goldfarb, D., Nanometer-Scale Distance Measurements in Proteins Using Gd<sup>3+</sup> Spin Labeling. *J. Am. Chem. Soc.* **2010**, *132*, 9040-9048.
14. Raitsimring, A. M.; Gunanathan, C.; Potapov, A.; Efremenko, I.; Martin, J. M. L.; Milstein, D.; Goldfarb, D., Gd<sup>3+</sup> Complexes as Potential Spin Labels for High Field Pulsed EPR Distance Measurements. *J. Am. Chem. Soc.* **2007**, *129*, 14138-14138.
15. Martorana, A.; Bellapadrona, G.; Feintuch, A.; Di Gregorio, E.; Aime, S.; Goldfarb, D., Probing Protein Conformation in Cells by EPR Distance Measurements using Gd<sup>3+</sup> Spin Labeling. *J. Am. Chem. Soc.* **2014**, *136*, 13458-13465.
16. Qi, M.; Gross, A.; Jeschke, G.; Godt, A.; Drescher, M., Gd(III)-PyMTA Label is Suitable for in-Cell EPR. *J. Am. Chem. Soc.* **2014**, *136*, 15366-15378.
17. Abdelkader, E. H.; Feintuch, A.; Yao, X.; Adams, L. A.; Aurelio, L.; Graham, B.; Goldfarb, D.; Otting, G., Protein Conformation by EPR Spectroscopy Using Gadolinium Tags Clicked to Genetically Encoded p-Azido-L-Phenylalanine. *Chem. Comm.* **2015**, *51*, 15898-15901.
18. Fleissner, M. R.; Brustad, E. M.; Kalai, T.; Altenbach, C.; Cascio, D.; Peters, F. B.; Hideg, K.; Peuker, S.; Schultz, P. G.; Hubbell, W. L., Site-Directed Spin Labeling of a Genetically Encoded Unnatural Amino Acid. *Proc. Natl. Acad. Sci. U S A* **2009**, *106*, 21637-21642.
19. Garbuio, L.; Bordignon, E.; Brooks, E. K.; Hubbell, W. L.; Jeschke, G.; Yulikov, M., Orthogonal Spin Labeling and Gd(III)-Nitroxide Distance Measurements on Bacteriophage T4-Lysozyme. *J. Phys. Chem. B* **2013**, *117*, 3145-3153.
20. Lueders, P.; Jeschke, G.; Yulikov, M., Double Electron-Electron Resonance Measured Between Gd<sup>3+</sup> Ions and Nitroxide Radicals. *J. Phys. Chem. Lett.* **2011**, *2*, 604-609.
21. Kaminker, I.; Tkach, I.; Manukovsky, N.; Huber, T.; Yagi, H.; Otting, G.; Bennati, M.; Goldfarb, D., W-Band Orientation Selective DEER Measurements on a Gd<sup>3+</sup>/Nitroxide Mixed-Labeled Protein Dimer with a Dual Mode Cavity. *J. Magn. Reson.* **2013**, *227*, 66-71.

22. Kaminker, I.; Yagi, H.; Huber, T.; Feintuch, A.; Otting, G.; Goldfarb, D., Spectroscopic Selection of Distance Measurements in a Protein Dimer with Mixed Nitroxide and Gd<sup>3+</sup> Spin Labels. *Phys. Chem. Chem. Phys.* **2012**, *14*, 4355-4358.
23. Meyer, A.; Schiemann, O., PELDOR and RIDME Measurements on a High-Spin Manganese(II) Bisnitroxide Model Complex. *J. Phys. Chem. A* **2016**, *120*, 3463-3472.
24. Akhmetzyanov, D.; Plackmeyer, J.; Endeward, B.; Denysenkov, V.; Prisner, T. F., Pulsed Electron-Electron Double Resonance Spectroscopy Between a High-Spin Mn<sup>2+</sup> Ion and a Nitroxide Spin Label. *Phys. Chem. Chem. Phys.* **2015**, *17*, 6760-6766.
25. Kaminker, I.; Bye, M.; Mendelman, N.; Gislason, K.; Sigurdsson, S. T.; Goldfarb, D., Distance Measurements Between Manganese(II) and Nitroxide Spin-Labels by DEER Determine a Binding Site of Mn<sup>2+</sup> in the HP92 Loop of Ribosomal RNA. *Phys. Chem. Chem. Phys.* **2015**, *17*, 15098-15102.
26. Joseph, B.; Tormyshev, V. M.; Rogozhnikova, O. Y.; Akhmetzyanov, D.; Bagryanskaya, E. G.; Prisner, T. F., Selective High-Resolution Detection of Membrane Protein-Ligand Interaction in Native Membranes Using Trityl-Nitroxide PELDOR. *Angew. Chem. Int. Ed.* **2016**, *55*, 11538-11542.
27. Shevelev, G. Y.; Krumkacheva, O. A.; Lomzov, A. A.; Kuzhelev, A. A.; Trukhin, D. V.; Rogozhnikova, O. Y.; Tormyshev, V. M.; Pyshnyi, D. V.; Fedin, M. V.; Bagryanskaya, E. G., Triarylmethyl Labels: Toward Improving the Accuracy of EPR Nanoscale Distance Measurements in DNAs. *J. Phys. Chem. B* **2015**, *119*, 13641-13648.
28. Astashkin, A. V., Mapping the Structure of Metalloproteins with RIDME. *Methods Enzymol.* **2015**, *563*, 251-284.
29. Jassoy, J. J.; Berndhäuser, A.; Duthie, F.; Kühn, S. P.; Hagelueken, G.; Schiemann, O., Versatile Trityl Spin Labels for Nanometer Distance Measurements on Biomolecules in Vitro and within Cells. *Angew. Chem. Int. Ed.* **2017**, *56*, 177-181.
30. Lueders, P.; Jager, H.; Hemminga, M. A.; Jeschke, G.; Yulikov, M., Distance Measurements on Orthogonally Spin-Labeled Membrane Spanning WALP23 Polypeptides. *J. Phys. Chem. B* **2013**, *117*, 2061-2068.
31. Yulikov, M.; Lueders, P.; Warsi, M. F.; Chechik, V.; Jeschke, G., Distance Measurements in Au Nanoparticles Functionalized with Nitroxide Radicals and Gd<sup>3+</sup>-DTPA Chelate Complexes. *Phys. Chem. Chem. Phys.* **2012**, *14*, 10732-10746.
32. Narr, E.; Godt, A.; Jeschke, G., Selective Measurements of a Nitroxide-Nitroxide Separation of 5 nm and a Nitroxide-Copper Separation of 2.5 nm in a Terpyridine-Based Copper(II) Complex by Pulse EPR Spectroscopy. *Angew. Chem. Int. Ed.* **2002**, *41*, 3907-3910.
33. Meyer, A.; Abdullin, D.; Schnakenburg, G.; Schiemann, O., Single and Double Nitroxide Labeled Bis(terpyridine)-Copper(II): Influence of Orientation Selectivity and Multispin Effects on PELDOR and RIDME. *Phys. Chem. Chem. Phys.* **2016**, *18*, 9262-9271.
34. After submitting the manuscript it has been brought to our attention that distance measurements between three spins, Cu<sup>2+</sup>, trityl, and a nitroxide in a model compound have been reported by Bowen, A. M.; Schuldais, R.; Plackmeyer, J.; Sigurdsson, S.; Prisner, T. F. at the EUROMAR 2017 conference ( book of abstracts, p. 147)
35. Müller, M.; Affolter, M.; Leupin, W.; Otting, G.; Wüthrich, K.; Gehring, W. J., Isolation and Sequence-Specific DNA Binding of the Antennapedia Homeodomain. *EMBO J.* **1988**, *7*, 4299-4304.

36. Shepherd, N. E.; Gamsjaeger, R.; Vandevenne, M.; Cubeddu, L.; Mackay, J. P., Site Directed Nitroxide Spin Labeling of Oligonucleotides for NMR and EPR Studies. *Tetrahedron* **2015**, *71*, 813-819.
37. Wu, Z.; Lee, M. D.; Carruthers, T. J.; Szabo, M.; Dennis, M. L.; Swarbrick, J. D.; Graham, B.; Otting, G., New Lanthanide Tag for the Generation of Pseudocontact Shifts in DNA by Site-Specific Ligation to a Phosphorothioate Group. *Bioconjugate Chem.* **2017**, *28*, 1741-1748.
38. Jeschke, G.; Chechik, V.; Ionita, P.; Godt, A.; Zimmermann, H.; Banham, J.; Timmel, C. R.; Hilger, D.; Jung, H., DeerAnalysis2006 - a Comprehensive Software Package for Analyzing Pulsed ELDOR Data. *Appl. Magn. Reson.* **2006**, *30*, 473-498.
39. Billeter, M.; Qian, Y. Q.; Otting, G.; Müller, M.; Gehring, W.; Wüthrich, K., Determination of the Nuclear-Magnetic-Resonance Solution Structure of an Antennapedia Homodomain-DNA Complex *J. Mol. Biol.* **1993**, *234*, 1084-1094.
40. Tkach, I.; Sicoli, G.; Hobartner, C.; Bennati, M., A Dual-Mode Microwave Resonator for Double Electron-Electron Spin Resonance Spectroscopy at W-band Microwave Frequencies. *J. Magn. Reson.* **2011**, *209*, 341-346.